

UC Riverside

Journal of Citrus Pathology

Title

Huanglongbing Surveillance Program Actions in the State of Bahia, Brazil

Permalink

<https://escholarship.org/uc/item/4tr8c5rx>

Journal

Journal of Citrus Pathology, 1(1)

Authors

Silva, S. X.B.
Andrade, E. C.
Nascimento, A. S.
et al.

Publication Date

2014

DOI

10.5070/C411024184

Copyright Information

Copyright 2014 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

1.6 P

Huanglongbing Surveillance Program Actions in the State of Bahia, Brazil

Silva, S.X.B.¹, Andrade, E.C.², Nascimento, A.S.², Barbosa, C.J.^{2,3}, Girardi, E.A.², Astúa, J.F.²,
and Laranjeira, F.F.²

¹Laboratory of Epidemiology and Health Information Technology / Bahia State Agency of Agricultural Defense (ADAB), Salvador/BA, Brazil

²Embrapa Cassava & Fruits, Cruz das Almas/BA, Brazil

³Centro de Laboratórios da Agropecuária (CLA), Empresa Baiana de Desenvolvimento Agrícola (EBDA), Salvador/BA, Brazil

Huanglongbing (HLB) is found in South/Southeastern states of Brazil, but citrus is grown all over the country. For that reason, surveillance procedures should be carried out frequently and contingency plans developed. This study reports the actions of the State Bureau of Agricultural Defense of Bahia (ADAB), for a commercial orchard in Bom Jesus da Lapa (state of Bahia, Northeastern Brazil) suspected of having HLB symptomatic plants. Besides having a Contingency Plan, a protocol that establishes actions to detect and eradicate the disease, ADAB is also a partner of the collaborative network HLB BioMath. In August 2012, an ADAB team inspected 22 hectares of three year old orchards of both Pêra sweet orange and Ponkan mandarin. Scouting was performed on total area, and 4.5% of the plants were marked as having HLB-like symptoms. Samples were tested with polymerase chain reaction (PCR) by the Phytopathological Clinic of Sylvio Moreira Citrus Center. The results were negative for HLB bacterium, but a *Phytoplasma* of 16SrIX Group was found. All symptomatic plants will be eliminated and the surveillance area will be extended. Despite this finding, Bom Jesus da Lapa is 700km away from the most important citrus regions in Bahia: Reconcavo and Litoral Norte. In those regions no suspected plant was ever reported.

Index terms: *Candidatus Liberibacter*, Free Area, Agricultural Defense.