

UC San Diego

UC San Diego Previously Published Works

Title

Profiles of brain metastases: Prioritization of therapeutic targets

Permalink

<https://escholarship.org/uc/item/4x06c00c>

Journal

International Journal of Cancer, 143(11)

ISSN

0020-7136

Authors

Ferguson, Sherise D
Zheng, Siyuan
Xiu, Joanne
et al.

Publication Date

2018-12-01

DOI

10.1002/ijc.31624

Peer reviewed

Profiles of brain metastases: Prioritization of therapeutic targets

Sherise D. Ferguson¹, Siyuan Zheng², Joanne Xiu³, Shouhao Zhou⁴, Mustafa Khasraw⁵, Priscilla K. Brastianos⁶, Santosh Kesari⁷, Jethro Hu⁸, Jeremy Rudnick⁸, Michael E. Salacz⁹, David Piccioni¹⁰, Suyun Huang¹, Michael A. Davies¹¹, Isabella C. Glitza¹¹, John V. Heymach¹², Jianjun Zhang¹², Nuhad K. Ibrahim¹³, John F. DeGroot¹⁴, Joseph McCarty¹, Barbara J. O'Brien¹⁴, Raymond Sawaya¹, Roeland G.W. Verhaak², Sandeep K. Reddy³, Waldemar Priebe¹⁵, Zoran Gatalica³, David Spetzler³ and Amy B. Heimberger¹

¹Departments of Neurosurgery, The University of Texas MD Anderson Cancer Center, Houston, TX

²Departments of Genome Medicine, The University of Texas MD Anderson Cancer Center, Houston, TX

³Caris Life Sciences, Phoenix, AZ

⁴Departments of Biostatistics, The University of Texas MD Anderson Cancer Center, Houston, TX

⁵NHMRC Clinical Trials Centre, University of Sydney, Sydney, Australia

⁶Division of Neuro-Oncology, Massachusetts General Hospital, Harvard Medical School, Boston, MA

⁷Pacific Neuroscience Institute and John Wayne Cancer Institute at Providence Saint John's Health Center, Santa Monica, CA

⁸Cedars-Sinai, Los Angeles, CA

⁹University of Kansas Cancer Center, Kansas City, KS

¹⁰Department of Neurosciences, University of California at San Diego Moores Cancer Center, La Jolla, CA

¹¹Departments of Melanoma Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

¹²Departments of Thoracic Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

¹³Departments of Breast Medical Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

¹⁴Departments of Neuro-Oncology, The University of Texas MD Anderson Cancer Center, Houston, TX

¹⁵Departments of Experimental Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, TX

We sought to compare the tumor profiles of brain metastases from common cancers with those of primary tumors and extracranial metastases in order to identify potential targets and prioritize rational treatment strategies. Tumor samples were collected from both the primary and metastatic sites of nonsmall cell lung cancer, breast cancer and melanoma from patients in locations worldwide, and these were submitted to Caris Life Sciences for tumor multiplatform analysis, including gene sequencing (Sanger and next-generation sequencing with a targeted 47-gene panel), protein expression (assayed by immunohistochemistry) and gene amplification (assayed by in situ hybridization). The data analysis considered differential protein expression, gene amplification and mutations among brain metastases, extracranial metastases and primary tumors. The analyzed population included: 16,999 unmatched primary tumor and/or metastasis samples: 8,178 nonsmall cell lung cancers (5,098 primaries; 2,787 systemic metastases; 293 brain metastases), 7,064 breast cancers (3,496 primaries; 3,469 systemic metastases; 99 brain metastases) and 1,757 melanomas (660 primaries; 996 systemic metastases; 101 brain metastases). TOP2A expression was increased in brain metastases from all 3 cancers, and brain metastases overexpressed multiple proteins clustering around functions critical to DNA synthesis and repair and implicated in chemotherapy resistance, including RRM1, TS, ERCC1 and TOPO1. cMET was overexpressed in melanoma brain metastases relative to primary skin specimens. Brain metastasis patients may particularly benefit from therapeutic targeting of enzymes associated with DNA synthesis, replication and/or repair.

Key words: brain metastases, molecular profiling, multiplatform analysis, DNA repair enzymes, TOP2A

Additional Supporting Information may be found in the online version of this article.

Grant sponsor: National Institutes of Health; Grant numbers: P30CA16672 and CA1208113; Grant sponsor: provost funds provided by Ethan Dmitrovsky.

Grant sponsor: National Institutes of Health; **Grant numbers:** CA1208113, P30CA16672; **Grant sponsor:** provost funds provided by Ethan Dmitrovsky

DOI: 10.1002/ijc.31624

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

History: Received 4 Jan 2018; Accepted 17 Apr 2018; Online 19 June 2018

Correspondence to: Sherise D. Ferguson, M.D., Department of Neurosurgery, The University of Texas MD Anderson Cancer Center, 1400 Holcombe Blvd., Houston, TX 77030, Tel.: 713-792-2400, E-mail: sdferguson@mdanderson.org

What's new?

Brain metastases are difficult to treat and generally lethal. Here, the authors sought clues to possible treatment targets by comparing the molecular characteristics of lung, breast, and skin cancers with those of brain metastases. They analyzed DNA sequence, protein expression, and gene amplification on over 17,000 samples, the largest metastases cohort ever studied. They found that the enzyme TOP2A, involved in DNA transcription, was enriched in the metastatic cells relative to the primary tumor cells. Other enzymes, involved in DNA synthesis, repair, and replication were also overexpressed in metastases, and could potentially be useful therapeutic targets.

Introduction

Metastasis is the most common source of tumors in the brain, with lung cancer, breast cancer and melanoma being the most common solid tumors spreading to the central nervous system (CNS).¹ Overall, it is estimated that 9–17% of cancer patients will develop brain metastases during the course of their disease and that approximately half will die within 3–27 months from initial diagnosis of CNS involvement.^{1–3} The reported median survival times for NSCLC, breast cancer and melanoma after the diagnosis of brain metastasis are 7.8 months, 15 months and 7–7.9 months, respectively.^{4–6} The poor survival associated with brain metastases is largely attributed to our incomplete understanding of the molecular mechanisms governing organotropism and the brain itself being a known sanctuary site. Additionally, most systemic therapeutics are not capable of crossing the blood–brain barrier (BBB). The need for additional treatment options is urgent; more than 100,000 patients per year develop brain metastases in the United States, and the current treatment cost exceeds 3 billion dollars.⁷ Current treatment options include surgical resection and various forms of irradiation, but these have only a modest impact on overall outcome and carry variable risks of neurological morbidity. There are no effective systemic therapies specifically approved for brain metastases. Unfortunately, once CNS involvement is present, effective treatment options are limited and the prognosis is dismal. These issues are pressing, as the incidence of brain metastasis may be increasing due to longer patient survival owing to newer systemic therapies that control extracranial disease, along with more frequent magnetic resonance imaging surveillance.^{3,5}

There has been a surge in clinical trials guided by genomics in order to better match patients with effective, targeted cancer treatments.⁸ The development of CNS metastases often results in exclusion of patients from clinical trials,⁹ particularly phase I trials of many novel therapeutics, mainly due to the known poor prognosis and limited efficacy of systemic therapy for these patients historically. Detailed molecular analyses of metastases across pathologies may help identify new therapeutic targets that can benefit these patients. Thus, we performed a multiplatform analysis of a large cohort of brain and systemic metastasis samples collected from the cancers most commonly predisposed to produce brain metastases in order to identify potentially actionable targets.

Materials and Methods**Study population**

From 2009 to 2015, tumors collected from both the primary and metastatic sites of non-small cell lung cancer (NSCLC), breast

cancer and melanoma from patients in locations worldwide were submitted to Caris Life Sciences for tumor multiplatform analysis. A total of 8,178 NSCLC (5,098 primaries; 2,787 systemic metastases; 293 brain metastases), 7,064 breast cancer (3,496 primaries; 3,469 systemic metastases; 99 brain metastases) and 1,757 melanoma (660 primaries; 996 systemic metastases; 101 brain metastases) samples (primary and metastases not matched to the same patient) were retrospectively analyzed. Our study is exempt per policy 45 CFR 46.101 (b); the data analyzed is from an existing commercial repository and subject information is de-identified.

Immunohistochemistry

All analysis by immunohistochemistry (IHC) was performed on full slides of formalin-fixed paraffin-embedded (FFPE) tumor specimens. Following the requirements of the Clinical Laboratory Improvement Amendments (CLIA)/Compliance Assistance Office (CAO) and International Organization for Standardization (ISO), staining conditions were performed per the manufacturer's instructions using automated staining techniques and were optimized and validated. The results were evaluated and confirmed by independent board-certified pathologists. Immunohistochemical scoring was based on staining percentage (0–100%) and intensity (0 = no staining; 1+ = weak staining; 2+ = moderate staining; 3+ = strong staining). Representative staining photos are shown in Supporting Information Figure S1. Results were categorized as positive or negative by defined thresholds specific to each marker based on published clinical literature that associates biomarker status with patient responses to therapeutic agents. Supporting Information Table S1 shows the thresholds for positive and negative categorization. Tumor cells were evaluated for proteins of interest. Analyzed proteins and antibodies used are listed in Supporting Information Table S2. PD-L1 staining results are specific for membranous staining, and PD-1 expression was evaluated by analysis of tumor-infiltrating lymphocytes as previously described.¹⁰ The expression of PD-L1 was analyzed with B7-H1 antibody (R&D systems) using automated immunohistochemical methods. Dilutions and conditions were based on package insert instructions, were optimized and validated and met the standards and requirements of the CLO/CAO and ISO.

In situ hybridization (ISH)

Gene amplification was assessed using fluorescence in situ hybridization (FISH) for *EGFR* (EGFR/CEP7 probe) and *TOP2A* (TOP2A/CEP17 probe). Both FISH and chromogenic

in situ hybridization (CISH) were used to detect gene amplification of *cMET* (*cMET/CEP7* probe and Ventana kit) and *Her2* (*HER-2/CEP17* probe and INFORM *HER-2* Dual ISH DNA Probe Cocktail). *EGFR* amplification was defined by the presence of an *EGFR/CEP7* ratio of ≥ 2 , or ≥ 15 *EGFR* copies per cell in $\geq 10\%$ of analyzed cells. A ratio of *TOP2:CEP17* ≥ 2.0 and an *HER-2/CEP17* ratio ≥ 2 was considered amplified; *cMET* was considered amplified if ≥ 5 copies per tumor cell were detected on average.

Mutational analysis

Using the Illumina MiSeq platform, sequencing was performed on genomic tumor DNA isolated from FFPE tumor tissue (germline DNA was not sequenced). Reference genome used was hg19 from the UCSC Genome Browser database (<http://hgdownload.cse.ucsc.edu/goldenPath/hg19/bigZips/>). Specific regions of 47 genes (panel of pan-cancer genes of interest related to cancer genomics based on current literature) were amplified using the customized Illumina TruSeq Amplicon Cancer Hotspot panel.¹¹ All variants reported were detected with $>99\%$ confidence based on the mutation frequency present. Mutations including *PTEN*, *TP53* and *BRAF* and were evaluated by Next-generation sequencing (NGS; Illumina MiSeq platform) (Supporting Information Table S3). The average depth of coverage of the hotspot panel was $>1,500\times$ and the tumor content requirement is $\geq 20\%$. This test has a sensitivity to detect as low as a 10% population of cells containing a mutation in a sequenced amplicon. All variants reported were detected with $>99\%$ confidence based on the frequency of the mutation present and the amplicon coverage. Note that as this is a commercial repository, mutational testing on individual samples was performed at the discretion of the ordering physician. Hence, not all samples were tested for all mutations. Additionally, a cohort of patients was profiled prior to the availability of NGS.

Statistical analysis

Comparisons of biomarker profiles, including protein expression (IHC), gene amplification (CISH/FISH) and mutation (Sanger sequencing and NGS) between tumor sites were performed using Fisher's exact test. To adjust for the multiple testing and limit the risk of false positive results, *p* values resulting from the Fisher's exact test were corrected using the Benjamini-Hochberg procedure.¹² All reported *p* values are two sided and corrected for multiple comparison. *P* values of less than 0.05 were declared as statistically significant. All analyses were performed with statistical software R v3.3.1.

Results

Study population

Patient characteristics, number of tumor specimens and distribution of systemic metastases (extracranial) are shown in Table 1 and Supporting Information Table S4. A total of 16,999 specimens were included for analysis; 9,252 primary lesions, 493 brain metastases and 7,252 systemic (extra-cranial

metastases). The majority of extracranial specimens were obtained from the lymph nodes. Sex was equally dichotomized among the NSCLC patients, breast cancer cases were mostly restricted to females and males were overrepresented in melanoma. Overall, the breast cancer patient population was younger than the NSCLC and melanoma cohorts.

TOP2A expression is a high frequency target in brain metastases relative to primary tumors

Using a CLIA-compliant cancer protein panel, differential expression was profiled in NSCLC, breast cancer and melanoma specimens (Fig. 1, Supporting Information Table S5). Across pathologies, TOP2A, which controls and alters the topologic states of DNA during transcription, was the protein expressed more frequently in the brain metastases relative to the primary tumors most consistently (NSCLC: 74.5% versus 55%, $p < 0.0001$; breast cancer: 78.8% versus 50.1%, $p = 0.0001$; melanoma: 76.4% versus 45.8%, $p = 0.0007$) (Fig. 2). Levels of two other enzymes involved in DNA synthesis and chemoresistance, RRM1 and TS, were also increased in NSCLC brain metastases relative to the primary (31.8% versus 20.9%, $p = 0.0049$; and 34.9% versus 21.5%, $p = 0.0006$, respectively), but this was only a trend for increased expression detected in breast cancer and melanoma brain metastases. Other DNA topoisomerases and enzymes essential for DNA replication/repair such as TOPO1 (77.9% versus 63.4%, $p = 0.0357$) and ERCC1 (61.8% versus 42.1%, $p = 0.0199$) were also more frequently expressed in breast cancer brain metastases. Her2/Neu expression was more common in breast cancer brain metastases (23.3%) compared with primary specimens (11.6%) ($p = 0.0227$), whereas the frequency of expression of all hormonal receptors (AR: 20.5% versus 51.3%, $p < 0.0001$; ER: 34.4% versus 54.1%, $p = 0.0049$; PR: 25.3% versus 43.6%, $p = 0.008$) was significantly reduced in brain metastases relative to the primary tumor. Melanoma brain metastases exhibited increased *cMET* expression frequency relative to the primary (36.4% versus 7.9%, $p = 0.0003$) but decreased TUBB3 expression (38.2% versus 76.2%, $p = 0.0007$).

Her2/Neu protein expression is more frequent in breast cancer brain metastases than in systemic metastases

To ascertain whether the differences in protein expression were a function of the process of metastasis in general or were more specific to brain metastasis, we next compared differential protein expression in brain and systemic metastases across tumor pathologies (Fig. 1, Supporting Information Table S5). Her2/Neu expression was significantly higher in breast brain metastases (23.3%) than in systemic metastases (9.9%) ($p = 0.0044$). Moreover, all hormone receptors were expressed less frequently in breast cancer brain metastases (AR, $p < 0.0001$; ER, $p = 0.0001$ and PR, $p = 0.019$)—similar to the comparison with primary sites. Systemic breast metastases showed an increased frequency of *PTEN* (52.7% versus 34%, $p = 0.0049$) and *MGMT* (68% versus 47.3%, $p = 0.0011$) relative to brain metastases (Fig. 2).

Table 1. Summary of characteristics of patients with brain and systemic metastases from non-small cell lung cancer, breast cancer and melanoma

Histology	Specimen site	Number of samples	Age (median)	Sex (% female)
Nonsmall cell lung cancer	Brain	293	61	54.9
	Lung	5,098	68	52
	Systemic Metastases	2,787	65	51
Breast cancer	Brain	99	51	100
	Breast	3,496	55	99.3
	Systemic Metastases	3,469	57	99
Melanoma	Brain	101	62	35.3
	Skin/(Primary)	660	66	35.6
	Systemic Metastases	996	64	62.5
Summary	Primary	9,254	63	69
	Brain	493	59	70
	Systemic Metastases	7,252	61	71.8
	Total	16,999	62	70

Breast cancer brain metastasis demonstrated enrichment of Her2 and EGFR gene amplifications relative to systemic metastases and primary lesions

Gene amplification results for NSCLC, breast cancer and melanoma are presented in Table 2. A total of 2,888 breast cancer specimens were evaluated for *Her2* amplification using FISH (2,808 breast primaries), with 28.8% (n = 80) of brain metastases displaying *Her2* amplification relative to 15.3% (n = 430) of the primary specimens (p = 0.0227). *Her2* amplification was also significantly higher in breast cancer brain metastases than in systemic metastases (13.1%) (p = 0.004), consistent with HER2-expressing cells having a greater propensity to spread to the brain.^{13,14} Furthermore, *EGFR* amplification was more common in breast cancer brain metastases (31.1%) than

in primaries (14.1%; p = 0.0341). Similarly, *EGFR* amplification was higher in brain metastases than in systemic lesions (14.2%; p = 0.0333). *EGFR* was also amplified in melanoma brain metastases (50%) and NSCLC brain metastases (36.3%) relative to the primary cancers (6.1% and 28%, respectively), but these differences were not statistically significant.

Minimal differences in mutational frequency between metastases and primary lesions

DNA sequencing data were analyzed for a panel of 47 well-annotated pan-cancer genes (listed in Supporting Information Table S6). In most cases, the most common mutations were found in both the primary cancers and brain metastases with




Figure 1. Differential protein expression by tumor pathology (NSCLC, breast cancer and melanoma). Figure displaying changes in expression of 20 cancer-related proteins in primary lesions, brain and systemic metastases. Numbers in the cells indicate rounded up fractions.




Figure 2. Bar graphs of protein expression frequency in brain metastases (BM), primary lesions and systemic metastases. The values used to generate this figure are absolute frequencies of expression i.e., the number of individual cases showing specific protein expression. Significant findings are displayed for (a) Non-small cell lung cancer; (b) breast cancer and (c) melanoma. Asterisks indicate comparisons that remain statistically significant after correction for multiple comparisons ($p < 0.05$). Black denotes brain metastases; blue, primary tumors; and green, systemic metastases.

similar frequencies. However, there were some notable exceptions in breast cancer. For example, *ABL1* mutations were more frequent in breast cancer brain metastases (11.1%; 2/18) than in primary breast cancers (0.4%; 3/822) ($p = 0.0319$) and *PIK3CA* mutations were more frequent in breast systemic metastases (30.3%; 415/1,370) than in brain metastases (10.3%; 4/39) ($p = 0.0426$) (Fig. 3; Supporting Information Table S7).

Discussion

Multiplatform analysis was performed on approximately 17,000 primary and metastatic cancers, which is the largest cohort described in the literature, in order to reveal potential therapeutic targets. We observed a significant association between TOP2A protein expression and brain metastases. TOP2A is a critical enzyme in DNA transcription and regulation that is highly expressed in proliferating cells, as such we cannot confirm this finding is not influenced by higher proliferative activity¹⁵ TOP2A overexpression is associated with an aggressive cancer phenotype and decreased patient survival,^{16–18} which makes it an attractive therapeutic target. Unfortunately, most currently available TOP2A inhibitors have limited BBB penetration, making them ineffective for treating brain metastases.¹⁹ We did not observe TOP2A amplification in brain metastases, however it is established that TOP2A protein expression is not secondary to gene amplification, which likely explains this discordance.²⁰

The proteins RRM1, ERCC1 and TS (involved in DNA replication, repair and chemotherapy resistance) were more frequently expressed in metastases than in primary lesions across all three tumor pathologies (NSCLC, breast cancer and melanoma). The DNA repair enzyme ERCC1, which plays a role in therapeutic resistance to platinum-based chemotherapy,^{21,22} has not been previously documented to be associated with




Figure 3. Schematic displaying the mutational frequency of cancer-related genes in primary lesions, brain and systemic metastases. Only genes with mutational frequency of >10% in at least one subgroup are shown in the figure.

Table 2. Frequency of gene amplification detected by fluorescence in situ hybridization (FISH) or chromogenic in situ hybridization (CISH) in brain metastases and in tumors taken from the primary and systemic sites of nonsmall cell lung cancer, breast cancer, and melanoma

Gene	NSCLC			BREAST CANCER			MELANOMA		
	Brain	Primary	Systemic	Brain	Primary	Systemic	Brain	Primary	Systemic
<i>EGFR</i>	36.3% (29/80)	28.0% (438/1,567)	34.5% (250/724)	31.1% ² (14/45)	14.4% (121/839)	14.2% ⁴ (138/970)	50% ¹ (4/8)	6.1% (2/33)	28.9% (26/90)
<i>TOP2A</i>	0% (0/14)	5% (9/181)	1.6% (2/128)	16.7% (7/42)	8.5% (112/1,315)	6.3% ³ (101/1,594)	0% (0/5)	0% (0/1)	0% (0/48)
<i>Her2</i>	5.8% (10/171)	3.6% (79/2,207)	3% (36/1,204)	28.8% ² (23/80)	15.3% (430/2,808)	13.1% ⁴ (361/2,767)	0% (0/41)	0% (0/205)	0% (0/247)
<i>cMET</i>	6.4% (8/125)	4% (61/1,517)	5.2% (47/906)	5.6% (1/18)	1.1% (11/1,027)	0.7% (7/987)	8.6% ¹ (3/35)	0.6% (1/168)	3.5% (8/230)

NSCLC, nonsmall cell lung cancer.

¹Bivariate analysis $p < 0.05$ (brain metastases relative to primary).

²Corrected for multiple testing $p < 0.05$ (brain metastases relative to primary).

³Bivariate analysis $p < 0.05$ (systemic metastases relative to brain metastases).

⁴Corrected for multiple testing $p < 0.05$ (systemic metastases relative to brain metastases)

breast cancer brain metastases. Similarly, the DNA synthesis protein RRM1, which is linked to chemosensitivity to fluoropyrimidine and gemcitabine-based chemotherapy,²³ and TS,²⁴ were shown to have increased expression in NSCLC brain metastases. This expression may be a result of the development of therapeutic resistance mechanisms to prior therapy or an intrinsic feature of rare tumor clones in the primary that have an increased propensity for aggressive behavior, including metastasis. A recent single institution study reported the genomic analysis of 500 metastatic cancer patients and also reported a notable incidence of defects in DNA repair in concordance with our findings.²⁵

Consistent with the literature, we identified less frequent expression of hormone receptor expression in breast cancer brain metastases^{26–28}—although in a much larger patient cohort and with the novel extension of this to androgen receptor (AR) status. Hormone receptor loss is a known marker of aggressive disease, negative prognosis and limited treatment options. It has been suggested that the loss of hormone receptor expression is due to dedifferentiation of the metastatic tumor²⁹ or the selection of cytogenetically unique clones that have an enhanced propensity to metastasize to distant sites.^{30,31} Alternatively, the loss of hormone receptor expression in breast cancer patients may represent selection bias as a result of treatment.³² Brain metastasis is a late event in hormone receptor-positive breast cancer. As a breast cancer loses its hormone sensitivity, it may behave more like triple-negative breast cancer, known to have a higher rate of brain metastasis and a worse prognosis.³³ Because our study did not employ paired primary tumor/metastasis samples, we cannot exclude the possibility that our brain metastasis cohort is enriched for triple-negative cases.

Specific to melanoma brain metastases was the increased expression of cMET, which is a receptor tyrosine kinase that binds HGF, thereby leading to the upregulation of oncogenic signaling cascades. The association of cMET with tumor

progression and brain metastasis has been previously documented in animal models using *cMET* knockdown demonstrating decreased occurrence of brain metastases by 70%.^{34–36} Therapies targeting *cMET* via the use of monoclonal antibodies are in various stages of clinical trial testing in multiple cancer histologies,³⁷ but the efficacy of such drugs in the central nervous system is unclear (NCT02414139, NCT02468661). Another therapeutic consideration in this setting would be cabozantinib, which is being evaluated in breast cancer patients with brain metastasis (NCT02260531).

PD-1 expression was very frequent in both brain metastases and the primaries of NSCLC and melanoma cancers. The interaction between the immune system, tumor and the tumor microenvironment has been the focus of intense investigation—specifically the role of the PD-1/PDL-1 signaling axis.³⁸ The efficacy of PD-1 inhibition (i.e., nivolumab and pembrolizumab) as a monotherapy or in combination therapy has shown favorable survival in clinical trials for patients with advanced melanoma and NSCLC.^{39–43} However, the role of immune checkpoint inhibitors in the treatment of CNS cancers, including metastasis, is currently being defined,^{44,45} and our current results provide further support for these agents in this context.

Although most genetic alterations were present at similar frequencies in the brain metastases and the primary tumors, there were some notable exceptions that were higher in the brain metastasis. *ABL1* mutations and *HER2* and *EGFR* amplification were common in breast cancer brain metastases, consistent with a prior study of whole-exome sequencing of 86 matched primary tumor/brain metastasis samples from multiple cancers.⁴⁶ In previous study, clinically actionable alterations not detected in matched primary tumors were found in more than 50% of brain metastases. Our study⁴⁶ also included matched samples and tumor pathologies that were not in our data set (e.g., renal cell carcinoma), some of which were highlighted for their alterations specific to brain metastases. It is also possible that in our

analysis, we were not powered to detect mutations present at a low frequency (subclonal events). Yet, our sample size was very large, which allowed for rigorous statistical analysis. Comparing the findings of the current study to smaller cohorts of published samples is challenging as the majority of studies focus on mutational changes/alterations in oncogenic signaling pathways while the major findings in our study focus on differential protein expression.^{47,48} For example, a study comparing marker profiles of the 43 tumors that had evaluable material from matched sets of primary breast cancer and brain metastases showed no statistically significant difference between the primary and metastatic breast carcinomas for any markers.⁴⁹ However, these authors did not assess for several of the markers evaluated in our study, which were found to be significantly higher in breast cancer brain metastases (i.e. TOP2A, TOPO1, ERCC1). An additional study by Thompson et al. compared 41 matched breast and brain metastases. These authors reported significant differences in p27kip1 and cyclin D1 expression between brain and primary lesions, however, they also did not assess the expression of the markers listed above.⁵⁰

Although our study is strengthened by its large sample size and wide array of markers analyzed, it has limitations. To collect such a large volume of data, a commercial database was utilized and clinical annotation is not available, including prior treatment regimens. Hence, we cannot account for the impact of treatment-related changes on the patterns of expression in the brain metastasis samples. Furthermore, this analysis does not clarify whether the differentially expressed proteins are the etiology for the development of brain metastasis or are a reflection of the unique brain microenvironment, which would be informative for preventive strategies. Additionally, we cannot exclude the possibility that tumors from different sites may represent variations in molecular subtype or different stages of the disease. Ideally matched samples would better resolve these questions; however, matched samples collected from multiple centers worldwide from a cohort size of this magnitude is not feasible and is a major challenge even at a single institution.²⁵ Yet, this is the largest data set with genomic/molecular characterization of brain metastases to date.

In conclusion, despite these limitations, we found enriched expression of a number of novel markers associated with

DNA synthesis, replication and repair in brain metastases compared with primary cancer specimens. This trend occurred across tumor histologies, supporting the prioritization of therapies that target these pathways. Although several of the therapeutic target candidates are not brain metastasis-specific genes, overexpressed proteins, or amplified genes, this data informs which targets would be suitable for therapeutic purposes during advanced stage metastasis, with the inclusion of brain metastasis patients.

Acknowledgements

We thank David M. Wildrick, Ph.D., for editorial assistance.

Conflict of interest

Amy B. Heimberger serves on the Caris Life Sciences Scientific Advisory Board and is a stockholder in the company. Santosh Kesari serves as a consultant to Caris Life Sciences. Joanne Xiu and Zoran Gatalica are employees of Caris Life Sciences. David Spetzler is President and Chief Scientific Officer of Caris Life Sciences. Sandeep K. Reddy was an employee of Caris Life Science at the time of manuscript preparation. Priscilla K. Brastianos serves as a consultant for Eli Lilly and Angiobond and honorarium for Merck and Genentech. Michael A. Davies serves on the advisory board for Novartis, Genentech, Bristol-Meyers Squibb, Vaccines, Synder, Sanofi-Aventis and is the PI of grants from Genentech, Bristol-Meyers Squibb, Myriad, Merck and Sanofi-Aventis. John F. DeGroot is a consultant for Celldex, Deciphera Pharmaceuticals, AbbVie, FivePrime Therapeutics, Inc., GW Pharma, Carthera, Eli Lilly, Boston Biomedical Inc, Kairos Venture Investments, Syneos Health and Monteris; serves on the advisory board for Genentech, Celldex, Foundation Medicine Inc., Novogen, Deciphera, Astrazeneca, Insys Therapeutics, Kadmon, Merck, Eli Lilly; has stock in Ziopharm Oncology and Gilead; has research support from Sanofi-Aventis, Astrazeneca, EMD-Serono, Eli Lilly, Novartis, Deciphera Pharmaceuticals, Mundipharma; has other financial/material interests in DSMB: VBL Therapeutics, DSMB: Novella, VBI Vaccines Inc. and his spouse is an employee of Ziopharm Oncology.

References

- Nayak L, Lee EQ, Wen PY. Epidemiology of brain metastases. *Curr Oncol Rep* 2012;14(1):48–54.
- Brastianos PK, Curry WT, Oh KS. Clinical discussion and review of the management of brain metastases. *J Natl Compr Canc Netw* 2013;11(9):1153–64.
- Eichler AF, Chung E, Kodack DP, et al. The biology of brain metastases—translation to new therapies. *Nat Rev Clin Oncol* 2011;8(6):344–56.
- Ali A, Goffin JR, Arnold A, et al. Survival of patients with non-small-cell lung cancer after a diagnosis of brain metastases. *Curr Oncol* 2013;20(4):e300–6.
- Rostami R, Mittal S, Rostami P, et al. Brain metastasis in breast cancer: a comprehensive literature review. *J Neurooncol* 2016;127(3):407–14.
- Spagnolo F, Picasso V, Lambertini M, et al. Survival of patients with metastatic melanoma and brain metastases in the era of MAP-kinase inhibitors and immunologic checkpoint blockade antibodies: A systematic review. *Cancer Treat Rev* 2016;45:38–45.
- Fox BD, Cheung VJ, Patel AJ, et al. Epidemiology of metastatic brain tumors. *Neurosurg Clin N Am* 2011;22(1):1–6.
- Davies MA. Targeted therapy for brain metastases. *Adv Pharmacol* 2012;65:109–42.
- Gounder MM, Nayak L, Sahebjam S, et al. Evaluation of the safety and benefit of phase I oncology trials for patients with primary CNS tumors. *J Clin Oncol* 2015;33(28):3186–92.
- Garber ST, Hashimoto Y, Weathers SP, et al. Immune checkpoint blockade as a potential therapeutic target: surveying CNS malignancies. *Neuro Oncol* 2016;18(10):1357–66.
- Xiu J, Piccioni D, Juarez T, et al. Multi-platform molecular profiling of a large cohort of

- glioblastomas reveals potential therapeutic strategies. *Oncotarget* 2016;7(16):21556–69.
12. Benjamini YH. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J Roy Statist Soc B* 1995;57:289–300.
 13. Palmieri D, Bronder JL, Herring JM, et al. Her-2 overexpression increases the metastatic outgrowth of breast cancer cells in the brain. *Cancer Res* 2007;67(9):4190–8.
 14. Zhang L, Ridgway LD, Wetzel MD, et al. The identification and characterization of breast cancer CTCs competent for brain metastasis. *Sci Transl Med* 2013;5(180):180ra148.
 15. Huang H, Liu J, Meng Q, et al. Multidrug resistance protein and topoisomerase 2 alpha expression in non-small cell lung cancer are related with brain metastasis postoperatively. *Int J Clin Exp Pathol* 2015;8(9):11537–42.
 16. Depowski PL, Rosenthal SI, Brien TP, et al. Topoisomerase IIalpha expression in breast cancer: correlation with outcome variables. *Mod Pathol* 2000;13(5):542–7.
 17. Kauffmann A, Rosselli F, Lazar V, et al. High expression of DNA repair pathways is associated with metastasis in melanoma patients. *Oncogene* 2008;27(5):565–73.
 18. Song L, Robson T, Doig T, et al. DNA repair and replication proteins as prognostic markers in melanoma. *Histopathology* 2013;62(2):343–50.
 19. Crivellari D, Pagani O, Veronesi A, et al. High incidence of central nervous system involvement in patients with metastatic or locally advanced breast cancer treated with epirubicin and docetaxel. *Ann Oncol* 2001;12(3):353–6.
 20. Panvichian R, Tantiwetrueangdet A, Angkathunyakul N, et al. TOP2A amplification and overexpression in hepatocellular carcinoma tissues. *BioMed Res Int* 2015;2015:381602.
 21. Altaha R, Liang X, Yu JJ, et al. Excision repair cross complementing-group 1: gene expression and platinum resistance. *Int J Mol Med* 2004; 14(6):959–70.
 22. Reed E. Platinum-DNA adduct, nucleotide excision repair and platinum based anti-cancer chemotherapy. *Cancer Treat Rev* 1998;24(5):331–44.
 23. Lee JJ, Maeng CH, et al. The immunohistochemical overexpression of ribonucleotide reductase regulatory subunit M1 (RRM1) protein is a predictor of shorter survival to gemcitabine-based chemotherapy in advanced non-small cell lung cancer (NSCLC). *Lung Cancer* 2010;70(2):205–10.
 24. Cascinu S, Aschele C, Barni S, et al. Thymidylate synthase protein expression in advanced colon cancer: correlation with the site of metastasis and the clinical response to leucovorin-modulated bolus 5-fluorouracil. *Clin Cancer Res* 1999;5(8):1996–9.
 25. Robinson DR, Wu YM, Lonigro RJ, et al. Integrative clinical genomics of metastatic cancer. *Nature* 2017;548(7667):297–303.
 26. Bachmann C, Grischke EM, Fehm T, et al. CNS metastases of breast cancer show discordant immunohistochemical phenotype compared to primary. *J Cancer Res Clin Oncol* 2013;139(4): 551–6.
 27. Curtit E, Nerich V, Mansi L, et al. Discordances in estrogen receptor status, progesterone receptor status, and HER2 status between primary breast cancer and metastasis. *Oncologist* 2013;18(6): 667–74.
 28. Shen Q, Sahin AA, Hess KR, et al. Breast cancer with brain metastases: clinicopathologic features, survival, and paired biomarker analysis. *Oncologist* 2015;20(5):466–73.
 29. Bachmann C, Grischke EM, Staebler A, et al. Receptor change-clinicopathologic analysis of matched pairs of primary and cerebral metastatic breast cancer. *J Cancer Res Clin Oncol* 2013; 139(11):1909–16.
 30. Heim S, Teixeira MR, Dietrich CU, et al. Cytogenetic polyclonality in tumors of the breast. *Cancer Genet Cytogenet* 1997;95(1):16–9.
 31. Kuukasjarvi T, Karhu R, Tanner M, et al. Genetic heterogeneity and clonal evolution underlying development of asynchronous metastasis in human breast cancer. *Cancer Res* 1997;57(8): 1597–604.
 32. Rasbridge SA, Gillett CE, Seymour AM, et al. The effects of chemotherapy on morphology, cellular proliferation, apoptosis and oncoprotein expression in primary breast carcinoma. *Br J Cancer* 1994;70(2):335–41.
 33. Sim HW, Morris PG, Patil S, et al. Brain metastases in breast cancer. *Expert Rev Anticancer Ther* 2014;14(2):173–83.
 34. Benedettini E, Sholl LM, Peyton M, et al. Met activation in non-small cell lung cancer is associated with de novo resistance to EGFR inhibitors and the development of brain metastasis. *Am J Pathol* 2010;177(1):415–23.
 35. Breindel JL, Haskins JW, Cowell EP, et al. EGF receptor activates MET through MAPK to enhance non-small cell lung carcinoma invasion and brain metastasis. *Cancer Res* 2013;73(16): 5053–65.
 36. Preusser M, Streubel B, Berghoff AS, et al. Amplification and overexpression of CMET is a common event in brain metastases of non-small cell lung cancer. *Histopathology* 2014;65(5):684–92.
 37. Blumenschein GR Jr, Mills GB, Gonzalez-Angulo AM. Targeting the hepatocyte growth factor-cMET axis in cancer therapy. *J Clin Oncol* 2012;30(26):3287–96.
 38. Dong H, Strome SE, Salomao DR, et al. Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion. *Nat Med* 2002;8(8):793–800.
 39. Borghaei H, Paz-Ares L, Horn L, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015; 373(17):1627–39.
 40. Brahmer J, Reckamp KL, Baas P, et al. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 2015;373(2): 123–35.
 41. Hamid O, Robert C, Daud A, et al. Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *N Engl J Med* 2013;369(2):134–44.
 42. Larkin J, Chiarion-Sileni V, Gonzalez R, et al. Combined nivolumab and ipilimumab or monotherapy in untreated melanoma. *N Engl J Med* 2015;373(1):23–34.
 43. Robert C, Schachter J, Long GV, et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med*. 2015; 372(26):2521–32.
 44. Goldberg SB, Gettinger SN, Mahajan A, et al. Pembrolizumab for patients with melanoma or non-small-cell lung cancer and untreated brain metastases: early analysis of a non-randomised, open-label, phase 2 trial. *Lancet Oncol* 2016;17(7): 976–83.
 45. Margolin K, Ernstoff MS, Hamid O, et al. Ipilimumab in patients with melanoma and brain metastases: an open-label, phase 2 trial. *Lancet Oncol* 2012;13(5):459–65.
 46. Brastianos PK, Carter SL, Santagata S, et al. Genomic characterization of brain metastases reveals branched evolution and potential therapeutic targets. *Cancer Discov* 2015;5(11):1164–77.
 47. Chen G, Chakravarti N, Aardalen K, et al. Molecular profiling of patient-matched brain and extracranial melanoma metastases implicates the PI3K pathway as a therapeutic target. *Clin Cancer Res* 2014;20(21):5537–46.
 48. Goswami RS, Patel KP, Singh RR, et al. Hotspot mutation panel testing reveals clonal evolution in a study of 265 paired primary and metastatic tumors. *Clin Cancer Res* 2015;21(11): 2644–51.
 49. Brogi E, Murphy CG, Johnson ML, et al. Breast carcinoma with brain metastases: clinical analysis and immunoprofile on tissue microarrays. *Ann Oncol* 2011;22(12):2597–603.
 50. Thomson AH, McGrane J, Mathew J, et al. Changing molecular profile of brain metastases compared with matched breast primary cancers and impact on clinical outcomes. *Br J Cancer* 2016;114(7):793–800.