

UCSF

UC San Francisco Previously Published Works

Title

A longitudinal study of persistent smoking among HIV-positive gay and bisexual men in primary relationships

Permalink

<https://escholarship.org/uc/item/4xv6g9f5>

Authors

Gamarel, Kristi E
Neilands, Torsten B
Conroy, Amy A
[et al.](#)

Publication Date

2017-03-01

DOI

10.1016/j.addbeh.2016.11.019

Peer reviewed



Published in final edited form as:

Addict Behav. 2017 March ; 66: 118–124. doi:10.1016/j.addbeh.2016.11.019.

A longitudinal study of persistent smoking among HIV-positive gay and bisexual men in primary relationships

Kristi E. Gamarel, PhD^{1,2}, Torsten B. Neilands, PhD³, Amy A. Conroy, PhD³, Samantha E. Dilworth, MS³, Nadra Lisha, PhD³, Jonelle M. Taylor, BA³, Lynae A. Darbes, PhD⁴, and Mallory O. Johnson, PhD³

¹Center for Alcohol and Addiction Studies, Brown University School of Public Health

²Department of Behavioral and Social Sciences, Brown University School of Public Health

³Department of Medicine, University of California, San Francisco

⁴Department of Health Behavior and Biological Sciences, School of Nursing, University of Michigan

Abstract

Introduction—We examined the stability of smoking behaviors, and factors associated with persistent smoking in a longitudinal study of HIV-positive gay and bisexual men in primary relationships.

Methods—A sample of 377 HIV-positive men on antiretroviral therapy and their same-sex partners completed five assessments over two years. Participants completed semi-structured interviews which assessed smoking status, sociodemographic factors, relationship dynamics, and HIV-related disease characteristics. Latent transition analysis estimated the amount of transition in smoking over time. Latent class analysis examined factors associated with smoking status across the study period.

Results—At baseline, 28.1% ($n = 106$) of participants reported current smoking. Over 90% of the HIV-positive men remained in the same smoking category over time (68.4% persistent non-smokers; 24.1% persistent smokers). Men whose partners smoked and men with lower income had higher odds of being persistent smokers, whereas older men and men who identified as Latino race/ethnicity had lower odds of being persistent smokers compared to non-smokers.

Conclusions—Despite efforts to reduce smoking among people living with HIV (PLWH), a substantial subset of men continued to smoke during their two years in the study. Findings suggest that primary partners who also smoke and low income were the strongest predictors of sustained smoking behaviors among HIV-positive men. Additional research is needed to better understand how to increase motivation and support for smoking cessation among PLWH and their primary partners, while attending to how socioeconomic status may inhibit access to and the sustained impact of existing smoking cessation programs.

1. Introduction

Tobacco use continues to be one of the leading causes of preventable morbidity and premature mortality in the United States and is a well-recognized risk factor for chronic diseases including cardiovascular disease, pulmonary disease, and cancer (Centers for Disease Control and Prevention, 2015). Among people living with HIV (PLWH) and in particular, HIV-positive gay, bisexual and other men who have sex with men (MSM), smoking is a highly prevalent behavior (Friis-Moller et al., 2003; Gritz, Vidrine, Lazev, Amick, & Arduino, 2004; Lifson et al., 2010; Mamary, Bahrs, & Martinez, 2002; Mdodo et al., 2015; O'Cleirigh, Dale, et al., 2015; O'Cleirigh, Valentine, et al., 2015; Vittecoq et al., 2003). Among HIV-positive individuals, smoking has been linked to an increased likelihood of HIV-related medical complications (Humfleet, Hall, Delucchi, & Dilley, 2013), and has been shown to negatively impact immune and virologic response (O'Cleirigh et al., 2014).

Despite advances in smoking cessation interventions, available data on smoking cessation interventions among PLWH indicate that cessation rates are low (Humfleet et al., 2013). In fact, fewer than 8% of HIV-positive smokers are actively engaged in any type of smoking-cessation program (Cioe, Crawford, & Stein, 2014), and engagement is even lower among HIV-positive gay and bisexual men (Webb, Vanable, Carey, & Blair, 2007). Thus, PLWH and in particular HIV-positive gay and bisexual men have been cited as a high-priority group for smoking cessation interventions; however, additional research is needed to better understand the relative stability of smoking over time among HIV-positive gay and bisexual men outside of the context of a controlled, intervention study, as well as which risk factors are associated with persistent smoking in order to develop better smoking prevention and cessation interventions.

Prior research has shown a range of factors associated with smoking behaviors among PLWH. With few exceptions (Pacek, Latkin, Crum, Stuart, & Knowlton, 2014), studies have largely focused on individual-level characteristics such as younger age, as well as comorbid depression and alcohol use which decreases the probability of successful smoking cessation (Gritz et al., 2004). Additionally, HIV-positive gay and bisexual men may also cope with unique minority stressors such as internalization of negative messages about one's sexual identity or expression, which can lead to maladaptive coping responses such as substance use (Meyer, 2003); these unique minority stressors have been associated with tobacco use among gay and bisexual men (Gamarel, Neilands, Dilworth, Taylor, & Johnson, 2015; Pachankis, Hatzenbuehler, & Starks, 2014). Moreover, recent evidence suggests that living with another smoker is an increased risk factor for sexual minority men specifically (Gamarel et al., 2016).

Close relationships are consistently linked to better health outcomes across a number of conditions (Lewis et al., 2006) and within couples affected by HIV, primary partners play a positive role in HIV-related health outcomes (Johnson et al., 2012). The mutual influence of partners on each other's health—or dyadic interdependence—can have positive and negative effects on health behaviors, such as smoking (Lewis et al., 2006). For example, individuals who are partnered with smokers are more likely to smoke themselves (Sutton, 1993), and less likely to quit if their partner also smokes (Chandola, Head, & Bartley, 2004; Cobb et al.,

2004; Holahan et al., 2012; Homish & Leonard, 2005; McBride et al., 1998). Conversely, positive influences related to relationship commitment and provision of partner encouragement to engage in healthy behaviors may contribute to health-enhancing behaviors such as quitting smoking to protect the longevity of one's relationship (Weiselquist, Rusbult, Foster, & Agnew, 1999). Thus, it is likely that relationship factors – such as commitment level and partner encouragement– may strongly influence smoking behaviors over time among HIV-positive men in same-sex relationships, given the previously observed associations with smoking behaviors among the general population.

Despite effective public health campaigns, public policy initiatives, and improvements in smoking cessation treatment programs, PLWH continue to be a group which warrants future study. To date, few studies have conducted longitudinal analyses to understand factors associated with smoking behaviors over time among HIV-positive men in same-sex relationships. Using data from a longitudinal study of HIV-positive gay and bisexual men and their same-sex primary partners, we explored the following questions: a) what is the stability of smoking status over a two-year period?, and b) to what extent is partner smoking status and relationship factors associated with smoking compared to non-smoking across a two-year period over and above existing individual and social factors? Based on existing literature showing low smoking cessation rates among PLWH in the context of intervention studies and unique minority stressors experienced by gay and bisexual men, we hypothesized that smoking status would be relatively stable across participants' two years in the study. Secondly, we hypothesized that partner smoking status and perceptions of partner encouragement would be robust predictors of smoking status over time, such that partners who also smoked and those who perceived their partners were a negative influence on their health would have greater odds of persistent smoking over time. Additionally, we hypothesized that relationship commitment would be protective such that higher levels of commitment would be predictive of non-smoking behaviors over time.

2. Methods

The data derive from the Duo Project, a longitudinal study of 266 same-sex male couples in which at least one partner was HIV-positive. While the primary purpose of the Duo Project was to examine relationship dynamics and ART adherence over time, we also collected one item on tobacco use. Details of the study procedures have been described elsewhere (Conroy, Gamarel, Neilands et al., 2016; Johnson et al., 2012). In total, 532 men participated in the baseline assessment, of which 377 were HIV-positive men on ART. Couples were enrolled for a two-year period and data collection began in January 2009 and ended in September 2014. Participants were recruited from the San Francisco Bay Area in the United States (U.S.) using passive recruitment strategies, and participant and provider referrals. Flyers were posted in clinics, community bulletin boards, AIDS service organizations, and at other community-based organizations. Media ads were placed online and in print publications targeting HIV-positive and gay/bisexual men. Interested individuals contacted study staff for more information on the study. Men were eligible for the study if they met the following criteria: (1) in a primary relationship, which was defined as “currently (for at least 3 months) in a relationship with someone you feel committed to above anyone else and with whom you have had a sexual relationship,” which is the definition used in many couples-

based HIV prevention studies (Beougher et al., 2015; Mitchell & Sullivan, 2015); (2) at least one partner in the relationship is HIV-positive and on an acknowledged ART regimen for at least 30 days; (3) at least 18 years old; (4) born male and currently identify as male; (5) English-speaking; and (6) able and willing to provide informed consent.

Partners were screened separately over the phone to assess eligibility and if both partners were eligible, couples were scheduled for an in-person interview at the study research center. Both partners were required to attend the assessment appointments together, but were separated during data collection. Data were collected using a combination of Computer Assisted Personal Interviewing (CAPI) and Audio Computer Assisted Self Interviewing (ACASI) methods, which optimize data integrity through the reduction of data entry errors while minimizing the effects of social desirability bias (Turner, Ku, Rogers et al., 1998). Couples were assessed at baseline and every six months thereafter for a total of five assessment waves. All HIV-positive participants had blood drawn for viral load tests at their baseline, 12-month, and 24-month visits.

Of the 377 HIV-positive participants on ART, 21% were lost during the course of the study to relationship breakup, 4% to relocation or travel, 2% to the death of a partner, and 6% withdrew or were withdrawn from the study due to inability to follow study protocols.

Ethical approval was obtained from the Committee on Human Research at the University of California, San Francisco. Written informed consent was obtained from all participants. Each partner of the couple was paid US \$50 for each survey completed and HIV-positive participants were paid an additional \$10 for each blood sample.

2.1. Measures

Smoking Status—At each assessment, participants were asked “Do you currently smoke cigarettes?” Answers to this question were used to classify participants as nonsmokers (reported not smoking at that time point) or current smokers (reported smoking at that time point). Smoking status at each time point was coded as smoking (1) versus non-smoking (0). Participants were also asked on average how many cigarettes they smoked per day during the past three months.

Partner Smoking Status—Partner smoking status was assessed by a variable indicating whether the partner was currently a smoker (yes/no).

Relationship Factors—We assessed relationship factors which may impact smoking persistence. First relationship commitment was assessed with a 4-item scale consisting of statements such as “I am committed to maintaining my relationship with my partner” (Kurdek, 1995). Response options ranged from 1 (not at all true) to 9 (extremely true) ($\alpha = 0.95$). Second, a lack of partner health behavior encouragement was assessed with one item that asked participants to endorse how much their partners contributed to their health (i.e., “How often does your partner encourage you to engage in healthy behaviors (e.g. exercising, eating healthier)”) Response options ranged from 1 (always encourages) to 5 (never encourages) with higher numbers indicating less support.

Sociodemographic Characteristics—Participants reported their age, race and ethnicity, education level, and income level. In our analyses race/ethnicity was represented as a series of indicator variables for Black, Latino, and other non-White race/ethnicity with White as the reference group. Because only 22 participants reported less than a GED or High School degree, education was coded as achieving high school graduation or equivalent (1) vs. less than high school education (0). This cutoff was also chosen given evidence that smokers with 12 years of schooling or less have the highest odds of persistent smoking as compared to smokers who completed college (Breslau, Johnson, Hiripi, & Kessler, 2001). Similarly, evidence indicates that low annual income levels are associated with persistent smoking (Slopen et al., 2013); thus, income level was dichotomized as less than \$20,000 (1) versus \$20,000 or more (0). This cut off was chosen given the small number of participants who reported earning less than \$10,000 annually (21.8%) and evidence that earning less than \$20,000 in San Francisco is considered low income. Participants also reported their relationship duration (in years, averaged across the couple) and HIV-positive partners reported the length of time since initiating ART (in years).

Depressive symptoms—The 20-item Center for Epidemiologic Studies Depression Scale (CES-D) was administered to measure depressed mood in the past week (Radloff, 1977). The CES-D consists of 20 items (e.g., “could not get going”). Participants responded on a 4-point scale ranging from 0 = “rarely or none of the time” to 3 = “most or all of the time.” After reverse scoring items 4, 8, 12, and 16, individual items were summed such that higher scores indicate greater depressive symptoms. Total possible scores range from 0 to 60. Internal consistency was acceptable within our sample ($\alpha = 0.91$).

Alcohol Use—Alcohol use was assessed with the 10-item Alcohol Use Disorders Identification Test (AUDIT) (Saunders, Aasland, Baber, De la Fuente, & Grant, 1993), which has been validated in samples of PLWH (Strauss & Rindskopf, 2009). The AUDIT contains three questions which assess alcohol use and seven items on alcohol problems/ alcohol dependence and subsequent problems. Each question has five responses which are scored from 0 to 4. Total scores ranged from 0 to 40. For analyses, we used a continuous variable such that higher values indicated greater alcohol use and potential problems.

Internalized Heterosexism—Internalized heterosexism, which is the extent to which individuals internalize negative societal attitudes about their sexual identity, was assessed via four items used in prior studies with HIV-positive gay and bisexual men ($\alpha = 0.75$) (Johnson, Carrico, Chesney, & Morin, 2008; Nungesser, 1983). Example items include: “I wish I were heterosexual,” “Whenever I think a lot about being gay, I feel critical about myself.” Each question has six responses which are scored from 0 to 5 and higher scores indicate greater internalized heterosexism.

HIV Medication Adherence and Disease Characteristics—For HIV-positive men, ART adherence was assessed using the Visual Analog Scale (VAS), which measures the proportion of ART medications taken in the past 30 days (Walsh, Mandalia, & Gazzard, 2002). The majority of participants in this sample self-reported nearly perfect ART adherence ($M = 94.27$, $Mdn = 98.00$, $SD = 11.61$); therefore, prior studies using these data

(Gamarel, Neilands, Dilworth, Taylor, & Johnson, 2015; Johnson et al., 2012) have dichotomized medication adherence as 100% adherence (1) versus less than 100% adherence (0). This cut off was also chosen to reflect the distinction between those who meet the public health goal of 100% medication adherence versus those who do not. HIV viral load assays were performed to detect HIV RNA at or above 20 copies/mL (Roche Molecular Systems, Inc.). Viral load was dichotomized as detectable (1) versus undetectable (0). Couple HIV serostatus was dichotomized as concordant HIV-positive (1) versus HIV-discordant (0).

2.2. Statistical Analyses

One-way frequency tables were computed for all analysis variables and measures of central tendency and variability were computed for continuous variables. Our primary data analyses were constructed to address our two primary research questions: 1) whether smoking status remained stable over time and 2) which of the pre-specified baseline variables of substantive interest described previously were predictive of smoking status over the study's follow-up period. To examine whether smoking status changed over time, latent transition analysis (LTA) with a first-order hidden Markov model was used to estimate transition probabilities from one wave to the next. The likelihood ratio test statistic (LRT) was computed to assess LTA model fit and nested LRTs were computed to compare nested LTA models. Because the LTA revealed that more than 90% of the participants maintained their original smoking status over all five assessment waves, latent class analysis (LCA) was used to examine associations between persistent smoking and baseline covariates. Due to the lack of complete smoking data across all assessment waves, persistent smoking was conceptualized as a latent variable measured by the available observed data so that cases with partial data could be included in the analysis along with cases with complete data. One-, two-, and three-class solutions were examined and compared using the Lo-Mendell-Rubin adjusted likelihood ratio test statistic to determine the optimal number of classes for the latent variable of persistent smoking (Nylund, Asparouhov, & Muthén, 2007). In addition, for all LTAs and LCAs containing two or more classes, the entropy statistic, a measure of classification quality, was computed. Entropy values range from 0 to 1, with larger values indicating better classification quality. *Mplus 7.4* was used for all LTA and LCA analyses.

Based on prior literature, we were particularly interested in the extent to which relationship factors were associated with persistent smoking, after accounting for individual-level covariates. Baseline covariates included in the LCA were age, high school graduation, Black race, Latino race, other non-White race, relationship duration, low income, CES-D score, AUDIT score, internalized heterosexism, length of time since initiating ART, 100% HIV medication adherence, detectable baseline viral load, and couple HIV-serostatus. Latent smoking status was regressed onto the baseline covariates using logistic regression. For age, high school graduation, race/ethnicity, low income, CES-D, AUDIT, internalized heterosexism, relationship commitment, and lack of health support, both actor and partner effects were included in line with methods developed in the dyadic research literature to elucidate partner influences on behavior after controlling for actors' own influences using the Actor-Partner Interdependence Model (APIM) approach (Kenny, Kashy, & Cook, 2006).

All inferential analyses employed a cluster-adjusted robust variance estimator (*Mplus* MLR estimator) to adjust inferences for nesting of participants within couples.

Following descriptive analyses to characterize the sample, our subsequent LTA and LCA analyses proceeded sequentially such that the LCA was informed by the results of the LTA. Specifically, first LTA addressed our research question assessing the stability of smoking status over time. Our second research question sought to examine which relationship dynamics are associated with persistent smoking, which was investigated using LCA once we established that latent smoking status did not change over the two years of the study in the initial LTA.

3. Results

Table 1 presents the baseline sample characteristics of the HIV-positive men on ART. Participants ranged in age from 22 to 69 years old ($M = 46.15$ years, $SD = 9.93$). The majority of the sample (95.8%) had earned a high school degree or higher, and over half earned less than \$20,000 annually. The majority of the men (91.5%) identified as gay, and slightly more than half (55.2%) identified as white race/ethnicity. Relationship duration averaged 6.52 years ($SD = 4.94$) and about two-thirds of the sample (68.7%) were in HIV-positive seroconcordant relationships. HIV-positive participants on ART had been diagnosed an average of 13.45 years ($SD = 8.04$), had been taking ART for 9.85 ($SD = 7.04$) years on average, and nearly half (44.8%) had a detectable HIV viral load. At baseline, over one-quarter of the 377 participants ($n = 106$, 28.1%) reported currently smoking cigarettes, with a range of 1 to 90 cigarettes smoked per day ($M = 11.70$, $SD = 13.61$) and remained fairly stable over time with a slight decrease to 22.1% at two years. Attrition was moderate: 8% at 6 months, 17% at 12 months, 27% at 18 months, and 32% at 24 months.

Latent Transition Analysis (LTA)

An initial LTA model was fitted, which allowed the transition probabilities to vary across comparisons of waves. The fit of the LTA model to the data was satisfactory ($\chi^2(20) = 26.80$, $p = .14$). As shown in Table 2, transition probabilities indicated that greater than 90% of the participants remained in the same class from one assessment visit to the next. The entropy for the model was .943, indicating very good classification quality. Due to the similarities in transition probabilities across time, a second, more restricted LTA, was estimated in which the transition matrices were held equal across time. The fit of this model to the data was also satisfactory ($\chi^2(26) = 30.16$, $p = .26$). The entropy for this model was .945, indicating similar or equivalent classification quality of the more parsimonious LTA. A nested likelihood ratio chi-square test indicated the simplified LTA with the equivalent transition probabilities fit the data equally as well as the LTA model which allowed the transition probabilities to differ across time ($\chi^2(6) = 3.15$, $p = .79$). In other words, there were no significant differences in the transition probabilities across the four transition points, implying that participants were equally likely to remain in or move out of the persistent smokers class into the persistent non-smokers class across the four transition points. The final estimated proportion of persistent non-smokers was 68.4% and the final estimated proportion of persistent smokers was 24.1%, resulting in 92.5% of participants exhibiting

persistent status. Based on these results, we proposed a two-class latent class analysis containing a persistent smoker class and a persistent non-smoker class.

Latent Class Analysis (LCA)

A series of initial unconditional LCAs were fitted to evaluate support for the hypothesized two-class model. These analyses considered models with one, two, and three latent classes, respectively. LCAs with k classes were compared with corresponding LCAs with $k-1$ classes using the Lo-Mendell-Rubin adjusted likelihood ratio test. Results indicated that the two-class LCA fit the data no worse than the three-class LCA ($\chi^2(6) = 40.54, p = .07$), but the two-class LCA was strongly preferred to the one-class LCA ($\chi^2(6) = 815.12, p < .0001$). The entropy for the two-class model was .985 whereas the entropy for the three-class model was .889, indicating that the two-class model had better classification quality. Therefore, we fit a conditional two-class LCA including all covariates. Four participants were missing viral load assay information, yielding an analysis N of 373. The entropy for this model was .951. Results of the logistic regression of latent class membership onto the pre-specified baseline covariates appear in Table 3. Participants with older age and Latino race/ethnicity had lower odds of being a persistent smoker compared to being a persistent non-smoker. However, low-income participants and those whose partners also smoked had higher odds of being persistent smokers compared to being a persistent non-smoker. There were no significant differences in commitment level and partner support.

4. Discussion

In this study, over one-quarter of HIV-positive men reported current smoking at the baseline assessment, with 92.5% of participants estimated to remain in the same smoking category across the two years of study, and 24.1% of the participants estimated to be persistent smokers. While the smoking prevalence was lower than other samples of HIV-positive adults (Cioe et al., 2016; Lifson et al., 2010; O'Cleirigh et al., 2015), these findings are consistent with prior research illustrating that people living with HIV are significantly less likely to quit smoking, despite the availability of pharmacologic and behavioral smoking cessation aids. Prior studies have reported relatively high motivation to quit smoking among people living with HIV and high rates of quit attempts (Benard et al., 2007); however, this readiness may not appear to translate into successful smoking cessation, as demonstrated in this study and others (Mdodo et al., 2015). Several smoking cessation trials to evaluate intensive counseling and cell phone interventions with PLWH suggest that these interventions can be efficacious, but studies were limited by short follow-up or a non-randomized design (Cioe, 2013). Another study demonstrated increased smoking cessation rates following the implementation of a training program for HIV clinicians (Huber et al., 2012). However, the limited success of smoking cessation interventions necessitates moving beyond individual-level factors to identify important interpersonal and structural barriers to uptake and sustained use of smoking cessation programs for PLWH that can be readily applied within current care models.

Consistent with previous findings among non-HIV-positive samples (Brath, Grabovac, Schalk, Degen, & Dorner, 2016; Reynolds, Neidig, & Wewers, 2014), we found that having

a partner who smoked was a robust predictor of persistent smoking compared to persistent non-smoking among HIV-positive men. Partners who also smoke have been cited as a significant barrier to cessation (Christakis & Fowler, 2008), and couples in which both members smoke have significantly lower quit rates and higher relapse rates (Ferguson, Bauld, Chesterman, & Judge, 2005; Garvey, Bliss, Hitchcock, Heinold, & Rosner, 1992; Palmer, Baucom, & McBride, 2000). Thus, health care providers should assess and address the smoking status of the partner when conducting smoking cessation programs to identify and help patients resolve barriers to cessation. For example, health care providers who invite and teach partners how to support their partners in quitting has been shown to be an effective strategy to promote smoking cessation (McBride et al. 2004).

Notably, we did not find that relationship factors such as commitment level and the provision of partner encouragement for engaging in healthy behaviors to be associated with persistent smoking. Longitudinal studies have provided evidence that higher support is related to less smoking in general but these studies have not specifically examined persistent smoking as an outcome (Mermelstein, Cohen, Lichtenstein, Baer, & Kamarck, 1986). Our lack of an association with partner encouragement is somewhat consistent with the disappointing results of other intervention studies using partner support to increase smoking cessation (Cohen et al., 1989; Lichtenstein, Glasgow, Lando, Ossip-Klein, & Boles, 1996; Park, Tudiver, & Campbell, 2012). Future research is warranted to better understand each partner's motivations for and barriers to smoking cessation to guide smoking cessation interventions with couples.

Socioeconomic disadvantage as measured by low income was associated with greater odds of persistent smoking compared to persistent non-smoking across the two years of the study. While some evidence suggests that the proportion of low-income smokers who want to quit is equivalent to the general population (Lebrun-Harris, Fiore, Tomoyasu, & Ngo-Metzger, 2015), other studies illustrate that few low-income smokers attempt to quit (Burkhalter, Springer, Chhabra, Ostroff, & Rapkin, 2005; Caleyachetty, Lewis, McNeill, & Leonard-Bee, 2012; Reid, Hammond, Boudreau, Fong, & Siahpush M., 2010). Additionally, when low-income smokers do try to quit, they are less likely to be successful than other smokers (Caleyachetty et al., 2012; Hiscock, Judge, & Bauld, 2011; Levy, Romano, & Mumford, 2005), which may be a result of limited insurance coverage, financial constraints and reduced access to quality healthcare (DeNavas-Walt, Proctor, & Smith, 2009; Hiscock R., Bauld, Amos, Fidler, & Munafò M., 2012). There is also evidence that low income neighborhoods and those with higher rates of same-sex couples have a higher density of tobacco retailers (Cantrell et al., 2015; Lee, Pan, Henriksen, Goldstein, & Ribisl, 2016). Thus, HIV-positive gay and bisexual men in lower socioeconomic positions may be more likely to share social environments with other smokers, such as romantic partners, which has been associated with smoking initiation, continued smoking, and relapse after quitting (Hiscock, et al., 2012). These findings highlight the need for smoking cessation programs to account for the socioeconomic constraints that may impede access to and sustained use of evidence-based treatments. Evidence suggests that incorporating tobacco treatment within community agencies may be particularly helpful for lower SES smokers (Christiansen, Reeder, TerBeek, Fiore, & Baker, 2015). Specifically, brief motivational interviewing interventions for unmotivated smoker delivered by community health workers have showed

particular promise (Christiansen et al., 2015). Furthermore, a lack of social support is a common reason for why lower SES smokers fail to quit smoking and other research suggests that having access to non-smokers in one's social network can be a powerful aid to quitting (Hiscock et al., 2012). Thus, future research may include the development and testing of social network interventions designed to provide peer support and enhance motivation, which could be adapted from effective interventions with HIV-positive drug-using adults of lower socioeconomic status (Latkin, Sherman, & Knowlton 2003).

Internalized stigma, poor mental health, and alcohol use have been shown to increase the risk of smoking among PLWH (Gamarel et al., 2015; Gritz et al., 2004; Pacek et al., 2014). In previous cross-sectional analyses of the Duo Project, internalized heterosexism was positively associated with smoking (Gamarel et al., 2015). However, in this study, internalized heterosexism was not associated with persistent smoking across the two years of the study. Additional research is needed to better understand the associations between stigma, mental health, substance use, and smoking behaviors over time among PLWH.

4.1. Limitations

Our study has several limitations. Our sample of HIV-positive gay and bisexual men on ART in San Francisco may be less diverse than other communities heavily impacted by HIV in the United States, which limits generalizability. For example, our sample was predominately white, linked to HIV care, self-reported high ART adherence, and in a committed relationship, which may explain the lower smoker rates in this study compared to others with PLWH. Second, this was a longitudinal study and therefore participant attrition could be a concern if those who completed the study were different from those who did not. For example, the results could be biased by the greater participation of high-functioning couples who stayed together over the study period, which has been noted as a limitation in other couples studies (Conroy et al., 2016). However, by conceptualizing persistent smoking status as a latent variable, our modeling approach was able to include all participants, including those who dropped out, rather than only those who completed all five assessment waves, thereby broadening the generalizability beyond the population who completed all assessments. Another limitation was our reliance on self-reported smoking status; however, self-report has been established as a fairly reliable indicator of smoking status (Vartiainen, Seppälä, Lillsunde, & Puska, 2002). Finally, the data analyzed here were collected as part of a larger study of relationship dynamics and ART adherence among same-sex male couples. Although tobacco use behaviors were assessed, these measures did not include standardized measures of smoking history, nicotine dependency, interest and self-efficacy for quitting, and factors maintaining smoking behaviors. Additionally, we were unable to examine differences between non-daily and daily smokers. While evidence suggests that non-daily smokers have increased health risks (Lindson-Hawley et al. 2016; Shane, Ling, & Glantz, 2010), non-daily smokers tend to be social or weekend smokers and are less likely to consider themselves “smokers” compared to daily smokers (Shiffman et al. 2012). Thus, future studies should include a more extensive measurement of tobacco use behaviors, including couple-level motivators and barriers to smoking cessation.

4.2. Conclusions

This study is among the first to examine whether partner smoking status and relationship factors are associated with smoking behaviors over time among PLWH. Study findings provide valuable information about the need to attend to relationship and structural barriers to smoking cessation. Specifically, partners' smoking behaviors and low income appear to negatively influence the likelihood of changes in smoking behaviors and should be monitored routinely in primary care settings. Findings from this study also emphasize that members of an individual's social network, particularly primary partners, may strongly influence one's smoking behaviors. Thus, smoking cessation efforts in dual-smoker couples may need to address both partners' smoking at the same time in order to change behavior.

References

- Benard A, Bonnet F, Tessier JF, Fossoux H, Dupon M, Mercie P, et al. Tobacco addiction and HIV infection: toward the implementation of cessation programs. ANRS CO3 Aquitaine Cohort. *AIDS Patient Care and STDs*. 2007; 21(7):458–468. [PubMed: 17651027]
- Beougher SC, Bircher AE, Chakravarty D, Darbes LA, Mandic CG, Neilands TB, Hoff CC. Couple-level motivations to test for HIV for gay men in relationships. *Archives of Sexual Behavior*. 2015; 44(2):499–508. [PubMed: 25550145]
- Brath H, Grabovac I, Schalk H, Degen O, Dorner TE. Prevalence and correlates of smoking and reading to quit in people living with HIV in Austria and Germany. *PLoS One*. 2016:e0150553. [PubMed: 26919722]
- Breslau N, Johnson EO, Hiripi E, Kessler R. Nicotine dependence in the United States: Prevalence, trends, and smoking persistence. *Archives of General Psychiatry*. 2001; 58(9):810–816. [PubMed: 11545662]
- Burkhalter JE, Springer CM, Chhabra R, Ostroff JS, Rapkin BD. Tobacco use and readiness to quit smoking in low-income HIV-infected persons. *Nicotine and Tobacco Research*. 2005; 7:511–522. [PubMed: 16085522]
- Caleyachetty A, Lewis S, McNeill A, Leonardi-Bee J. Struggling to make ends meet: exploring pathways to understand why smokers in financial difficulties are less likely to quit successfully. *European Journal of Public Health*. 2012; 22(Suppl 1):41–48. [PubMed: 22294784]
- Cantrell J, Anesetti-Rothermel A, Pearson JL, Xiao H, Vallone D, Kirchner TR. The impact of tobacco retail outlet environment on adult cessation and differences by neighborhood poverty. *Addiction*. 2015; 119(1):152–161.
- Caleyachetty A, Lewis S, McNeill A, Leonardi-Bee J. Struggling to make ends meet: exploring pathways to understand why smokers in financial difficulties are less likely to quit successfully. *European Journal of Public Health*. 2012; 22(Suppl 1):41–8. [PubMed: 22294784]
- Centers for Disease Control and Prevention. Current Cigarette Smoking Among Adults—United States, 2005–2014. *Morbidity and Mortality Weekly Report*. 2015; 64(44):1233–1240. [PubMed: 26562061]
- Chandola T, Head J, Bartley M. Socio-demographic predictors of quitting: How important are household factors? *Addiction*. 2004; 99(6):770–777. [PubMed: 15139875]
- Chaturvedi AK, Pfeiffer RM, Chang L, Goedert JJ, Biggar RJ, Engels EA. Elevated risk of lung cancer among people with AIDS. *AIDS*. 2007; 21:207–213. [PubMed: 17197812]
- Christiansen BA, Reeder KM, TerBeek EG, Fiore MC, Baker TB. Motivating low socioeconomic smokers to accept evidence-based smoking cessation treatment: A brief intervention for the community agency setting. *Nicotine and Tobacco Research*. 2015; 17(8):1002–1011. [PubMed: 26180226]
- Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *New England journal of medicine*. 2008; 358(21):2249–2258. [PubMed: 18499567]
- Cioe PA. Smoking cessation interventions in HIV-infected adults in North America: A literature review. *Journal of Addictive Behaviors, Therapy, and Rehabilitation*. 2013; 2(3):100–112.

- Cioe PA, Crawford SL, Stein MD. Cardiovascular risk-factor knowledge and risk perception among HIV-infected adults. *Journal of the Association of Nurses in AIDS Care*. 2014; 25(1):60–69. [PubMed: 24070645]
- Cioe PA, Gamarel KE, Pantalone DW, Monti PM, Mayer KH, Kahler CW. Cigarette smoking and antiretroviral therapy (ART) adherence in a sample of heavy drinking HIV-infected men who have sex with men (MSM). *AIDS and Behavior*. 2016
- Cobb LK, McAdams-DeMarco MA, Huxley RR, Woodward M, Koton S, Coresh J, Anderson CAM. The association of spousal smoking status with the ability to quit smoking: The Atherosclerosis Risk in Communities Study. *American Journal of Epidemiology*. 2004; 179(10):1182–1187.
- Cohen S, Lichtenstein E, Prochaska JO, Rossi JS, Gritz ER, Carr CR, Abrams D. Debunking myths about self-quitting: Evidence from 10 prospective studies of persons who attempt to quit smoking by themselves. *American Psychologist*. 1989; 44(11):1355. [PubMed: 2589730]
- Conroy AA, Gamarel KE, Neilands TB, Dilworth SE, Darbes LA, Johnson MO. Relationship dynamics and partner beliefs about viral suppression: A longitudinal study of male couples living with HIV/AIDS (The Duo Project). *AIDS and Behavior*. 2016; 20(7):1572–83. [PubMed: 27150895]
- Crothers K, Butt AA, Gibert CL, Rodriguez-Barradas MC, Crystal S, Justice AC. Veterans Aging Cohort 5 Project Team. Increased COPD among HIV-positive compared to HIV-negative veterans. *Chest*. 2006; 130:1326–1333. [PubMed: 17099007]
- DeNavas-Walt, C., Proctor, BD., Smith, JC. *Income, Poverty, and Health Insurance Coverage in the United States: 2008*. Washington, DC: US Government Printing Office; 2009.
- Ferguson J, Bauld L, Chesterman J, Judge K. The English smoking treatment services: one-year outcomes. *Addiction*. 2005; 100(Suppl 2):59–69. [PubMed: 15755262]
- Friis-Moller N, Weber R, Reiss P, Thiebaut R, Kirk O, d'Arminio Monforte A, et al. Cardiovascular disease risk factors in HIV patients-association with antiretroviral therapy. Results from the DAD study. *AIDS*. 2003; 17(8):1179–1193. [PubMed: 12819520]
- Gamarel KE, Kahler CW, Lee JH, Reisner SL, Mereish EH, Matthews AK, Operario D. Sexual orientation disparities by smoking vary by sex and household smoking among US adults: Findings from the 2003–2012 National Health and Nutrition Examination Surveys. *Preventive Medicine*. 2016; 82:1–6. [PubMed: 26598804]
- Gamarel KE, Neilands TB, Dilworth SE, Taylor JM, Johnson MO. Smoking, internalized heterosexism, and HIV disease management among male couples. *AIDS Care*. 2015; 26(5):649–654.
- Garvey AJ, Bliss RE, Hitchcock JL, Heinold JW, Rosner B. Predictors of smoking relapse among self-quitters: a report from the Normative Aging Study. *Addictive Behaviors*. 1992; 17(4):367–377. [PubMed: 1502970]
- Gritz ER, Vidrine DJ, Lazev AB, Amick BC, Arduino RC. Smoking behavior in a low-income multiethnic HIV/AIDS population. *Nicotine Tob Res*. 2004; 6(1):71–77. [PubMed: 14982690]
- Hiscock R, Judge K, Bauld L. Social inequalities in quitting smoking: what factors mediate the relationship between socioeconomic position and smoking cessation? *Journal of Public Health*. 2011; 33:39–47. [PubMed: 21178184]
- Hiscock R, Bauld L, Amos A, Fidler JA, Munafò M. Socioeconomic status and smoking: A review. *Annals of the New York Academy Science*. 2012; 1248:107–123.
- Holahan CJ, North RJ, Holahan CK, Hayes RB, Powers DA, Okckene JK. Social influences on smoking in middle-aged and older women. *Psychology of Addictive Behaviors*. 2012; 26(3):519–526.
- Homish GG, Leonard KE. Spousal influence on smoking behaviors in a US community sample of newly married couples. *Social Sciences & Medicine*. 2005; 61(12):2557–2567.
- Huber M, Ledergerber B, Sauter R, Young J, Fehr J, Cusini A, et al. Outcome of smoking cessation counselling of HIV-positive persons by HIV care physicians. *HIV Medicine*. 2012; 13(7):387–397. [PubMed: 22257025]
- Humfleet GL, Hall SM, Delucchi KL, Dille JW. A randomized clinical trial of smoking cessation treatments provided in HIV clinical care settings. *Nicotine and Tobacco Research*. 2013; 15(8):1436–1445. [PubMed: 23430708]

- Johnson MO, Carrico AW, Chesney MA, Morin SF. Internalized heterosexism among HIV-positive, gay-identified men: Implications for HIV prevention and care. *Journal of Consulting and Clinical Psychology*. 2008; 76(5):829–839. [PubMed: 18837600]
- Johnson MO, Dilworth SE, Taylor JM, Darbes LA, Comfort ML, Neilands TB. Primary relationships, HIV treatment adherence, and virologic control. *AIDS and Behavior*. 2012; 16(6):1511–1521. [PubMed: 21811842]
- Kenny, DA., Kashy, DA., Cook, WL. *Dyadic Data Analysis*. New York, NY: Guilford Press; 2006.
- Kurdek LA. Assessing multiple determinants of relationship commitment in cohabitating gay, cohabitating lesbian, dating heterosexual, and married heterosexual couples. *Family Relations*. 1995; 44(3):261–266.
- Latkin C, Sherman S, Knowlton A. HIV prevention among drug users: Outcome of a network-oriented peer outreach intervention. *Health Psychology*. 2003; 22(4):322–339.
- Lebrun-Harris LA, Fiore MC, Tomoyasu N, Ngo-Metzger Q. Cigarette smoking, desire to quit, and tobacco-related counseling among patients at adult health centers. *American Journal of Public Health*. 2015; 105:180–188. [PubMed: 24625147]
- Lee JGL, Pan WK, Henriksen L, Goldstein AO, Ribisl KM. Is there a relationship between the concentration of same-sex couples and tobacco retailer density? *Nicotine and Tobacco Research*. 2016; 18(2):147–155. [PubMed: 25744959]
- Levy DT, Romano E, Mumford E. The relationship of smoking cessation to sociodemographic characteristics, smoking intensity, and tobacco control policies. *Nicotine and Tobacco Research*. 2005; 7:387–396. [PubMed: 16085506]
- Lewden C, Salmon D, Morlat P, Bévilacqua S, Jouglu E, Bonnet F. Mortality 2000 study group. Causes of death among human immunodeficiency virus (HIV)-infected adults in the era of potent antiretroviral therapy: emerging role of hepatitis and cancers, persistent role of AIDS. *International Journal of Epidemiology*. 2005; 34:121–130. [PubMed: 15561752]
- Lewis MA, McBride CM, Pollak KI, Puleo E, Butterfield RM, Emmons KM. Understanding health behavior change among couples: An interdependence and communal coping approach. *Social Science & Medicine*. 2006; 62:1369–1380. [PubMed: 16146666]
- Lichtenstein E, Glasgow RE, Lando HA, Ossip-Klein DJ, Boles SM. Telephone counseling for smoking cessation: rationales and meta-analytic review of evidence. *Health Education Research*. 1996; 11(2):243–257. [PubMed: 10163409]
- Lifson AR, Neuhaus J, Arribas JR, van den Berg-Wolf M, Labriola AM, Read TR. Smoking-related health risks among persons with HIV in the Strategies for Management of Antiretroviral Therapy clinical trial. *American Journal of Public Health*. 2010; 100(10):1896–1903. [PubMed: 20724677]
- Lindson-Hawley N, Hartmann-Boyce J, Fanshawe TR, Begh R, Farley A, Lancaster T. Interventions to reduce harm from continued tobacco use. *Cochrane Database of Systematic Reviews*. 2016; 10 CD005231.
- Mamary EM, Bahrs D, Martinez S. Cigarette smoking and the desire to quit among individuals living with HIV. *AIDS Patient Care and STDs*. 2002; 16(1):39–42. [PubMed: 11839217]
- McBride CM, Curry SJ, Grothaus LC, Nelson JC, Lando H, Pirie PL. Partner smoking status and pregnant smoker's perceptions of support for and likelihood of smoking cessation. *Health Psychology*. 1998; 17(1):63–69. [PubMed: 9459072]
- McBride CM, Baucom DH, Peterson BL, Pollak KI, Palmer C, Westman E, Lyna P. Prenatal and postpartum smoking abstinence a partner-assisted approach. *American Journal of Preventative Medicine*. 2004; 27:232–238.
- Mdodo R, Frazier EL, Dube SR, Mattson CL, Sutton MY, Brooks JT, et al. Cigarette Smoking Prevalence Among Adults With HIV Compared With the General Adult Population in the United States: Cross-sectional Surveys. *Annals of Internal Medicine*. 2015; 162(5):335–344. [PubMed: 25732274]
- Mermelstein R, Cohen S, Lichtenstein E, Baer JS, Kamarck T. Social support and smoking cessation and maintenance. *Journal of Consulting and Clinical Psychology*. 1986; 54:447–453. [PubMed: 3745596]
- Meyer IH. Prejudice, social stress, and mental health in lesbian, gay, and bisexual populations: Conceptual issues and research evidence. *Psychological Bulletin*. 2003; 129(4):674–697.

- Mitchell JW, Sullivan PS. HIV-negative partnered men's attitudes toward using in-home rapid HIV test and associated factors among a sample of US HIV-negative and HIV-discordant male couples. *Sexually Transmitted Diseases*. 2015; 42(3):123–128. [PubMed: 25668643]
- Nungesser, L. *Homosexual Acts, Actors, and Identities*. New York, NY: Praeger; 1983.
- Nylund KL, Asparouhov T, Muthén BO. Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Structural Equation Modeling*. 2007; 14(4):535–569.
- O'Cleirigh C, Dale S, Elsesser S, Pantalone DW, Mayer KH, Bradford JB, Safren SA. Sexual minority specific and related traumatic experiences are associated with increased risk for smoking among gay and bisexual men. *Journal of Psychosomatic Research*. 2015; 78(5):472–477. [PubMed: 25754971]
- O'Cleirigh C, Valentine SE, Pinkston M, Herman D, Bedoya CA, Gordan JR, Safren SA. The unique challenges facing HIV-positive patients who smoke cigarettes: HIV viremia, ART adherence, engagement in HIV care, and concurrent substance use. *AIDS and Behavior*. 2015; 19:178–185. [PubMed: 24770984]
- O'Cleirigh C, Valentine SE, Pinkston M, Herman D, Bedoya CA, Gordon JR, Safren SA. The Unique Challenges Facing HIV-Positive Patients Who Smoke Cigarettes: HIV Viremia, Art Adherence, Engagement in HIV care, and Concurrent Substance Use. *AIDS and behavior*. 2014:1–8. [PubMed: 23321946]
- Pacek LR, Latkin C, Crum RM, Stuart EA, Knowlton AR. Current cigarette smoking among HIV-positive current and former drug users: Associations with individual and social characteristics. *AIDS and Behavior*. 2014; 18(7):1368–1377. [PubMed: 24287787]
- Pachankis PE, Hatzenbuehler ML, Starks TJ. The influence of structural stigma and rejection sensitivity on young sexual minority men's daily tobacco and alcohol use. *Social Science & Medicine*. 2014; 103:67–75. [PubMed: 24507912]
- Palmer, CA., Baucom, DH., McBride, CM. Couple approaches to smoking cessation. In: Schmalting, KB., editor. *The Psychology of Couples and Illness: Theory, Research, & Practice*. Washington, D.C.: American Psychological Association; 2000. p. 407xvi
- Park EW, Tudiver FG, Campbell T. Enhancing partner support to improve smoking cessation. *Cochrane Database Systematic Review*. 2012; 7
- Radloff LS. The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1(3):385–401.
- Reid JL, Hammond D, Boudreau C, Fong GT, Siahpush M. Socioeconomic disparities in quit intentions, quit attempts, and smoking abstinence among smokers in four western countries: findings from the International Tobacco Control Four Country Survey. *Nicotine and Tobacco Research*. 2010; 12(Supple 1):S20.
- Reynolds NR, Neidig JL, Wewers ME. Illness representation and smoking behavior: a focus group study of HIV-positive men. *Journal of the Association of Nurses AIDS Care*. 2014; 15(4):37–47.
- Saunders JB, Aasland OG, Baber TF, De la Fuente JR, Grant M. Development of the Alcohol Use Disorders Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-III. *Addiction*. 1993; 88(6):791–804. [PubMed: 8329970]
- Schane RE, Ling PM, Glantz SA. Health effects of light and intermittent smoking a review. *Circulation*. 2010; 121(13):1518–1522. [PubMed: 20368531]
- Shiffman S, Tindle H, Li X, Scholl S, Dunbar M, Mitchell-Miland C. Characteristics and smoking patterns of intermittent smokers. *Experimental and Clinical Psychopharmacology*. 2012; 20(4): 264–277. [PubMed: 22390208]
- Slopen N, Kontos EZ, Ryff CD, Ayanian JZ, Alpert MA, Williams DR. Psychosocial stress and cigarette smoking persistence, cessation, and relapse over 9–10 years: A prospective study of middle-aged adults in the United States. *Cancer Causes Control*. 2013; 24(10):1849–1863. [PubMed: 23860953]
- Sutton GC. Do Men Grow to Resemble Their Wives, or Vice Versa? *Journal of Biosocial Science*. 1993; 25(1):25–29. [PubMed: 8425882]

- Turner CF, Ku L, Rogers SM, Lindberg LD, Pleck JH, Sonenstein FL. Adolescent sexual behavior, drug use, and violence: Increased reporting with computer survey technology. *Science*. 1998; 280(5365):867–873. [PubMed: 9572724]
- Vartiainen E, Seppälä T, Lillsunde P, Puska P. Validation of self reported smoking by serum cotinine measurement in a community-based study. *Journal of Epidemiological Community Health*. 2002; 56:167–170.
- Vittecoq D, Escaut L, Chironi G, Teicher E, Monsuez JJ, Andrejak M, et al. Coronary heart disease in HIV-infected patients in the highly active antiretroviral treatment era. *AIDS*. 2003; 17(Suppl 1):S70–76. [PubMed: 12870533]
- Walsh JC, Mandalia S, Gazzard BG. Responses to a 1 month self-report on adherence to antiretroviral therapy are consistent with electronic data and virological treatment outcome. *AIDS*. 2002; 16(2): 269–277. [PubMed: 11807312]
- Webb MS, Vanable PA, Carey MP, Blair DC. Cigarette smoking among HIV+ men and women: examining health, substance use, and psychosocial correlates across the smoking spectrum. *Journal of Behavioral Medicine*. 2007; 30(5):371–383. [PubMed: 17570050]
- Weiselquist J, Rusbult CE, Foster CA, Agnew CR. Commitment, pro-relationship behavior, and trust in close relationships. *Journal of Personality and Social Psychology*. 1999; 77(5):942–966. [PubMed: 10573874]

Table 1

Sample Characteristics of HIV-positive men on ART (N = 377)

<i>Demographics</i>	Mean (SD), N (%)
Age (years) (mean, SD)	46.15 (9.93)
Education Level, (N, %)	
Less than High school	16 (4.2)
High school or higher	361 (95.8)
Annual Income (N, %)	
Less than \$20,000	197 (52.3)
\$20,000 or higher	180 (47.4)
Couple HIV status (N, %)	
Serodiscordant	118 (31.3)
Seroconcordant	259 (68.7)
Sexual Identity (N, %)	
Gay	345 (91.5)
Bisexual	28 (7.4)
Other	4 (1.1)
Race/ethnicity (N, %)	
Black	62 (16.4)
White	208 (55.2)
Latino	73 (19.4)
Other	34 (9.0)
Time living with HIV, years (mean, SD)	13.45 (8.04)
<i>Relationship Characteristics</i>	
Relationship duration, years (mean, SD)	6.52 (4.94)
Commitment (mean, SD)	32.49 (5.92)
Partner health support (mean, SD)	1.91 (1.16)
<i>Health Status</i>	
Depressive Symptoms (mean, SD)	15.02 (10.42)
100% Adherence to ART (N, %)	297 (78.8)
Viral Suppression (N=373, %)	169 (44.8)
<i>Substance Use</i>	
Alcohol Use (mean, SD)	3.92 (5.67)
Current Smoker: Baseline (Wave 1) (N, %)	106 (28.12)
Current Smoker: 6 months (Wave 2) (N=346, %)	100 (28.90)
Current Smoker: 12 months (Wave 3) (N=313, %)	83 (26.52)
Current Smoker: 18 months (Wave 4) (N=277, %)	68 (24.55)
Current Smoker: 24 months (Wave 5) (N =258, %)	57 (22.09)

Table 2

Latent Transition Analysis: Transition Probabilities (N = 377)

	Latent Class 1	Latent Class 2
Model 1		
Wave 1 vs. Wave 2		
Latent Class 1	.988	.012
Latent Class 2	.020	.980
Wave 2 vs. Wave 3		
Latent Class 1	.919	.081
Latent Class 2	.004	.996
Wave 3 vs. Wave 4		
Latent Class 1	.961	.039
Latent Class 2	.005	.995
Wave 4 vs. Wave 5		
Latent Class 1	.938	.062
Latent Class 2	.010	.990
Model 2		
Latent Class 1	.993	.007
Latent Class 2	.050	.950

Notes: Latent transition analyses were performed using *Mplus* 7.4. Model 1 and Model 2 both feature hidden Markov first-order latent transitions assuming measurement invariance over time. Model 1 allows differing transition matrices over time whereas Model 2 assumes a single transition matrix across all time points.

Table 3

Latent Class Analysis: Logistic Regression Odds Ratios (OR) and 95% Confidence Intervals (CIs) for the Regressions of Persistent Smoking onto Baseline Covariates (N = 373).

Covariate	OR	95% CI	p
Actor Age (years)	0.934	0.875, 0.998	.044
Partner Age (years)	1.023	0.985, 1.063	.238
Actor High School Education	0.256	0.044, 1.491	.130
Partner High School Education	0.769	0.161, 3.670	.742
Actor Black Race	1.129	0.452, 2.825	.795
Actor Latino Ethnicity	0.200	0.069, 0.575	.003
Actor Other Non-White Race	0.831	0.295, 2.340	.725
Partner Black Race	1.541	0.569, 4.168	.395
Partner Latino Ethnicity	0.915	0.341, 2.459	.861
Partner Other Non-White Race	1.874	0.601, 5.839	.279
Relationship Length (years)	0.994	0.917, 1.077	.879
Actor Low Income	3.895	1.866, 8.130	<.001
Partner Low Income	1.606	0.797, 3.238	.185
Actor Depression (CES-D)	0.992	0.961, 1.024	.609
Partner Depression (CES-D)	0.976	0.944, 1.009	.147
Actor Alcohol Use (AUDIT)	1.023	0.960, 1.090	.485
Partner Alcohol Use (AUDIT)	1.015	0.962, 1.071	.586
Actor Internalized Heterosexism	1.018	0.931, 1.114	.694
Partner Internalized Heterosexism	1.048	0.968, 1.135	.248
Actor Relationship Commitment	0.998	0.920, 1.082	.957
Partner Relationship Commitment	0.953	0.889, 1.021	.171
Actor Perception of Health Support	1.170	0.865, 1.582	.308
Partner Perception of Health Support	1.064	0.791, 1.432	.681
Actor Time Living with HIV	1.000	0.995, 1.005	.910
Actor Detectable Viral Load	0.418	0.172, 1.018	.055
Actor 100% HIV Medication Adherence	0.556	0.186, 1.664	.294
Couple Concordant HIV-positive	1.405	0.563, 3.509	.466
Partner is a Current Smoker (Baseline)	12.857	5.196, 31.814	<.001