

Lawrence Berkeley National Laboratory

Recent Work

Title

THE SPECTROSCOPIC UTILITY OF THE TWO-PROTON PICK-UP ($6M, |B$) REACTION

Permalink

<https://escholarship.org/uc/item/4zs7n13k>

Author

Weisenmiller, R.B.

Publication Date

1975-10-01

0 0 4 4 0 5 1 8 5

Submitted to Physical Review Letters

LBL-4304
Preprint c.1

RECEIVED
LAWRENCE
BERKELEY LABORATORY

NOV 5 1975

LIBRARY AND
DOCUMENTS SECTION


THE SPECTROSCOPIC UTILITY OF THE TWO-PROTON
PICK-UP (${}^6\text{Li}$, ${}^8\text{B}$) REACTION

R. B. Weisenmiller, N. A. Jelley, K. H. Wilcox,
G. J. Wozniak, and Joseph Cerny

October 1975

Prepared for the U. S. Energy Research and
Development Administration under Contract W-7405-ENG-48

For Reference
Not to be taken from this room



LBL-4304
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE SPECTROSCOPIC UTILITY OF THE TWO-PROTON PICK-UP (${}^6\text{Li}, {}^8\text{B}$) REACTION*

R. B. Weisenmiller, N. A. Jelley[†], K. H. Wilcox,
G. J. Wozniak, and Joseph Cerny

Department of Chemistry and
Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

October 1975

Abstract:

Data for the (${}^6\text{Li}, {}^8\text{B}$) reaction on targets of ${}^{10}\text{B}$, ${}^{12}\text{C}$, ${}^{13}\text{C}$, and ${}^{16}\text{O}$ at 80 and 93 MeV are presented. Spectroscopic selectivity is observed and evidence for the dominance of spatially symmetric transfer of the two protons is shown. On the $T_z = 0$ targets a close similarity is found to the analogous (p,t) reaction, and the ${}^{13}\text{C}$ results indicate the location of low-lying lp-shell states in ${}^{11}\text{Be}$.

- - -

Although light-ion induced two-nucleon transfer reactions have been used extensively to study two-particle and two-hole configurations, the difficulties inherent in observing the ($n, {}^3\text{He}$) reaction have hindered the study of two-proton hole states. However, the development of heavy-ion beams has made available several possible reactions capable of probing such states in neutron-excess nuclei. We wish to report on results, obtained in the first survey of a two-proton pick-up reaction on lp-shell targets, which demonstrate the feasibility of the (${}^6\text{Li}, {}^8\text{B}$) reaction as a means of studying these configurations.

Among the two-proton pick-up reactions reported^{1,2} to date - (${}^6\text{Li}, {}^8\text{B}$), (${}^{11}\text{B}, {}^{13}\text{N}$), and (${}^{18}\text{O}, {}^{20}\text{Ne}$) - the first has several experimental advantages. Both ${}^7\text{B}$ and ${}^9\text{B}$ are particle-unbound, thus allowing clean separation of the ${}^8\text{B}$ particles from other boron isotopes by particle identification techniques with solid-state detector telescopes. Since ${}^8\text{B}$ has no bound excited states, its energy spectra lack the shadow peak ambiguity of the (${}^{18}\text{O}, {}^{20}\text{Ne}$) reaction, though this advantage is also shared by the (${}^{11}\text{B}, {}^{13}\text{N}$) reaction. However, the (${}^6\text{Li}, {}^8\text{B}$) reaction is the lightest of the broadly feasible two-proton pick-up reactions, and the kinematic contribution to the energy resolution is therefore smaller, particularly for light targets.

This study focussed on lp-shell targets because this region has been investigated thoroughly with other two-nucleon transfer reactions and the coefficients of fractional parentage (c.f.p.) relevant to two-nucleon transfer have been calculated³; furthermore, there is an obvious symmetry in the (mirror) final states populated in the (p,t) and (${}^6\text{Li}, {}^8\text{B}$) reactions on $T_z = 0$ targets and we report results from three such targets: ${}^{12}\text{C}$, ${}^{16}\text{O}$, and ${}^{10}\text{B}$. The two-particle c.f.p.⁴ for ${}^8\text{B}(2^+) \rightarrow {}^6\text{Li}(1^+) + 2p$ show that the two protons can be transferred from a spatially symmetric (${}^1\text{D}$) state or an anti-symmetric (${}^3\text{P}$) state relative to the ${}^6\text{Li}$ core. The simplest cluster transfer mechanism corresponds to an internal ${}^1\text{S}$ state (as in the (p,t) reaction) for the transferred nucleons, which for the (${}^6\text{Li}, {}^8\text{B}$) reaction can only arise from the ${}^1\text{D}$ component. However, there is a much larger amplitude for the ${}^3\text{P}$ than for the ${}^1\text{D}$ component, so if anti-symmetric transfer⁵ is important the expected symmetry between the (${}^6\text{Li}, {}^8\text{B}$) and (p,t) reactions might be distorted. Although most of the established two-proton hole states in the lp-shell can be populated by both transfer symmetries, there are a few known levels which would be fed predominantly by spatially anti-symmetric transfer,^{3,5} and two examples of these are discussed below.

The general techniques for the production of lithium beams at the Lawrence Berkeley Laboratory 88-inch cyclotron and the identification of ^8B reaction products have been described previously.² Because these reactions have highly-negative Q-values, beam energies of 80 and 93 MeV were used. Two telescopes, consisting of two transmission (ΔE) detectors, typically 15 and 10 μm thick, a 200- μm E detector and a 1000- μm reject detector, were employed. Each subtended 0.29 msr; a typical energy resolution was 300 keV. Background reduction in this moderately low yield reaction ($\sim 1\text{-}15 \mu\text{b/sr c.m.}$) was accomplished by requiring a conformity in the particle identification signals generated by each of the two ΔE detectors. Two ^{16}O targets were used: a 16% oxidized (by atom) 0.34 mg/cm^2 Li target and a 0.21 mg/cm^2 silicon dioxide target. Other targets were 0.22 mg/cm^2 natural carbon, 0.14 mg/cm^2 ^{10}B (96%), and 0.14 mg/cm^2 ^{13}C (90%). Kinematic shifts were utilized in the analysis to discern the levels of interest from those arising from target contaminants.

A comparison between a $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}$ spectrum and a $^{12}\text{C}(p,t)^{10}\text{C}$ spectrum,⁶ Figs. 1(a) and 1(b), shows that a close similarity exists between these reactions. In both spectra the 0^+ ground state and the first excited 2^+ state (2_1^+) are strongly populated, in agreement with the calculated two-particle transition strengths.³ The next higher peak, observed at 5.96 MeV in ^{10}Be , could be expected largely to consist of the 2^+ member of the $2^+, 1^-$ doublet at this energy,⁷ since lowest order shell model configurations and a simple pick-up reaction mechanism prohibit forming the 1^- state; indeed the angular distribution of the analog of this state in the (p,t) data⁶ is consistent with $L = 2$. The population of this 2_2^+ level is less than the 2_1^+ level, although transitions to the former have a much greater theoretical strength.³ Possibly some of this missing strength is contained in the 2_3^+ level at 7.54 MeV in ^{10}Be which, although thought⁸ to have a dominant sd-shell character, is observably populated.

Comparison with the (p,t) data, shown in Fig. 1(b), implies that the analog of this 2_3^+ state might be the 6.6 MeV level in ^{10}C , which is again consistent with the (p,t) angular distribution.⁶ Finally, evidence is seen at several angles for the weak population of a probable⁷ 2^+ state at 9.4 MeV and a state at 11.8 MeV.

Figures 2(a)-2(b) provide another example of the similarity between the ($^6\text{Li}, ^8\text{B}$) and (p,t)⁹ reactions, now employing ^{16}O targets. In both spectra, the lp-shell two-hole states, such as the 0^+ ground state and the two lowest 2^+ states, are populated the strongest, while the non p-shell states, such as the 1^- and $^8 0^+$ levels at 6.09 and 6.59 MeV in ^{14}C , respectively, are only weakly populated. (These 1^- and 0^+ states are obscured in Fig. 2(a) because of a carbon contaminant peak, but were seen using the SiO_2 target.) The large theoretical spectroscopic amplitude for the 2_1^+ configuration may well be shared among the 7.01, 8.32, and 10.44 MeV states,¹⁰ all of which are fed by the ($^6\text{Li}, ^8\text{B}$) reaction.

Another interesting comparison can be made between these two reactions induced on a ^{10}B target. In this case, Cohen and Kurath³ predict that a low-lying excited state in the product nucleus should be populated more strongly than its ground state. This is borne out in the $^{10}\text{B}(\text{}^6\text{Li}, ^8\text{B})^8\text{Li}$ data, shown in Fig. 3(a), and also in the $^{10}\text{B}(\text{p,t})^8\text{B}$ data,¹¹ which are not shown. One sees the expected strong population of the 3^+ level at 2.26 MeV relative to that of the $^8\text{Li } 2^+$ ground state [the known⁷ ^8Li level at 6.53 MeV was also observed].

The weak population of two specific final states in the above data indicates that spatially anti-symmetric transfer of two protons in the ($^6\text{Li}, ^8\text{B}$) reaction is probably not an important transfer mode. Little yield at all angles of the 1^+ level at 0.98 MeV in ^8Li , see Fig. 3(a), was observed; this is particularly significant since this level is essentially solely connected to the ^{10}B ground state via spatially anti-symmetric transfer. Similarly, transitions from ^{16}O to the 1^+ state¹⁰ at 11.29 MeV in ^{14}C are only possible via this transfer

mode^{3,5} and again are not observed (reactions on target contaminants obscure this region in Fig. 2(a), but this fact was established with the SiO₂ target). Also noteworthy in this context is the general similarity observed in the (p,t) and (⁶Li, ⁸B) reactions on the T_z = 0 targets.

The (⁶Li, ⁸B) reaction on a ¹³C target provides a good test of the spectroscopic selectivity of this reaction, because transitions to the ¹¹Be 1/2⁺ ground state¹² are forbidden in first order (see earlier ¹⁰Be, 1⁻ level discussion). This unusual level ordering in ¹¹Be, consisting of an sd-shell 1/2⁺ ground state with the lowest lp-shell 1/2⁻ state lying at 0.32 MeV, was explained by Talmi and Unna¹³ as a consequence of the differing interaction energies of the 2s_{1/2} and lp_{1/2} neutron with the lp_{3/2} protons. An energy spectrum of the ¹³C(⁶Li, ⁸B)¹¹Be reaction is shown in Fig. 3(b). Using the energy scale readily determined from the ¹²C contaminant, we find that the observed strength is predominantly to the 1/2⁻ state at 0.32 MeV, thereby additionally confirming its assignment as the lowest lp-shell level.¹²

Also seen in the ¹³C(⁶Li, ⁸B)¹¹Be energy spectrum in Fig. 3(b) is a state at 2.69 MeV and a peak at ~ 4.0 MeV (which may contain both of the known¹² states at 3.89 and 3.96 MeV). Observation at several angles determined that the known levels at 1.79 and 3.41 MeV are populated weakly, if at all. Each of these levels was observed in the ⁹Be(t,p)¹¹Be reaction,¹² but it can populate both positive and negative parity states. The fact that the states at 2.69 and ~ 4.0 MeV were seen in the (⁶Li, ⁸B) reaction strongly indicates that these states are lp-shell states and thus would have negative parity; this is consistent with the predicted³ locations of the first three negative parity states in ¹¹Be (given a 1/2⁻ state at 0.32 MeV, a 3/2⁻ is expected at 2.6 MeV [but with a small transition strength], and a 5/2⁻, at 4.98 MeV).

The differential cross sections for these (${}^6\text{Li}, {}^8\text{B}$) reactions decrease monotonically with angle (see Fig. 4). This is consistent with what one might expect for a high-energy heavy-ion reaction on light targets when one is far from the grazing angle. Typical cross sections ($\theta_{\text{c.m.}} \sim 20^\circ$) for the dominant transitions were $\sim 15 \mu\text{b/sr c.m.}$ for reactions on ${}^{10}\text{B}$, ${}^{12}\text{C}$, and ${}^{16}\text{O}$ targets and $\sim 2 \mu\text{b/sr}$ on the ${}^{13}\text{C}$ target. The highly-negative Q -values of these reactions cause a mismatch of the dynamic properties of the ${}^6\text{Li}$ and ${}^8\text{B}$ nuclei, which contributes to the small cross sections observed.

While it may be difficult to extract quantitative spectroscopic information from this reaction owing to the large kinematic mismatch (even if the relevant optical model parameters existed), considerable qualitative information has been gained. Furthermore, no evidence for the spatially anti-symmetric transfer of the two protons in this heavy-ion reaction was observed. Although the (${}^6\text{Li}, {}^8\text{B}$) reaction has other possible complications compared to the counterpart (p, t) reaction, it clearly offers excellent possibilities for the spectroscopic study of two-proton hole states in neutron-excess nuclei.

We would like to acknowledge our gratitude to Dr. D. Kurath for providing some two-nucleon fractional parentage coefficients and for some useful discussions on anti-symmetric transfer, to Dr. D. Ashery for help with some of the experiments, and to Drs. D. K. Scott and M. S. Zisman for some valuable discussions.

FOOTNOTES AND REFERENCES

*Work performed under the auspices of the U. S. Energy Research and Development Administration.

†Present Address: Nuclear Physics Laboratory, University of Oxford, England.

1. R. H. Siemssen, C. L. Fink, L. R. Greenwood, and H. J. Körner, Phys. Rev. Lett. 28, 626 (1972); P. R. Christensen, V. I. Manko, F. D. Becchetti, and R. J. Nickles, Nucl. Phys. A207, 33 (1973); D. K. Scott, B. G. Harvey, D. L. Hendrie, L. Kraus, C. F. Maguire, J. Mahoney, Y. Terrien, and K. Yagi, Phys. Rev. Lett. 33, 1343 (1974).
2. N. A. Jelley, K. H. Wilcox, R. B. Weisenmiller, G. J. Wozniak, and J. Cerny, Phys. Rev. C 9, 2067 (1974).
3. S. Cohen and D. Kurath, Nucl. Phys. A141, 145 (1970).
4. D. Kurath, private communication.
5. D. Kurath, Comments on Nuclear and Particle Physics V2, 55 (1972); O. Lkhagva and I. Rotter, Yad. Fiz. 11, 1037 (1970) [Sov. J. Nucl. Phys. 11, 576 (1970)].
6. D. Ashery, M. S. Zisman, G. Goth, G. J. Wozniak, R. B. Weisenmiller and J. Cerny, unpublished data; see also W. Benenson, G. M. Crawley, J. D. Dreisbach, and W. P. Johnson, Nucl. Phys. A97, 510 (1967).
7. F. Ajzenberg-Selove and T. Lauritsen, Nucl. Phys. A227, 1 (1974) and references therein.
8. D. E. Alburger, E. K. Warburton, A. Gallmann, and D. H. Wilkinson, Phys. Rev. 185, 1242 (1969).
9. D. G. Fleming, J. C. Hardy, and J. Cerny, Nucl. Phys. A162, 225 (1971) and unpublished data.
10. G. Kaschl, G. Mairle, H. Mackh, D. Hartwig, and U. Schwinn, Nucl. Phys. A178, 275 (1971).

11. G. T. A. Squier, A. R. Johnston, E. W. Spiers, S. A. Harbison, and N. M. Stewart, Nucl. Phys. A141, 158 (1970).
12. F. Ajzenberg-Selove, Nucl. Phys. A248, 1 (1975).
13. I. Talmi and I. Unna, Phys. Rev. Lett. 4, 469 (1960).

FIGURE CAPTIONS


Fig. 1(a) A composite spectrum of the $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}$ reaction ($E ^6\text{Li} = 80 \text{ MeV}$) between $\theta_{\text{lab.}} = 12.8^\circ$ and 16.8° in which the data were kinematically shifted to $\theta_{\text{lab.}} = 15.8^\circ$. (b) The $^{12}\text{C}(p,t)^{10}\text{C}$ reaction induced by 54 MeV protons at 19.5° (D. Ashery et al.⁶). (This angle lies at the first minimum in the ^{10}C g.s. angular distribution).

Fig. 2(a) An energy spectrum from a partially oxidized Li target for the $^{16}\text{O}(^6\text{Li}, ^8\text{B})^{14}\text{C}$ reaction. These data were collected at 13.5° with a 93 MeV ^6Li beam. Carbon contamination gave rise to the ^{10}Be states. (b) The $^{16}\text{O}(p,t)^{14}\text{O}$ reaction induced by 54.1 MeV protons at 27° (D. G. Fleming et al.⁹).

Fig. 3(a) A composite spectrum of the $^{10}\text{B}(^6\text{Li}, ^8\text{B})^8\text{Li}$ reaction ($E ^6\text{Li} = 80 \text{ MeV}$) between $\theta_{\text{lab.}} = 9.7^\circ$ and 20.3° in which the data were kinematically shifted to $\theta_{\text{lab.}} = 9.7^\circ$. (b) A composite spectrum of the $^{13}\text{C}(^6\text{Li}, ^8\text{B})^{11}\text{Be}$ reaction ($E ^6\text{Li} = 80 \text{ MeV}$) between $\theta_{\text{lab.}} = 9.4^\circ$ and 20.3° for a total of 32,900 μC in which the data were kinematically shifted to $\theta_{\text{lab.}} = 14.3^\circ$. Oxygen contamination gave rise to the ^{14}C states.


Fig. 4 Angular distributions for reactions induced by an 80 MeV ^6Li beam:

- (a) $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}$ g.s.; (b) $^{12}\text{C}(^6\text{Li}, ^8\text{B})^{10}\text{Be}^*$ (3.37 MeV, 2^+);
 (c) $^{16}\text{O}(^6\text{Li}, ^8\text{B})^{14}\text{C}$ g.s., and (d) $^{13}\text{C}(^6\text{Li}, ^8\text{B})^{11}\text{Be}^*$ (0.32 MeV, $1/2^-$).



XBL 759-3890

Fig. 1



XBL 759-3891

Fig. 3




Fig. 4

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research and Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720