

# UC San Diego

## UC San Diego Previously Published Works

### Title

DNA Delivery by Virus-Like Nanocarriers in Plant Cells

### Permalink

<https://escholarship.org/uc/item/5004v69h>

### Journal

Nano Letters, 24(26)

### ISSN

1530-6984

### Authors

Islam, Reyazul  
Youngblood, Marina  
Kim, Hye-In  
[et al.](#)

### Publication Date

2024-07-03

### DOI

10.1021/acs.nanolett.3c04735

Peer reviewed

# 1 DNA Delivery by Virus-Like Nanocarriers in Plant Cells

2 Md Reyazul Islam,<sup>#</sup> Marina Anderson-Youngblood,<sup>#</sup> Hye-In Kim, Ivonne González-Gamboa,  
3 Andrea Gabriela Monroy-Borrego, Adam A. Caparco, Gregory V. Lowry, Nicole F. Steinmetz,\*  
4 and Juan Pablo Giraldo\*



Cite This: <https://doi.org/10.1021/acs.nanolett.3c04735>



Read Online

ACCESS |



Metrics & More




Article Recommendations



Supporting Information

5 **ABSTRACT:** Tobacco mild green mosaic virus (TMGMV)-like nano-  
6 carriers were designed for gene delivery to plant cells. High aspect ratio  
7 TMGMVs were coated with a polycationic biopolymer, poly(allylamine)  
8 hydrochloride (PAH), to generate highly charged nanomaterials (TMGMV-  
9 PAH;  $56.20 \pm 4.7$  mV) that efficiently load (1:6 TMGMV:DNA mass ratio)  
10 and deliver single-stranded and plasmid DNA to plant cells. The TMGMV-  
11 PAH were taken up through energy-independent mechanisms in *Arabidopsis*  
12 protoplasts. TMGMV-PAH delivered a plasmid DNA encoding a green  
13 fluorescent protein (GFP) to the protoplast nucleus (70% viability), as  
14 evidenced by GFP expression using confocal microscopy and Western blot  
15 analysis. TMGMV-PAH were inactivated (iTGMV-PAH) using UV cross-  
16 linking to prevent systemic infection in intact plants. Inactivated iTGMV-  
17 PAH-mediated pDNA delivery and gene expression of GFP *in vivo* was  
18 determined using confocal microscopy and RT-qPCR. Virus-like nano-  
19 carrier-mediated gene delivery can act as a facile and biocompatible tool for advancing genetic engineering in plants.

20 **KEYWORDS:** virus, nanoparticles, gene delivery, protoplasts, plant genetics, agriculture



21 **T**he rapid increase in the human global population is  
22 projected to require a 35 to 55% increase in food  
23 production by 2050.<sup>1</sup> Addressing this challenge during a  
24 changing climate and without sustainable conventional  
25 agricultural practices raises concerns about food security.<sup>2</sup>  
26 Plant genetic engineering has been widely employed to  
27 generate crops with increased yield,<sup>3</sup> improved quality,  
28 enhanced resistance to herbicides,<sup>4</sup> insects,<sup>5</sup> diseases,<sup>6,7</sup> and  
29 biotic and abiotic stresses.<sup>8,9</sup> Genetically modified plants for  
30 biomanufacturing also hold immense potential for synthesizing  
31 small-molecule drugs,<sup>10</sup> recombinant protein therapeutics,<sup>11,12</sup>  
32 and vaccines.<sup>13,14</sup> Despite numerous biotechnological advance-  
33 ments over the past few decades, the genetic transformation of  
34 many plant species still poses considerable challenges. The  
35 delivery of transgenes into plant species mainly relies on two  
36 transformation methods: *Agrobacterium tumefaciens*-mediated  
37 transformation system<sup>15</sup> and particle bombardment.<sup>16</sup> How-  
38 ever, the *Agrobacterium*-mediated system has some significant  
39 drawbacks such as uncontrollable target gene integration into  
40 the host chromosomes causing positional effects on gene  
41 expression, and many plant species are inherently resistant to  
42 *Agrobacterium* infection<sup>17</sup> or showed low transformation  
43 efficiency (~5% to 33%).<sup>18,19</sup> Biolistics has been utilized in  
44 various plant species, as a random gene delivery system into  
45 the host nucleus, mitochondria, and chloroplast.<sup>4</sup> Particle  
46 bombardment is performed by high-pressure gene gun delivery  
47 that damages host genomic DNA and results in random


insertions of multiple copies of the gene.<sup>20</sup> The particle  
48 bombardment system is also expensive, requires labor-intensive  
49 tissue culture and selection, has low transformation efficiency  
50 often requiring hundreds of transformation attempts to  
51 generate a transgenic line,<sup>20,21</sup> and has not been successfully  
52 implemented in diverse plant species.<sup>22</sup> Therefore, there is a  
53 pressing need for a versatile, plant-species-independent, and  
54 easy-to-use tool for plant genetic transformation, allowing for  
55 efficient delivery of exogenous genes.

56 Recent advancements in nanotechnology have revealed the  
57 potential of nanomaterials in facilitating the delivery of genetic  
58 materials, such as plasmid DNA<sup>23–25</sup> and siRNA,<sup>26,27</sup> as well as  
59 biomacromolecules like functional proteins,<sup>28</sup> active ingre-  
60 dients,<sup>29,30</sup> nutrients,<sup>31</sup> and therapeutics<sup>32</sup> in plants. Single-  
61 walled carbon nanotubes (SWCNTs),<sup>23,24,33</sup> mesoporous silica  
62 nanoparticles (MSNs),<sup>34,35</sup> layered double hydroxide (LDH)  
63 clay nanosheets,<sup>26</sup> and functional peptide–DNA com-  
64 plexes<sup>25,36</sup> have demonstrated delivery of functional DNA/  
65 RNA cargoes into plant cells without mechanical assistance. 66

**Received:** December 4, 2023

**Revised:** June 10, 2024

**Accepted:** June 12, 2024




**Figure 1.** Intracellular DNA delivery in *Arabidopsis* plant cells mediated by virus-like nanocarriers. Negatively charged TMGMVs or inactivated (iTGMVs) were coated with a biopolymer, poly(allylamine) hydrochloride (PAH), imparting them with positive charge (TMGMV-PAH). The TMGMV-PAH were loaded by electrostatics with a DNA oligo (GT<sub>15</sub>, 30 bp ssDNA) that was covalently linked to a Cy3 organic dye (TMGMV-PAH-GT<sub>15</sub>-Cy3), or a plasmid DNA (pDNA) encoding a reporter gene of a green fluorescent protein (GFP). The nanocarriers and DNA cargoes spontaneously enter plant cell membranes without mechanical aid through energy-independent uptake mechanisms. Inactivated iTGMV-PAH mediated the delivery and expression of pDNA in *Arabidopsis* epidermal cells.

67 Several studies have demonstrated the possibility of carbon  
68 nanotube-mediated gene delivery in plant nucleus,<sup>23,36</sup>  
69 chloroplast,<sup>24,33</sup> and mitochondrial<sup>25</sup> genomes. However,  
70 there is a need to develop high aspect ratio nanomaterials  
71 for plant transformation that are degradable, biocompatible,  
72 and manufactured with controlled aspect ratios on a large scale.  
73 We turned toward plant virus nanoparticles as a biodegradable,  
74 cost-effective, and easily scalable nanotechnology with tunable  
75 surface chemistry.<sup>29,30,37</sup>

76 Tobacco mild green mosaic virus (TMGMV)<sup>38</sup> is a plant  
77 virus within the tobamovirus genus, also known as the U2  
78 strain of tobacco mosaic virus (TMV), approved by the U.S.  
79 Environmental Protection Agency (EPA) for use in bio-  
80 herbicides.<sup>39</sup> The nucleoprotein components of TMGMV are  
81 self-assembled from 2130 identical copies of a coat protein and  
82 ssRNA to form a 300 × 18 nm soft matter rod-shaped  
83 structure with a 4 nm wide hollow interior channel.<sup>29,38,40</sup> The  
84 nanocarriers derived from TMGMV are of interest for delivery  
85 applications due to their unique physio-chemical properties,  
86 such as biodegradability (protein-based particles), the ability to  
87 self-assemble into identical and high aspect ratio structures,  
88 and large-scale economical production with high purity and  
89 reproducibly.<sup>29,41</sup> The chemical design space is well under-  
90 stood and TMGMV can be functionalized with cargo through  
91 covalent chemistry<sup>42</sup> or encapsulation.<sup>29</sup> There are also well-  
92 established methods of TMGMV RNA inactivation through  
93 UV cross-linking or chemical treatments for use in plant  
94 species susceptible to infection.<sup>43</sup> TMGMV particles have been  
95 utilized as a carrier for active ingredients such as a porphyrin-  
96 based photosensitizer drugs (500 Zn-porphyrin molecules/

TMGMV) for cancer cell abolition of melanoma and cervical 97  
cancer models,<sup>40</sup> as well as ivermectin (10% mass loading 98  
efficiency to TMGMV) to treat plants infected with parasitic 99  
nematodes.<sup>29,30,44</sup> Plant virus-derived vectors (plasmids with 100  
virus genetic elements) have been extensively used for genetic 101  
engineering in plants through the mechanical inoculation of 102  
plasmid DNA, biolistics, vascular puncture, agroinoculation, or 103  
insect-mediated vector delivery.<sup>45,46</sup> These applications 104  
focused on delivery of RNA packaged inside the capsid.<sup>47</sup> To 105  
date, plant virus coat proteins have not been engineered as 106  
carriers for facile plasmid DNA delivery in plant cells. 107

In this study, we developed native and inactivated TMGMV- 108  
based nanomaterials as a platform for the nuclear delivery of 109  
DNA in *Arabidopsis thaliana* protoplasts and intact plants, 110  
respectively (Figure 1). Although PEG-mediated protoplast 111  
transformations achieve high transient transformation efficien- 112  
cies (50–90% in viable cells),<sup>48</sup> protoplast systems are crucial 113  
for developing genetic transformation tools and understanding 114  
nanoparticle–plant cell interaction processes.<sup>23,33,49</sup> Because 115  
plant protoplasts lack a cell wall, this study also included DNA 116  
delivery analysis *in vivo* using *Arabidopsis* leaf epidermal cells. 117  
We functionalized TMGMV by covalently coating a poly- 118  
cationic biopolymer, poly(allylamine) hydrochloride (PAH), 119  
on the TMGMV surface (TMGMV-PAH). The PAH imparts a 120  
positive charge to TMGMV-PAH for binding to DNA through 121  
electrostatic interactions. PAH has been extensively used for 122  
pharmaceutical and drug delivery applications due to its high 123  
water-solubility and biodegradable properties.<sup>50,51</sup> To deter- 124  
mine whether TMGMV-PAH delivered single-stranded DNA 125  
(ssDNA) into protoplast cells without using mechanical aid 126




**Figure 2.** Characterization of TMGMV-PAH coated with single-stranded DNA. (a) Transmission electron microscopy of negative-stained TMGMV and TMGMV-PAH. Yellow arrows indicate PAH coated on the surface of TMGMV. Scale bars 100 nm. (b) Denaturing Nu-PAGE gel electrophoresis under white light followed by Coomassie staining, 1: TMGMV, 2: TMGMV-PAH, 3: TMGMV-PAH-GT<sub>15</sub>-Cy3, M: prestained molecular weight standards. The arrow indicates the position of the TMGMV coat protein (CP) at 17.5 kDa (lower arrow) and PAH conjugated TMGMV-PAH CP at 35 kDa (upper arrow) or higher molecular weight. (c) UV-vis absorbance and (d) zeta potential (10 mM MES, pH 6.0) of TMGMV before and after coating with PAH and GT<sub>15</sub>-Cy3. The data are the means  $\pm$  SD ( $n = 4$ ). Statistical analysis was performed by one-way analysis of variance (ANOVA) with Tukey's *posthoc* multiple comparison analysis (GraphPad Prism 6); \* $P < 0.05$ ; \*\*\*\* $P < 0.0001$ .

127 while maintaining biocompatibility, we employed confocal  
128 microscopy to track the ssDNA cargo covalently bonded to a  
129 fluorophore (Cy3) and protoplast bioavailability assays. We  
130 also demonstrated the high loading capacity of plasmid DNA  
131 (pDNA) onto the TMGMV-PAH, and assessed the pDNA  
132 delivery, uptake mechanism, and transgene expression in  
133 protoplasts. Finally, we used inactivated iTMGMV-PAH to  
134 demonstrate pDNA delivery and expression in *Arabidopsis* leaf  
135 epidermal cells *in vivo*. Using virus-like nanocarriers for DNA  
136 delivery in plant cells offers a promising solution for plant  
137 genetic transformations that is scalable and biocompatible with  
138 high manufacturing quality and reproducibility.

139 The selection of polymer coating for TMGMV focused on  
140 cationic biopolymers capable of binding electrostatically with  
141 negatively charged pDNA. Among various options, PAH,  
142 polylysine, and polyarginine were prioritized due to their  
143 higher  $pK_a$  values (above pH 8) and FDA approval for other  
144 applications. TMGMV coated with polylysine and polyarginine  
145 were negatively charged, making them unsuitable for pDNA  
146 coating (Figure S1). In contrast, PAH TMGMVs were  
147 positively charged, and therefore, PAH was chosen as the  
148 coating for TMGMV in this study. We characterized TMGMV,  
149 TMGMV-PAH, and GT<sub>15</sub>-Cy3-loaded TMGMV-PAH  
150 (TMGMV-PAH-GT<sub>15</sub>-Cy3) by UV-vis, dynamic light scatter-  
151 ing (DLS), zeta potential ( $\zeta$ ), transmission electron micros-  
152 copy (TEM), Nu-PAGE protein analysis, and fluorescence  
153 emission spectra. TEM imaging of TMGMV and TMGMV-  
154 PAH shows high aspect ratio, rod-shaped nanostructures  
155 (Figure 2a) consistent with previous studies using TMGMV  
156 for pesticide delivery.<sup>29,42</sup> The TMGMV-PAH had a rough

157 surface, which is different from native TMGMV (Figure 2a),  
158 indicating coating of the PAH polymer on the TMGMV  
159 surface. We utilized a carbodiimide coupling reaction to  
160 covalently bond the amine functional groups of PAH to the  
161 carboxyl groups in TMGMV (Figure S2),<sup>42</sup> and the chemical  
162 conjugation was confirmed by Fourier-transform infrared  
163 spectroscopy (FTIR; Figure S3). Based on TEM analysis, the  
164 average lengths of TMGMV and TMGMV-PAH were  
165 nonsignificantly different,  $129.9 \pm 57.7$  and  $191.3 \pm 95$  nm,  
166 respectively. Notably, broken nanomaterials were also observed  
167 in both uncoated TMGMV and TMGMV-PAH, which can  
168 occur during preparation or imaging of the TMGMV TEM  
169 samples.<sup>29,42</sup> Furthermore, the conjugation of PAH ( $\sim 17.5$   
170 kDa) to TMGMV coat protein (CP) was confirmed by  
171 denatured Nu-PAGE protein analysis, which indicated the  
172 presence of higher molecular weight bands at  $\sim 35$  kDa, in  
173 addition to the TMGMV CP band at  $\sim 17.5$  kDa (Figure 2b).  
174 The smeared protein bands were observed due to the high  
175 positive charge of TMGMV-PAH CP ( $56.20 \pm 4.7$  mV) that  
176 hinders the relative mobility toward the electrode in the Nu-  
177 PAGE system. Both TEM and Nu-PAGE analysis indicate that  
178 PAH is coated onto the TMGMV-PAH.

179 To investigate DNA delivery by TMGMV-PAH in  
180 protoplasts, we used confocal microscopy to track ssDNA  
181 oligonucleotide (GT)<sub>15</sub> covalently linked to the Cy3  
182 fluorescent dye (GT<sub>15</sub>-Cy3). Cy3 is bright, photostable, and  
183 its emission range does not overlap with chloroplast  
184 autofluorescence.<sup>24</sup> GT<sub>15</sub>-Cy3 has been previously employed  
185 for coating positively charged carbon nanotubes for determin-  
186 ing subcellular localization in plants.<sup>24,33,52</sup> The UV-vis




**Figure 3.** Delivery of single-stranded DNA by TMGMV-PAH in plant protoplasts. (a) Confocal images of isolated mesophyll protoplasts with chlorophyll autofluorescence (magenta) exposed to TMGMV-PAH-GT<sub>15</sub>-Cy3 (0.1 mg/mL). The GT<sub>15</sub>-Cy3 was detected in protoplast membranes (white arrows) and nuclei (yellow arrows). Scale bars 30  $\mu$ m. (b) After treatment with TMGMV-PAH-GT<sub>15</sub>-Cy3, protoplasts were stained either with a nuclear marker, Hoechst, or cell membrane staining dye, FM-4-64 for confocal microscopy imaging. Scale bars 5  $\mu$ m. (c) Orthogonal projections from z-stacks of different planes ( $x/y$ ,  $x/z$ , or  $y/z$ ) of confocal microscopy images indicating localization of GT<sub>15</sub>-Cy3 with Hoechst nuclear marker. Scale bars 30  $\mu$ m. (d) Quantitative analysis of subcellular localization of GT<sub>15</sub>-Cy3 with Hoechst nuclear marker and FM-4-64 cell membrane dye. The data are means  $\pm$  SD ( $n = 3$ ).

187 absorbance spectra of TMGMV, TMGMV-PAH, and  
 188 TMGMV-PAH-GT<sub>15</sub>-Cy3 indicated characteristic absorption  
 189 peaks at 260 nm (Figure 2c). TMGMV-PAH-GT<sub>15</sub>-Cy3  
 190 showed distinct absorption peaks at 550 nm that corresponded  
 191 to the Cy3 dye on TMGMV-PAH (Figure 2c). To validate the  
 192 binding of GT<sub>15</sub>-Cy3 to TMGMV-PAH and confirm the  
 193 absence of unbound dye, the sample was purified using a  
 194 centrifugal filter unit (100 K MWCO). Following the second  
 195 wash step, no absorbance corresponding to GT<sub>15</sub>-Cy3 was  
 196 detected in the eluent (Figure S4a), whereas TMGMV-PAH-  
 197 GT<sub>15</sub>-Cy3 exhibited fluorescence emission peaks at 567 nm,  
 198 attributed to the attachment of GT<sub>15</sub>-Cy3 on TMGMV-PAH  
 199 (Figure S4b). DLS analysis indicated well dispersed nanoma-  
 200 terials with increasing hydrodynamic diameter from 267  $\pm$  1.6  
 201 nm for TMGMV to 310  $\pm$  1.3 nm for TMGMV-PAH and 361

$\pm$  3.2 nm for TMGMV-PAH-GT<sub>15</sub>-Cy3 ( $P < 0.005$ ; Figure 202  
 S4c). We observed a significant change of  $\zeta$  potential after 203  
 conjugation of PAH from negative charged TMGMV ( $-22.37$  204  
 $\pm 2.3$  mV) to highly positive charged TMGMV-PAH ( $56.20 \pm$  205  
 $4.7$  mV;  $P < 0.0001$ ; 10 mM MES buffer, pH 6.0; Figure 2d), 206  
 indicating binding of polycationic PAH to the TMGMV 207  
 surface. As expected, the  $\zeta$  potential for TMGMV-PAH slightly 208  
 decreased from  $56.20 \pm 4.7$  to  $47.69 \pm 4.4$  mV when loading 209  
 GT<sub>15</sub>-Cy3 ( $P < 0.05$ ; Figure 2d) due to the electrostatic 210  
 bonding between the negatively charged GT<sub>15</sub> and the 211  
 positively charged TMGMV-PAH. 212


To examine *in vitro* DNA delivery and subcellular local- 213  
 ization in plant cells using TMGMV-PAH as a nanocarrier, 214  
*Arabidopsis* protoplasts were isolated and incubated with 215  
 TMGMV-PAH coated with GT<sub>15</sub>-Cy3. Protoplasts are model 216



**Figure 4.** Plasmid DNA delivery and expression mediated by virus-like nanocarriers in isolated plant protoplasts. (a) DNA loading analysis by agarose gel electrophoresis of pDNA (p35S-eGFP) bound to TMGMV-PAH at mass ratios 1:1 to 1:30. M: DNA ladder. Black arrows indicate supercoiled (below) and circular (upper) pDNA bands. The red arrow indicates pDNA bound to TMGMV-PAH that prevents its mobility through the gel. (b) Zeta potential measurements of virus-like nanocarriers with or without pDNA (10 mM MES, pH 6.0). Data are means  $\pm$  SD ( $n = 3-4$ ). Statistical analysis was performed by one-way ANOVA and *Dunnett's* multiple comparisons posthoc test; \* $P < 0.01$ ; \*\*\* $P < 0.0001$ . (c) Representative TEM images of TMGMV, TMGMV-PAH, and pDNA-loaded at 1:6 mass ratios to TMGMV-PAH. Scale bar 100 nm. Arrows indicate pDNA attachment to TMGMV-PAH. (d) pDNA delivery and expression mediated by TMGMV-PAH in isolated plant protoplasts determined by confocal microscopy. Scale bar 10  $\mu$ m. (e) GFP expression analysis by Western blotting. The arrow indicates 27 kDa of GFP protein and asterisks indicate nonspecific bands. M, protein ladder.

217 systems for gene expression analysis that have been used in  
 218 numerous plant nanoparticle studies of uptake and gene  
 219 delivery.<sup>23,33,49</sup> To assess the delivery of GT<sub>15</sub>-Cy3 bound to  
 220 TMGMV-PAH and their subcellular localization using  
 221 confocal microscopy, isolated protoplasts (Figure S5) were  
 222 incubated with 0.1 mg/mL of TMGMV-PAH-GT<sub>15</sub>-Cy3 at  
 223 room temperature for 2 h before imaging. Confocal  
 224 fluorescence microscopy images indicated a significant level

of GT<sub>15</sub>-Cy3 fluorescence signal in protoplast cell membranes,  
 225 and nuclei when treated with TMGMV-PAH-GT<sub>15</sub>-Cy3  
 226 (Figure 3a). In contrast, control confocal images of protoplasts  
 227 treated with GT<sub>15</sub>-Cy3 did not show GT<sub>15</sub>-Cy3 fluorescence  
 228 signal indicating that GT<sub>15</sub>-Cy3 alone cannot be taken up by  
 229 protoplasts under these exposure conditions (Figure 3a). To  
 230 confirm TMGMV-PAH-GT<sub>15</sub>-Cy3 interaction with protoplast  
 231 cell membranes and GT<sub>15</sub>-Cy3 nuclear delivery by TMGMV-  
 232



**Figure 5.** Plasmid DNA delivery and expression mediated by iTMGMV-PAH-pDNA in *Arabidopsis* leaves. Green fluorescence protein (GFP) (a) confocal microscopy images and (b) and fluorescence intensity ( $n = 3$ ) indicating GFP expression in leaf epidermal cells infiltrated with iTMGMV-PAH-pDNA. Three-week-old *Arabidopsis* leaves were abaxially infiltrated with (1:6) 0.1 mg/mL iTMGMV-PAH: 0.6 mg/mL pDNA and analyzed 2 days post infiltration ( $n = 3$ ). Scale bars 30  $\mu\text{m}$ . One-way ANOVA with Tukey's *posthoc* multiple comparison analysis; \*\*\*\* $P < 0.0001$ . (c) RT-qPCR analysis of GFP mRNA expression levels after 2 days of iTMGMV-PAH-pDNA infiltration in *Arabidopsis* leaves. Statistical analysis was performed by one-way ANOVA with Tukey's *posthoc* multiple comparison analysis; \*\* $P < 0.005$  ( $n = 3$ ).

233 PAH, protoplasts were stained with a cell membrane marker  
234 FM-4-64 and a nuclear staining marker Hoechst. The GT<sub>15</sub>-  
235 Cy3 fluorescence was observed localized with FM-4-64 and  
236 Hoechst fluorescence signals in protoplasts cell membrane and  
237 nucleus, respectively (Figures 3b and S6). Orthogonal  
238 projections from Z-stacks of different planes ( $x/y$ ,  $x/z$ , or  $y/z$ )  
239 of the confocal microscope images confirmed nuclear  
240 uptake of GT<sub>15</sub>-Cy3 using TMGMV-PAH as shown by the  
241 colocalization with Hoechst fluorescence dye (Figure 3c).  
242 Quantitative subcellular localization analysis indicated that  
243 approximately 38%  $\pm$  1.5 of the GT<sub>15</sub>-Cy3 fluorescence signal  
244 was observed in protoplast cell membranes, while 11%  $\pm$  3.0  
245 localized with a nuclear marker (Hoechst; Figure 3d).  
246 Together, our results indicate that high aspect ratio and highly  
247 positive charged TMGMV-PAH allow penetration through  
248 plant cell membranes and facilitate ssDNA delivery (GT<sub>15</sub>-  
249 Cy3) into the nucleus, similar to inorganic high aspect ratio  
250 nanomaterials with positive charge.<sup>23</sup>

251 To elucidate the mechanism of DNA delivery into plant cells  
252 by TMGMV-PAH, we conducted a cell uptake assay with  
253 TMGMV-PAH-GT<sub>15</sub>-Cy3 at 4 °C to inhibit energy-dependent  
254 uptake mechanisms, including endocytosis.<sup>53</sup> We observed a  
255 similar percentage of protoplasts with GT<sub>15</sub>-Cy3 delivery by  
256 TMGMV-PAH at 4 °C (10%  $\pm$  1.6) and 25 °C (11%  $\pm$  3.2)  
257 (Figure S7). Thus, DNA delivered by TMGMV-PAH passively  
258 traverses the protoplast membrane by an energy-independent  
259 mechanism. This is consistent with previous studies demon-  
260 strating that highly charged inorganic nanomaterials sponta-  
261 neously penetrate plant cells, by creating temporary pores in  
262 their lipid membranes.<sup>23,24,33,54,55</sup> To determine the specific  
263 endocytosis pathways involved in nanoparticle uptake, a variety  
264 of endocytosis inhibitors can be employed.<sup>56</sup> However,  
265 temperature dependent assays block all endocytosis pathways,  
266 thus giving unequivocal evidence that the nanocarriers are not  
267 taken up through energy dependent mechanisms.

268 We investigated the TMGMV-PAH loading of pDNA,  
269 encoding a green fluorescent protein (GFP) in a transient  
270 expression vector (p35S-eGFP) (Figure S8), and delivery in  
271 *Arabidopsis* protoplasts. The TMGMV-PAH-pDNA were  
272 loaded at various concentrations of pDNA (TMGMV-

273 PAH:pDNA mass ratios 1:1 to 1:20  $w/w$ ). The gel  
274 electrophoresis of pDNA mobility shift assay (EMSA) showed  
275 no unbound or free pDNA running into the agarose gel at a  
276 mass ratio of TMGMV-PAH/pDNA = 1:1 to 1:12 ( $w/w$ ),  
277 meaning that pDNA loading was 100% up to a 1:12 ( $w/w$ )  
278 mass ratio (Figure 4a). The 1:12 TMGMV-PAH to pDNA  
279 mass loading ratio is multiple times higher than the 1:2 and  
280 10:1 nanomaterial/pDNA loading ratio reported in previous  
281 studies using inorganic nanomaterials for DNA delivery in  
282 plant cells.<sup>23,57</sup> Oversaturated and unbound free pDNA bands  
283 were observed at TMGMV-PAH/pDNA mass ratios of 1:20  
284 ( $w/w$ ) and higher in EMSA (Figure 4a). The loading of pDNA  
285 gradually reduced  $\zeta$  potential as the loading ratio of pDNA  
286 increased from 1:1 to 1:12 (Figure 4b) due to the electrostatic  
287 bonding between the negatively charged pDNA and the  
288 positively charged TMGMV-PAH. The highest decrease in  $\zeta$   
289 potential was observed after pDNA loading to TMGMV-PAH  
290 at a mass ratio of 1:12, dropping from the initial  $+7.53 \pm 5.2$   
291 mV for TMGMV-PAH to  $+9.57 \pm 10.6$  mV ( $P < 0.0001$ ;  
292 Figure 4b). At the loading mass ratio of 1:20, the  $\zeta$  potential  
293 became negative,  $-31.17 \pm 6.4$  mV, representing the  
294 oversaturation of the nanocarriers and free pDNA in the  
295 suspension. This finding indicates maximum pDNA loading at  
296 a 1:12 mass ratio and is consistent with our EMSA analysis. We  
297 confirmed morphological integrity of TMGMV-PAH loaded  
298 with pDNA from 1:1 to 1:12 mass ratios by TEM (Figures 4c  
299 and S9). In addition, we also assessed pDNA stability by an *in*  
300 *vitro* pDNA degradation assay using DNase I (nuclease), which  
301 showed that pDNA molecules, when loaded onto TMGMV-  
302 PAH, were protected from DNase I nuclease activity (Figure  
303 S10).

304 To demonstrate pDNA delivery and expression in plant  
305 cells, we incubated isolated protoplasts with TMGMV-PAH-  
306 pDNA complexes at 1:6 mass ratio having a high positive  
307 charge ( $+42.16 \pm 5.1$  mV) and loading of pDNA (Figure 4b)  
308 to promote uptake through lipid membranes<sup>49</sup> and increase the  
309 amount of pDNA delivery, respectively. We used 25  $\mu\text{g}$  of  
310 pDNA for TMGMV-PAH-mediated protoplast transformation,  
311 a standard concentration of pDNA (5–30  $\mu\text{g}$ ) established for  
312 PEG-mediated protoplast transformation.<sup>58</sup> Therefore, we

313 adjusted the TMGMV-PAH concentration to 0.04 mg/mL to  
314 keep a 1:6 mass ratio of the pDNA loading. Protoplasts were  
315 incubated with TMGMV-PAH-pDNA, and gene expression  
316 was determined after 24 h by confocal fluorescence microscopy  
317 imaging. We observed GFP expression in protoplasts when  
318 incubated with TMGMV-PAH-pDNA (Figure 4d) at a 16%  $\pm$   
319 3.0 ( $P < 0.001$ ) transformation efficiency. This transformation  
320 efficiency is lower than what is reported for PEG-mediated  
321 transformation in *Arabidopsis* plant protoplasts (50% to 90% in  
322 viable cells).<sup>48</sup> However, this demonstrates that virus-like  
323 nanocarriers can be engineered to deliver DNA to the plant  
324 nuclear genome. Further optimization of plant virus type or the  
325 nanocarrier charge, size, and aspect ratio properties may result  
326 in higher transformation efficiencies. Nevertheless, GFP  
327 expression was observed using TMGMV-PAH-pDNA, but  
328 was not detected when protoplasts were incubated with pDNA  
329 alone and TMGMV-PAH alone (Figure 4d). To further  
330 confirm GFP expression in protoplasts treated with TMGMV-  
331 PAH-pDNA, we performed a Western blot analysis on total  
332 soluble protein using an anti-GFP antibody, which detected an  
333  $\sim 27$  kDa GFP-specific protein band (Figure 4e).

334 For GFP expression analysis *in vivo*, we inactivated TMGMV  
335 to prevent plant infection using UV light exposure as reported  
336 previously.<sup>43</sup> The TEM size of inactivated iTMGMV (110.73  
337  $\pm 30.22$  nm) is similar to those of active TMGMV (129.9  $\pm$   
338 57.7 nm) ( $P > 0.05$ ) (Figure S11). In contrast, the zeta  
339 potential of iTMGMV is more negative ( $-36.29 \pm 4.23$  mV)  
340 compared to that of active TMGMV ( $-22.4 \pm 2.3$  mV) (10  
341 mM MES Buffer, pH 6.0) ( $P < 0.0001$ ). This resulted in  
342 iTMGMV-PAH-pDNA having a higher zeta potential ( $58.53 \pm$   
343  $0.50$  mV) than TMGMV-PAH-pDNA ( $42.16 \pm 5.1$  mV;  $P <$   
344  $0.001$ ). We abaxially infiltrated the inactivated iTMGMV-PAH  
345 coated in pDNA into 3-week-old *Arabidopsis* leaves at the  
346 previously established 1:6 mass loading ratio. Confocal  
347 microscopy analysis indicated that 0.1 mg/mL of iTMGMV-  
348 PAH bound to 0.6 mg/mL of pDNA can enable GFP  
349 expression into leaf epidermal cells (Figure 5a). Buffer or  
350 iTMGMV-PAH infiltrated leaves did not exhibit GFP  
351 fluorescence (Figure S12). Leaves infiltrated with iTMGMV-  
352 PAH-pDNA had a high GFP fluorescence intensity (Figure  
353 5b). RT-qPCR analysis quantifying GFP mRNA fold change  
354 expression supported GFP expression mediated by 0.1 mg/mL  
355 iTMGMV-PAH:0.6 mg/mL pDNA (Figure 5c). Together,  
356 these analyses show that (i) TMGMV-PAHs have the highest  
357 pDNA mass loading ratio for nanocarriers reported to date,  
358 preserve and protect the pDNA integrity from degradation,  
359 and facilitate spontaneous pDNA translocation across the plant  
360 plasma membrane and cell wall, enabling transgene expression  
361 in the nucleus *in vitro* and *in vivo*.

362 Maintaining cell viability after exposure to nanocarriers with  
363 DNA is crucial for enabling biocompatible gene delivery tools  
364 for plants.<sup>59</sup> We evaluated protoplast viability of TMGMV-  
365 PAH coated with GT<sub>15</sub>-Cy3 (0.1–0.5 mg/mL) or pDNA (0.04  
366 mg/mL) using fluorescein diacetate (FDA),<sup>60</sup> a lipophilic  
367 fluorescent dye that is permeable to membranes of living cells.  
368 Following endogenous esterase-mediated enzymatic activity,  
369 nonfluorescent FDA is transformed to fluorescein, a green  
370 fluorescence compound. Broken cells lack esterases, rendering  
371 them devoid of fluorescein signal. The FDA-treated protoplast  
372 cells were analyzed by confocal microscopy imaging, and viable  
373 cell percentages were calculated based on the fluorescein  
374 presence. Both TMGMV-PAH-GT<sub>15</sub>-Cy3 or TMGMV-PAH-  
375 pDNA treated and control (untreated) protoplasts showed

bright green fluorescence characteristic of fluorescein and  
376 normal morphology (Figure S13a,b). Approximately 71%  $\pm$   
377 3.5 of cells remained viable after exposure to TMGMV-PAH-  
378 GT<sub>15</sub>-Cy3 (0.1 mg/mL), while increased concentrations  
379 resulted in a gradual reduction in fluorescein signal and  
380 increased number of broken cells (Figure S13c). A dramatic  
381 reduction in the fluorescein signal in protoplasts was observed  
382 after exposure to TMGMV-PAH-GT<sub>15</sub>-Cy3 (0.5 mg/mL), in  
383 which almost no viable cells were observed (Figure S13c). For  
384 protoplasts exposed to the TMGMV-PAH:pDNA mass ratio  
385 (1:3), approximately 74%  $\pm 3.0$  of cells remained viable, which  
386 is not significantly different from the viability of untreated  
387 protoplasts (Figure S13d). In contrast, when TMGMV-PAH  
388 was loaded with pDNA at the mass ratios of 1:6 and 1:12,  
389 significant decreases were observed in cell viability, approx-  
390 imately 65%  $\pm 5.5$  ( $P < 0.039$ ) at the 1:6 ratio and 43%  $\pm 8.5$   
391 ( $P < 0.0003$ ) at the 1:12 ratio cells were viable when compared  
392 to the protoplasts-only cells (Figure S13d). The TMGMV-  
393 PAH-pDNA concentration in this protoplast viability assay was  
394 kept similar to that used in the transformation analysis (0.04  
395 mg/mL). These findings suggest that an increased loading of  
396 pDNA onto TMGMV-PAH can affect plant cell viability.  
397 Biocompatibility of iTMGMV-PAH-pDNA in *Arabidopsis*  
398 leaves was determined using propidium iodide, a fluorescent  
399 dye that stains the nucleus of dead cells (Figure S14). Confocal  
400 microscopy images of leaf cells infiltrated with our chosen  
401 concentration for GFP expression analysis of 0.1 mg/mL  
402 iTMGMV-PAH: 0.6 mg/mL pDNA showed a similar  
403 percentage of dead cells ( $4.5 \pm 1.7\%$ ) to leaves treated with  
404 buffer control ( $7.9 \pm 3.4\%$ ;  $P > 0.5$ ; Figure S14a,b). Higher  
405 concentrations of 0.15 mg/mL iTMGMV-PAH: 0.9 mg/mL  
406 pDNA significantly increased the percentage of dead cells  
407 ( $15.8 \pm 2.2\%$ ;  $P < 0.01$ ). Overall, our results indicate that  
408 DNA coated TMGMV-PAH are highly biocompatible with  
409 plant cells both *in vitro* in plant protoplasts and *in vivo* in leaf  
410 cells. 411

We engineered plant virus coat protein nanocarriers  
412 (TMGMV-PAH) for facile plasmid DNA delivery into the  
413 plant cell nucleus without mechanical or biological aid, with  
414 high biocompatibility and the highest loading of DNA  
415 nanocarriers for plant cells reported to date. We demonstrated  
416 this approach using TMGMV-PAH that spontaneously  
417 delivered a transgene (GFP) encoded in an expression vector  
418 (pDNA) into plant protoplasts and epidermal cell nuclei. GFP  
419 gene delivery and expression in plant cells has been mediated  
420 by high aspect ratio carbon nanotubes.<sup>23–25,33</sup> In this work, we  
421 used high aspect ratio protein-based nanomaterials, native  
422 TMGMV in protoplasts, and inactivated iTMGMVs *in vivo* to  
423 prevent plant infection.<sup>43</sup> TMGMV's ability to move across  
424 plant cell barriers in numerous plant species<sup>43,61</sup> suggests that  
425 these nanocarriers could mediate DNA delivery to protoplasts  
426 or leaf cells from different plant species. 427

Future research will assess if pDNA mediated delivery by  
428 TMGMV-PAHs in plant cells results in transient expression of  
429 transgenes, similar to what has been reported in previous  
430 studies about pDNA delivery using inorganic nanomaterials,  
431 <sup>23,24,33</sup> or enable stable plant transformation and genome  
432 editing with higher efficiency compared to current DNA  
433 delivery protocols using biological or mechanical aid. TMGMV  
434 may prove to be a promising tool for the delivery of genes,  
435 small-interfering RNA (siRNA), and clustered regularly  
436 interspaced short palindromic repeats (CRISPR) in plants  
437 for gene editing applications. Targeted delivery approaches 438



could be implemented for TMGMV-mediated gene delivery into plastid genomes including coating with targeting peptides<sup>62</sup> for gene delivery to plant chloroplasts,<sup>24,62</sup> and mitochondria.<sup>25</sup> Our nanotechnology approach utilizing TMGMV-PAH for DNA delivery paves the way for developing plant virus-based nanocarriers with tunable and well-controlled properties,<sup>41,42,63</sup> cost-effectiveness, scalability,<sup>64,65</sup> degradability,<sup>63</sup> and high biocompatibility,<sup>63,66</sup> which enable more sustainable agriculture and advanced plant bioengineering.

## ■ ASSOCIATED CONTENT

### SI Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acs.nanolett.3c04735>.

Detailed experimental procedures, including nanocarrier synthesis and characterization, microscopy, protoplast isolation, abaxial infiltration, RT-qPCR, gel electrophoresis, and biocompatibility assays (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Authors

**Nicole F. Steinmetz** – Department of NanoEngineering, University of California, San Diego, La Jolla, California 92093, United States; Department of Bioengineering, Department of Radiology, Center for Nano-Immuno Engineering, Shu and K.C. Chien and Peter Farrell Collaboratory, Institute for Materials Discovery and Design, Moores Cancer Center, and Center for Engineering in Cancer, Institute for Engineering in Medicine, University of California, San Diego, La Jolla, California 92093, United States; [orcid.org/0000-0002-0130-0481](https://orcid.org/0000-0002-0130-0481); Email: [nsteinmetz@ucsd.edu](mailto:nsteinmetz@ucsd.edu)

**Juan Pablo Giraldo** – Department of Botany and Plant Sciences, University of California, Riverside, California 92507, United States; [orcid.org/0000-0002-8400-8944](https://orcid.org/0000-0002-8400-8944); Email: [juanpablo.giraldo@ucr.edu](mailto:juanpablo.giraldo@ucr.edu)

### Authors

**Md Reyazul Islam** – Department of Botany and Plant Sciences, University of California, Riverside, California 92507, United States; [orcid.org/0000-0001-9494-3197](https://orcid.org/0000-0001-9494-3197)

**Marina Anderson-Youngblood** – Department of Botany and Plant Sciences, University of California, Riverside, California 92507, United States; [orcid.org/0009-0004-9588-8168](https://orcid.org/0009-0004-9588-8168)

**Hye-In Kim** – Department of Botany and Plant Sciences, University of California, Riverside, California 92507, United States; [orcid.org/0000-0002-5203-7361](https://orcid.org/0000-0002-5203-7361)

**Ivonne González-Gamboa** – Department of NanoEngineering, University of California, San Diego, La Jolla, California 92093, United States; Department of Molecular Biology, University of California, San Diego, La Jolla, California 92093, United States; [orcid.org/0000-0003-1617-8252](https://orcid.org/0000-0003-1617-8252)

**Andrea Gabriela Monroy-Borrego** – Department of NanoEngineering, University of California, San Diego, La Jolla, California 92093, United States

**Adam A. Caparco** – Department of NanoEngineering, University of California, San Diego, La Jolla, California 92093, United States; [orcid.org/0000-0002-8545-8349](https://orcid.org/0000-0002-8545-8349)

**Gregory V. Lowry** – Department of Civil and Environmental Engineering and Center for Environmental Implications of NanoTechnology (CEINT), Carnegie Mellon University,

Pittsburgh, Pennsylvania 15213, United States; [orcid.org/0000-0001-8599-008X](https://orcid.org/0000-0001-8599-008X)

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/acs.nanolett.3c04735>

### Author Contributions

<sup>#</sup>These authors contributed equally to this work (M.R.I. and M.A.-Y.). J.P.G. and N.F.S. conceived the idea and designed experiments with M.R.I. M.R.I. performed nanomaterial synthesis and characterization, *in vitro* DNA loading and delivery, gene expression analysis, cell viability, endocytosis, and confocal microscopy assays. J.P.G. designed *in vivo* experiments with M.A.-Y. who performed inactivated nanomaterial synthesis, *in vivo* pDNA delivery and gene expression analysis using RT-qPCR and confocal microscopy, and biocompatibility assays. N.F.S. and A.A.C. designed the iTMGMV formulation for the *in vivo* studies, which was prepared and characterized for quality control by A.A.C. G.V.L. contributed with data analysis. H.K. performed polymer coating design and synthesis of nanocarriers, TEM, zeta potential, and FTIR analysis of nanomaterials. I.G.-G. purified and lyophilized native TMGMV. A.G.M.-B. performed TEM of nanomaterials loaded with plasmid DNA and analysis with I.G.-G. All authors contributed to writing the manuscript.

### Notes

The authors declare the following competing financial interest(s): A pending patent entitled Compositions and Methods for Delivery of Nucleic Acids is based on this work. J.P.G., M.R.I., H.K. (University of California, Riverside), and N.F.S. (University of California, San Diego) are inventors in this patent. Specific aspects of the manuscript covered in the patent disclosure include compositions and methods for delivery of DNA in plant cells. N.F.S. is a cofounder of, has equity in, and has a financial interest in Mosaic ImmunoEngineering Inc. N.F.S. is a cofounder and serves as manager of Pokometz Scientific LLC, under which she is paid as a consultant to Mosaic ImmunoEngineering Inc., Flagship Laboratories 95 Inc., and Arana Biosciences Inc. The other authors declare no potential conflict of interest.

## ■ ACKNOWLEDGMENTS

This material is based on work mainly supported by the National Science Foundation under Grant FMSG: Bio: 2134535. M.A.Y. received funding from NSF National Research Traineeship Program Grant DBI-1922642. A.G.M.-B. was supported by a CONAHCYT doctoral studies scholarship (CVU 1062156). A.A.C. was supported through NIFA-2022-67012-36698. We also thank partial support from UC San Diego Materials Research Science and Engineering Center (MRSEC) through NSF Grant DMR-2011924 to N.F.S. The authors thank the University of California San Diego - Cellular and Molecular Medicine Electron Microscopy Core (UCSD-CMM-EM Core, RRID:SCR\_022039) for equipment access and technical assistance. The UCSD-CMM-EM Core is supported in part by the National Institutes of Health Award Number S10OD023527.

## ■ REFERENCES

(1) van Dijk, M.; Morley, T.; Rau, M. L.; Sanghai, Y. A Meta-Analysis of Projected Global Food Demand and Population at Risk of Hunger for the Period 2010–2050. *Nat. Food* **2021**, *2* (7), 494–501.

- (2) Intergovernmental Panel on Climate Change (IPCC). Food Security. *Climate Change and Land: IPCC Special Report on Climate Change, Desertification, Land Degradation, Sustainable Land Management, Food Security, and Greenhouse Gas Fluxes in Terrestrial Ecosystems*; Cambridge University Press, 2022; pp 437–550.
- (3) Sedeek, K. E. M.; Mahas, A.; Mahfouz, M. Plant Genome Engineering for Targeted Improvement of Crop Traits. *Front. Plant Sci.* **2019**, *10*, 114.
- (4) Daniell, H.; Datta, R.; Varma, S.; Gray, S.; Lee, S. B. Containment of Herbicide Resistance through Genetic Engineering of the Chloroplast Genome. *Nat. Biotechnol.* **1998**, *16* (4), 345–348.
- (5) Liu, Y.; Wu, H.; Chen, H.; Liu, Y.; He, J.; Kang, H.; Sun, Z.; Pan, G.; Wang, Q.; Hu, J.; Zhou, F.; Zhou, K.; Zheng, X.; Ren, Y.; Chen, L.; Wang, Y.; Zhao, Z.; Lin, Q.; Wu, F.; Zhang, X.; Guo, X.; Cheng, X.; Jiang, L.; Wu, C.; Wang, H.; Wan, J. A Gene Cluster Encoding Lectin Receptor Kinases Confers Broad-Spectrum and Durable Insect Resistance in Rice. *Nat. Biotechnol.* **2015**, *33* (3), 301–305.
- (6) van Esse, H. P.; Reuber, T. L.; van der Does, D. Genetic Modification to Improve Disease Resistance in Crops. *New Phytol.* **2020**, *225* (1), 70–86.
- (7) Liu, X.; Ao, K.; Yao, J.; Zhang, Y.; Li, X. Engineering Plant Disease Resistance against Biotrophic Pathogens. *Curr. Opin. Plant Biol.* **2021**, *60*, No. 101987.
- (8) Zhang, H.; Zhu, J.; Gong, Z.; Zhu, J.-K. Abiotic Stress Responses in Plants. *Nat. Rev. Genet.* **2022**, *23* (2), 104–119.
- (9) Saijo, Y.; Loo, E. P.-I. Plant Immunity in Signal Integration between Biotic and Abiotic Stress Responses. *New Phytol.* **2020**, *225* (1), 87–104.
- (10) Srivastav, V. K.; Egbuna, C.; Tiwari, M. Chapter 1 - Plant Secondary Metabolites as Lead Compounds for the Production of Potent Drugs. In *Phytochemicals as Lead Compounds for New Drug Discovery*; Egbuna, C., Kumar, S., Ifemeje, J. C., Ezzat, S. M., Kaliyaperumal, S., Eds.; Elsevier, 2020; pp 3–14.
- (11) Wolfson, W. Grow Your Own: Protalix BioTherapeutics Produces Drugs in Carrot Cells. *Chem. Biol.* **2013**, *20* (8), 969–970.
- (12) Daniell, H.; Nair, S. K.; Esmaili, N.; Wakade, G.; Shahid, N.; Ganesan, P. K.; Islam, M. R.; Shepley-McTaggart, A.; Feng, S.; Gary, E. N.; Ali, A. R.; Nuth, M.; Cruz, S. N.; Graham-Wooten, J.; Streatfield, S. J.; Montoya-Lopez, R.; Kaznica, P.; Mawson, M.; Green, B. J.; Ricciardi, R.; Milone, M.; Harty, R. N.; Wang, P.; Weiner, D. B.; Margulies, K. B.; Collman, R. G. Debulking SARS-CoV-2 in Saliva Using Angiotensin Converting Enzyme 2 in Chewing Gum to Decrease Oral Virus Transmission and Infection. *Mol. Ther.* **2022**, *30* (5), 1966–1978.
- (13) Paolino, K. M.; Regules, J. A.; Moon, J. E.; Ruck, R. C.; Bennett, J. W.; Remich, S. A.; Mills, K. T.; Lin, L.; Washington, C. N.; Fomillos, G. A.; Lindsey, C. Y.; O'Brien, K. A.; Shi, M.; Mark Jones, R.; Green, B. J.; Tottey, S.; Chichester, J. A.; Streatfield, S. J.; Yusibov, V. Safety and Immunogenicity of a Plant-Derived Recombinant Protective Antigen (rPA)-Based Vaccine against Bacillus Anthracis: A Phase 1 Dose-Escalation Study in Healthy Adults. *Vaccine* **2022**, *40* (12), 1864–1871.
- (14) Charland, N.; Gobeil, P.; Pillet, S.; Boulay, I.; Séguin, A.; Makarkov, A.; Heizer, G.; Bhutada, K.; Mahmood, A.; Trépanier, S.; Hager, K.; Jiang-Wright, J.; Atkins, J.; Saxena, P.; Cheng, M. P.; Vinh, D. C.; Boutet, P.; Roman, F.; Van Der Most, R.; Ceregido, M. A.; Dionne, M.; Tellier, G.; Gauthier, J.-S.; Essink, B.; Libman, M.; Haffizulla, J.; Fréchette, A.; D'Aoust, M.-A.; Landry, N.; Ward, B. J. Safety and Immunogenicity of an AS03-Adjuvanted Plant-Based SARS-CoV-2 Vaccine in Adults with and without Comorbidities. *NPJ Vaccines* **2022**, *7* (1), 142.
- (15) Herrera-Estrella, L.; Depicker, A.; Van Montagu, M.; Schell, J. Expression of Chimaeric Genes Transferred into Plant Cells Using a Ti-Plasmid-Derived Vector. *Nature* **1983**, *303* (5914), 209.
- (16) Klein, T. M.; Harper, E. C.; Svab, Z.; Sanford, J. C.; Fromm, M. E.; Maliga, P. Stable Genetic Transformation of Intact Nicotiana Cells by the Particle Bombardment Process. *Proc. Natl. Acad. Sci. U. S. A.* **1988**, *85* (22), 8502–8505.
- (17) Shrawat, A. K.; Lörz, H. Agrobacterium-Mediated Transformation of Cereals: A Promising Approach Crossing Barriers. *Plant Biotechnol. J.* **2006**, *4* (6), 575–603.
- (18) Ye, X.; Shrawat, A.; Moeller, L.; Rode, R.; Rivlin, A.; Kelm, D.; Martinell, B. J.; Williams, E. J.; Paisley, A.; Duncan, D. R.; Armstrong, C. L. Agrobacterium-Mediated Direct Transformation of Wheat Mature Embryos through Organogenesis. *Front. Plant Sci.* **2023**, *14*, No. 1202235.
- (19) Hayta, S.; Smedley, M. A.; Clarke, M.; Forner, M.; Harwood, W. A. An Efficient Agrobacterium-Mediated Transformation Protocol for Hexaploid and Tetraploid Wheat. *Curr. Protoc* **2021**, *1* (3), No. e58.
- (20) Liu, J.; Nannas, N. J.; Fu, F.-F.; Shi, J.; Aspinwall, B.; Parrott, W. A.; Dawe, R. K. Genome-Scale Sequence Disruption Following Biolistic Transformation in Rice and Maize. *Plant Cell* **2019**, *31* (2), 368–383.
- (21) Frame, B. R.; Zhang, H.; Cacciolone, S. M.; Sidorenko, L. V.; Dietrich, C. R.; Pegg, S. E.; Zhen, S.; Schnable, P. S.; Wang, K. Production of Transgenic Maize from Bombarded Type II Callus: Effect of Gold Particle Size and Callus Morphology on Transformation Efficiency. *In Vitro Cellular & Developmental Biology - Plant* **2000**, *36* (1), 21–29.
- (22) Baltes, N. J.; Gil-Humanes, J.; Voytas, D. F. Genome Engineering and Agriculture: Opportunities and Challenges. *Prog. Mol. Biol. Transl. Sci.* **2017**, *149*, 1–26.
- (23) Demirer, G. S.; Zhang, H.; Matos, J. L.; Goh, N. S.; Cunningham, F. J.; Sung, Y.; Chang, R.; Aditham, A. J.; Chio, L.; Cho, M.-J.; Staskawicz, B.; Landry, M. P. High Aspect Ratio Nanomaterials Enable Delivery of Functional Genetic Material without DNA Integration in Mature Plants. *Nat. Nanotechnol.* **2019**, *14* (5), 456–464.
- (24) Santana, I.; Jeon, S.-J.; Kim, H.-I.; Islam, M. R.; Castillo, C.; Garcia, G. F. H.; Newkirk, G. M.; Giraldo, J. P. Targeted Carbon Nanostructures for Chemical and Gene Delivery to Plant Chloroplasts. *ACS Nano* **2022**, *16* (8), 12156–12173.
- (25) Law, S. S. Y.; Liou, G.; Nagai, Y.; Giménez-Dejoo, J.; Tateishi, A.; Tsuchiya, K.; Kodama, Y.; Fujigaya, T.; Numata, K. Polymer-Coated Carbon Nanotube Hybrids with Functional Peptides for Gene Delivery into Plant Mitochondria. *Nat. Commun.* **2022**, *13* (1), 2417.
- (26) Mitter, N.; Worrall, E. A.; Robinson, K. E.; Li, P.; Jain, R. G.; Taochy, C.; Fletcher, S. J.; Carroll, B. J.; Lu, G. Q. M.; Xu, Z. P. Clay Nanosheets for Topical Delivery of RNAi for Sustained Protection against Plant Viruses. *Nat. Plants* **2017**, *3*, No. 16207.
- (27) Li, S.; Li, J.; Du, M.; Deng, G.; Song, Z.; Han, H. Efficient Gene Silencing in Intact Plant Cells Using siRNA Delivered By Functional Graphene Oxide Nanoparticles. *Angew. Chem., Int. Ed.* **2022**, *61* (40), No. e202210014.
- (28) Wang, J. W.; Cunningham, F. J.; Goh, N. S.; Boozarpour, N. N.; Pham, M.; Landry, M. P. Nanoparticles for Protein Delivery in Planta. *Curr. Opin. Plant Biol.* **2021**, *60*, No. 102052.
- (29) Chariou, P. L.; Steinmetz, N. F. Delivery of Pesticides to Plant Parasitic Nematodes Using Tobacco Mild Green Mosaic Virus as a Nanocarrier. *ACS Nano* **2017**, *11* (5), 4719–4730.
- (30) Caparco, A. A.; González-Gamboa, I.; Hays, S. S.; Pokorski, J. K.; Steinmetz, N. F. Delivery of Nematocides Using TMGMV-Derived Spherical Nanoparticles. *Nano Lett.* **2023**, *23* (12), 5785–5793.
- (31) Solanki, P.; Bhargava, A.; Chhipa, H.; Jain, N.; Panwar, J. Nano-Fertilizers and Their Smart Delivery System. In *Nanotechnologies in Food and Agriculture*; Rai, M., Ribeiro, C., Mattoso, L., Duran, N., Eds.; Springer International Publishing: Cham, 2015; pp 81–101.
- (32) Borgatta, J.; Shen, Y.; Tamez, C.; Green, C.; Hedlund Orbeck, J. K.; Cahill, M. S.; Protter, C.; Deng, C.; Wang, Y.; Elmer, W.; White, J. C.; Hamers, R. J. Influence of CuO Nanoparticle Aspect Ratio and Surface Charge on Disease Suppression in Tomato (*Solanum Lycopersicum*). *J. Agric. Food Chem.* **2023**, *71*, 9644.
- (33) Kwak, S.-Y.; Lew, T. T. S.; Sweeney, C. J.; Koman, V. B.; Wong, M. H.; Bohmert-Tatarev, K.; Snell, K. D.; Seo, J. S.; Chua, N.-H.; Strano, M. S. Chloroplast-Selective Gene Delivery and Expression in

- 692 Planta Using Chitosan-Complexed Single-Walled Carbon Nanotube  
693 Carriers. *Nat. Nanotechnol.* **2019**, *14* (5), 447–455.
- 694 (34) Chang, F.-P.; Kuang, L.-Y.; Huang, C.-A.; Jane, W.-N.; Hung,  
695 Y.; Hsing, Y.-I. C.; Mou, C.-Y. A Simple Plant Gene Delivery System  
696 Using Mesoporous Silica Nanoparticles as Carriers. *J. Mater. Chem. B*  
697 *Mater. Biol. Med.* **2013**, *1* (39), 5279–5287.
- 698 (35) Torney, F.; Trewyn, B. G.; Lin, V. S.-Y.; Wang, K. Mesoporous  
699 Silica Nanoparticles Deliver DNA and Chemicals into Plants. *Nat.*  
700 *Nanotechnol.* **2007**, *2* (5), 295–300.
- 701 (36) Li, J.; Li, S.; Du, M.; Song, Z.; Han, H. Nuclear Delivery of  
702 Exogenous Gene in Mature Plants Using Nuclear Location Signal and  
703 Cell-Penetrating Peptide Nanocomplex. *ACS Appl. Nano Mater.* **2023**,  
704 *6* (1), 160–170.
- 705 (37) Jones, I.; Roy, P. Small Is Beautiful: Virus-like Particles as  
706 Vaccines. *Biochem.* **2021**, *43* (4), 18–21.
- 707 (38) Pattanayek, R.; Stubbs, G. Structure of the U2 Strain of  
708 Tobacco Mosaic Virus Refined at 3.5 Å Resolution Using X-Ray Fiber  
709 Diffraction. *J. Mol. Biol.* **1992**, *228* (2), 516–528.
- 710 (39) Charudattan, R. Use of Plant Viruses as Bioherbicides: The  
711 First Virus-Based Bioherbicide and Future Opportunities. *Pest Manag.*  
712 *Sci.* **2024**, *80*, 103.
- 713 (40) Chariou, P. L.; Wang, L.; Desai, C.; Park, J.; Robbins, L. K.; von  
714 Recum, H. A.; Ghiladi, R. A.; Steinmetz, N. F. Let There Be Light:  
715 Targeted Photodynamic Therapy Using High Aspect Ratio Plant Viral  
716 Nanoparticles. *Macromol. Biosci.* **2019**, *19* (5), No. e1800407.
- 717 (41) Fan, X. Z.; Pomerantseva, E.; Gnerlich, M.; Brown, A.;  
718 Gerasopoulos, K.; McCarthy, M.; Culver, J.; Ghodssi, R. Tobacco  
719 Mosaic Virus: A Biological Building Block for Micro/nano/bio  
720 Systems. *J. Vac. Sci. Technol. A* **2013**, *31* (5), No. 050815.
- 721 (42) González-Gamboa, I.; Caparco, A. A.; McCaskill, J. M.;  
722 Steinmetz, N. F. Bioconjugation Strategies for Tobacco Mild Green  
723 Mosaic Virus. *ChemBioChem* **2022**, *23* (18), No. e202200323.
- 724 (43) Chariou, P. L.; Ma, Y.; Hensley, M.; Rosskopf, E. N.; Hong, J.  
725 C.; Charudattan, R.; Steinmetz, N. F. Inactivated Plant Viruses as an  
726 Agrochemical Delivery Platform. *ACS Agric. Sci. Technol.* **2021**, *1* (3),  
727 124–130.
- 728 (44) González-Gamboa, I.; Caparco, A. A.; McCaskill, J.;  
729 Fuenlabrada-Velázquez, P.; Hays, S. S.; Jin, Z.; Jokerst, J. V.;  
730 Pokorski, J. K.; Steinmetz, N. F. Inter-Coat Protein Loading of  
731 Active Ingredients into Tobacco Mild Green Mosaic Virus through  
732 Partial Dissociation and Reassembly of the Virion. *Sci. Rep.* **2024**, *14*  
733 (1), 7168.
- 734 (45) Abrahamian, P.; Hammond, R. W.; Hammond, J. Plant Virus-  
735 Derived Vectors: Applications in Agricultural and Medical Bio-  
736 technology. *Annu. Rev. Virol.* **2020**, *7* (1), 513–535.
- 737 (46) Scholthof, K.-B. G.; Mirkov, T. E.; Scholthof, H. B. Plant Virus  
738 Gene Vectors: Biotechnology Applications in Agriculture and  
739 Medicine. *Genet. Eng.* **2002**, *24*, 67–85.
- 740 (47) Rössner, C.; Lotz, D.; Becker, A. VIGS Goes Viral: How VIGS  
741 Transforms Our Understanding of Plant Science. *Annu. Rev. Plant*  
742 *Biol.* **2022**, *73*, 703–728.
- 743 (48) Yoo, S.-D.; Cho, Y.-H.; Sheen, J. Arabidopsis Mesophyll  
744 Protoplasts: A Versatile Cell System for Transient Gene Expression  
745 Analysis. *Nat. Protoc.* **2007**, *2* (7), 1565–1572.
- 746 (49) Lew, T. T. S.; Wong, M. H.; Kwak, S.-Y.; Sinclair, R.; Koman,  
747 V. B.; Strano, M. S. Rational Design Principles for the Transport and  
748 Subcellular Distribution of Nanomaterials into Plant Protoplasts.  
749 *Small* **2018**, *14* (44), No. e1802086.
- 750 (50) Jian, W.; Xu, S.; Wang, J.; Feng, S. Layer-by-Layer Assembly of  
751 Poly(allylamine Hydrochloride)/polyurethane and Its Loading and  
752 Release Behavior for Methylene Orange. *J. Appl. Polym. Sci.* **2013**, *129*  
753 (4), 2070–2075.
- 754 (51) Kreke, M. R.; Badami, A. S.; Brady, J. B.; Akers, R. M.;  
755 Goldstein, A. S. Modulation of Protein Adsorption and Cell Adhesion  
756 by Poly(allylamine hydrochloride) Heparin Films. *Biomaterials* **2005**,  
757 *26* (16), 2975–2981.
- 758 (52) Giraldo, J. P.; Landry, M. P.; Kwak, S.-Y.; Jain, R. M.; Wong, M.  
759 H.; Iverson, N. M.; Ben-Naim, M.; Strano, M. S. A Ratiometric Sensor  
Using Single Chirality Near-Infrared Fluorescent Carbon Nanotubes: 760  
Application to In Vivo Monitoring. *Small* **2015**, *11* (32), 3973–3984. 761  
(53) He, Z.; Liu, K.; Manaloto, E.; Casey, A.; Cribaro, G. P.; Byrne, 762  
H. J.; Tian, F.; Barcia, C.; Conway, G. E.; Cullen, P. J.; Curtin, J. F. 763  
Cold Atmospheric Plasma Induces ATP-Dependent Endocytosis of 764  
Nanoparticles and Synergistic U373MG Cancer Cell Death. *Sci. Rep.* 765  
**2018**, *8* (1), 5298. 766  
(54) Giraldo, J. P.; Landry, M. P.; Faltermeier, S. M.; McNicholas, T. 767  
P.; Iverson, N. M.; Boghossian, A. A.; Reuel, N. F.; Hilmer, A. J.; Sen, 768  
F.; Brew, J. A.; Strano, M. S. Plant Nanobionics Approach to Augment 769  
Photosynthesis and Biochemical Sensing. *Nat. Mater.* **2014**, *13* (4), 770  
400–408. 771  
(55) Wong, M. H.; Misra, R. P.; Giraldo, J. P.; Kwak, S.-Y.; Son, Y.; 772  
Landry, M. P.; Swan, J. W.; Blankschtein, D.; Strano, M. S. Lipid 773  
Exchange Envelope Penetration (LEEP) of Nanoparticles for Plant 774  
Engineering: A Universal Localization Mechanism. *Nano Lett.* **2016**, 775  
*16* (2), 1161–1172. 776  
(56) Rennick, J. J.; Johnston, A. P. R.; Parton, R. G. Key Principles 777  
and Methods for Studying the Endocytosis of Biological and 778  
Nanoparticle Therapeutics. *Nat. Nanotechnol.* **2021**, *16* (3), 266–276. 779  
(57) Liu, Q.; Li, Y.; Xu, K.; Li, D.; Hu, H.; Zhou, F.; Song, P.; Yu, Y.; 780  
Wei, Q.; Liu, Q.; Wang, W.; Bu, R.; Sun, H.; Wang, X.; Hao, J.; Li, H.; 781  
Li, C. Clay Nanosheet-Mediated Delivery of Recombinant Plasmids 782  
Expressing Artificial miRNAs via Leaf Spray to Prevent Infection by 783  
Plant DNA Viruses. *Hortic Res.* **2020**, *7* (1), 179. 784  
(58) Lee, D. W.; Hwang, I. Transient Expression and Analysis of 785  
Chloroplast Proteins in Arabidopsis Protoplasts. In *Chloroplast* 786  
*Research in Arabidopsis: Methods and Protocols*; Jarvis, R. P., Ed.; 787  
Humana Press: Totowa, NJ, 2011; Vol. 1, pp 59–71. 788  
(59) Wang, X.; Xie, H.; Wang, P.; Yin, H. Nanoparticles in Plants: 789  
Uptake, Transport and Physiological Activity in Leaf and Root. 790  
*Materials* **2023**, *16* (8), 3097. 791  
(60) Wang, H.; Zhu, X.; Li, H.; Cui, J.; Liu, C.; Chen, X.; Zhang, W. 792  
Induction of Caspase-3-like Activity in Rice Following Release of 793  
Cytochrome-F from the Chloroplast and Subsequent Interaction with 794  
the Ubiquitin-Proteasome System. *Sci. Rep.* **2014**, *4*, 5989. 795  
(61) de Andrés-Torán, R.; Guidoum, L.; Zamfir, A. D.; Mora, M. Á.; 796  
Moreno-Vázquez, S.; García-Arenal, F. Tobacco Mild Green Mosaic 797  
Virus (TMGMV) Isolates from Different Plant Families Show No 798  
Evidence of Differential Adaptation to Their Host of Origin. *Viruses* 799  
**2023**, *15* (12), 2384. 800  
(62) Santana, I.; Wu, H.; Hu, P.; Giraldo, J. P. Targeted Delivery of 801  
Nanomaterials with Chemical Cargoes in Plants Enabled by a 802  
Biorecognition Motif. *Nat. Commun.* **2020**, *11* (1), 2045. 803  
(63) Koudelka, K. J.; Pitek, A. S.; Manchester, M.; Steinmetz, N. F. 804  
Virus-Based Nanoparticles as Versatile Nanomachines. *Annu. Rev.* 805  
*Virol.* **2015**, *2* (1), 379–401. 806  
(64) Chung, Y. H.; Church, D.; Koellhoffer, E. C.; Osota, E.; Shukla, 807  
S.; Rybicki, E. P.; Pokorski, J. K.; Steinmetz, N. F. Integrating Plant 808  
Molecular Farming and Materials Research for next-Generation 809  
Vaccines. *Nat. Rev. Mater.* **2022**, *7* (5), 372–388. 810  
(65) Rybicki, E. P. Plant Molecular Farming of Virus-like 811  
Nanoparticles as Vaccines and Reagents. *Wiley Interdiscip. Rev.* 812  
*Nanomed. Nanobiotechnol.* **2020**, *12* (2), No. e1587. 813  
(66) Kim, K. R.; Lee, A. S.; Kim, S. M.; Heo, H. R.; Kim, C. S. Virus- 814  
like Nanoparticles as a Theranostic Platform for Cancer. *Front Bioeng* 815  
*Biotechnol.* **2023**, *10*, No. 1106767. 816