

UC Irvine

UC Irvine Previously Published Works

Title

Sources of upper tropospheric HOx over the South Pacific Convergence Zone: A case study

Permalink

<https://escholarship.org/uc/item/5027n3tn>

Journal

Journal of Geophysical Research, 107(D2)

ISSN

0148-0227

Authors

Mari, Céline
Saüt, Carine
Jacob, Daniel J
[et al.](#)

Publication Date

2003

DOI

10.1029/2000jd000304

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Sources of upper tropospheric HO_x over the South Pacific Convergence Zone: A case study

Céline Mari,¹ Carine Saüt,¹ Daniel J. Jacob,² François Ravetta,^{2,3} Bruce Anderson,⁴ Melody A. Avery,⁵ Donald R. Blake,⁶ William H. Brune,⁷ Ian Faloon,⁸ G. L. Gregory,⁴ Brian G. Heikes,⁹ Glen W. Sachse,⁴ Scott T. Sandholm,¹⁰ Hanwant B. Singh,¹¹ Robert W. Talbot,¹² David Tan,¹⁰ and Stephanie Vay⁴

Received 27 December 2000; revised 2 July 2001; accepted 5 July 2001; published 12 November 2002.

[1] A zero-dimensional (0-D) model has been applied to study the sources of hydrogen oxide radicals (HO_x = HO₂ + OH) in the tropical upper troposphere during the Pacific Exploratory Mission in the tropics (PEM-Tropics B) aircraft mission over the South Pacific in March–April 1999. Observations made across the Southern Pacific Convergence Zone (SPCZ) and the southern branch of the Intertropical Convergence Zone (ITCZ) provided the opportunity to contrast the relative contributions of different sources of HO_x in a nitrogen oxide radical (NO_x)-limited regime, in relatively pristine tropical air. The primary sources of HO_x vary significantly along the flight track, in correlation with the supply of water vapor. The latitudinal variation of HO_x sources is found to be controlled also by the levels of NO_x and primary HO_x production rates $P(\text{HO}_x)$. Budget calculations in the 8- to 12-km altitude range show that the reaction $\text{O}(^1\text{D}) + \text{H}_2\text{O}$ is a major HO_x source in the cloud region traversed by the aircraft, including SPCZ and the southern branch of the ITCZ. Production from acetone becomes significant in drier region south of 20°S and can become dominant where water vapor mixing ratios lie under 200 ppmv. Over the SPCZ region, in the cloud outflow, CH₃ OOH transported by convection accounts for 22% to 64% of the total primary source. Oxidation of methane amplifies the primary HO_x source by 1–1.8 in the dry regions. **INDEX TERMS:** 0320 Atmospheric Composition and Structure: Cloud physics and chemistry; 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; 0368 Atmospheric Composition and Structure: Troposphere—constituent transport and chemistry; 3314 Meteorology and Atmospheric Dynamics: Convective processes; **KEYWORDS:** HO_x, convection, SPCZ, upper troposphere, PEM-TROPICS

Citation: Mari, C., et al., Sources of upper tropospheric HO_x over the South Pacific Convergence Zone: A case study, *J. Geophys. Res.*, 107, 8229, doi:10.1029/2000JD000304, 2002. [printed 108(D2), 2003]

¹Laboratoire d'Aérodynamique, UMR CNRS/Université Paul Sabatier, Toulouse, France.

²Division of Engineering and Applied Science, Harvard University, Cambridge, Massachusetts, USA.

³Service d'Aéronomie, Paris, France.

⁴NASA Langley Research Center, Hampton, Virginia, USA.

⁵Department of Physics, Hampton University, Hampton, Virginia, USA.

⁶Department of Chemistry, University of California, Irvine, California, USA.

⁷Department of Meteorology, Pennsylvania State University, University Park, Pennsylvania, USA.

⁸National Center for Atmospheric Research, Boulder, Colorado, USA.

⁹Center for Atmospheric Chemistry Studies, University of Rhode Island, Narragansett, Rhode Island, USA.


¹⁰Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA.

¹¹NASA Ames Research Center, Moffett Field, California, USA.

¹²CSRC/EOS, University of New Hampshire, Durham, New Hampshire, USA.

1. Introduction

[2] Tropical regions play an important role in determining the global oxidizing power of the atmosphere because of the high UV radiation and humidity, promoting the formation of hydrogen oxide radicals (HO_x = OH + peroxy radicals). Surprising results have emerged from the first measurements of HO_x radicals in the upper troposphere: observed HO_x levels were frequently 2–4 times higher than expected based on the commonly assumed primary source:



To explain the elevated levels of HO_x in the upper troposphere, new sources have been suggested. Acetone (CH₃C(O)CH₃) was found to be a major source at low levels of water vapor in the upper troposphere [Singh *et al.*, 1995; Arnold *et al.*, 1997; McKeen *et al.*, 1997; Wennberg *et*




Figure 1. GEOS-10 IR satellite image for 21 March, at 0300 UTC with track of flight 10, during PEM-Tropics B experiment. Flight 10 started on 20 March, 2230 UTC, from Fiji, returning on 21 March, 0615 UTC. Median values of observed pressure (in hPa), ozone (in ppbv), CH₃I (in pptv), NO_x (in pptv) and water vapor mixing ratios (ppmv) are listed for four latitudinal regions: south of SPCZ (24°–20°S), SPCZ (20°–17°S), North of SPCZ (17°–13°S) and ITCZ (13°–10°S).

al., 1998]. Recent studies have suggested that deep convection may be an effective process in lifting HO_x precursors from the boundary layer to the middle and upper troposphere [Prather and Jacob, 1997]. Several of these precursors have been identified including methylhydroperoxide (CH₃OOH) [Jaeglé *et al.*, 1997; Folkins *et al.*, 1998; Cohan *et al.*, 1999], formaldehyde (CH₂O) and other aldehydes [Müller and Brasseur, 1999]. Photolysis of convected peroxides and formaldehyde, together with acetone seemed to provide a strong enough source to account for the observed levels of HO_x. The importance of these new sources was tested successfully for northern midlatitudes [Jaeglé *et al.*, 2001] and tropical northern and southern Pacific oceans [Crawford *et al.*, 1999], in air masses impacted by natural (biomass burning) and anthropogenic pollution. In this paper we examine the question of the origin of HO_x radicals and the mechanism controlling the fast photochemistry of upper tropospheric HO_x in the clean tropical southern Pacific ocean during a season when biomass burning and advection of continental emitted species were low.

[3] In particular, we focus the study on the convection outflow and surrounding air from the South Pacific Convergence Zone (SPCZ). Our analysis is based on extensive observations of HO_x radicals, their precursors, and related species obtained during the Pacific Exploratory Mission in the tropics (PEM-Tropics B). This mission was designed to provide a detailed picture of tropospheric chemistry over the

South Pacific Basin during the wet season (March–April 1999) [Raper *et al.*, 2001]. Our analysis is based on data from the DC-8 aircraft northeast of Fiji (17°S, 177°E) as the aircraft traversed an extensive and strong band of SPCZ convection convective outflow at 10–12 km altitude (section 2).

[4] The SPCZ is a major dynamical feature within the South Pacific. It is characterized by strong upward and downward motions and convective precipitations, and represents a barrier for horizontal atmospheric transport [Gregory *et al.*, 1999]. Air north and south of the SPCZ has distinct chemical signatures, reflective of the impact of long range transport, dry air and subsidence south of the SPCZ and strong ascending motions and precipitations related to convective clouds north of the SPCZ. The SPCZ flight during PEM-Tropics B offers us the opportunity to focus on HO_x sources and chemistry in the tropical upper troposphere along a strong gradient of convective influence.

[5] Our approach is based on a zero-dimensional (0-D) model (section 3) constrained by the ensemble of PEM-Tropics B aircraft observations. This model allows us to identify the dominant HO_x sources controlling the HO_x concentration in the upper troposphere along the flight track. Budget calculations are then performed and dominant sources are discussed as a function of latitude (section 4).

2. Experimental Conditions

[6] Flight 10 was dedicated to the study of SPCZ-related convection. The flight traversed the extensive and strong band of convection, sampling air masses on the edges and the outflow of the convective system (Figure 1). The flight took place during daytime, starting from Fiji on 20 March, 2230 UTC (20 March, 0930 local solar time), ending on 21 March at 0615 UTC (20 March, 1715 local solar time). Figure 2 shows the altitude of the aircraft and the measured values of NO₂ photolysis rate as a function of time.




Figure 2. Photolysis rate of NO₂, in s⁻¹ and altitude of the aircraft, in km, as a function of local solar date and time, during flight 10. The entire flight took place during daytime, crossing the SPCZ twice during the outbound and returning part of the flight.




Figure 3. Three-day kinematic backward trajectories from the HYSPLIT4 trajectory model, analyzed meteorological fields are from NCEP/FNL. Trajectories start at 10 km altitude and are calculated for equally spaced points along flight 10 track. Starting points are marked with diamonds, ending points with triangles.

[7] Chemical variations observed across the SPCZ at 200–300 hPa (Figure 1) showed the transition between subtropical and tropical air masses as well as strong convective influence within the SPCZ convection

area, a near-zonal band of deep convection was present between 5° and 15°S at the northern type of the flight track, commensurate with the southern branch of the ITCZ [Fuelberg *et al.*, 2001]. The satellite image of Figure 1 displays two distinct regions: a region north of the SPCZ, characterized by a strong convective activity and a cloud free region south of the SPCZ characterized by dry air and subsidence. We calculated 3-day kinematic backward trajectories using the Hybrid Single-Particle Lagrangian Integrated Trajectory (HYSPLIT4) 1997 model (Web address: <http://www.arl.noaa.gov/ready/hysplit4.html>, NOAA Air Resources Laboratory, Silver Spring, MD) starting at 10 km altitude along the flight track (Figure 3). Most of the trajectories showed a weak northeasterly flow of tropical maritime air, with those South of 20°S traversing Australia within the past 2 weeks.

[8] In Figure 1 and Table 1, median concentrations at 8–12 km altitude observed along the flight track are reported for four distinct regions. In the southern part of the track (south of 20°S), relatively high concentrations of ozone (mean value 32 ppbv), NO (mean value 29 pptv) and small enhancements of CO (up to 58 ppbv) were measured (Figure 4), suggesting some continental influence. Indeed, back trajectories indicate that air arriving on the southwest side of the SPCZ in the upper troposphere originated from Australia (Figure 3). Pickering *et al.* [2001] also concluded that air arriving on the southwest side of the SPCZ at 11 km likely consisted of outflow from deep convection over northern Australia 3 days earlier. Observed PAN and HNO₃ concentrations were also relatively high in this region (median values 59 pptv and 109 pptv, respectively). High mixing ratios of acetone and methanol were measured in the southwest-portion of the flight suggesting a continental biogenic origin of the sampled air mass [Jacob *et al.*,

Table 1. Median Observations in the Upper Troposphere (8 to 12 km) for the Four Latitudinal Regions Along the Flight Track in Figure 1^a

Observations	24°–20°S South of SPCZ	20°–17°S SPCZ	17°–13°S North of SPCZ	13°–10°S ITCZ
Pressure, hPa	262 (215–350)	216 (215–348)	238 (216–341)	245 (216–350)
Temperature, K	233 (224–250)	225 (223–250)	229 (224–250)	232 (224–251)
Ozone column, DU	254 (253–261)	253 (250–254)	251 (250–253)	258 (253–259)
H ₂ O, ppmv	86 (36–393)	296 (65–1268)	165 (97–1469)	354 (136–2304)
RH (water),	22 (5–43)	77 (9–154)	61 (20–93)	86 (26–106)
RH (ice),	19 (4–44)	76 (7–163)	58 (17–91)	85 (23–107)
OH, pptv	0.24 (0.12–0.39)	0.23 (0.13–0.34)	0.18 (0.09–0.27)	0.24 (0.13–0.53)
HO ₂ , pptv	6 (3–10)	7 (3–13)	7 (5–9)	9 (7–14)
H ₂ O ₂ , ppbv	146 (97–502)	160 (80–352)	174 (101–263)	201 (114–481)
CH ₃ OOH, ppbv	67 (34–158)	116 (30–304)	100 (37–212)	116 (45–181)
CH ₂ O, ppbv	80 (56–306)	110 (56–182)	96 (52–253)	118 (80–201)
NO, pptv	29 (5–59)	10 (7–38)	10 (6–19)	12 (6–78)
HNO ₃ , pptv	109 (51–145)	37 (5–135)	26 (9–52)	39 (33–42)
O ₃ , ppbv	32 (17–55)	14 (10–41)	15 (13–21)	15 (12–19)
CH ₃ I, pptv	0.08 (0.02–0.17)	0.21 (0.08–0.40)	0.11 (0.08–0.19)	0.13 (0.07–0.16)
CO, pptv	48 (42–58)	46 (41–54)	49 (46–51)	49 (47–54)
PAN, pptv	59 (10–117)	10 (8–113)	8 (7–14)	9 (8–12)
Methanol, pptv	2706 (402–2866)	895 (402–1780)	1841 (903–2740)	786 (395–2589)
Ethane, pptv	228 (195–278)	229 (173–285)	254 (234–261)	258 (247–263)
Propane, pptv	6 (4–11)	5 (3–7)	6 (4–7)	6 (4–8)
CH ₄ , pptv	1715 (1706–1742)	1713 (1700–1733)	1714 (1708–1718)	1715 (1706–1721)
Acetone, pptv	509 (303–533)	372 (304–454)	423 (388–544)	351 (337–373)
Aerosol surface area (liquid and ice), $\mu\text{m}^2 \text{cm}^{-3}$	3 (1–4)	1135 (1–149352)	3 (0–39201)	741 (0–6479)

^a The values are medians. The minimum and maximum are listed in parentheses.




Figure 4. CO (crosses) and O₃ (diamonds) mixing ratios, respectively in pptv and ppbv, as a function of latitude, for both outbound and return parts of flight 10.

2002]. In contrast, ethane and propane, emitted preferentially by anthropogenic sources, have constant mixing ratios as a function of latitude. For the regions further north, PAN and HNO₃ median mixing ratios were lower than 10 and 40 pptv, respectively. The region south of the SPCZ was characterized by strong subsidence associated with the upper tropospheric subtropical jet. Consequently in this region, water vapor concentrations were low (median value 86 ppmv).

[9] The flight segment between 17° and 20°S was characterized by deep convection associated with the SPCZ as evidenced by high concentrations of water vapor, CH₃I and CH₃OOH and low concentrations of ozone and NO. Between 13°S and 17°S, the air mass was drier, but NO and ozone concentrations remained low. Finally, the northernmost part of the flight took place in ITCZ convective outflow with chemical composition very similar to that of the SPCZ outflow.

3. Model Description

[10] We use a steady state photochemical box model [Crassier et al., 1999] including the Regional Atmospheric Chemistry Model (RACM) mechanism [Stockwell et al., 1997]. This mechanism is suitable for representation of remote to polluted urban conditions. The model is constrained with the observed 1-minute averages of the concentrations along the flight track for species other than HO_x. The constrained species are NO_x (NO + NO₂), O₃, H₂O, CO, CH₄, acetone, propane, methanol, ethanol, ethane, PAN, HNO₃, H₂O₂ and CH₃OOH. When measurements were unavailable, median values shown in Table 1 were used for model calculation. The constrained parameters are held constant throughout each simulation. NO is allowed to vary diurnally and is constrained so that the NO concentration matches the measurement at the time of the day of the observation. Temperature, pressure and aerosol surface area are also determined from the aircraft observations. Photolysis rates are calculated for clear sky conditions with the NCAR Tropospheric Ultraviolet-Visible radiative transfer code [Madronich and Weller, 1990], scaled to the values

of J(NO₂) observed aboard the aircraft [Shetter et al., 2002] to account for cloud effects using the correction factor J(NO₂)/J(NO₂)_{clear sky}. For wavelengths smaller than those relevant for J(NO₂), this approach can lead to a slight overcorrection of the photolysis rate (for example J(O¹D)).

[11] Heterogeneous reactions of HO₂, NO₂, NO₃ and N₂O₅ in aerosols are included with reaction probabilities of 0.1, 0.0001, 0.01, and 0.1 respectively [Jacob, 2000] and assuming aerosols to be aqueous. Aerosol surface areas measured during flight 10 remained around 3 μm² cm⁻³ except in cirrus clouds associated with convective outflow, where surface areas were considerably higher (see Table 1). In these cirrus clouds, a reaction probability of 0.025 was used for HO₂ following Cooper and Abbatt [1996].

[12] The photochemical model is integrated to diurnal steady state as defined by reproducibility of concentrations over a one-day cycle. CH₂O is a prognostic variable in photochemical steady state in the model; however, the model systematically underestimates the observations by a factor of 2 to 6. A similar difficulty in simulating CH₂O has been pointed out for the ensemble of PEM-Tropics B data [Ravetta et al., 2000] and in previous studies [Schultz et al., 1999; Jaeglé et al., 2001]. According to Singh et al. [2000], heterogeneous conversion of methanol to CH₂O on aerosols might provide an additional source of formaldehyde. During Flight 10, however, no clear correlation was found between methanol and aerosol surface area. Constraining CH₂O with observed values increases of HO₂ mixing ratios by 25%.

[13] Peroxides in the upper troposphere may be produced locally, serving as reservoirs for HO_x, or they may be convected from lower altitudes, providing a primary source of HO_x to the upper troposphere [Jaeglé et al., 1997; Prather and Jacob, 1997; Cohan et al., 1999; Mari et al., 2000]. The impact of convective injection can be estimated by comparison of the photochemical steady state concentrations computed from the model with observed values (results of both nonlocal and local contributions). Local contribution was determined from a steady state photochemical calculation with all the normally fixed species except CH₃OOH and H₂O₂. The ratio of measured-to-calculated concentrations in the upper troposphere then provides a good indicator of nonlocal contribution if higher than unity. A second criteria was applied to determine where nonlocal contribution coincide with convective injection. We looked at air masses for which CH₃I and relative humidity were in the top octile of measurements at 8–12 km altitude (i.e., CH₃I > 0.2 pptv and relative humidity > 89%) [Cohan et al., 1999]. Thus, in the following, convected peroxides are defined as the nonlocal portion determined from a steady state calculation when correlated with high CH₃I and relative humidity. Both criteria lead to the conclusion that convection increased CH₃OOH by factor of 1.9–5.5, essentially in the SPCZ cloud region (Table 2), suggesting that, in the case of CH₃OOH, convective transport could be important. By contrast, observed median H₂O₂ mixing ratios were within 30% of the photochemical steady state computed by the model, revealing that local photochemistry could sustain the observed H₂O₂ mixing ratios without invoking transport. This is also consistent with vertical transport of H₂O₂ being limited in the con-

Table 2. Median Values for the Ratio of Measured-to-Calculated CH₃OOH and H₂O₂ Total Production and Amplification Factor in the Upper Troposphere (8 to 12 km) for the Four Regions Along the Flight Track in Figure 1^a

	24°–20°S South of SPCZ	20°–17°S SPCZ	17°–13°S North of SPCZ	13°–10°S ITCZ
H ₂ O ₂ (meas/calc)	0.98 (0.63–3.6)	0.78 (0.13–1.73)	1.03 (0.23–1.38)	0.87 (0.4–1.60)
CH ₃ OOH (meas/calc)	1.03 (0.32–1.97)	1.44 (0.23–5.55)	1.33 (0.37–3.63)	1.23 (0.43–3.14)
Total HO _x production pptv/d	1577 (589–4121)	1270 (563–4271)	1238 (537–2619)	2349 (1347–6890)
Amplification factor ^b	1.32 (1.09–1.80)	1.14 (0.95–1.56)	1.22 (1.01–1.29)	1.12 (0.86–1.54)

^a The values are medians. The minimum and maximum values are listed in parentheses.

^b Amplification factor for methane oxidation A_{CH_4} is defined as the relative increase in the primary source of HO_x due to oxidation of CH₄ by OH.

vective updrafts by scavenging by convective precipitations [Crawford *et al.*, 1999; Mari *et al.*, 2000; Pickering *et al.*, 2001].

[14] In the following analysis, we consider the primary sources of HO_x from photolysis of convected CH₃OOH and H₂O₂, photolysis of acetone and photolysis of ozone with subsequent reaction of O(¹D) with water vapor. In order to quantify the relative importance of these primary sources of HO_x, we adapted the method described in previous investigations to our chemical mechanism [Arnold *et al.*, 1997; Folkins and Chatfield, 2000; Jaeglé *et al.*, 2001]. This method is based on the calculation of HO_x yields (i.e., the total number of HO_x molecules generated for each molecule of precursor photolyzed). The primary HO_x production, $P(\text{HO}_x)$ can then be expressed following Jaeglé *et al.* [2001]:

$$P(\text{HO}_x) = 2k[\text{O}(\text{D})][\text{H}_2\text{O}] + \sum_i Y_i J_i [\text{X}_i]$$

where X_i is either CH₃OOH, H₂O₂, or acetone, J_i is the associated photolysis rate, and Y_i is the corresponding yield.

[15] Oxidation of methane is considered as a secondary source of HO_x because it depends on the initial concentration of available OH [Wennberg *et al.*, 1998; Jaeglé *et al.*, 2001]. A measure of the amplification of HO_x production by this autocatalytic source is given by the amplification factor calculated following Jaeglé *et al.* [2001] and defined as the relative increase in the primary source of HO_x due to oxidation of CH₄ by OH:

$$A_{CH_4} = \frac{P(\text{HO}_x) + y_{CH_4} k[\text{OH}][\text{CH}_4]}{P(\text{HO}_x)}$$

[16] In Figure 5, we examine the trend of HO_x yields from CH₃OOH, acetone and CH₄ as a function of NO_x mixing ratios and $P(\text{HO}_x)$ calculated by the box model. The HO_x yields from methane and acetone increase with increasing NO_x. The HO_x yield from CH₄ increases with NO_x from about 0.25 at NO_x = 10 pptv to 0.5 at NO_x = 60 pptv and decreases with increasing HO_x production rate from 0.5 to nearly zero. These variations are consistent with previous studies [Folkins and Chatfield, 2000; Jaeglé *et al.*, 2001]. The HO_x yield from acetone increases with increasing NO_x and decreasing primary production. This variation reflects the greatest impact of acetone on tropospheric chemistry for high NO_x mixing ratios. The HO_x yield from CH₃OOH decreases with increasing NO_x and increasing primary production from about 1.7 at NO_x = 10 pptv and $P(\text{HO}_x)$ =

100 pptv/d down to 1.3 at NO_x = 60 pptv and $P(\text{HO}_x)$ = 1500 pptv/d similar to the values calculated by Folkins and Chatfield [2000].

4. Comparison Between Model and Observations

[17] Figure 6 compares simulated and observed HO₂ and OH mixing ratios for each individual point of the 1-minutes merge data set in the upper troposphere above 8 km. Also




Figure 5. Calculated HO_x yields from methane (diamonds), methylhydroperoxide (crosses) and acetone (triangles) during flight 10 for each simulated point as a function of (a) NO_x mixing ratios and (b) total primary production rates, $P(\text{HO}_x)$. Regression lines (solid) are also plotted for each yield.




Figure 6. Comparison between observations and model calculations for (a) HO₂ mixing ratios, in pptv, (b) OH mixing ratios, in pptv and (c) HO₂/OH mixing ratio in the upper troposphere (8 km to 12 km). One minutes averages corresponding to PEM-Tropics B Flight 10 are shown by crosses. Also listed are the slope (s) of the linear fit, the least squares correlation coefficient (r^2) between model and observations, and the median ratio (R) of simulated-to-observed mixing ratios.

shown in this plot are the least squares correlation coefficient (r^2) between model and observations, the slope (s) of the linear fit obtained by the multilinear regression method [Bevington, 1969], and the median ratio (R) of simulated-to-observed mixing ratios. We find that the model captures 80% of the variance of the observed HO₂ concentrations. The median simulated-to-observed ratio is 1.1 and the slope

of the linear fit is 0.85. The simulation of OH mixing ratio is poorer with 38% of the variance of observed OH captured by the model. The median simulated-to-observed ratio is 0.86 and the slope is 0.67. Figure 7 shows HO_x mixing ratios from the model and the observations as a function of latitude. A maximum is both measured and simulated in the SPCZ region (13.5 pptv) at 19°S. In the northern edge of the SPCZ region, simulated HO_x mixing ratios decrease sharply under 1 pptv due to high aerosol surface areas representative of cirrus clouds. There were no observations under these conditions. Two maxima are calculated and observed in the southern branch of the ITCZ with HO_x levels rising up to 13 and 15 pptv, respectively. Between the two cloud bands, HO_x mixing ratios are halved and the dry intermediate region is characterized by a minimum of HO_x abundances. The origin of these latitudinal variations is studied in the following section. As seen in Figure 6, the HO₂/OH ratio varies from 20 to 90, with the median model to observed ratio for HO₂/OH of 1.26. The model represents the observed ratio with a correlation coefficient $r^2 = 0.37$. However, this low correlation coefficient is better than the one obtained for the entire set of data during PEM-Tropics




Figure 7. Comparison of simulated (diamonds) and observed (crosses) (a) HO_x mixing ratios, in pptv, and (b) HO₂/OH ratio, as a function of latitude, for both outbound and return parts of flight 10.




Figure 8. HO_x yields of (triangles) CH₄, CH₃OOH (crosses) and acetone (diamonds) as a function of latitude. Solid segments represent averaged values of CH₄, CH₃OOH, and acetone yields in the four latitudinal regions in Figure 1. $P(\text{HO}_x)$ stands for total primary production rate.

B ($r^2 = 0.01$ in the 6- to 12-km altitude range; Ravetta et al., submitted). In Figure 7, the variation of the HO₂/OH ratio with latitude shows a region of enhanced values south of 20°S, where NO concentrations were also largest. The model simulates the observed peaks of HO₂/OH ratio along the flight track, although, it tends to slightly overestimate the ratio.

5. Primary Sources of HO_x

[18] In order to quantify and better understand the processes involved in the variation of the HO_x levels as a function of latitude, we perform a budget calculation following the methodology described above. Figure 8 shows the variation of HO_x yields as a function of latitude for methylhydroperoxide, acetone, and methane in the four regions described previously. The HO_x yield of acetone is maximum south of 20°S as could be expected in a region of relatively high NO_x mixing ratios and low $P(\text{HO}_x)$. The yield is minimum north of 10°S where the impact of high NO_x mixing ratios is counteracted by high $P(\text{HO}_x)$. HO_x yield CH₃OOH is maximum in the cloud free region between SPCZ and the zonal cloud band, characterized by low NO_x mixing ratio and low $P(\text{HO}_x)$. In contrast, minimum values are calculated above 13°S where mean NO_x mixing ratio and $P(\text{HO}_x)$ are higher. The upper troposphere in the tropical region sampled during PEM-Tropics B is characterized by high HO_x production rates (around 2000 pptv/d) and low NO_x concentrations (<60 pptv). Under these conditions, negative values of yield from HO_x can be predicted as shown in Figure 8. During this flight, we found a maximum amplification of the primary source due to the oxidation of methane ($A_{\text{CH}_4} = 1.80$) in the southern-

most region for points combining maximum observed NO levels (40 < NO < 60 pptv) and low $P(\text{HO}_x)$ (<200 pptv/d). Only 6% of the simulated points reveal a small quenching (A_{CH_4}). These points correspond to observed NO levels lower than 20 pptv and $P(\text{HO}_x)$ higher than 1000 pptv/d. Thus, the amplification is largest when is low $P(\text{HO}_x)$ and NO_x mixing ratio is high. This result is concordant with previous findings showing that the dependence of A_{CH_4} on $P(\text{HO}_x)$ acts as a weak buffer for HO_x concentrations [Crawford et al., 1999; Jaeglé et al., 2001].

[19] Dominant primary production terms are plotted in Figure 9 in the 10- to 12-km altitude range where convective injection of CH₃OOH was identified. A large fraction of the total HO_x production is attributed to water vapor, photolysis of CH₃OOH pumped by convection from the marine boundary layer and acetone. North of 20°S, HO_x production is driven principally by the O¹D + H₂O reaction (Table 3); this contribution is strongest in the cloudy regions corresponding to the SPCZ and ITCZ cloud bands (Figure 1) where it accounts for 90% of the total primary production of HO_x. We found that 75% of the simulated HO_x variance is produced by water vapor. In the drier regions south of 20°S, production from water vapor and acetone are comparable. Production from acetone contributes to 18–81% of the HO_x primary production in the 8- to 10-km altitude range (mean contribution 33%) and to 36–77% in the 10- to 12-km altitude range (mean contribution 49%). We found that production from acetone becomes significant and can be dominant over production from water vapor for water vapor mixing ratios lower than 200 ppmv. Impact of convected CH₃OOH on the HO_x production is limited to the SPCZ cloud band. The effect of convected CH₃OOH is sensitive at 10–12 km. In Table 3, the primary production rate from




Figure 9. HO_x production rates, in pptv/d, from water vapor (plusses), convected CH₃OOH (stars) and acetone (diamonds) as a function of latitude in the 10- to 12-km altitude range, for the four latitudinal regions in Figure 1.

CH₃OOH accounts for 22% to 64% of the total primary source in this altitude range. Clearly, this impact is spatially restricted to the outflow of the cloud.

6. Conclusion

[20] A 0-D model is applied to study the photochemistry of HO_x (HO₂ + OH) in the tropical upper troposphere, during the PEM-Tropics B experiment. In particular, airborne observations across the SPCZ-related convection gave the opportunity to quantify the relative contribution of the sources of HO_x in a remote nonpolluted tropical region, characterized by NO_x-limited conditions and influenced by marine deep convective clouds. The predominant primary sources of HO_x are identified as reaction O(¹D) + H₂O, photolysis of convected CH₃OOH and photolysis of acetone. HO_x chemistry shows a latitudinal dependence, reflecting the different origins of the air masses traversed by

the aircraft. The southernmost region (<20°S) is characterized by a strong subsidence associated with the upper tropospheric subtropical jet and advection of air from Australia. Because of the subsidence, water vapor mixing ratios are low and advection of continental air masses induces relatively high NO_x and ozone mixing ratios. In this region, HO_x primary production from acetone and water vapor are comparable, in particular in the 10- to 12-km altitude range where production from acetone can account for 36% to 77% of the total primary production rate. Amplification of the primary source due to the oxidation of methane shows maximum values (1.1–1.8) in the southernmost dry region. The SPCZ region is characterized by elevated water vapor mixing ratios and convective injection of low soluble species like CH₃I and CH₃OOH in the cloud outflow. Total primary production rates are higher than in the southernmost dry region and controlled mainly by water vapor abundances. When enhancement of CH₃OOH by convective transport is predicted, production from convected CH₃OOH becomes important and contributes for 22% to 64% of the primary production rate. The northern cloud band, corresponding to the southern branch of the ITCZ, is characterized by higher NO_x mixing ratios and HO_x production rates compared to the SPCZ cloud band. No impact of acetone or convected CH₃OOH is predicted and the reaction O(¹D) + H₂O is largely prevailing. This region is also characterized by some occurrences of low NO_x mixing ratios and elevated primary production rates leading to a diminution of HO_x primary sources by oxidation of methane. Between, the two cloud bands the aircraft passed through a dry region characterized by relatively low NO_x mixing ratios, low primary production rates, and minimum HO_x mixing ratios. In this region, production from acetone is important, although rarely dominant, and methane oxidation amplifies the primary source production by 1.0–1.3. The general good agreement in the clean tropical Pacific upper troposphere should not hide the remaining uncertainties regarding HO_x in the upper troposphere. Problems remain for very high NO_x at which systematic differences between observed and simulated HO_x have been found, reflecting gaps in our understanding of the coupling between HO_x and NO_x [Brune *et al.*, 1999;

Table 3. Mean Relative Contribution of Primary Sources in the Upper Troposphere in the 8–10 km and 10–12 km Altitude Ranges and for the Four Latitudinal Regions Along the Flight Track in Figure 1

	24°–20°S	20°–17°S	17°–13°S	13°–10°S
<i>10–12 km Altitude</i>				
$P(\text{HO}_x)^a$, pptv/d	322 (95–528) ^b	374 (76–1727)	254 (82–598)	549 (320–1223)
$P(\text{HO}_x)(\text{O}^1\text{D} + \text{H}_2\text{O})$, %	51 (23–64)	59 (13–91)	66 (44–82)	77 (64–92)
$P(\text{HO}_x)(\text{CH}_3\text{OOH})$, %	0 (0–0) ^c	11 (0–64)	0 (0–0)	0 (0–0)
$P(\text{HO}_x)(\text{acetone})$, %	49 (36–77)	30 (9–55)	34 (18–56)	23 (8–36)
<i>8–10 km Altitude</i>				
$P(\text{HO}_x)$, pptv/d	715 (96–1581)	1141 (375–2442)	681 (171–1193)	1482 (510–3832)
$P(\text{HO}_x)(\text{O}^1\text{D} + \text{H}_2\text{O})$, %	67 (19–82)	84 (59–95)	79 (54–91)	93 (79–99)
$P(\text{HO}_x)(\text{CH}_3\text{OOH})$, %	0 (0–0)	0 (0–0)	0 (0–0)	0 (0–0)
$P(\text{HO}_x)(\text{acetone})$, %	33 (18–81)	16 (5–41)	21 (9–46)	7 (1–21)

^aPrimary sources of HO_x, $P(\text{HO}_x)$ are photolysis of convected CH₃OOH and H₂O₂, photolysis of acetone, and photolysis of ozone with subsequent reaction of O(¹D) with water vapor. During flight 10, no convective injection of H₂O₂ was identified by the model.

^bThe values are means. The minimum and maximum are listed in parentheses.

^cZero values mean no contribution from photolysis of convected CH₃OOH to the total primary source of HO_x.

Faloona et al., 2000]. Chemistry of HO_x at night and sunlight/sunset is still poorly understood [Wennberg et al., 1999; Jaeglé et al., 1999] and the fate of inside cirrus clouds certainly needs further study.

References

- Arnold, F., V. Burger, B. Droste-Franke, F. Grimm, A. Krieger, J. Schneider, and T. Stimp, Acetone in the upper troposphere and lower stratosphere: Impact on trace gases and aerosols, *Geophys. Res. Lett.*, **24**, 3017–3020, 1997.
- Bevington, P., *Data Reduction and Error Analysis for the Physical Sciences*, McGraw-Hill, New York, 1969.
- Brune, W. H., et al., OH and HO₂ chemistry in the north Atlantic free troposphere, *Geophys. Res. Lett.*, **25**, 3077–3080, 1999.
- Cohan, D. S., M. G. Schultz, D. J. Jacob, B. G. Heikes, and D. R. Blake, Convective injection and photochemical decay of peroxides in the tropical upper troposphere: Methyl iodide as a tracer of marine convection, *J. Geophys. Res.*, **104**, 5717–5724, 1999.
- Cooper, P. L., and J. P. D. Abbatt, Heterogeneous interactions of OH and HO₂ radicals with surfaces characteristic of atmospheric particulate matter, *J. Phys. Chem.*, **100**, 2249–2254, 1996.
- Crassier, V., K. Suhre, P. Tulet, and R. Rosset, Development of a reduced chemical scheme for use in mesoscale meteorological models, *Atmos. Environ.*, **34**, 2633–2644, 1999.
- Crawford, J., et al., Assessment of upper tropospheric HO_x sources over the tropical Pacific based on NASA GTE/PEM data: Net effect on HO_x and other photochemical parameters, *J. Geophys. Res.*, **104**, 16,255–16,273, 1999.
- Faloona, et al., Observations of HO_x and its relationship with NO_x in the upper troposphere during SONEX, *J. Geophys. Res.*, **105**, 3771–3783, 2000.
- Folkens, I., R. Chatfield, H. Singh, Y. Chen, and B. Heikes, Ozone production efficiencies of acetone and peroxides in the upper troposphere, *Geophys. Res. Lett.*, **25**, 1305–1308, 1998.
- Folkens, I., and R. Chatfield, Impact of acetone on ozone production and OH in the upper troposphere at high NO_x, *Geophys. Res. Lett.*, **105**, 11,585–11,599, 2000.
- Gregory, G. L., et al., Chemical characteristics of Pacific tropospheric air in the region of the ITCZ and SPCZ, *J. Geophys. Res.*, **104**, 5697–5716, 1999.
- Jacob, D. J., Heterogeneous chemistry and tropospheric ozone, *Atmos. Environ.*, **34**, 2131–2159, 2000.
- Jacob, D. J., B. D. Field, E. Jin, I. Bey, Q. Li, J. A. Logan, and R. M. Yantosca, Atmospheric budget of acetone, *J. Geophys. Res.*, **107**(D9), 4100, doi:10.1029/2001JD000694, 2002.
- Jaeglé, L., et al., Observed OH and HO₂ in the upper troposphere suggest a major source from convective injection of peroxides, *Geophys. Res. Lett.*, **24**, 3181–3184, 1997.
- Jaeglé, L., et al., Ozone production in the upper troposphere and the influence of aircraft during SONEX: Approach of NO_x-saturated conditions, *Geophys. Res. Lett.*, **26**, 3081–3084, 1999.
- Jaeglé, L., D. J. Jacob, W. H. Brune, and P. O. Wennberg, Chemistry of HO_x radicals in the upper troposphere, *Atmos. Environ.*, **35**, 469–489, 2001.
- Lefer, B. L., S. R. Hall, L. Cinquini, R. E. Shetter, J. D. Barrick, and J. H. Crawford, Comparison of airborne NO₂ photolysis frequency measurements during PEM-Tropics B, *J. Geophys. Res.*, **106**, 32,645–32,656, 2001.
- Madronich, S., and G. Weller, Numerical integration errors in calculated tropospheric photodissociation rate coefficients, *J. Atmos. Sci.*, **10**, 289–300, 1990.
- Mari, C., D. J. Jacob, and P. Bechtold, Transport and scavenging of soluble gases in a deep convective cloud, *J. Geophys. Res.*, **105**(D17), 22,255–22,267, 2000.
- McKeen, S. A., T. Gierczak, J. B. Burholder, P. O. Wennberg, T. F. Hanisco, E. R. Keim, R.-S. Gao, S. C. Liu, A. R. Ravishankara, and D. W. Fahey, The photochemistry of acetone in the upper troposphere: A source of odd-hydrogen radicals, *Geophys. Res. Lett.*, **24**, 3177–3180, 1997.
- Müller, J.-F., and G. Brasseur, Sources of upper tropospheric HO_x: A three-dimensional study, *J. Geophys. Res.*, **104**, 1705–1715, 1999.
- Pickering, K. E., et al., Trace gas transport and scavenging in PEM Tropics-B SPCZ convection, *J. Geophys. Res.*, **106**(D23), 32,591–32,607, 2001.
- Prather, M. J., and D. J. Jacob, A persistent imbalance in HO_x and NO_x photochemistry of the upper troposphere driven by deep tropical convection, *Geophys. Res. Lett.*, **24**, 3189–3192, 1997.
- Raper, J. L., M. M. Kleb, D. J. Jacob, D. D. Davis, R. E. Newell, H. E. Fuelberg, J. M. Bendura, J. M. Hoell, and R. J. McNeal, Pacific Exploratory Mission in the Tropical Pacific: PEM-Tropics B, March–April 1999, *J. Geophys. Res.*, **106**(D23), 32,401–32,425, 2001.
- Ravetta, F., et al., Experimental evidence for the importance of convected methylhydroperoxide as a source of hydrogen oxide (HO_x) radicals in the tropical upper troposphere, *J. Geophys. Res.*, **106**, 32,709–32,716, 2001.
- Schultz, M. G., et al., On the origin of tropospheric ozone and NO_x over the tropical South Pacific, *J. Geophys. Res.*, **104**, 5829–5843, 1999.
- Singh, H. B., M. Kanakidou, P. Crutzen, and D. Jacob, High concentrations and photochemistry of carbonyls and alcohols in the global troposphere, *Nature*, **378**, 50–54, 1995.
- Singh, et al., Distribution and fate of selected oxygenated organic species in the troposphere and lower stratosphere over the Atlantic, *J. Geophys. Res.*, **105**, 3795–3806, 2000.
- Stockwell, W. R., F. Kirchner, M. Kuhn, and S. Seefeld, A new mechanism for regional atmospheric chemistry modeling, *J. Geophys. Res.*, **22**, 25,847–25,879, 1997.
- Wennberg, P. O., et al., Hydrogen radicals, nitrogen radicals and the production of O₃ in the upper troposphere, *Science*, **279**, 49–53, 1998.
- Wennberg, P. O., et al., Twilight observations suggest unknown sources of HO_x, *Geophys. Res. Lett.*, **26**, 1373–1376, 1999.
- B. Anderson, G. L. Gregory, G. W. Sachse, and S. Vay, NASA Langley Research Center, Hampton, VA 23681, USA.
- D. Blake, Department of Chemistry, University of California, Irvine, CA 92717, USA.
- W. H. Brune, Department of Meteorology, The Pennsylvania State University, University Park, PA 16802-5013, USA.
- I. Faloona, Atmospheric Chemistry Division, National Center for Atmospheric Research, Boulder, CO 80305, USA.
- B. Heikes, Center for Atmospheric Chemistry Studies, University of Rhode Island, Narragansett, RI 02882, USA.
- D. J. Jacob, Division of Engineering and Applied Science, Harvard University, Cambridge, MA 02138, USA.
- C. Mari and C. Saüt, Laboratoire d'Aérodynamique, UMR 5560 CNRS/Université Paul Sabatier, Toulouse, F-31400, France. (marc@aero.obs-mip.fr)
- M. Avery, Department of Physics, Hampton University, Hampton, VA 29668, USA.
- S. T. Sandholm, Earth and Atmospheric Sciences Baker Building, Georgia Institute of Technology, Atlanta, GA 30332, USA.
- H. B. Singh, NASA Ames Research Center, Moffett Field, CA 94035, USA.
- F. Ravetta, Service d'Aéronomie du CNRS, Université Paris 6, Boite 102, 4 Place Jussieu, 75252, Paris Cedex 05, France.
- R. W. Talbot, CSRC/EOS, University of New Hampshire, Durham, NH 30824, USA.
- D. Tan, Georgia Institute of Technology, Atlanta, GA 30332-0340, USA.