

Lawrence Berkeley National Laboratory

Recent Work

Title

OK THE MECHANIC OF RADIOLYSIS IN AQUEOUS FORMIC ACID SOLUTIONS

Permalink

<https://escholarship.org/uc/item/5037888b>

Authors

Garrison, Warren M.
Bennett, Winifred
Jayko, Michael.

Publication Date

1955-11-01

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-486

ON THE MECHANISM OF RADIOLYSIS IN AQUEOUS FORMIC ACID SOLUTIONS

Warren M. Garrison, Winifred Bennett and Michael Jayko

November, 1955

1

ON THE MECHANISM OF RADIOLYSIS IN AQUEOUS FORMIC ACID SOLUTIONS

(1) This work was performed under the auspices of the Atomic Energy Commission

Warren M. Garrison, Winifred Bennett and Michael Jayko


Crocker Laboratory, Radiation Laboratory, University of California
Berkeley, California

November, 1955

In detailed kinetic studies of the indirect action of γ -radiation on dilute formic acid solutions, Hart ² has shown that H and OH free radicals

(2) E. J. Hart, (a) J. Phys. Chem. 56, 594 (1952); (b) J. Am. Chem. Soc. 76, 4198 (1954); (c) ibid 76, 4312 (1954).

formed by radiation decomposition of water react with formic acid according to



Radical removal then occurs via



or through the alternate to 3,4, i.e.,



The γ -ray data are consistent with the conclusion that no net reduction of formic acid occurs in dilute solution at pH values below 3. However, at concentrations above .05 M, evidence for the reduction of formic acid has been obtained. ^{2c} On the basis of the reaction scheme, 1-5, it is seen that reduction cannot involve a competing reaction of the type



since both 2 and 2a have the same concentration dependency. Some other intermediate is apparently involved in reaction with the HCOOH species. It has been proposed (a) that this intermediate is the secondary electron formed in the ionization process and (b) that electron capture by formic acid ($\text{HCOOH} + e = \text{HCO} + \text{OH}^-$) competes with electron capture by water ($\text{H}_2\text{O} + e = \text{H} + \text{OH}^-$) at the higher formic acid concentration.

Results recently obtained in this Laboratory indicate, however, that an alternative mechanism may be involved. It has been shown that irradiation

(3) W. M. Garrison, W. Bennett, M. Jayko, and S. Cole, presented at 128th American Chemical Society Meeting, Minneapolis, Minn., 1955. Univ. of California Radiation Lab. Report, 3208 (1955).


of dilute oxygen-free formic acid solutions with cyclotron-produced helium ions results in the formation of higher organic acids and carbonyl compounds as major products. Both glyoxal and glyoxylic acid, together with smaller amounts of oxalic, glycolic, tartronic, mesoxalic, and tartaric acids are formed at radiation doses which correspond to the removal of less than one percent of the formic acid initially present in the target solution.

Evidence that the glyoxylic acid observed in these studies is not formed through secondary processes initiated by H reduction of oxalic acid can be seen in the experimental data published several years ago by Fricke, Hart, and Smith. In this earlier work it was observed that hydrogen production



(4) H. Fricke, E. J. Hart, and H. P. Smith, J. Chem. Phys. 6, 229 (1938).

in the x-ray irradiation of formic acid-oxalic acid mixtures did not decrease with increasing oxalic acid concentrations until the ratio $C(\text{COOH})_2 / C\text{HCOOH}$ approached a value of 1. The corresponding ratio in the cyclotron irradiated solutions was approximately 10^{-3} .

On the basis of a consideration of the γ -ray and cyclotron data, it is concluded that radiolysis of formic acid solutions under both conditions can be interpreted without the assumption of competing electron capture by the solute if it is assumed that H adds reversibly to formic acid according to




If reaction 1 above is fast as compared to 2, then the reduced formic acid radical $HC(OH)_2$ could combine with $COOH$ to give glyoxylic acid or dimerize to form glyoxal at the high radical concentrations obtained in cyclotron irradiations, i.e.,



(5) W. M. Garrison, W. Bennett, S. Cole, H. R. Raymond, and B. M. Weeks, J. Am. Chem. Soc. 77, 2720 (1955).

In the γ -ray irradiations, the radical concentrations are lower, and in the more dilute formic acid solutions H would be preferentially removed via 2. The concentration effect observed by Hart is then accounted for on the assumption that reaction 8 occurs as a competing process for removal of $HC(OH)_2$



A detailed study of the proposed mechanism is now in progress in this laboratory. Information on the properties of free radical intermediates involved in the radiolysis of other aquo-organic systems is being obtained through study of radical combination reactions induced by cyclotron irradiation.

We wish to thank the staff of the 60-inch Cyclotron at Crocker Laboratory for their continued cooperation in this work.