

UCSF

UC San Francisco Previously Published Works

Title

Organizational Predictors of Colonoscopy Follow-up for Positive Fecal Occult Blood Test Results: An Observational Study

Permalink

<https://escholarship.org/uc/item/50j9q992>

Journal

Cancer Epidemiology Biomarkers & Prevention, 24(2)

ISSN

1055-9965

Authors

Partin, Melissa R
Burgess, Diana J
Burgess, James F
[et al.](#)

Publication Date

2015-02-01

DOI

10.1158/1055-9965.epi-14-1170

Peer reviewed



Published in final edited form as:

Cancer Epidemiol Biomarkers Prev. 2015 February ; 24(2): 422–434. doi:
10.1158/1055-9965.EPI-14-1170.

Organizational predictors of colonoscopy follow-up for positive fecal occult blood test results: an observational study

Melissa R. Partin^{1,2}, Diana J. Burgess^{1,2}, James F. Burgess Jr.^{3,4}, Amy Gravelly¹, David Haggstrom^{5,6}, Sarah E Lillie^{1,2}, Sean Nugent¹, Adam A Powell¹, Aasma Shaukat^{1,2}, Louise C. Walter⁷, and David B. Nelson^{1,2}

¹Center for Chronic Disease Outcomes Research, Minneapolis Veterans Affairs Health Care System

²Department of Medicine, University of Minnesota

³Center for Healthcare Organization and Implementation Research, Boston Veterans Affairs Health Care System

⁴Department of Health Policy and Management, Boston University School of Public Health

⁵VA Health Services Research & Development Center for Health Information and Communication, Roudebush VAMC, Indianapolis, IN

⁶Division of General Internal Medicine and Geriatrics, Department of Medicine, Indiana University School of Medicine

⁷Division of Geriatrics, San Francisco VA Medical Center and University of California, San Francisco

Abstract

Background—This study assessed the contribution of organizational structures and processes identified from facility surveys to follow-up for positive Fecal Occult Blood Tests (FOBT+).

Methods—We identified 74,104 patients with FOBT+ results from 98 Veterans Health Administration (VHA) facilities between 8/16/09-3/20/11 and followed them until 9/30/11 for completion of colonoscopy. We identified patient characteristics from VHA administrative records, and organizational factors from facility surveys completed by Primary Care and Gastroenterology Chiefs. We estimated predictors of colonoscopy completion within 60 days and 6 months using hierarchical logistic regression models.

Results—30% of patients with FOBT+ results received colonoscopy within 60 days and 49% within 6 months. Having Gastroenterology or Laboratory staff notify Gastroenterology providers directly about FOBT+ cases was a significant predictor of 60-day (odds ratio (OR)=1.85, p=0.01) and 6-month follow-up (OR 1.25, p=0.008). Additional predictors of 60-day follow-up included adequacy of colonoscopy appointment availability (OR 1.43, p=0.01) and frequent individual

Corresponding author contact information: Melissa R. Partin, PhD., Minneapolis Veterans Affairs Medical Center, 1 Veterans Drive, Minneapolis MN 55417, Melissa.partin@va.gov.

Potential competing interests: None

feedback to Primary Care providers about FOBT+ referral timeliness (OR 1.79, $p=0.04$). Additional predictors of 6-month follow-up included using guideline-concordant surveillance intervals for low-risk adenomas (OR 1.57, $p=0.01$) and using group appointments and combined verbal-written methods for colonoscopy preparation instruction (OR 1.48, $p=0.0001$).

Conclusion—Directly notifying Gastroenterology providers about FOBT+ results, employing guideline-concordant adenoma surveillance intervals, and using colonoscopy preparations instruction methods that provide both verbal and written information may increase overall follow-up rates. Enhancing follow-up within 60-days may require increased colonoscopy capacity and feedback to Primary Care providers.

Impact—These findings may inform organizational-level interventions to improve FOBT+ follow-up.

Keywords

Early Detection of Cancer; Diagnostic Services; Mass Screening; Colorectal Neoplasms; Colonoscopy; Health Services Accessibility; Veterans Health

INTRODUCTION

Colorectal cancer is the third most common cancer and the third leading cause of cancer death among men and women in the United States.(1;2) The best known defense against colorectal cancer is early detection and prevention through routine screening. Current guidelines endorse multiple colorectal cancer screening methods,(3–5) but fecal occult blood testing (FOBT) and colonoscopy are the most widely used.(6) Two of the largest integrated health care systems in the United States (Kaiser Permanente and the Veterans Health Administration) have achieved high colorectal cancer screening rates using screening programs emphasizing FOBT.(7;8) Although randomized controlled trials have demonstrated that FOBT can be a highly efficacious screening method if FOBT positive (FOBT+) results are followed by diagnostic colonoscopy,(9–11) many FOBT-based screening programs document challenges assuring that FOBT+ results receive follow-up colonoscopy in a timely manner.(12–15) Proportions of FOBT+ cases failing to receive follow-up colonoscopy reported in prior studies range from 35–63%,(12–15) and the median waiting times from FOBT+ to colonoscopy range from 105–202 days.(12–14;16)

Both the Veterans Health Administration (VHA) and the Canadian Association of Gastroenterology Wait Time Consensus Group recommend performing a colonoscopy within 60 days of FOBT+ results.(17;18) However, recent data from the VHA documenting that 50% of FOBT+ cases fail to receive follow-up colonoscopy within this window,(19) and from a Canadian survey of Gastroenterologists documenting an average waiting time from FOBT+ results to colonoscopy of 105 days(16) suggest significant gaps remain in assuring timely follow-up. Closing these gaps will require identifying modifiable contributors to persistent FOBT+ follow-up delays.

Most prior studies examining contributors to FOBT+ follow-up have focused on non-modifiable individual-level factors,(20;21) identifying significant associations with patient age,(22–25) gender,(14;24;26) race,(25) comorbidity,(23) personal history of bowel disease,

(27) family history of colorectal cancer,(28) and recent colonoscopy.(25) A few have identified modifiable individual-level contributors, including patient fears and worries about colorectal cancer(27;28) and provider awareness of guidelines(29;30) and intentions to order diagnostic testing for FOBT+ results.(26) However, very few prior studies have examined the contribution of modifiable organizational-level factors to FOBT+ follow-up patterns,(21) despite growing recognition that understanding the role that the care environment plays in assuring quality is essential to designing effective interventions and making further improvements in cancer care.(31;32)

We conducted a study to assess the contribution of modifiable organizational-level factors to FOBT+ follow-up rates. We hypothesized that higher follow-up rates would be associated with: (1) organizational structures designed to facilitate quality improvement (i.e., leadership support, resource alignment, feedback, and incentives), and (2) organizational processes that: control system-wide demand for colonoscopy (i.e., “demand efficiency” processes); minimize wasted appointments and the number of steps required to complete a colonoscopy (i.e., “supply efficiency” processes), and address patient barriers to colonoscopy completion (i.e., “patient-centered” processes). We assessed the contribution of these organizational-level factors while controlling for individual-level factors demonstrated to be associated with FOBT+ follow-up in prior studies.

MATERIALS AND METHODS

Setting and participants

We identified a cohort of patients who had outpatient FOBT+ results from a Veterans Health Administration (VHA) facility between 8/2009-3/2011 (one year prior and six months after the start date for the organizational survey, described below) and followed them until 9/2011 (six months after the last FOBT date) for completion of follow-up colonoscopy. To identify patients with FOBT+ results, we identified from VHA laboratory records all outpatient FOBT procedures performed at VHA facilities during the sample accrual period, using the codes provided in Supplementary Material 1. We then defined FOBT+ cases as any individual card test with a positive result, or any multiple card series with one or more cards with positive results. If an eligible patient had more than one FOBT+ result in the sample selection window, the first FOBT+ result was selected for the sample. We then excluded patients if they: did not receive their FOBT+ results from one of 125 VHA facilities that conducted at least 1,400 FOBTs in 2009; were age <18 or >100 at the time the FOBT+ result was recorded; had a prior diagnosis of colorectal cancer in VHA medical records; or received their FOBT+ from a VHA community-based outpatient clinic that refers less than 70% of colonoscopies to one of the 125 VHA facilities included in the sampling frame, leaving 86,926 eligible FOBT+ patients available for analysis. We linked this patient sample to facility-level data on organizational structures and processes obtained from web-based surveys (described previously(33) and in Supplementary Material 2) administered to the Chiefs of Primary Care (73% response rate) and Gastroenterology (81% response rate) beginning in 8/2010, yielding 74,014 patients from 98 facilities responding to one or both surveys (Figure 1). We excluded 43 facilities completing <1,400 FOBT procedures in 2009

to ensure an adequate sample (> 100) of FOBT+ patients from each facility was available for estimating the association between organizational factors and follow-up rates.

Conceptual framework

Our conceptualization of key organizational-level contributors to FOBT+ follow-up is informed by Donabedian's framework for understanding the quality of care,(34) and prior research documenting the association between specific organizational structures, organizational processes, individual-level factors, and quality outcomes. Organizational structures associated with quality outcomes in prior studies include: *leadership support*, (35;36) personal and frequent *feedback*, (37;38) *incentives*,(39) and *resource alignment* to improvement goals(35;36;40) (such as tracking systems,(41) and quality improvement training.(15)) Organizational processes refer to approaches used to complete each step required to assure FOBT+ follow-up (i.e., notification, referral, scheduling, patient education). For our analysis, we categorized organizational processes into three groups: (1) those that control system-wide demand for colonoscopy, which we refer to as “*demand efficiency*” processes; (2) those that minimize wasted appointments and the number of steps required to complete a colonoscopy, which we refer to as “*supply efficiency*” processes; and (3) those that address patient barriers to colonoscopy completion, which we refer to as “*patient-centered*” processes. Individual-level factors we control for in our hypothesis tests related to organizational-level factors include characteristics of patients and FOBT procedures found to be associated with FOBT+ follow-up rates in prior studies (i.e., age, race, residence, comorbidities, personal history of polyps, and ordering provider characteristics).

Data Sources and Measures

Our primary outcome was follow-up colonoscopy completion, identified from VHA administrative records using the codes in Supplementary Material 3. We separately examined correlates of colonoscopy completion within 60 days (the VHA recommended follow-up interval) and 6 months.

Predictors—Table 1 provides the survey question wording, response options, and coding for analysis for all organizational predictors we examined.

Organizational structures: We measured *leadership support* using two items: (1) a question from the Primary Care survey asking the extent to which “not a priority to leadership” is a barrier to providing timely FOBT+ follow-up, and (2) an identical question from the Gastroenterology survey. We measured *resource alignment* with two items: (1) “tracking”, a question on the Primary Care Chief survey about how frequently their program tracks what happens to patients with FOBT+ results, and (2) a question on the Gastroenterology Chief survey asking the extent to which “colonoscopy appointment availability” is a barrier to providing timely FOBT+ follow-up. *Feedback* was assessed with two measures: (1) “Primary Care feedback”, which combined two measures on the type and frequency of feedback (see Table 1) into a single measure, and (2) “Gastroenterology feedback” which combined similar measures from the Gastroenterology survey. *Incentives* were assessed with two items: (1) “Primary Care Incentives”, and (2) “Gastroenterology

Incentives”, both of which asked Chiefs “which of the following do [providers in your Primary Care program/staff in your Gastroenterology program] receive for their performance on assuring timely follow-up of positive FOBT results” (see Table 1 for response options and coding).

Organizational processes: Measures of *demand efficiency processes* included: (1) number of contraindications the facility’s colorectal cancer screening clinical reminder asks about; (2) information on the facility’s colonoscopy consult template includes contraindications; and (3) the typical surveillance interval for patients with 1–2 adenomas <1cm at the facility is guideline-consistent (i.e., 5–10 years). (4) Measures of *supply efficiency processes* included: (1) Gastroenterology providers are directly notified of FOBT+ cases (either by Lab or by Gastroenterology Staff who take responsibility for identifying FOBT+ cases); (2) patient colonoscopy prep instruction does not require a separate appointment; (3) a pre-op appointment is not required for colonoscopy; and (4) overbooking is used to minimize wasted appointments. Measures of *patient-centered processes* included: (1) patient notification of FOBT+ results includes phone contact; (2) colonoscopy appointment times are negotiated (using a scheduler or letter requesting the patient call the clinic to set up an appointment) rather than pre-assigned to patients in a mailed letter; (3) patient prep instruction procedures include opportunities for questions (i.e., some verbal instruction); and (4) patients receive colonoscopy appointment reminders that include prep instructions.

We used VHA administrative data to identify the following individual-level factors controlled for in our predictive models: age (<50, 50–64, 65–84, 85); race (Non-Hispanic White, Hispanic, African American, American Indian, Asian or Pacific Islander, Unknown); residence (urban, rural); drive time to the nearest VHA specialty care facility (<60, >60 minutes); Charlson comorbidity score for the 1 year prior to the FOBT+ result; mental health diagnoses (psychiatric only, substance abuse only, or dual diagnosis, none); personal history of colorectal polyps or benign neoplasms (ICD-9 211.3-4, 569.0, v12.72); whether the FOBT was ordered by their primary care provider; what type of provider ordered their FOBT (physician, nurse practitioner or physician assistant, resident, nurse or other staff); and what type of facility ordered their FOBT (VHA specialty care facility or community-based outpatient clinic).

Analysis—Because the limited number of facilities in the sample precluded the simultaneous inclusion of a large number of facility-level predictors, we pursued the following steps to select predictors for inclusion in the final model. We initially fit separate bivariate hierarchical logistic regression models (with random effects for facility of care) for each organizational structure and process measure. We then included in a base multivariable model all measures with p-values < 0.10, or with more than a 5% difference between model estimated completion rates (among the levels of a categorical measure or between the mean and one standard deviation shift from the mean for a continuous measure). We also included in the base model month of FOBT+ result and any patient-level predictors associated with colonoscopy completion within the respective timeframe (see Supplementary Table 1 for bivariate estimates derived from this step). We then reduced the number of predictors in this model in a stepwise fashion, retaining explanatory measures with p-values < 0.10. Using this

final multivariable model, we constructed model-based odds ratios and least square mean completion rate estimates (using the observed marginal distributions of the other covariates) for each of the organizational structure and process measures included in the model. To explore the impact of potentially valid reasons for not completing a follow-up colonoscopy at a VHA facility on our estimates, we fit a final set of models excluding patients who may not have been appropriate for colorectal cancer screening (i.e., age <45 or >85, with documentation of limited life expectancy in the medical record, or a colonoscopy in the prior 10 years), and treating patients who refused colonoscopy or chose to pursue colonoscopy in the private sector as having adequate follow-up. Supplementary Material 4 provides details on how we estimated the prevalence of each of these reasons for not completing a colonoscopy.

Human subjects approval

The study was reviewed and approved by the Institutional Review Boards at the Minneapolis VA Medical Center (approval 9/15/2009), and the Boston VA Medical Center (approval 2/10/2010).

RESULTS

Patients included in the analysis were primarily non-Hispanic white (65%) married (53%) men (96%) over the age of 50 (95%) (Table 2). Roughly half (52%) lived in urban areas, and most (61%) lived less than 60 minutes from a VHA medical center providing specialty services. A total of 20% had a personal history of colorectal polyps or benign neoplasms, 68% had a mental health or substance abuse diagnosis, and the mean Charlson Comorbidity Score was 1.9. Most participants (58%) had their FOBT procedures ordered at a VHA hospital, by their Primary Care provider (70%), and most ordering providers were physicians (72%).

Few facilities cited lack of leadership support as a barrier to improving FOBT+ follow-up rates, but 38 (49%) considered colonoscopy appointment availability a barrier (Table 3). The majority (64%) reported tracking what happens to patients with FOBT+ results on a weekly or more frequent basis, but only 11% reported providing individual, frequent feedback to Primary Care Staff about their FOBT+ referral practices. A higher percentage (22%) reported providing verbal, frequent feedback to Gastroenterology Staff about colonoscopy follow-up rates. The majority reported no incentives for Primary Care (56%) or Gastroenterology Staff (63%) tied to FOBT+ follow-up performance. About half (48%) reported including some information on contraindications on their colorectal cancer screening clinical reminder, and 47% reported including information on contraindications on their colonoscopy consult template. The modal surveillance interval for patients with 1–2 adenomas <1cm was 5 years (84%). Most (70%) relied on Primary Care to notify Gastroenterology of FOBT+ cases, and most (70%) did not require a separate appointment for colonoscopy prep instruction. The majority (62%) did not require a pre-op appointment for colonoscopy, and 54% reported using overbooking to minimize wasted colonoscopy appointments. Most (67%) used patient notification procedures that included some phone contact, but scheduling procedures were highly variable. The approach used to instruct

patients about colonoscopy preparation was highly variable, with 36% using written methods only, 23% using verbal methods delivered by phone or an individual appointment, and 32% using group appointments or some other method involving both written and verbal instruction. Finally, most (69%) did not review prep procedures in their colonoscopy appointment reminders.

The cumulative proportion of patients with FOBT+ results receiving a colonoscopy at a VHA facility within 60 days was 30% (range 10–57% across facilities). Organizational structures significantly associated with 60-day follow-up rates in the multivariable model included: *colonoscopy appointment availability* is not a key barrier (OR 1.43, CI 1.09–1.90, $p=0.01$); and monthly or more frequent *Primary Care feedback* (OR 1.79, CI 1.02–3.16, $p=0.04$). Organizational processes significantly associated with 60-day follow-up rates in the multivariable model included: *colonoscopy consult information* does not ask about indication or contraindications for colonoscopy (OR 1.48, CI 1.07–2.05, $p=0.02$), or asks about indication only (OR 1.49, CI 1.10–2.02, $p=0.01$) (a finding counter to our demand efficiency hypothesis), and *Gastroenterology notification* directly of FOBT+ results by Gastroenterology Staff (OR 1.85, CI 1.17–2.91, $p=0.01$) (Table 4). No patient-centered process measures were significantly associated with the 60-day outcomes in the adjusted model. After excluding potentially inappropriate FOBTs, and treating refusals and private sector colonoscopies completed after the FOBT+ as adequately followed up, the estimated follow-up rates increase markedly, and the odds ratio estimates for colonoscopy appointment availability and Gastroenterology notification remain significant. However, the odds ratio estimates for Primary Care feedback, and colonoscopy consult information attenuate and are no longer statistically significant.

The cumulative proportion of patients with FOBT+ results receiving a colonoscopy within 6 months was 49% (range 30–70% across facilities). No organizational structures were significantly associated with the 6-month outcome in the multivariable model. Organizational processes significantly associated with 6-month follow-up rates in the multivariable model included: using an *adenoma surveillance interval* of at least 5 years (5 years OR 1.32, CI 1.02–1.71, $p=0.04$; 7–10 years OR 1.57, CI 1.11–2.20, $p=0.01$); having *Gastroenterology notification* directly from lab about FOBT+ cases (OR 1.25, CI 1.06–1.47, $p=0.008$); and using group appointments or other combined verbal and written methods to provide patient *colonoscopy prep instruction* (OR 1.48, CI 1.22–1.79, $p=0.05$) (Table 5). After excluding potentially inappropriate FOBTs, and treating refusals and private sector colonoscopies completed after the FOBT+ as adequately followed up (last 2 columns of Table 5), the estimated follow-up rates increase to more than 67% in all subgroups, and only the estimate of group appointment for colonoscopy prep instructions remained significant (OR 1.50, CI 1.16–1.95, $p=0.003$).

DISCUSSION

Consistent with our hypothesis that organizational structures supporting quality improvement would be positively associated with follow-up rates, we found that adequacy of colonoscopy appointment availability, and providing primary care providers with individual, frequent feedback about the timeliness of FOBT+ referrals were positively

associated with receiving follow-up colonoscopy within 60 days of an FOBT+. These findings are consistent with prior research documenting that limited colonoscopy capacity is the most common barrier to reducing FOBT+ follow-up delay,(41) and that individual, frequent feedback can improve adherence to clinical practice guidelines.(43;44) However, the fact that no organizational structures were significantly associated with 6 month follow-up rates suggests that these factors have more influence on how quickly, rather than whether, patients with FOBT+ results receive colonoscopy follow-up.

We found partial support for our hypotheses that higher follow-up rates would be associated with organizational processes enhancing demand efficiency, supply efficiency, and patient-centered processes. Specifically, our findings suggest that: using surveillance intervals for low-risk adenomas that are not more aggressive than recommended by guidelines(42) (a demand efficiency measure); assigning responsibility for identifying FOBT+ cases to Lab or Gastroenterology staff (a supply efficiency measure); and employing group and other combined verbal and written colonoscopy prep instruction processes (a patient-centered process) are positively associated with overall follow-up rates.

One organizational process measure (Gastroenterology providers are notified by lab or Gastroenterology Staff about FOBT+ results) was significantly associated with both 60-day and 6-month follow-up rates. These results are consistent with findings from a prior randomized trial conducted in four VHA facilities, which found 30-day, 90-day and 6-month follow-up rates improved significantly (by 9–31%, $p < 0.03$) in facilities that implemented an electronic intervention to directly notify Gastroenterology staff of FOBT+ results, but did not significantly change in the usual care comparison facilities.(2)

Counter to our hypothesis that organizational processes designed to reduce unnecessary demand for colonoscopy would be positively associated with follow-up rates, we found facilities that asked about contraindications on colonoscopy consult templates had lower rather than higher 60-day follow-up rates. This finding might be explained by the fact that patients with documented limited life expectancy, recent colonoscopy, and refusal to complete follow-up colonoscopy were not initially excluded from our sample. Indeed, after excluding these cases from the analysis, the association of consult template characteristics with follow-up rates was no longer statistically significant. Thus, including information on contraindications on the colonoscopy consult template may identify individuals who should not have been screened, and some of these individuals appropriately do not have follow-up colonoscopy.

We hypothesized that colonoscopy prep instruction processes that involve some verbal instruction would be associated with higher follow-up rates than methods that involved only written instruction because verbal instruction processes would provide more opportunities to address patient questions. However, our findings suggest that some forms of verbal instruction (i.e., phone and individual appointments) were associated with lower 6-month follow-up rates than written only instruction methods. Because we did not collect information on the specific content of the prep instruction provided, we can only speculate about why group preparation instruction and other combined verbal and written methods were superior to verbal phone and individual appointment instruction. One possibility is that

group prep appointments and other combined methods may employ a more structured approach than other verbal instruction methods, and so are more likely to encourage patients to identify and clarify aspects of the preparation they do not understand. Alternatively the group/peer setting and other combined approaches may prompt greater patient engagement. The resulting enhanced clarification and/or engagement may increase the proportion of patients that attend their scheduled colonoscopy appointment and present with adequate bowel preparation, thereby reducing delays associated with needing to reschedule colonoscopy appointments. A final possibility is that instruction approaches that require patients to complete a group class or other formal instruction before scheduling a colonoscopy lead to self-selection of individuals that are more likely to adhere to their colonoscopy appointment. All of these explanations are consistent with findings from one prior randomized trial, which found that patients participating in a nurse-led group colonoscopy prep education program had higher colonoscopy completion rates and lower cancellation rates due to poor bowel preparation than patients who received an educational brochure only.(45) Our finding that other measures of patient-centered processes (phone results notification, negotiated appointment scheduling, appointment reminders that review prep instructions) were not associated with follow-up rates contrasts with previous studies attributing high endoscopy attendance to patient-centered processes such as education(46) and reminder systems,(47;48) may be unique to this patient population trained in the hierarchical traditions of the military, and may not generalize to other health care settings.

This study has a number of strengths, including the large sample size of patients and medical facilities, the rigorous methodology used to adjust our estimates for reasons a colonoscopy was not completed, and the fact that it identifies several modifiable organizational predictors of FOBT+ follow-up rates. However, our findings should be qualified by several limitations. First, we may be underestimating actual follow-up rates because some patients with FOBT+ results may have pursued colonoscopy outside of the VHA. Indeed secondary analyses we conducted on this cohort suggest that up to 15% of patients with FOBT+ results who did not receive a colonoscopy in VHA within 6 months had documentation in their chart notes that they were pursuing colonoscopy in the private sector. However, sensitivity analyses treating patients with documentation of pursuing colonoscopy in the private sector as adequately followed up did not significantly alter the pattern of associations between organizational factors and follow-up rates reported here. A second possible limitation is that our measures of organizational structures and processes may include some measurement error. Structure and process reports from Chiefs were measured at one point in time, in most cases with single-item measures, and may therefore be inaccurate (given that facilities may make periodic adjustments to structures and processes) or insufficiently sensitive (from oversimplification of the underlying processes). Further, lack of variability in our sample forced us to collapse potentially distinct categories for several measures. Future studies should examine whether more detailed measures in more variable facility samples yield different results. Additionally, our analysis excluded 12,822 FOBT+ patients from 25 facilities with incomplete facility survey data, and 10,806 FOBT+ from 43 facilities conducting fewer than 1,400 FOBTs in 2009, which may raise concerns about whether our findings can be generalized to FOBT+ patients from other VHA facilities in the sampling frame. However, our previous analysis of the survey data found no significant variation in

facility FOBT+ follow-up rates or characteristics by survey response status,(33) and facilities excluded based on FOBT volume represented not only smaller facilities with FOBT-based screening programs, but also larger facilities with colonoscopy-based screening programs. Finally, the VHA is a unique context, characterized by a predominantly male, low income population with higher than average comorbidity burden, including high rates of mental health and substance abuse diagnoses. Therefore, our findings may not generalize to other health care contexts. Given that VHA is the largest integrated health care system in the United States, however, our findings have important implications for a substantial population of health providers and consumers in this country.

Despite these limitations, the insights gleaned from this study regarding the role organizational structures and processes can play in assuring patients with FOBT+ results receive timely colonoscopy will be helpful in guiding future efforts to improve FOBT+ follow-up rates. Specifically, our most robust findings suggest that Gastroenterology clinics may be able to significantly increase the proportion of FOBT+ results that receive follow-up colonoscopy by assuming responsibility for identifying FOBT+ results, and employing prep education processes that include both written and verbal information, but in order to increase the proportion of FOBT+ patients that receive follow-up colonoscopy within 60 days, it may be necessary to increase colonoscopy appointment availability. Given that the significant organizational-level predictors of follow-up rates we identified all had modest effects (i.e., resulting in at most 5–14% differences in follow-up rates), multifaceted strategies designed not only to increase colonoscopy follow-up for FOBT+ results, but also to reduce FOBT use in patients who would not complete colonoscopy follow-up due to contraindications or personal preference, may be needed to close remaining gaps. A fruitful area for future research would be the evaluation of such multifaceted strategies.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support: This research was funded by the Department of Veterans Affairs Health Services Research & Development (VA HSR&D) grant # IIR 08-334-2 (Partin), VA HSR&D Research Career Scientist award #RCS 10-185 (Partin), VA HSR&D Career Development Award (CDA) #CDA 07-016 (Haggstrom) VA HSR&D CDA #CDA 08-024 (Powell), VA Clinical Science Research & Development Career Development Award CDA-2 (Shaukat), and National Institute on Aging at the National Institutes of Health grant # K24AG041180 (Walter). The views expressed in this article are those of the authors and do not necessarily represent the position or policy of the Department of Veterans Affairs or the National Institutes of Health.

Reference List

1. American Cancer Society. Cancer Facts and Figures 2013. Atlanta: American Cancer Society; 2013.
2. Humphrey LL, Shannon J, Partin MR, O'Malley J, Chen Z, Helfand M. Improving the follow-up of positive hemocult screening tests: an electronic intervention. *J Gen Intern Med.* 2011 Jul; 26(7): 691–7. [PubMed: 21327529]
3. Levin B, Lieberman DA, McFarland B, Andrews KS, Brooks D, Bond J, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal

- Cancer, and the American College of Radiology. *Gastroenterology*. 2008 May; 134(5):1570–95. [PubMed: 18384785]
4. Rex DK, Johnson DA, Anderson JC, Schoenfeld PS, Burke CA, Inadomi JM. American College of Gastroenterology guidelines for colorectal cancer screening 2009 [corrected]. *Am J Gastroenterol*. 2009 Mar; 104(3):739–50. [PubMed: 19240699]
 5. USPSTF. Screening for colorectal cancer: U.S. Preventive Services Task Force recommendation statement. *Ann Intern Med*. 2008 Nov 4; 149(9):627–37. [PubMed: 18838716]
 6. Yabroff KR, Klabunde CN, Yuan G, McNeel TS, Brown ML, Casciotti D, et al. Are Physicians' Recommendations For Colorectal Cancer Screening Guideline-Consistent? *Journal of General Internal Medicine*. 2010 Oct 14; 26(2):177–84. [PubMed: 20949328]
 7. Levin TR, Jamieson L, Burley DA, Reyes J, Oehrli M, Caldwell C. Organized colorectal cancer screening in integrated health care systems. *Epidemiol Rev*. 2011 Jul; 33(1):101–10. [PubMed: 21709143]
 8. El Serag HB, Petersen L, Hampel H, Richardson P, Cooper G. The use of screening colonoscopy for patients cared for by the Department of Veterans Affairs. *Arch Intern Med*. 2006 Nov 13; 166(20):2202–8. [PubMed: 17101937]
 9. Hardcastle JD, Chamberlain JO, Robinson MH, Moss SM, Amar SS, Balfour TW, et al. Randomised controlled trial of faecal-occult-blood screening for colorectal cancer. *Lancet*. 1996 Nov 30; 348(9040):1472–7. [PubMed: 8942775]
 10. Kronborg O, Fenger C, Olsen J, Jorgensen OD, Sondergaard O. Randomised study of screening for colorectal cancer with faecal-occult-blood test. *Lancet*. 1996 Nov 30; 348(9040):1467–71. [PubMed: 8942774]
 11. Mandel JS, Bond JH, Church TR, Snover DC, Bradley GM, Schuman LM, et al. Reducing mortality from colorectal cancer by screening for fecal occult blood. Minnesota Colon Cancer Control Study. *N Engl J Med*. 1993 May 13; 328(19):1365–71. [PubMed: 8474513]
 12. Gellad ZF, Almirall D, Provenzale D, Fisher DA. Time from positive screening fecal occult blood test to colonoscopy and risk of neoplasia. *Dig Dis Sci*. 2009 Nov; 54(11):2497–502. [PubMed: 19093199]
 13. Larson MF, Ko CW, Dominitz JA. Effectiveness of a provider reminder on fecal occult blood test follow-up. *Dig Dis Sci*. 2009 Sep; 54(9):1991–6. [PubMed: 19255849]
 14. Paszat L, Rabeneck L, Kiefer L, Mai V, Ritvo P, Sullivan T. Endoscopic follow-up of positive fecal occult blood testing in the Ontario FOBT Project. *The Canadian Journal of Gastroenterology*. 2007 Jun; 21(6):379–82.
 15. Powell AA, Nugent S, Ordin DL, Noorbaloochi S, Partin MR. Evaluation of a VHA Collaborative to Improve Follow-up After a Positive Colorectal Cancer Screening Test. *Medical Care*. 2011; 49(41):897–903. [PubMed: 21642875]
 16. Leddin D, Armstrong D, Borgaonkar M, Bridges RJ, Fallone CA, Telford JJ, et al. The 2012 SAGE wait times program: Survey of Access to GastroEnterology in Canada. *Can J Gastroenterol*. 2013 Feb; 27(2):83–9. [PubMed: 23472243]
 17. Paterson WG, Depew WT, Pare P, Petrunia D, Switzer C, Veldhuyzen van Zanten SJ, et al. Canadian consensus on medically acceptable wait times for digestive health care. *Canadian Journal of Gastroenterology*. 2006 Jun; 20(6):411–23. [PubMed: 16779459]
 18. Veterans Health Administration. VHA Directive 2007–004: Colorectal Cancer Screening. Washington DC: Department of Veterans Affairs; 2007 Jan 12.
 19. Combined Assessment Program Summary Report: Evaluation of Colorectal Cancer Screening and Follow-Up in Veterans Health Administration Facilities. Department of Veterans Affairs Office of Inspector General; 2013.
 20. Zapka J, Taplin SH, Price RA, Cranos C, Yabroff R. Factors in quality care--the case of follow-up to abnormal cancer screening tests--problems in the steps and interfaces of care. *J Natl Cancer Inst Monogr*. 2010; 2010(40):58–71. [PubMed: 20386054]
 21. Zapka JM, Edwards HM, Chollette V, Taplin SH. Follow-up to Abnormal Cancer Screening Tests: Considering the Multilevel Context of Care. *Cancer Epidemiol Biomarkers Prev*. 2014 Oct; 23(10):1965–73. [PubMed: 25073625]

22. Choi KS, Lee HY, Jun JK, Shin A, Park EC. Adherence to follow-up after a positive fecal occult blood test in an organized colorectal cancer screening program in Korea, 2004–2008. *J Gastroenterol Hepatol*. 2012 Jun; 27(6):1070–7. [PubMed: 22004224]
23. Fisher DA, Zullig LL, Grambow SC, Abbott DH, Sandler RS, Fletcher RH, et al. Determinants of medical system delay in the diagnosis of colorectal cancer within the Veteran Affairs Health System. *Dig Dis Sci*. 2010 May; 55(5):1434–41. [PubMed: 20238248]
24. Miglioretti DL, Rutter CM, Bradford SC, Zauber AG, Kessler LG, Feuer EJ, et al. Improvement in the diagnostic evaluation of a positive fecal occult blood test in an integrated health care organization. *Med Care*. 2008 Sep; 46(9 Suppl 1):S91–S96. [PubMed: 18725839]
25. Rao SK, Schilling TF, Sequist TD. Challenges in the management of positive fecal occult blood tests. *J Gen Intern Med*. 2009 Mar; 24(3):356–60. [PubMed: 19130147]
26. Turner B, Myers RE, Hyslop T, Hauck WW, Weinberg D, Brigham T, et al. Physician and patient factors associated with ordering a colon evaluation after a positive fecal occult blood test. *J Gen Intern Med*. 2003 May; 18(5):357–63. [PubMed: 12795734]
27. Ishikawa Y, Zheng YF, Nishiuchi H, Suda T, Hasumi T, Saito H. Classification tree analysis to enhance targeting for follow-up exam of colorectal cancer screening. *BMC Cancer*. 2013; 13:470. [PubMed: 24112563]
28. Shields HM, Weiner MS, Henry DR, Lloyd JA, Ransil BJ, Lamphier DA, et al. Factors that influence the decision to do an adequate evaluation of a patient with a positive stool for occult blood. *Am J Gastroenterol*. 2001 Jan; 96(1):196–203. [PubMed: 11197252]
29. Lurie JD, Welch HG. Diagnostic testing following fecal occult blood screening in the elderly. *J Natl Cancer Inst*. 1999 Oct 6; 91(19):1641–6. [PubMed: 10511591]
30. Nadel MR, Berkowitz Z, Klabunde CN, Smith RA, Coughlin SS, White MC. Fecal Occult Blood Testing Beliefs and Practices of U.S. Primary Care Physicians: Serious Deviations from Evidence-Based Recommendations. *Journal of General Internal Medicine*. 2010 Apr 10.
31. Taplin SH, Clauser S, Rodgers AB, Breslau E, Rayson D. Interfaces across the cancer continuum offer opportunities to improve the process of care. *J Natl Cancer Inst Monogr*. 2010; 2010(40): 104–10. [PubMed: 20386059]
32. Clauser SB, Taplin SH, Foster MK, Fagan P, Kaluzny AD. Multilevel intervention research: lessons learned and pathways forward. *J Natl Cancer Inst Monogr*. 2012 May; 2012(44):127–33. [PubMed: 22623606]
33. Partin MR, Powell AA, Burgess DJ, Haggstrom D, Gravely AA, Halek K, et al. Adding Postal Follow-Up to a Web-Based Survey of Primary Care and Gastroenterology Clinic Physician Chiefs Improved Response Rates but not Response Quality or Representativeness. *Eval Health Prof*. 2014 published online 5 December 2013.
34. Donabedian, A. *Explorations in Quality Assessment and Monitoring*. Vol. I. Ann Arbor, MI: Health Administration Press; 1980.
35. Lukas C, Holmes S, Cohen A, Restuccia J, Cramer I, Shwartz M, et al. Transformational change in health care systems: An organizational model. *Health Care Management Review*. 2007; 32(4): 309–20. [PubMed: 18075440]
36. Anhang PR, Zapka J, Edwards H, Taplin SH. Organizational factors and the cancer screening process. *J Natl Cancer Inst Monogr*. 2010; 2010(40):38–57. [PubMed: 20386053]
37. Mandelblatt JS, Yabroff KR. Effectiveness of interventions designed to increase mammography use: a meta-analysis of provider-targeted strategies. *Cancer Epidemiol Biomarkers Prev*. 1999 Sep; 8(9):759–67. [PubMed: 10498394]
38. Myers RE, Turner B, Weinberg D, Hyslop T, Hauck WW, Brigham T, et al. Impact of a physician-oriented intervention on follow-up in colorectal cancer screening. *Prev Med*. 2004 Apr; 38(4):375–81. [PubMed: 15020170]
39. Engelstad LP, Stewart SL, Nguyen BH, Bedeian KL, Rubin MM, Pasick RJ, et al. Abnormal Pap smear follow-up in a high-risk population. *Cancer Epidemiol Biomarkers Prev*. 2001 Oct; 10(10): 1015–20. [PubMed: 11588126]
40. Yano EM, Soban LM, Parkerton PH, Etzioni DA. Primary care practice organization influences colorectal cancer screening performance. *Health Services Research*. 2007 Jun; 42(3 Pt 1):1130–49. [PubMed: 17489907]

41. Powell AA, Gravelly AA, Ordin DL, Schlosser JE, Partin MR. Timely follow-up of positive fecal occult blood tests strategies associated with improvement. *Am J Prev Med.* 2009 Aug; 37(2):87–93. [PubMed: 19524390]
42. Winawer SJ, Zauber AG, Fletcher RH, Stillman JS, O'Brien MJ, Levin B, et al. Guidelines for colonoscopy surveillance after polypectomy: a consensus update by the US Multi-Society Task Force on Colorectal Cancer and the American Cancer Society. *Gastroenterology.* 2006; 130:1872–85. [PubMed: 16697750]
43. Hysong SJ. Meta-analysis: audit and feedback features impact effectiveness on care quality. *Med Care.* 2009 Mar; 47(3):356–63. [PubMed: 19194332]
44. Ivers N, Jamtvedt G, Flottorp S, Young JM, Odgaard-Jensen J, French SD, et al. Audit and feedback: effects on professional practice and healthcare outcomes. *Cochrane Database Syst Rev.* 2012; 6:CD000259. [PubMed: 22696318]
45. Abuksis G, Mor M, Segal N, Shemesh I, Morad I, Plaut S, et al. A patient education program is cost-effective for preventing failure of endoscopic procedures in a gastroenterology department. *Am J Gastroenterol.* 2001 Jun; 96(6):1786–90. [PubMed: 11419830]
46. Denberg TD, Coombes JM, Byers TE, Marcus AC, Feinberg LE, Steiner JF, et al. Effect of a mailed brochure on appointment-keeping for screening colonoscopy: a randomized trial. *Ann Intern Med.* 2006 Dec 19; 145(12):895–900. [PubMed: 17179058]
47. Gurudu SR, Fry LC, Fleischer DE, Jones BH, Trunkenbolz MR, Leighton JA. Factors contributing to patient nonattendance at open-access endoscopy. *Dig Dis Sci.* 2006 Nov; 51(11):1942–5. [PubMed: 17009114]
48. Lee CS, McCormick PA. Telephone reminders to reduce non-attendance rate for endoscopy. *J R Soc Med.* 2003 Nov; 96(11):547–8. [PubMed: 14594962]




Figure 1.
Subject Flowchart

Table 1
Question wording, response categories, and coding for analysis of Organizational Structure and Process measures.

Measure	Source	Question Wording	Coding for analysis														
Primary Care Leadership support	Primary Care Survey	<p>Please rate each of the following potential barriers to providing timely follow-up for positive FOBTs, where 1 is not a barrier and 5 is a key barrier to providing timely follow-up for positive FOBTs.</p> <p>Not a barrier</p> <table border="1"> <tr> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>Key barrier</td> <td>Don't Know</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>Not a priority to leadership</p>	1	2	3	4	5	Key barrier	Don't Know	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	1="key barrier" (category 5) 0=all other responses
1	2	3	4	5	Key barrier	Don't Know											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											
Gastroenterology Leadership support	Gastroenterology Survey	<p>Please rate each of the following potential barriers to providing timely follow-up for positive FOBTs, where 1 is not a barrier and 5 is a key barrier to providing timely follow-up for positive FOBTs.</p> <p>Not a barrier</p> <table border="1"> <tr> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>Key barrier</td> <td>Don't Know</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>Not a priority to leadership</p>	1	2	3	4	5	Key barrier	Don't Know	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	1="key barrier" (category 5) 0=all other responses
1	2	3	4	5	Key barrier	Don't Know											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											
Resource alignment: Tracking	Primary Care Survey	<p>Has your PRIMARY CARE PROGRAM assigned anyone the responsibility of tracking what happens to patients with positive FOBT results? (Select One)</p> <p><input type="checkbox"/> Yes</p> <p>Approximately how frequently does this person/s check these outcomes? (Select One)</p> <p><input type="checkbox"/> As needed on an ongoing basis</p> <p><input type="checkbox"/> Weekly</p> <p><input type="checkbox"/> Monthly</p> <p><input type="checkbox"/> Quarterly</p> <p><input type="checkbox"/> Don't know</p> <p><input type="checkbox"/> No</p> <p><input type="checkbox"/> Don't know</p>	1= someone is assigned to track "weekly or more frequently" 0=all other responses														
Resource alignment: Colonoscopy appointment availability	Gastroenterology Survey	<p>Please rate each of the following potential barriers to providing timely follow-up for positive FOBTs.</p> <p>Not a barrier</p> <table border="1"> <tr> <td>1</td> <td>2</td> <td>3</td> <td>4</td> <td>5</td> <td>Key barrier</td> <td>Don't Know</td> </tr> <tr> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> <td><input type="checkbox"/></td> </tr> </table> <p>Limited availability of colonoscopy appointments</p>	1	2	3	4	5	Key barrier	Don't Know	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	1="key barrier" (category 5) 0=all other responses
1	2	3	4	5	Key barrier	Don't Know											
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>											
Primary Care Feedback	Primary Care Survey	<p>Do primary care providers receive feedback about the amount of time it takes them to refer patients with positive FOBT results for colonoscopy?</p> <p><input type="checkbox"/> Yes</p> <p><input type="checkbox"/> No</p> <p><input type="checkbox"/> Don't know</p> <p>How are primary care providers given this feedback? (Check all that apply)</p>	0=no feedback 1=aggregate feedback at any frequency 2=individual infrequent feedback (less frequently than monthly)														

Measure	Source	Question Wording	Coding for analysis
Gastroenterology Feedback	Gastroenterology survey	<p><input type="checkbox"/> Individualized information at provider level</p> <p><input type="checkbox"/> Aggregate information at the team or clinic level</p> <p><input type="checkbox"/> Aggregate information at the facility level</p> <p><input type="checkbox"/> Other (specify):</p> <p><input type="checkbox"/> Don't know</p> <p>How frequently are primary care providers given this feedback? (Select One)</p> <p><input type="checkbox"/> Weekly</p> <p><input type="checkbox"/> Monthly</p> <p><input type="checkbox"/> Quarterly</p> <p><input type="checkbox"/> Annually</p> <p><input type="checkbox"/> Other (specify):</p> <p><input type="checkbox"/> Don't know</p>	3=individual frequent feedback (at least monthly)
Gastroenterology Feedback	Gastroenterology survey	<p>How is feedback on the timeliness of follow-up for patients with positive FOBT results given to GI/Endoscopy staff? (Check all that apply)</p> <p><input type="checkbox"/> Verbally in a staff meeting</p> <p><input type="checkbox"/> In writing in the form of an aggregate team, clinic or facility report</p> <p><input type="checkbox"/> Other (specify):</p> <p><input type="checkbox"/> No feedback provided to GI/Endoscopy staff on this aspect of care</p> <p><input type="checkbox"/> Don't know</p> <p>Approximately how frequently are GI/Endoscopy staff given this feedback? (Select One)</p> <p><input type="checkbox"/> Weekly</p> <p><input type="checkbox"/> Monthly</p> <p><input type="checkbox"/> Quarterly</p> <p><input type="checkbox"/> Annually</p> <p><input type="checkbox"/> Don't know</p>	0=written less than monthly 1=written feedback at least monthly 2=verbal feedback less than monthly 3=verbal feedback at least monthly (hypothesized as the optimal category, based on prior research) (43-44)
Primary Care Incentives	Primary Care Survey	<p>Which of the following do providers in your primary care program receive from leadership for their performance on assuring TIMELY FOLLOW-UP OF POSITIVE FOBT RESULTS? (Check all that apply)</p> <p><input type="checkbox"/> Recognition for good performance</p> <p><input type="checkbox"/> Monetary rewards for good performance</p> <p><input type="checkbox"/> Counseling or reprimands for poor performance</p> <p><input type="checkbox"/> None of the above</p> <p><input type="checkbox"/> Don't know</p>	0=no rewards or reprimands 1=Recognition only 2=Monetary rewards (alone or in combination with other incentives) 3=Reprimands (alone or in combination with recognition)
Gastroenterology Incentives	Gastroenterology Survey	<p>Which of the following rewards or reprimands do staff in your GI/Endoscopy program receive for their performance on assuring timely follow-up of positive FOBT results? (Check all that apply)</p> <p><input type="checkbox"/> Recognition from leadership for good performance</p> <p><input type="checkbox"/> Monetary rewards for good performance</p> <p><input type="checkbox"/> Counseling or reprimands for poor performance</p> <p><input type="checkbox"/> Other (specify):</p> <p><input type="checkbox"/> None of the above</p> <p><input type="checkbox"/> Don't know</p>	0=no rewards or reprimands 1=Recognition only 2=Monetary rewards (alone or in combination with other incentives) 3=Reprimands (alone or in combination with recognition)
Demand Efficiency: Number of contraindications on screening reminder	Primary Care Survey	<p>Does your facility's colorectal cancer screening clinical reminder ask about contraindications?</p> <p><input type="checkbox"/> Yes</p> <p><input type="checkbox"/> No</p> <p><input type="checkbox"/> Don't know</p> <p>Which of the following contraindications are asked about in the reminder? (Check all that apply)</p>	0=none 1=1-2 2=3 3=4 or more

Measure	Source	Question Wording	Coding for analysis
Demand Efficiency: Does colonoscopy consult template include contraindications?	Primary Care Survey	<ul style="list-style-type: none"> <input type="checkbox"/> Life limiting comorbidities/limited life expectancy <input type="checkbox"/> Health issues that increase risk of complications of colonoscopy <input type="checkbox"/> Recent colonoscopy <input type="checkbox"/> Patient not willing to undergo colonoscopy if screen is positive <input type="checkbox"/> Other (specify) <p>Does your Primary Care program use a consult template for referrals to Gastroenterology for follow-up of positive FOBT results?</p> <ul style="list-style-type: none"> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know <p>Which of the following items are included in the template? (Check all that apply)</p> <ul style="list-style-type: none"> <input type="checkbox"/> Anticoagulant use <input type="checkbox"/> Anti-platelet use <input type="checkbox"/> Diabetic <input type="checkbox"/> Anemia/Iron deficiency <input type="checkbox"/> Life expectancy or comorbidities related to life expectancy <input type="checkbox"/> Previous colonoscopy results <input type="checkbox"/> Physical/cognitive impairments that would make difficult to follow prep instructions <input type="checkbox"/> Lab values <input type="checkbox"/> Other (specify): <input type="checkbox"/> None <input type="checkbox"/> Don't know 	0=some contraindications 1=indications only 2=neither
Demand Efficiency: Guideline-concordant surveillance	Gastroenterology Survey	How soon is a repeat colonoscopy typically scheduled at your facility for the following types of patients: <input type="checkbox"/> 1-2 small adenomas (<1cm) on last colonoscopy: _____years	0=not guideline-concordant (<5 years) 1=5 years 2=7-10 years
Supply Efficiency: How Gastroenterology providers are notified of FOBT+	Gastroenterology Survey	How is the GI/Endoscopy clinic first notified of an FOBT positive result? (Check all that apply) <input type="checkbox"/> Lab sends notification directly to GI/Endoscopy <input type="checkbox"/> Primary care notifies using consult template/referral <input type="checkbox"/> Other(specify):(n=5 said Gastroenterology identifies FOBT+) <input type="checkbox"/> Don't know	0=Primary care notifies 1=Lab notifies 2=Gastroenterology notifies
Supply Efficiency: Does prep instruction require a separate appointment?	Gastroenterology Survey	What is the most typical way that patients receive their colonoscopy preparation instructions at your facility? (Select One) <input type="checkbox"/> Written instructions provided with prep kit <input type="checkbox"/> Written instructions mailed separate from prep kit <input type="checkbox"/> Verbal instructions provided over the phone <input type="checkbox"/> Individual appointment <input type="checkbox"/> Group appointment <input type="checkbox"/> Other (specify): <input type="checkbox"/> Don't know	0=individual or group appointment (including those indicated in specified "other" responses) 1=all other categories
Supply Efficiency: Pre-procedure appointment required?	Gastroenterology Survey	Do you require a pre-procedure clinic appointment prior to the colonoscopy appointment? <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know	0=Yes 1=No

Measure	Source	Question Wording	Coding for analysis
Supply Efficiency: Overbooking is used	Gastroenterology Survey	In the past six months, did your GI program implement any of the following processes to keep up with colonoscopy demand? (Check all that apply) <ul style="list-style-type: none"> <input type="checkbox"/> Added additional clinic hours <input type="checkbox"/> Implemented make up clinics <input type="checkbox"/> Added additional staff <input type="checkbox"/> Increased use of contract providers <input type="checkbox"/> Increased use of fee basis providers <input type="checkbox"/> Overbooked appointments <input type="checkbox"/> Other (specify): <input type="checkbox"/> No change implemented <input type="checkbox"/> Don't know 	1=Overbooking appointments is used 0=all other responses
Patient-Centered Process: Patient notification of FOBT+ includes phone contact	Primary Care Survey	How are patients seen in your primary care program typically first notified of a positive FOBT result? (Select One) <ul style="list-style-type: none"> <input type="checkbox"/> Letter from primary care clinic <input type="checkbox"/> Letter from GI clinic <input type="checkbox"/> Phone call from primary care or GI clerk <input type="checkbox"/> Phone call from primary care or GI nurse <input type="checkbox"/> Phone call from primary care or GI provider <input type="checkbox"/> Email or secure messaging <input type="checkbox"/> Other (specify): <input type="checkbox"/> Don't know 	0=notification does not include phone contact 1=notification includes some phone contact
Patient-Centered Process: Are colonoscopy appointment times negotiated with patients?	Gastroenterology Survey	Which of the following options best describes the most common way that patients are scheduled for the first appointment to follow-up on a positive FOBT at your facility? (Select One) <ul style="list-style-type: none"> <input type="checkbox"/> Patient is sent a letter instructing them to come to the clinic at a specific date/time <input type="checkbox"/> Patient is first sent a letter instructing them to call the clinic, and then an appointment date/time is negotiated over the phone <input type="checkbox"/> A scheduler calls the patient to arrange an appointment date/time <input type="checkbox"/> Other (specify): <input type="checkbox"/> Don't know 	1=Second and third response options, and other responses that include phone contact with patient 0=All other responses
Patient-Centered Process: Do prep instruction methods include opportunities for questions?	Gastroenterology Survey	What is the most typical way that patients receive their colonoscopy preparation instructions at your facility? (Select One) <ul style="list-style-type: none"> <input type="checkbox"/> Written instructions provided with prep kit <input type="checkbox"/> Written instructions mailed separate from prep kit <input type="checkbox"/> Verbal instructions provided over the phone <input type="checkbox"/> Individual appointment <input type="checkbox"/> Group appointment <input type="checkbox"/> Other (specify): <input type="checkbox"/> Don't know 	0=Written only 1=Verbal phone and individual appointment 2=Group appointment and "other" responses that included written and verbal combinations
Patient-Centered Process: Do appointment reminders review prep instruction?	Gastroenterology Survey	Are preparation procedures reviewed as part of the [colonoscopy appointment] reminder? <ul style="list-style-type: none"> <input type="checkbox"/> Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know 	0=No reminders or reminders do not review prep 1=Reminders review prep

Table 2

Characteristics of patients included in the analysis (N=74,104)

Characteristic	N	%
Race/ethnicity		
Native American	638	1
African American	12,866	17
Asian/Pacific Islander	1,058	1
White, Non-Hispanic	48,018	65
Hispanic	4,453	6
Unknown	7,071	10
Marital Status		
Married	38,887	53
Widowed	4,571	6
All Others	30,391	41
Gender		
Male	71,067	96
Female	3,037	4
Age		
< 50	3,352	5
50 – 64	40,513	55
65 – 84	27,725	37
85	2,514	3
Residence		
Urban	38,713	52
Rural/Highly Rural	35,118	48
Drive time to VA Medical Center		
> 60 minutes	28,661	39
30–60 minutes	17,694	24
6–29 minutes	25,094	34

Characteristic	N	%
<6 minutes	2,387	3
Polyps or benign neoplasms	14,770	20
Mental health diagnoses		
no mental health diagnosis	23,958	32
psychiatric only	19,383	26
substance abuse only	11,341	15
both psychiatric and substance abuse	19,422	26
Charlson Comorbidity Score (mean and range)	1.9 (0–19)	
FOBT ordering facility is a VHA hospital	42,627	58
FOBT was ordered by patient's Primary Care Provider	51,813	70
FOBT ordering provider type		
Physician	53,661	72
Nurse Practitioner, Physician Assistant	16,203	22
Resident	2,686	4
Other	1,535	2

Table 3

Distribution of facilities and patients across organizational structures and processes examined

Characteristic	Facilities (N=98) ¹		Patients (N=74,104)	
	N	%	N	%
ORGANIZATIONAL STRUCTURES				
Leadership support ^{GI}				
Primary Care leadership support ^b Is a key barrier	0	0	0	0
Is not a key barrier	75	100	56,897	100
Missing	23		17,207	
Gastroenterology leadership support ^c - Is a key barrier	2	2	671	1
Is not a key barrier	78	98	61,171	99
Missing	18		12,262	
Resources				
Colonoscopy appointment availability ^c - Is a key barrier	38	49	37,658	63
Is not a key barrier	39	51	22,021	37
Missing	21		17,207	
Tracking ^b > weekly	15	20	13,340	21
At least weekly	27	36	26,513	42
No tracking	33	44	23,429	37
Missing	23		10,822	
Feedback				
Primary Care feedback ^b - Aggregate	5	6	2,655	4
None	47	62	34,001	59
Individual, infrequent	8	11	6,520	11
Individual, frequent	16	21	14,394	25
Missing	22		16,534	
Gastroenterology feedback ^c - None	22	27	15,014	24
Written, infrequent	18	22	14,720	23
Verbal, infrequent	12	15	12,146	19
Verbal, frequent	18	22	13,432	21
Written, frequent	11	14	7,970	13
Missing	17		10,822	
Incentives				
Primary Care incentives ^b - None	42	56	28,439	50
Recognition only	6	8	4,789	8
Monetary reward (alone or in combination with other strategies)	6	8	4,452	8
Reprimand (alone or in combination with recognition)	21	28	19,217	34
Missing	23		17,207	

Characteristic	Facilities (N=98) ¹		Patients (N=74,104)	
	N	%	N	%
Gastroenterology incentives^c – None	52	64	37,204	59
Recognition only	15	19	12,893	20
Monetary reward (alone or in combination with other strategies)	7	9	6,257	10
Reprimand (alone or in combination with recognition)	7	9	6,599	10
Missing	17		10,822	
ORGANIZATIONAL PROCESSES				
Demand Efficiency Processes				
Contraindications on colorectal Cancer Screening Reminder^b – None	39	52	29,996	53
1–2	21	28	14,327	25
3	7	9	4,669	8
4+	8	11	7,905	14
Missing	23		17,207	
Information on colonoscopy consult^b – Contraindications	35	47	29,167	51
Indication	22	29	16,781	29
Neither	18	24	10,949	19
Missing	23		17,207	
Surveillance for 1–2 Adenomas <1 cm^c – <5 years	6	8	2,519	4
5 years	67	84	56,645	90
7–10 years	7	9	3,693	6
Missing	18		11,247	
Supply Efficiency Processes				
Gastroenterology notification^c – by primary care	55	70	42,829	69
by Lab	19	24	14,005	22
by Gastroenterology	5	6	5,512	9
Missing	19		11,758	
Colonoscopy prep instruction^c – separate appointment required	24	30	21,079	33
Separate appointment not required	57	70	42,203	67
Missing	17		10,822	
Pre-op Appointment^c – required	31	38	28,060	44
Not required	50	62	35,222	56
Missing	17		10,822	
Overbooking^c – is used to meet colonoscopy demand	43	54	38,087	60
Is not used	24	30	17,427	28
No changes to meet colonoscopy demand	13	16	7,494	12
Missing	18		11,096	
Patient-centered Processes				
Patients notification^b – Written contact only	25	33	21,649	38

Characteristic	Facilities (N=98) ¹		Patients (N=74,104)	
	N	%	N	%
Some phone contact	50	67	35,248	62
Missing	23		17,207	
Colonoscopy Scheduling ^c – call from scheduler	25	31	19,560	31
Letter requesting patient call for appointment	11	14	7,887	12
Letter with assigned appointment	28	35	23,895	38
Other	17	21	11,940	19
Missing	17		10,822	
Colonoscopy prep instruction ^c – Verbal phone and individual appointment	19	23	15,965	25
Written only	36	44	23,442	37
Verbal Group or other combined verbal/written method	16	32	23,875	38
Missing	17		10,822	
Appointment reminders ^c – do not review prep/no reminder	25	31	24,385	39
Do review prep	56	69	38,897	61
Missing	17		10,822	

^a Distribution from 81 facilities responding to Gastroenterology Survey, or 76 responding to Primary care survey, depending on measure (98 facilities responded to one or both of the surveys).

^b Primary Care survey item

^c Gastroenterology survey item

Table 4

Odds ratios, 95% confidence intervals, and follow-up percentage estimates for organizational structures and processes derived from the original and sensitivity-adjusted^a multivariable regression models for 60-day follow-up rate outcome measure.^{b,c}

Characteristic	Original Multivariable estimates		Sensitivity-adjusted ^a Multivariable estimates.	
	OR (95% CI)	Follow-up % (95% CI)	OR (95% CI)	Follow-up % (95% CI)
ORGANIZATIONAL STRUCTURES				
Resources				
Colonoscopy appointment availability - Is a key barrier	1.00	26 (23–30)	1.00	48 (43–52)
Is not a key barrier	1.43 (1.09–1.90)	34 (30–38)	1.48 (1.14–1.92)	58 (53–62)
Feedback				
Primary Care feedback - Aggregate	1.00	24 (16–34)	1.00	48 (37–60)
None	1.09 (0.65–1.82)	26 (22–29)	0.96 (0.59–1.58)	47 (43–51)
Individual, infrequent	1.52 (0.85–2.71)	32 (25–40)	1.32 (0.75–2.31)	55 (47–63)
Individual, frequent	1.79 (1.02–3.16)	36 (30–43)	1.71 (0.99–2.97)	61 (54–68)
ORGANIZATIONAL PROCESSES				
Demand Efficiency Processes				
Information on colonoscopy consult - Contraindications	1.00	25 (22–29)	1.00	48 (44–53)
Indication	1.49 (1.10–2.02)	33 (28–40)	1.30 (0.96–1.74)	55 (49–61)
Neither	1.48 (1.07–2.05)	33 (27–40)	1.19 (0.86–1.63)	53 (46–60)
Supply Efficiency Processes				
Gastroenterology notification – by Primary Care	1.00	26 (24–30)	1.00	48 (45–52)
by lab	1.36 (0.97–1.90)	33 (26–40)	1.24 (0.89–1.73)	54 (46–61)
by Gastroenterology	1.85 (1.17–2.91)	40 (30–50)	1.82 (1.17–2.83)	63 (53–72)

^aExcluding patients who may not have been appropriate for colorectal cancer screening (age <45 or >85, documentation of limited life expectancy in the medical record, or colonoscopy in the prior 10 years), and treating patients who refused colonoscopy or chose to pursue colonoscopy in the private sector as having adequate follow-up.

^bBold odds ratios are significant at $p < 0.05$

^cOdds ratios for the individual-level factors controlled for in the 60-day model (age, race, residence, drive time to the nearest VHA specialty care facility, personal history of colorectal polyps or benign neoplasms, Charlson comorbidity score for the 1 year prior to the FOBT+ result, mental health diagnoses, whether the FOBT was ordered by their primary care provider, and what type of facility ordered their FOBT) are provided in Supplementary Table 2.

Table 5

Odds ratios, follow-up percentage estimates, and 95% confidence intervals for organizational processes significantly associated with 6-month follow-up rates in original and sensitivity-adjusted^a multivariable logistic regression models.^{b,c}

Characteristic	Original Multivariable estimates		Sensitivity-Adjusted ^a Multivariable estimates	
	OR (95% CI)	Follow-up % (95% CI)	OR (95% CI)	Follow-up % (95% CI)
Demand Efficiency Processes				
Surveillance for 1–2 Adenomas <1 cm – <5 years	1.00	42 (36–48)	1.00	70 (63–76)
5 years	1.32 (1.02–1.71)	49 (47–50)	1.14 (0.80–1.62)	73 (71–75)
7–10 years	1.57 (1.11–2.20)	53 (47–58)	1.33 (0.84–2.12)	76 (70–82)
Supply Efficiency Processes				
Gastroenterology notification - by primary care	1.00	47 (45–49)	1.00	72 (69–74)
by lab	1.25 (1.06–1.47)	52 (49–56)	1.18 (0.94–1.48)	75 (71–78)
by Gastroenterology	1.31 (0.99–1.73)	53 (47–60)	1.43 (0.98–2.10)	78 (72–84)
Patient-Centered Processes				
Colonoscopy prep instruction – verbal phone or individual appointment	1.00	43 (40–47)	1.00	68 (64–72)
Written only	1.18 (1.00–1.40)	48 (45–50)	1.22 (0.97–1.53)	72 (69–75)
Verbal group appointment or other combined verbal/written method	1.48 (1.22–1.79)	53 (50–56)	1.50 (1.16–1.95)	76 (73–79)

^aExcluding patients who may not have been appropriate for colorectal cancer screening (age <45 or >85, documentation of limited life expectancy in the medical record, or colonoscopy in the prior 10 years), and treating patients who refused colonoscopy or chose to pursue colonoscopy in the private sector as having adequate follow-up.

^bBold odds ratios are significant at $p < 0.05$

^cOdds ratios for the individual-level factors controlled for in the 6-month model (age, race, residence, drive time to the nearest VHA specialty care facility, personal history of colorectal polyps or benign neoplasms, Charlson comorbidity score for the 1 year prior to the FOBT+ result, mental health diagnoses, whether the FOBT was ordered by their primary care provider, what type of provider ordered their FOBT, and what type of facility ordered their FOBT) are provided in Supplementary Table 2.