

UC San Diego

UC San Diego Previously Published Works

Title

Gene Expression Signature in Adipose Tissue of Acromegaly Patients

Permalink

<https://escholarship.org/uc/item/5203g4jt>

Journal

PLOS ONE, 10(6)

ISSN

1932-6203

Authors

Hochberg, Irit
Tran, Quynh T
Barkan, Ariel L
et al.

Publication Date

2015

DOI

10.1371/journal.pone.0129359

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RESEARCH ARTICLE

Gene Expression Signature in Adipose Tissue of Acromegaly Patients

Irit Hochberg^{1,2*}, Quynh T. Tran³, Ariel L. Barkan^{4,5}, Alan R. Saltiel^{2,4}, William F. Chandler⁵, Dave Bridges^{6,7,2*}

1 Institute of Endocrinology, Diabetes and Metabolism, Rambam Health Care Campus, Haifa, Israel, **2** Life Sciences Institute, University of Michigan, Ann Arbor, MI, United States of America, **3** Department of Preventive Medicine, University of Tennessee Health Science Center, Memphis, TN, United States of America, **4** Department of Internal Medicine, University of Michigan, Ann Arbor, MI, United States of America, **5** Department of Neurosurgery, University of Tennessee Health Science Center, Memphis, TN, United States of America, **6** Department of Physiology, University of Tennessee Health Science Center, Memphis, TN, United States of America, **7** Children's Foundation Research Institute, Le Bonheur Children's Hospital, Memphis, TN, United States of America

* i_hochberg@rambam.health.gov.il (IH); dbridge9@uthsc.edu (DB)



OPEN ACCESS

Citation: Hochberg I, Tran QT, Barkan AL, Saltiel AR, Chandler WF, Bridges D (2015) Gene Expression Signature in Adipose Tissue of Acromegaly Patients. PLoS ONE 10(6): e0129359. doi:10.1371/journal.pone.0129359

Academic Editor: Martin Young, University of Alabama at Birmingham, UNITED STATES

Received: December 25, 2014

Accepted: May 7, 2015

Published: June 18, 2015

Copyright: © 2015 Hochberg et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was supported by Motor City Golf Classic (MCGC) Grant # G010640. This work utilized Metabolomics Core Services supported by grant U24 DK097153 of the NIH Common Fund to the University of Michigan. This work was also supported by a Le Bonheur Grant # 650 700 to DB.

Competing Interests: The authors have declared that no competing interests exist.

Abstract

To study the effect of chronic excess growth hormone on adipose tissue, we performed RNA sequencing in adipose tissue biopsies from patients with acromegaly (n = 7) or non-functioning pituitary adenomas (n = 11). The patients underwent clinical and metabolic profiling including assessment of HOMA-IR. Explants of adipose tissue were assayed *ex vivo* for lipolysis and ceramide levels. Patients with acromegaly had higher glucose, higher insulin levels and higher HOMA-IR score. We observed several previously reported transcriptional changes (*IGF1*, *IGFBP3*, *CISH*, *SOCS2*) that are known to be induced by GH/IGF-1 in liver but are also induced in adipose tissue. We also identified several novel transcriptional changes, some of which may be important for GH/IGF responses (*PTPN3* and *PTPN4*) and the effects of acromegaly on growth and proliferation. Several differentially expressed transcripts may be important in GH/IGF-1-induced metabolic changes. Specifically, induction of *LPL*, *ABHD5*, and *NRIP1* can contribute to enhanced lipolysis and may explain the elevated adipose tissue lipolysis in acromegalic patients. Higher expression of *TCF7L2* and the fatty acid desaturases *FADS1*, *FADS2* and *SCD* could contribute to insulin resistance. Ceramides were not different between the two groups. In summary, we have identified the acromegaly gene expression signature in human adipose tissue. The significance of altered expression of specific transcripts will enhance our understanding of the metabolic and proliferative changes associated with acromegaly.

Introduction

Acromegaly, i.e. excessive growth hormone (GH) production secondary to a pituitary adenoma, is a rare condition with an annual incidence of 3 patients per million [1]. The excess GH has important metabolic effects; the two most significant effects of GH on metabolism in

adipose tissue are insulin resistance and lipolysis [2]. Insulin resistance, presenting as diabetes or impaired glucose tolerance, is found in most acromegalic patients [3], and contributes to the enhanced morbidity [4]. Growth hormone induces the expression and secretion of IGF-1, so phenotypes associated with acromegaly may be due to either GH signaling, IGF-1 signaling or a combination of both [5,6].

There are few studies addressing the effect of GH specifically on the subcutaneous adipose tissue. Induction of STAT5 tyrosine phosphorylation and IGF1 mRNA expression has been detected in human subcutaneous adipose tissue biopsies taken after acute GH administration [7]. Subcutaneous adipocytes extracted from acromegalic patients are insulin resistant *ex vivo*, and after a glucose tolerance test there was 50% less insulin binding to its receptor and markedly decreased insulin-related anti-lipolytic activity [8]. *In vivo* measurement in humans detected GH-induced lipolysis in subcutaneous adipose tissue [9]. Pharmacologic inhibition of lipolysis reduced GH-induced insulin resistance, suggesting that some of this resistance is dependent on higher abundance of free fatty acids [10]. Microarray of gene expression has been published for subcutaneous adipose tissue biopsies before and after one year of GH treatment in GH deficient patients [11].

To study the effects of chronic excess GH, we used unbiased RNA sequencing in adipose tissue from acromegaly patients and controls. We found a distinctive pattern of changes in many transcripts that are highly associated with acromegaly. Many of these alterations may contribute to the metabolic effect of GH and reveal novel mechanisms of GH-induced insulin resistance and lipolysis in adipose tissue.

Materials and Methods

Patient Recruitment

The study was approved by the institutional review board of the University of Michigan Medical System. Written informed consent was obtained from all patients. Patients were recruited consecutively from a cohort undergoing transsphenoidal adenomectomy at the University of Michigan Medical Center for acromegaly or non-functioning pituitary adenoma over a 12 month period. All but one patient were newly diagnosed, none had previous surgery and only one previously diagnosed patient was treated with a somatostatin analog but IGF1 was still high without remission. None of the patients were on insulin, but one patient from each group was treated with metformin. Two patients with non-secreting adenomas were treated with beta blockers. Exclusion criteria were age <18 years old, current hormone treatment including glucocorticoids, malignancy, inflammatory disease, diabetes type 1 and established pituitary hormone deficiencies. For each patient, a data sheet was completed including, age, sex, anthropometric measurements, diagnosis of hypertension, diabetes, results of blood tests and medications. Fasting blood samples were assayed for glucose (Siemens Advia 1800) and insulin (Life Technologies) as instructed by the manufacturers.

Subcutaneous Fat Biopsy

During the course of pituitary surgery a routine subcutaneous fat graft is utilized to seal the surgical field upon completion of the procedure. A total of 500 mg of this fat graft was used for the study. ~200 mg were utilized for *ex vivo* lipolysis assay, ~300 mg was snap frozen in liquid nitrogen and stored at -80 degrees for RNA preparation and ceramide analysis.

Ex vivo Lipolysis

Twenty five mg pieces of adipose tissue were pre-incubated for 15 minutes in KRBH buffer (sigma) at 37°C and then incubated for 1 hour at 37°C in 300 µl KRBH in the presence or absence of isoproterenol 30nM in duplicate. Glycerol was assayed in supernatants using a glycerol assay kit (sigma) as instructed by manufacturer.

Ceramide Determinations

Ceramide analysis of tissue samples was performed by liquid chromatography-triple quadrupole mass spectrometry (LC-QQQ) according to a modified version of the protocol reported in [12]. Briefly, frozen tissue samples were pulverized under liquid nitrogen, then 20 mg portions were extracted using 1.6 mL of a 2:1:0.8 mixture of chloroform:methanol:water containing internal standards (50 ng each of C17 and C25 ceramide and C12 glucosylceramide per sample) [13]. The organic layer of the extract was dried under nitrogen gas and reconstituted in 100 µL of 60:40 acetonitrile: isopropanol. The re-constituted extract was analyzed by electrospray ionization LC-MS/MS on an Agilent (Santa Clara, CA) 6410 triple quadrupole instrument operating in positive ion multiple reaction monitoring mode. The LC column used was a Waters (Milford, MA) Xbridge C18 2.5 µ, 50 mm x 2.1 mm i.d. Mobile phase A was 5mM ammonium acetate, adjusted to pH 9.9 with ammonium hydroxide; mobile phase B was 60:40 acetonitrile:isopropanol. The gradient consisted of a linear ramp from 50 to 100%B over 5 minutes, a 20 minute hold at 100%B, and re-equilibration at 50%B for 10 minutes. Injection volume was 25 µL. Ceramides and glucosylceramides were identified by retention time and by MS/MS fragmentation parameters, and were quantitated by peak area relative to the closest-matching internal standard using Agilent MassHunter Quantitative Analysis software.

Transcriptomic Analysis

Total RNA was extracted from adipose tissue using the RNEasy kit (Qiagen) and its quality was verified using the Agilent 2100 Bioanalyzer (Agilent Technologies). At the University of Michigan DNA Sequencing Core, cDNA libraries from polyA mRNA were prepared using TruSeq cDNA synthesis kit and sequenced using a HiSeq 2000 (Illumina). Samples were run on 2 lanes of a HiSeq 2000 (Illumina) generating 8 612 682 to 16 469 501 single-ended 50 bp reads per sample. These were aligned to the human genome (Ensembl GRCh37.74, Genbank Assembly ID GCA_000001405.14) using TopHat version 2.0.10 [14], Bowtie 2 version 2.1.0 [15] and Samtools version 0.1.18. Reads were mapped to known genes using HTseq [16]. Gene expression was analyzed using DESeq2 version 1.2.10 [17]. To account for potential age-dependent changes in the subjects, we separated the patients into two groups, based on the median value, under 60 years of age versus 60 and above as has been previously reported for acromegaly studies [18]. We provide here both the non-age adjusted (S1 Table) and age-adjusted gene expression changes (S2 Table).

We then added this age group as a covariate along with the disease state. We tested for interactions between the age group and the disease state for each gene and did not identify any interaction term after adjusting for multiple observations ($q < 0.05$). All fold changes provided in this manuscript are age-adjusted fold change values calculated from this regression.

We used Gene Set Enrichment Analysis (GSEA v2.0.13 [19,20]) to determine whether our rank-ordered gene list for the comparison of acromegaly vs control patients is enriched in genes from gene ontology, KEGG, transcription factor or microRNA target gene sets. The gene list was ranked based on the shrunken log based 2 fold change and the statistical significance of the enrichment score was determined by performing 1000 phenotype permutations and setting

Table 1. Clinical characteristics.

	Control (n = 11)		Acromegaly (n = 9)		p
Age (years)	63.4	+/- 2.7	48.3	+/- 4.9	0.011
Height (cm)	170.0	+/- 2.4	180.1	+/- 4.0	0.036
Weight (kg)	89.4	+/- 6.7	103.9	+/- 9.3	0.21
BMI (kg/m ²)	30.7	+/- 1.8	31.7	+/- 2.1	0.69
Abdominal Circumference (cm)	100.7	+/- 4.6	104.9	+/- 6.3	0.59

Data represents mean +/- standard error.

doi:10.1371/journal.pone.0129359.t001

the enrichment statistics to classic. Other settings for GSEA pre-ranked were left by the software default.

These subjects for whom RNAseq was performed corresponded to the patients described in [Table 1](#), with the exception of subjects 29 and 31 (both acromegaly patients), which had clinical data but no RNAseq data. These data are available through the Gene Expression Omnibus (GSE57803).

Statistics

All statistical tests were performed using the R (version 3.0.2, [21]). To correct for multiple hypotheses, p-values were adjusted by the method of Benjamini and Hochberg [22] and referred to in this manuscript as q-values. The age adjusted, p-value corrected for multiple is denoted as q. Statistical significance was set at $p/q < 0.05$ for most comparisons except for GSEA analysis in which a $p < 0.25$ was used.

Descriptive statistics such as means and standard deviations were determined for clinical measurements. Student's t-test was used to test the difference in means of these measurements between control and acromegaly patients. Normality assumption was checked by Shapiro-Wilk test. Wilcoxon rank sum test was performed for HOMA-IR score, insulin levels and the 14:0, 16:0, 20:0 ceramides and the C16:0 glucosylceramide species as these data were not normally distributed. Welch's t-test was used for basal lipolysis since the equal variance assumption was rejected by Levene's test (car package version 2.0–19 [23]). Correlation coefficients were calculated by Pearson's product-moment.

Results and Discussion

Patient Characteristics

Clinical and metabolic measurements were obtained for 9 acromegaly patients and 11 controls. Patient characteristics are shown in [Table 1](#). There was no statistically significant difference in body mass index (BMI), abdominal circumference or weight. Acromegaly patients were younger ($p = 0.011$) and taller than their controls ($p = 0.036$).

Acromegaly Patients Were More Insulin Resistant and Had Higher Lipolysis

Acromegaly patients had elevated fasting glucose levels ($p = 0.013$) and higher fasted insulin ($p = 0.012$, [Fig 1A and 1B](#)). When combined, we observed higher HOMA-IR scores in the acromegalic patients than in the controls ($p = 0.001$, [Fig 1C](#)), reflecting a significant decrease in insulin sensitivity in the acromegaly patients, consistent with previous clinical findings [3].




Fig 1. Acromegalic patients have reduced insulin sensitivity and trend for higher lipolytic activity than their controls. A) Fasting blood glucose levels. B) Fasting insulin levels. C) HOMA-IR score from Control or Acromegaly subjects. D) *ex vivo* lipolysis as measured by glycerol release from excised white adipose tissue from control or acromegaly patients left untreated (Basal) or after stimulation with 30 nM isoproterenol (Iso). Data is presented as mean \pm standard error of the mean. Asterisk indicates $q < 0.05$.

doi:10.1371/journal.pone.0129359.g001

Subcutaneous adipose tissue chunks for lipolysis assay were available from 6 acromegaly patients and 9 controls. The results suggested that acromegaly patients may have higher basal lipolysis ($p = 0.11$), and higher lipolysis in the presence of isoproterenol ($p = 0.058$) even though they did not achieve statistical significance (Fig 1D). These data are consistent with previous reports linking GH signaling with increased lipolysis [24].

Transcriptomic Analysis

To determine which genes are altered in adipose tissue in acromegaly subjects, we performed a transcriptomic analysis of subcutaneous adipose tissue mRNA from 7 acromegalic patients and 11 controls. Patients separated along the first principal component approximately based on their age, with 10/11 controls in one group and 7/8 of the acromegalic patients on the other group (S1 Fig). This suggests that the major molecular differences between these groups can be explained by the presence or absence of acromegaly.

After correcting for age, we identified 418 genes that had significantly different expression in acromegaly. Of these, 198 genes were down-regulated and 290 were up-regulated in adipose tissue from the acromegalic patients. These transcripts form a signature identifying transcriptional differences in adipose tissue in response to long-term exposure to GH or indirectly to IGF-1 (Fig 2A and S1 and S2 Tables).

In general, gene expression changes in acromegalic patients over 60 were smaller than in patients under 60 (Fig 2B). Among genes that had significantly different expression, the fold change for a gene was $25 \pm 2.3\%$ higher in the younger cohort than the older cohort. This effect was statistically significant via a Wilcoxon-Rank Sum test ($p = 1.4 \times 10^{-11}$).

Gene set enrichment analysis testing KEGG pathways [25,26] showed enrichment of genes in the categories involved in metabolism, including upregulation of genes involved in the TCA cycle, fatty acid metabolism, biosynthesis of unsaturated fatty acids as well as genes which




Fig 2. Differential expression of genes in white adipose tissue from subjects with acromegaly compared to controls. A) Heatmap of the differentially expressed genes in white adipose tissue. Individual values are colored as the log fold change for a particular gene in a particular subject compared to the average expression of that gene across all cohorts, with brown indicating less expression and green indicating more expression (designated in the key as Row Z-score). The bar across the top indicates the subject's diagnosis, red for acromegaly and blue for controls. B) Scatterplot showing the log2 fold change for genes which had a statistically significant difference ($q < 0.05$) between acromegaly and control subjects. Each dot represents the log2 fold change for acromegaly for a gene in the under 60 and 60 or over cohorts. The solid line represents a slope of 1, which would imply no difference in fold change between age groups. The red line is a best fit line with a lower slope, showing that on average the fold change for older patients is smaller than the fold change for the under-60 patients.

doi:10.1371/journal.pone.0129359.g002




Fig 3. GH targets are differentially expressed in acromegaly subjects. A) mRNA Expression of A) *IGF1* and B) *IGFBP3* transcript levels in adipose tissue from control and acromegalic patients (C) Comparison between *IGF1* mRNA and IGF-1 serum levels in patients with acromegaly (D and E) Expression of mRNA for suppressors of growth hormone signaling (F) and Expression of tyrosine phosphatases associated with growth hormone signaling. Asterisks indicate $q < 0.05$ for the separated under 60 and 60 or over cohorts for panels A, B, D and E and for the age adjusted combined analysis for panel F. Barplots are presented as mean \pm standard error of the mean.

doi:10.1371/journal.pone.0129359.g003

regulate the cell cycle. We also observed downregulation of genes involved in pathways of GPCR signaling, MAPK signaling, inflammation and protein synthesis (S3 Table). We also examined the transcription factor networks that may underlie these changes in mRNA levels. We identified an up-regulation of several candidate transcription factors and microRNA's (S4 Table) whose targets are significantly altered in acromegalic subcutaneous white adipose tissue after adjusting for age. Notably among these are an up-regulation of E2F, GATA-1, MEF-2 and CREB targets and a down-regulation of AP1, STAT1, 3, 4 and 6, PPAR α , NF- κ B and SRF responsive genes.

Established GH Responsive Genes Are Up-Regulated in Adipose Tissue from Acromegaly Patients

Since acromegaly is caused by an overproduction of GH, we first analyzed known GH responsive genes. We found that expression of previously reported GH responsive genes, including *IGF1* (3.5 fold $q = 1.65 \times 10^{-6}$), and *IGFBP3* (2.3 fold, $q = 0.0002$) are elevated in acromegalic patients (Fig 3A and 3B). IGF-1 has been shown to be induced in adipocytes exposed to GH [27], while there were no previous reports regarding *IGFBP3* induction in adipose tissue. The confirmation of these previously reported acromegaly or GH-induced transcriptional changes strengthens our interpretation of other transcriptional changes. Neither the growth hormone receptor (GHR) nor the IGF-1 receptors (*IGF1R*, *IGF2R*) was significantly altered in acromegalic adipose tissue.

There was a correlation between *IGF1* mRNA and levels of IGF-1 in serum in the acromegaly patients ($R^2 = 0.51$, $p = 0.043$; Fig 3C), reflecting that increased induction of *IGF1* in adipose tissue is similar in its extent to serum IGF1 induction. Serum IGF1 is primarily thought to be derived from liver tissue due to the observation that serum IGF-1 levels are reduced 75% in a liver specific IGF-1 knockout [28]. Our data demonstrates that expression of the adipose tissue *IGF1* gene correlates well with that of serum IGF-1. Note that the older subjects had lower serum

IGF-1 than the younger subjects, indicating that circulating IGF-1 levels may correlate with generally reduced transcriptional changes observed in the older acromegalic patients (Fig 3C).

A Novel Negative Feedback Loop Is Induced by Chronic Exposure to High GH/IGF-1 Levels

SOCS2 AND *CISH*, both suppressors of cytokine signaling that have been shown to be important in down-regulating GH signaling, are up-regulated in acromegaly (1.7 and 2.3 fold respectively ($q = 0.003$ and $q = 0.00014$, Fig 3D and 3E). These data suggest that feedback mechanisms may be more active in younger patients, potentially either due to improved flexibility or reduced duration of the disease. These have been shown to be induced in liver and muscle by GH [29], and *SOCS2* has also been reported to be induced in adipocytes by GH [27,30]. We observed no significant differences in any PIAS genes.

We observed induction of the tyrosine phosphatases, *PTPN3* (also called PTP-H1, 2.5 fold higher $q = 0.0028$), *PTPN4* (1.6 fold $q = 0.00014$) and *PTPN13* (1.3 fold, $q = 0.038$) in acromegaly (Fig 3F). *PTPN3* has been reported to bind GH receptor *in vitro* in the presence of GH [31], and its overexpression reduces STAT5 signaling in response to GH [32]. *Ptpn3* Knockout mice have excessive GH activity, as demonstrated by excessive growth accompanied by a strong induction of liver *IGF1* mRNA and serum IGF-1 [32]. This is the first report of enhanced abundance of *PTPN3* mRNA in response to GH/IGF-1 exposure. The increased expression of *PTPN3* that we have observed in acromegaly suggests that this may be an additional negative feedback pathway induced by GH and reducing GH/IGF-1 signaling.

Genes Controlling DNA Replication, Proliferation and Apoptosis

We observed a difference in expression of several different genes regulating cellular proliferation in acromegalic subjects. The KEGG category containing DNA replication was enriched in acromegalic white adipose tissue, (S3 Table). Expression of Cyclin C (*CCNC*; 1.2 fold $q = 0.022$), and Cyclin E (*CCNE1*; 2.9 fold, $q = 6.5 \times 10^{-5}$) which are important for transition from G1 to S, were increased in acromegalic patients, and the negative regulator, cyclin dependent kinase inhibitor B (*CDKN2B*) was decreased (40% reduced fold $q = 0.016$, S2A Fig). *CDKN2B* has also been identified as a diabetes susceptibility gene identified in GWA studies [33,34].

Additional DNA replication genes that were induced were nucleosome assembly protein 1-like 1 (*NAP1L1*, 1.3 fold $q = 0.025$) and origin recognition complex, subunit 2 (*ORC2*, 1.7 fold $q = 0.0044$), which are important for DNA replication, and the anti-apoptotic regulators *BAG4* (BCL2-associated athanogene 4, 1.7 fold $q < 10^{-4}$) and *CAPN6* (calpain 6, 3.7 fold $q = 0.0011$) were also induced (S2 Fig). Together these implicate increased cell division, potentially of immune, vascular or pre-adipocyte cells in adipose tissue depots.

Apoptosis signal-regulating kinase 1 (*MAP3K5*) expression is higher in acromegaly (2.8 fold $q < 0.0004$), and there is also higher expression of its downstream substrates p38 α (*MAPK14* 1.2 fold $q = 0.012$), p38 δ (*MAPK13*, 2.7 fold $q < 10^{-4}$). The effect of GH or IGF-1 on these transcripts has not been reported before, and they could account for the effects of enhanced cell proliferation and apoptosis in response to GH/IGF-1 [35].

Transcriptional Changes Regulating Lipid Metabolism and Localization that May Contribute to Enhanced Lipolysis

To determine the potential causes of the increased lipolysis observed in Fig 1D, we examined the expression of human lipases in these adipose tissues. We observed no significant difference




Fig 4. Expression changes of selected genes, potentially responsible for lipolytic or insulin sensitivity alterations in acromegaly patients. mRNA Expression profile of genes potentially involved in lipid catabolism (A), regulation of lipolysis (B), fatty acid synthesis (C), fatty acid desaturation (D), *TCF7L2* (E), and glucocorticoid signaling (F). Asterisks indicate $q < 0.05$. Data indicates mean \pm standard error of the mean.

doi:10.1371/journal.pone.0129359.g004

in expression of the three classical triglyceride lipolysis enzymes hormone sensitive lipase (*LIPE*), adipose triglyceride lipase (*PNPLA*) or monoglycerol lipase (*MGLL*; Fig 4A). Lipoprotein lipase (*LPL*), the lipase important for lipolysis of triglycerides in apolipoproteins, was significantly more highly expressed in acromegaly patients (2.0 fold, $q = 0.004$). A strong induction of *LPL* expression in response to GH and absence of change in *HSL* were demonstrated before in a preadipocyte cell line [36–38] and in adipose tissue biopsies from GH deficient patients after treatment with GH [11]. Studies that addressed *LPL* enzymatic activity and not expression have found either no change or even a reduction in *LPL* activity in response to GH treatment of human adipocytes *in vitro* [39,40].

Although neither Hormone Sensitive Lipase (*LIPE*) nor ATGL (*PNPLA2*) were altered, two direct regulators of *HSL* and *ATGL* activity in adipocytes, abhydrolase domain containing 5 (*ABHD5*, also called CGI58 [41]), and nuclear receptor interacting protein 1 (*NRIP* (also called RIP140 [42]) were expressed at higher levels in adipose tissue from acromegaly patients (2.3 fold $q = 0.0016$ and 1.7 fold, $q = 0.043$ Fig 4B). CGI58 is an allosteric activator of lipolytic activity and RIP140 regulates CGI58's activity, therefore these data suggest that their transcriptional up-regulation could contribute to the induction of lipolysis by GH/IGF-1. *NRIP1* has also been proposed to be a transcription regulator of genes involved in lipid and glucose metabolism [42] and its induction could contribute to additional metabolic effects of GH/IGF-1 including disrupted glucose metabolism. We also examined *CIDEA/B/C* and *GOS2*, which have also been proposed to be positive regulators of lipolysis [43]. While there were no changes in the former, we did observe a non-significant elevation in *GOS2* (1.53 fold, $q = 0.246$; S2C Fig).

We next examined the expression of G-protein coupled receptors that induce lipolysis. The β_3 adrenergic (*ADRB3*) was more highly expressed in acromegaly patients compared to the

controls (5.2 fold, $q = 0.0064$). The $\beta 1$ receptor was also more highly expressed in acromegalic adipose tissue though it did not reach statistical significance (1.5 fold, $q = 0.20$; Fig 4B) suggesting a potential sensitization of these patients to adrenergic stimuli may underlie the enhanced *ex vivo* lipolysis.

In contrast to the lipolytic phenotype of acromegalic patients, several fatty acid and triglyceride synthesis genes were expressed at higher levels in acromegaly patients (Fig 4C). These include *ACSS2* (1.6 fold, $q = 0.044$) and *ACSS3* (1.7 fold, $q = 0.064$), which catalyze the activation of acetate for use in lipid synthesis. We also observed elevations in Acetoacetyl Co-A synthase (*AACS*; 1.9 fold, $q = 0.0066$), Acetyl Co-A carboxylase (*ACACA*, 1.7 fold $q = 0.039$), and Acyl-CoA synthetase long-chain family member 3 *ACSL3* (1.3 fold, $q = 0.045$), which convert long-chain fatty acids into fatty acyl-CoA esters. In addition, we also observed an elevation in fatty acid synthase *FASN* (1.7 fold, $q = 0.01$) in the acromegaly patients. The first two steps in triglyceride synthesis from fatty acids are catalyzed by glycerol-3-phosphate acyltransferase 9 (*AGPAT1,2,3* and 5) and 1-acyl-sn-glycerol-3-phosphate acyltransferase (*AGPAT9*). We observed transcriptional up-regulation of both *AGPAT2* (42% increased, $q = 0.02$) and *AGPAT9* (7.6 fold increased, $q = 2.7 \times 10^{-4}$) in acromegaly patients. Via pathway analysis, both fatty acid metabolism and unsaturated fatty acid biosynthetic pathways were up-regulated in the acromegaly patients (see S2 Table and below). This up-regulation may represent compensation by the adipose tissue due to enhanced lipid breakdown and oxidation in this and other tissues in acromegaly patients.

With respect to glycogen metabolism, expression of muscle glycogen phosphorylase (*PYGM*) was 3.2 fold higher ($q = 0.00078$) in the acromegaly patients (S2D Fig). The significance of glycogen in adipose tissue, or changes in glycogen content in acromegalic adipose tissue have not been characterized, but these findings would predict that the rates of glycogenolysis in acromegalic white adipose tissue would be elevated.

Transcripts Altered in Acromegaly that May Contribute to Insulin Resistance

One of the most pronounced differences between the acromegaly patients and the controls was a greater than 4 fold higher expression ($q < 1 \times 10^{-5}$; Fig 4D) of the three fatty acid desaturases—stearoyl-CoA desaturase (*SCD*, delta-9-desaturase), fatty acid desaturase 1 (*FADS1*, delta-5-desaturase) and fatty acid desaturase 2 (*FADS2*, delta-6-desaturase). *SCD* products and *FADS2* mRNA have also recently been shown to be induced by GH in mice [44]. The change in expression of these enzymes could be a possible link between acromegaly and insulin resistance due to an elevation of unsaturated fatty acids, as activity of *FADS1* and *FADS2* are associated with metabolic syndrome [45,46].

We observed no decrease in expression of canonical transcripts important for insulin signaling and response to insulin in adipocytes, including insulin receptor (*INSR*), *IRS1*, *IRS2*, *AKT1-3*, or GLUT4 (*SLC2A4*; see S3A Fig). This indicates that the observed insulin resistance is not caused by mRNA decreases in these genes. In fact, the KEGG category containing insulin signaling genes was generally up-regulated in these tissues (S3 Table). *AKT1* was significantly higher (1.3 fold, $q = 0.006$) and the remainder of these genes trended to be more highly expressed in the adipose tissue from the acromegalic and insulin resistant patients, potentially underlying a transcriptional up-regulation that compensates for an alternative insulin resistance mechanism.

One previously identified candidate is the phosphoinositide-3-kinase, regulatory subunit 1 (*PIK3RI*, also called p85 α), which was induced by GH in mouse adipose tissue and thought to contribute to GH-induced insulin resistance [47]. In our study *PIK3RI* expression was not

significantly different in the acromegaly patients, though it was modestly increased (25% increased, $q = 0.23$).

The ERK kinase pathway was down-regulated in the acromegaly patients including a 2–4 fold lower expression of the downstream transcription factors *FOS*, *JUN*, *JUNB* ($q < 0.04$, [S2 Table](#) and [S3B Fig](#)). Jun and Fos form the transcription factor AP1, which drives transcription of many targets involved in differentiation, proliferation and apoptosis [48]. Globally, we also found that AP1 targets were down-regulated. (ie, NES = -3.30, $q < 1 \times 10^{-4}$ for V\$AP1_Q4_01, [S4 Table](#)).

The cytokine modulators *STAT6* and the pro-inflammatory protein kinase IKK β (*IKKBK*) are expressed at lower levels (~18% reduced, $q = 0.0034$ for *STAT6* and $q = 0.009$ for *IKKBK*). Furthermore, the pro-inflammatory cytokines *IL1B*, *IL6* and *CCL2* (MCP-1) and the pro-inflammatory protein kinase *IKBKE* all trend towards lower expression, and genes from several KEGG pathways involved in inflammation were expressed at lower levels in the acromegaly cohort ([S2 Table](#) and [S3 Table](#) and [S3C Fig](#)). These data support the hypothesis that insulin resistance in these patients is not due to enhanced inflammatory signaling.

To test biochemically whether ceramides may play a role in the acromegaly associated insulin resistance, we took a lipidomics approach to analyze ceramide and glucosylceramide species from the adipose tissue explants of these patients. Elevated ceramides have been proposed to mediate insulin resistance by several models [49–52]. We observed a modest down-regulation of the mRNA levels of glycosylsphingolipid metabolic genes in our data (normalized enrichment score = -0.86 $q = 0.71$). We then measured ceramide species from 7 acromegaly patients and 11 control patients directly and observed no statistically significant changes in any ceramide species ([S4 Fig](#), q -values all > 0.25). We did however, detect modest elevations of C16:0, C18:0 and C24:0 ceramide species in subcutaneous adipose tissue from acromegalic patients. We therefore do not have strong evidence to support the hypothesis that ceramide elevations are causative of insulin resistance in acromegalic white adipose tissue.

TCF7L2, a transcription factor regulating many metabolism genes that is also a diabetes susceptibility gene [53] is up-regulated in the acromegaly patients (1.5 fold, $q = 0.00045$, [Fig 4E](#)). Mice with liver specific knockout of *Tcf7l2* are hypoglycemic, while transgenic mice overexpressing liver *Tcf7l2* are hyperglycemic [54]. *TCF7L2* in subcutaneous fat is higher and expression of splice isoforms is reduced in subcutaneous fat and in liver following bariatric surgery [55]. Higher expression of *TCF7L2* could also therefore be linked to insulin resistance in acromegaly.

Glucocorticoid Regulation

11 β Hydroxysteroid dehydrogenase 1 (*HSD11B1*), the enzyme that activates transformation of cortisone to cortisol, was reduced over 4 fold in acromegaly patients ($q = 0.0048$, [Fig 4F](#)). The down-regulation of expression and activity of this enzyme by GH/IGF1 has been confirmed *in vitro* [56], in GH deficient patients treated with GH [57] and in acromegaly patients [58,59]. In addition, we found higher expression of the glucocorticoid receptor (*NR3C1*, 1.5 fold $q = 0.00013$) in acromegaly patients ([Fig 2F](#)). Glucocorticoid receptor expression is repressed by cortisol [60], so the higher expression may be due to the reduced local cortisol levels caused by lowered *HSD11B1*.

Summary

In this study we have described a transcriptional signature in adipose tissue from subjects with acromegaly. We identified 418 adipose tissue genes altered in acromegaly patients. Some of

these genes may be direct targets of increased GH or IGF-1 signaling in adipose tissue, whereas others may be secondary adaptations to this condition.

Interestingly, we observed more modest gene expression changes in general for older acromegalic patients than for younger patients. We are unable to determine from our study how long patients were acromegalic prior to our study, so one possibility is that the older patients have had longer to adapt to elevated GH levels. Alternatively, elevated GH/IGF-1 signaling may play a stronger role in younger patients. It should be noted, however that this exploratory finding was limited since we only had 2 acromegalic patients over 60 in our study, so these age-dependent findings will need to be reproduced in a larger cohort.

The fact that the patients consistently had a relatively uniform change of expression of these genes suggests that we are able to draw valid conclusions about adipose tissue in acromegalic patients even from this small cohort. Furthermore, as mentioned throughout, our data agree with several previous studies in animal models and patients. The confirmation of these previously reported GH-dependent transcriptional changes strengthens our interpretation of other transcriptional changes. One potential caveat to our approach is the use of patients with a non-secreting adenoma as the control group. To avoid the possible effects of hypopituitarism on adipose tissue we excluded patients with pituitary hormone deficiencies. We chose to include this as the control group as these samples not only collected in an identical manner from the same surgeons and processed identically, but also controls for potential non-secreting effects of pituitary tumor growths in the acromegaly subjects.

A potential caveat is the potential confounding effect of anti-diabetic or anti-growth hormone medications. Only one acromegalic patient was on somatostatin, and his IGF-1 levels were non-responsive. Our exclusion criteria included any glucocorticoid treatment. One patient in each group was on metformin as an antidiabetic medication, so we do not feel that this affected our overall conclusions. Another potential limitation was our inability to reanalyze the samples by a secondary method for gene expression or to validate our findings at the protein level, due to a lack of sample. We could not analyze a second cohort due to the rarity of this disease.

These data provide a variety of novel transcriptional changes that may be causative of the co-morbidities associated with acromegaly. Further studies in animals and cells using knockout or overexpression of specific transcripts may verify which of the changes is crucial in metabolic effects of GH in adipose tissue.

Supporting Information

S1 Fig. Principal component analysis of normalized transcript levels. Normalized counts were transformed via a regularized log transformation then principal components were calculated. Samples were then colored based on age and diagnosis. The dotted line indicates the grouping of samples into groups based on their disease state.

(TIF)

S2 Fig. Expression changes of selected other transcripts. mRNA Expression profile of differentially expressed genes involved in A) MAPK Signaling, B) Cell cycle control, C) Lipolysis and D) Glycogen metabolism. Asterisks indicate $q < 0.05$. Barplots are presented as mean \pm standard error of the mean. mRNA Expression is in units of RPKM (reads per kilobase per million reads).

(TIF)

S3 Fig. Expression changes of selected insulin signal transduction genes. mRNA Expression profile of genes involved in insulin signaling (A) and inflammation (B). Asterisks indicate

$q < 0.05$. Barplots are presented as mean \pm standard error of the mean. mRNA Expression is in units of RPKM (reads per kilobase per million reads).

(TIF)

S4 Fig. Ceramide levels in adipose tissue from control and acromegalic patients. Ceramide (Cer) or glucosylceramide levels (GluCer) were determined as described in the methods and materials section. The number indicates the fatty acid species associated with each lipid. All values are normalized such that control values are equal to 1. Data indicates mean \pm standard error of the mean.

(TIF)

S1 Table. Expression changes between control and acromegaly subjects. Calculated expression, and expression changes for each gene are shown along with raw p-value, adjusted p-values and the fold change.

(CSV)

S2 Table. Age adjusted gene expression changes between control and acromegaly subjects. Patients were grouped into under-60 and 60 and over as the age and genes were first analyzed with age group as a covariate then after that adjustment, the disease state was taken into account. Three log₂ fold changes and p-values are presented. The age adjusted effect of Acromegaly (AcrovsControl), the effects if acromegaly in only the under 60 group (Acrovs-Con_0.60) and the effects of acromegaly in only the above 60 group (AcrovsCon_60.100).

(XLS)

S3 Table. Gene set enrichment analysis of GO and KEGG pathways. Size is the total size of the KEGG category, NES is the normalized enrichment score, NOM p-value is the raw p-value and FDR q-value is corrected for multiple observations. Gene details lists the specific genes which led to the enrichment of this category in our data. A negative enrichment score indicates down-regulation of the category in acromegaly.

(XLS)

S4 Table. Gene set enrichment analysis of transcription factor and miRNA pathways. These categories indicate that target genes regulated by these factors are altered in acromegalic white adipose tissue. Size is the total size of the category, NES is the normalized enrichment score, NOM p-value is the raw p-value and FDR q-value is corrected for multiple observations. Gene details lists the specific genes which led to the enrichment of this category in our data. A negative enrichment score indicates down-regulation of the category in acromegaly.

(XLS)

Acknowledgments

We thank Charlotte Gunden, Elizabeth Walkowiak and Eric Vasbinder for their valuable help in the study. The authors would like to thank Ian Brooks and the UTHSC-ORNL Center for Biomedical Informatics for provisioning the Rstudio server used in this analysis. We would also like to thank Solomon S. Solomon and the members of the Bridges and Reiter laboratories (at UTHSC) for helpful suggestions.

Author Contributions

Conceived and designed the experiments: IH ALB WFC. Performed the experiments: IH WFC. Analyzed the data: IH QTT DB. Contributed reagents/materials/analysis tools: ARS. Wrote the paper: IH QTT DB. Participated in conception and design: ARS.

References

1. Holdaway IM, Rajasoorya C (1999) Epidemiology of acromegaly. *Pituitary* 2: 29–41. PMID: [11081170](#)
2. Vijayakumar A, Novosyadlyy R, Wu Y, Yakar S, LeRoith D (2010) Biological effects of growth hormone on carbohydrate and lipid metabolism. *Growth Horm IGF Res* 20: 1–7. doi: [10.1016/j.ghir.2009.09.002](#) PMID: [19800274](#)
3. Ezzat S, Forster MJ, Berchtold P, Redelmeier DA, Boerlin V, Harris AG (1994) Acromegaly. Clinical and biochemical features in 500 patients. *Medicine (Baltimore)* 73: 233–240. PMID: [7934807](#)
4. Colao A, Baldelli R, Marzullo P, Ferretti E, Ferone D, Gargiulo D, et al. (2000) Systemic hypertension and impaired glucose tolerance are independently correlated to the severity of the acromegalic cardiomyopathy. *J Clin Endocrinol Metab* 85: 193–199. PMID: [10634386](#)
5. Heinrich UE, Schalch DS, Koch JG, Johnson CJ (1978) Nonsuppressible insulin-like activity (NSILA). II. Regulation of serum concentrations by growth hormone and insulin. *J Clin Endocrinol Metab* 46: 672–678. doi: [10.1210/jcem-46-4-672](#) PMID: [573281](#)
6. Ayuk J, Sheppard MC (2006) Growth hormone and its disorders. *Postgrad Med J* 82: 24–30. doi: [10.1136/pgmj.2005.036087](#) PMID: [16397076](#)
7. Jørgensen JOL, Jessen N, Pedersen SB, Vestergaard E, Gormsen L, Lund SA, et al. (2006) GH receptor signaling in skeletal muscle and adipose tissue in human subjects following exposure to an intravenous GH bolus. *Am J Physiol Endocrinol Metab* 291: E899–E905. doi: [10.1152/ajpendo.00024.2006](#) PMID: [16757551](#)
8. Bolinder J, Ostman J, Werner S, Amer P (1986) Insulin action in human adipose tissue in acromegaly. *J Clin Invest* 77: 1201–1206. doi: [10.1172/JCI112422](#) PMID: [3514677](#)
9. Gravhølt CH, Schmitz O, Simonsen L, Bülow J, Christiansen JS, Møller N, et al. (1999) Effects of a physiological GH pulse on interstitial glycerol in abdominal and femoral adipose tissue. *Am J Physiol* 277: E848–E854. PMID: [10567011](#)
10. Nielsen S, Møller N, Christiansen JS, Jørgensen JO (2001) Pharmacological antilipolysis restores insulin sensitivity during growth hormone exposure. *Diabetes* 50: 2301–2308. PMID: [11574412](#)
11. Khalfallah Y, Sassolas G, Borson-Chazot F, Vega N, Vidal H (2001) Expression of insulin target genes in skeletal muscle and adipose tissue in adult patients with growth hormone deficiency: effect of one year recombinant human growth hormone therapy. *J Endocrinol* 171: 285–292. PMID: [11691648](#)
12. Kasumov T, Huang H, Chung Y-M, Zhang R, McCullough AJ, Kirwan JP, et al. (2010) Quantification of ceramide species in biological samples by liquid chromatography electrospray ionization tandem mass spectrometry. *Anal Biochem* 401: 154–161. doi: [10.1016/j.ab.2010.02.023](#) PMID: [20178771](#)
13. Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37: 911–917. PMID: [13671378](#)
14. Kim D, Perteu G, Trapnell C, Pimentel H, Kelley R, Salzberg SL, et al. (2013) TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol* 14: R36. doi: [10.1186/gb-2013-14-4-r36](#) PMID: [23618408](#)
15. Langmead B, Trapnell C, Pop M, Salzberg SL (2009) Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 10: R25. doi: [10.1186/gb-2009-10-3-r25](#) PMID: [19261174](#)
16. Anders S, Pyl PT, Huber W (2014) HTSeq—A Python framework to work with high-throughput sequencing data. *Bioinformatics*: 1–4. doi: [10.1093/bioinformatics/btu638](#)
17. Love MI, Huber W, Anders S (2014) Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2. doi: [10.1101/002832](#)
18. Tanimoto K, Hizuka N, Fukuda I, Takano K, Hanafusa T (2008) The influence of age on the GH-IGF1 axis in patients with acromegaly. *Eur J Endocrinol* 159: 375–379. doi: [10.1530/EJE-08-0243](#) PMID: [18653547](#)
19. Mootha VK, Lindgren CM, Eriksson K-F, Subramanian A, Sihag S, Lehar J, et al. (2003) PGC-1alpha-responsive genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes. *Nat Genet* 34: 267–273. doi: [10.1038/ng1180](#) PMID: [12808457](#)
20. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA (2005) Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* 102: 15545–15550. doi: [10.1073/pnas.0506580102](#) PMID: [16199517](#)
21. R Development Core Team, R Core Team (2011) R: A language and environment for statistical computing.
22. Benjamini Y, Hochberg Y (1995) Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *J R Stat Soc Ser B* 57: 289–300.

23. Fox J, Weisberg S (2011) *An {R} Companion to Applied Regression*. Second. Thousand Oaks {CA}: Sage.
24. Moller L, Norrelund H, Jessen N, Flyvbjerg A, Pedersen SB, Gaylinn DB, et al. (2009) Impact of growth hormone receptor blockade on substrate metabolism during fasting in healthy subjects. *J Clin Endocrinol Metab* 94: 4524–4532. doi: [10.1210/jc.2009-0381](https://doi.org/10.1210/jc.2009-0381) PMID: [19820031](https://pubmed.ncbi.nlm.nih.gov/19820031/)
25. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. (2000) Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. *Nat Genet* 25: 25–29. doi: [10.1038/75556](https://doi.org/10.1038/75556) PMID: [10802651](https://pubmed.ncbi.nlm.nih.gov/10802651/)
26. Young MD, Wakefield MJ, Smyth GK, Oshlack A (2010) Gene ontology analysis for RNA-seq: accounting for selection bias. *Genome Biol* 11: R14. doi: [10.1186/gb-2010-11-2-r14](https://doi.org/10.1186/gb-2010-11-2-r14) PMID: [20132535](https://pubmed.ncbi.nlm.nih.gov/20132535/)
27. Fleenor D, Arumugam R, Freemark M (2006) Growth hormone and prolactin receptors in adipogenesis: STAT-5 activation, suppressors of cytokine signaling, and regulation of insulin-like growth factor I. *Horm Res* 66: 101–110. doi: [10.1159/000093667](https://doi.org/10.1159/000093667) PMID: [16735796](https://pubmed.ncbi.nlm.nih.gov/16735796/)
28. Haluzik M, Yakar S, Gavrilova O, Setser J, Boisclair Y, LeRoith D (2003) Insulin resistance in the liver-specific IGF-1 gene-deleted mouse is abrogated by deletion of the acid-labile subunit of the IGF-binding protein-3 complex: relative roles of growth hormone and IGF-1 in insulin resistance. *Diabetes* 52: 2483–2489. PMID: [14514630](https://pubmed.ncbi.nlm.nih.gov/14514630/)
29. Clasen BFF, Krusenstjerna-Hafstrøm T, Vendelbo MH, Thorsen K, Escande C, Møller N, et al. (2013) Gene expression in skeletal muscle after an acute intravenous GH bolus in human subjects: identification of a mechanism regulating ANGPTL4. *J Lipid Res* 54: 1988–1997. doi: [10.1194/jlr.P034520](https://doi.org/10.1194/jlr.P034520) PMID: [23606725](https://pubmed.ncbi.nlm.nih.gov/23606725/)
30. Huo JS, McEachin RC, Cui TX, Duggal NK, Hai T, States DJ, et al. (2006) Profiles of growth hormone (GH)-regulated genes reveal time-dependent responses and identify a mechanism for regulation of activating transcription factor 3 by GH. *J Biol Chem* 281: 4132–4141. doi: [10.1074/jbc.M508492200](https://doi.org/10.1074/jbc.M508492200) PMID: [16326703](https://pubmed.ncbi.nlm.nih.gov/16326703/)
31. Pasquali C, Curchod M-L, Wälchli S, Espanel X, Guerrier M, Arigoni F, et al. (2003) Identification of protein tyrosine phosphatases with specificity for the ligand-activated growth hormone receptor. *Mol Endocrinol* 17: 2228–2239. doi: [10.1210/me.2003-0011](https://doi.org/10.1210/me.2003-0011) PMID: [12907755](https://pubmed.ncbi.nlm.nih.gov/12907755/)
32. Pilecka I, Patrignani C, Pescini R, Curchod M-L, Perrin D, Xue Y, et al. (2007) Protein-tyrosine phosphatase H1 controls growth hormone receptor signaling and systemic growth. *J Biol Chem* 282: 35405–35415. doi: [10.1074/jbc.M705814200](https://doi.org/10.1074/jbc.M705814200) PMID: [17921143](https://pubmed.ncbi.nlm.nih.gov/17921143/)
33. Sladek R, Rocheleau G, Rung J, Dina C, Shen L, Serre D, et al. (2007) A genome-wide association study identifies novel risk loci for type 2 diabetes. *Nature* 445: 881–885. doi: [10.1038/nature05616](https://doi.org/10.1038/nature05616) PMID: [17293876](https://pubmed.ncbi.nlm.nih.gov/17293876/)
34. Saxena R, Voight BF, Lyssenko V, Burtt NP, de Bakker PIW, Chen H, et al. (2007) Genome-wide association analysis identifies loci for type 2 diabetes and triglyceride levels. *Science* (80-) 316: 1331–1336. doi: [10.1126/science.1142358](https://doi.org/10.1126/science.1142358) PMID: [17463246](https://pubmed.ncbi.nlm.nih.gov/17463246/)
35. Waters MJ, Brooks AJ (2012) Growth hormone and cell growth. *Endocr Dev* 23: 86–95. doi: [10.1159/000341761](https://doi.org/10.1159/000341761) PMID: [23182823](https://pubmed.ncbi.nlm.nih.gov/23182823/)
36. Pradines-Figueres A, Barcellini-Couget S, Dani C, Baudoin C, Ailhaud G (1990) Inhibition by serum components of the expression of lipoprotein lipase gene upon stimulation by growth hormone. *Biochem Biophys Res Commun* 166: 1118–1125. PMID: [2306231](https://pubmed.ncbi.nlm.nih.gov/2306231/)
37. Barcellini-Couget S, Vassaux G, Negrel R, Ailhaud G (1994) Rise in cytosolic Ca²⁺ abolishes in preadipose cells the expression of lipoprotein lipase stimulated by growth hormone. *Biochem Biophys Res Commun* 199: 136–143. PMID: [8123005](https://pubmed.ncbi.nlm.nih.gov/8123005/)
38. Padines-Figuères A, Barcellini-Couget S, Dani C, Vannier C, Ailhaud G (1990) Transcriptional control of the expression of lipoprotein lipase gene by growth hormone in preadipocyte Ob1771 cells. *J Lipid Res* 31: 1283–1291. PMID: [2401859](https://pubmed.ncbi.nlm.nih.gov/2401859/)
39. Simsolo RB (1995) Effects of acromegaly treatment and growth hormone on adipose tissue lipoprotein lipase. *J Clin Endocrinol Metab* 80: 3233–3238. doi: [10.1210/jc.80.11.3233](https://doi.org/10.1210/jc.80.11.3233) PMID: [7593431](https://pubmed.ncbi.nlm.nih.gov/7593431/)
40. Richelsen B, Pedersen SB, Kristensen K, Børglum JD, Nørrelund H, Christiansen JS, et al. (2000) Regulation of Lipoprotein Lipase and Hormone-Sensitive Lipase Activity and Gene Expression in Adipose and Muscle Tissue by Growth Hormone Treatment During Weight Loss in Obese Patients. 49: 906–911. doi: [10.1053/mt.2000.6738](https://doi.org/10.1053/mt.2000.6738) PMID: [10910003](https://pubmed.ncbi.nlm.nih.gov/10910003/)
41. Lass A, Zimmermann R, Haemmerle G, Riederer M, Schoiswohl G, Schweiger M, et al. (2006) Adipose triglyceride lipase-mediated lipolysis of cellular fat stores is activated by CGI-58 and defective in Charnarin-Dorfman Syndrome. *Cell Metab* 3: 309–319. doi: [10.1016/j.cmet.2006.03.005](https://doi.org/10.1016/j.cmet.2006.03.005) PMID: [16679289](https://pubmed.ncbi.nlm.nih.gov/16679289/)

42. Ho P-C, Chuang Y-S, Hung C-H, Wei L-N (2011) Cytoplasmic receptor-interacting protein 140 (RIP140) interacts with perilipin to regulate lipolysis. *Cell Signal* 23: 1396–1403. doi: [10.1016/j.cellsig.2011.03.023](https://doi.org/10.1016/j.cellsig.2011.03.023) PMID: [21504789](https://pubmed.ncbi.nlm.nih.gov/21504789/)
43. Nielsen TS, Jessen N, Jørgensen JOL, Møller N, Lund S (2014) Dissecting adipose tissue lipolysis: Molecular regulation and implications for metabolic disease. *J Mol Endocrinol* 52. doi: [10.1530/JME-13-0277](https://doi.org/10.1530/JME-13-0277)
44. Oberbauer AM, German JB, Murray JD (2011) Growth hormone enhances arachidonic acid metabolites in a growth hormone transgenic mouse. *Lipids* 46: 495–504. doi: [10.1007/s11745-011-3548-y](https://doi.org/10.1007/s11745-011-3548-y) PMID: [21442273](https://pubmed.ncbi.nlm.nih.gov/21442273/)
45. Kröger J, Schulze MB (2012) Recent insights into the relation of $\Delta 5$ desaturase and $\Delta 6$ desaturase activity to the development of type 2 diabetes. *Curr Opin Lipidol* 23: 4–10. doi: [10.1097/MOL.0b013e32834d2dc5](https://doi.org/10.1097/MOL.0b013e32834d2dc5) PMID: [22123669](https://pubmed.ncbi.nlm.nih.gov/22123669/)
46. Mayneris-Perxachs J, Guerendiain M, Castellote AI, Estruch R, Covas MI, Fito M, et al. (2013) Plasma fatty acid composition, estimated desaturase activities, and their relation with the metabolic syndrome in a population at high risk of cardiovascular disease. *Clin Nutr* 33: 90–97. doi: [10.1016/j.clnu.2013.03.001](https://doi.org/10.1016/j.clnu.2013.03.001) PMID: [23591154](https://pubmed.ncbi.nlm.nih.gov/23591154/)
47. Del Rincon J- P, Iida K, Gaylann BD, McCurdy CE, Leitner JW, Barbour LA, et al. (2007) Growth hormone regulation of p85alpha expression and phosphoinositide 3-kinase activity in adipose tissue: mechanism for growth hormone-mediated insulin resistance. *Diabetes* 56: 1638–1646. doi: [10.2337/db06-0299](https://doi.org/10.2337/db06-0299) PMID: [17363744](https://pubmed.ncbi.nlm.nih.gov/17363744/)
48. Shaulian E, Karin M (2002) AP-1 as a regulator of cell life and death. *Nat Cell Biol* 4: E131–E136. doi: [10.1038/ncb0502-e131](https://doi.org/10.1038/ncb0502-e131) PMID: [11988758](https://pubmed.ncbi.nlm.nih.gov/11988758/)
49. Holland WL, Brozinick JT, Wang L-PP, Hawkins ED, Sargent KM, Liu Y, et al. (2007) Inhibition of ceramide synthesis ameliorates glucocorticoid-, saturated-fat-, and obesity-induced insulin resistance. *Cell Metab* 5: 167–179. doi: [10.1016/j.cmet.2007.01.002](https://doi.org/10.1016/j.cmet.2007.01.002) PMID: [17339025](https://pubmed.ncbi.nlm.nih.gov/17339025/)
50. Adams JM, Pratipanawatr T, Berria R, Wang E, DeFronzo RA, Sullards MC, et al. (2004) Ceramide content is increased in skeletal muscle from obese insulin-resistant humans. *Diabetes* 53: 25–31. PMID: [14693694](https://pubmed.ncbi.nlm.nih.gov/14693694/)
51. Aerts JM, Ottenhoff R, Powlson AS, Grefhorst A, van Eijk M, Dubbelhuis PF, et al. (2007) Pharmacological inhibition of glucosylceramide synthase enhances insulin sensitivity. *Diabetes* 56: 1341–1349. doi: [10.2337/db06-1619](https://doi.org/10.2337/db06-1619) PMID: [17287460](https://pubmed.ncbi.nlm.nih.gov/17287460/)
52. Chavez JA, Knotts T a, Wang L-P, Li G, Dobrowsky RT, Florant GL, et al. (2003) A role for ceramide, but not diacylglycerol, in the antagonism of insulin signal transduction by saturated fatty acids. *J Biol Chem* 278: 10297–10303. doi: [10.1074/jbc.M212307200](https://doi.org/10.1074/jbc.M212307200) PMID: [12525490](https://pubmed.ncbi.nlm.nih.gov/12525490/)
53. Ip W, Chiang Y-TA, Jin T (2012) The involvement of the wnt signaling pathway and TCF7L2 in diabetes mellitus: The current understanding, dispute, and perspective. *Cell Biosci* 2: 28. doi: [10.1186/2045-3701-2-28](https://doi.org/10.1186/2045-3701-2-28) PMID: [22892353](https://pubmed.ncbi.nlm.nih.gov/22892353/)
54. Boj SF, van Es JH, Huch M, Li VSW, José A, Hatzis P, et al. (2012) Diabetes risk gene and Wnt effector Tcf7l2/TCF4 controls hepatic response to perinatal and adult metabolic demand. *Cell* 151: 1595–1607. doi: [10.1016/j.cell.2012.10.053](https://doi.org/10.1016/j.cell.2012.10.053) PMID: [23260145](https://pubmed.ncbi.nlm.nih.gov/23260145/)
55. Hindle AK, Brody F, Tevar R, Kluk B, Hill S, McCaffrey T, et al. (2009) TCF7L2 expression in diabetic patients undergoing bariatric surgery. *Surg Endosc* 23: 700–704. doi: [10.1007/s00464-008-0001-2](https://doi.org/10.1007/s00464-008-0001-2) PMID: [18806947](https://pubmed.ncbi.nlm.nih.gov/18806947/)
56. Napolitano a Voice MW, Edwards CR Seckl JR, Chapman KE (1998) 11Beta-hydroxysteroid dehydrogenase 1 in adipocytes: expression is differentiation-dependent and hormonally regulated. *J Steroid Biochem Mol Biol* 64: 251–260. PMID: [9618026](https://pubmed.ncbi.nlm.nih.gov/9618026/)
57. Paulsen SK, Pedersen SB, Jørgensen JOL, Fisker S, Christiansen JS, Flyvbjerg A, et al. (2006) Growth hormone (GH) substitution in GH-deficient patients inhibits 11beta-hydroxysteroid dehydrogenase type 1 messenger ribonucleic acid expression in adipose tissue. *J Clin Endocrinol Metab* 91: 1093–1098. doi: [10.1210/jc.2005-1694](https://doi.org/10.1210/jc.2005-1694) PMID: [16368752](https://pubmed.ncbi.nlm.nih.gov/16368752/)
58. Frajese GV, Taylor NF, Jenkins PJ, Besser GM, Monson JP (2004) Modulation of cortisol metabolism during treatment of acromegaly is independent of body composition and insulin sensitivity. *Horm Res* 61: 246–251. doi: [10.1159/000077135](https://doi.org/10.1159/000077135) PMID: [15004437](https://pubmed.ncbi.nlm.nih.gov/15004437/)
59. Moore JS, Monson JP, Kaltsas G, Putignano P, Wood PJ, Sheppard MC, et al. (1999) Modulation of 11beta-hydroxysteroid dehydrogenase isozymes by growth hormone and insulin-like growth factor: in vivo and in vitro studies. *J Clin Endocrinol Metab* 84: 4172–4177. PMID: [10566668](https://pubmed.ncbi.nlm.nih.gov/10566668/)
60. Kalinyak JE, Dorin RI, Hoffman AR, Perlman AJ (1987) Tissue-specific regulation of glucocorticoid receptor mRNA by dexamethasone. *J Biol Chem* 262: 10441–10444. PMID: [2440867](https://pubmed.ncbi.nlm.nih.gov/2440867/)