

UC Irvine

UC Irvine Previously Published Works

Title

The Aquaporin Gene Family of the Yellow Fever Mosquito, *Aedes aegypti*

Permalink

<https://escholarship.org/uc/item/52k798c3>

Journal

PLoS ONE, 5(12)

ISSN

1932-6203

Authors

Drake, Lisa L.
Boudko, Dmitri Y.
Marinotti, Osvaldo
[et al.](#)

Publication Date

2010-12-29

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

The Aquaporin Gene Family of the Yellow Fever Mosquito, *Aedes aegypti*

Lisa L. Drake¹, Dmitri Y. Boudko⁴, Osvaldo Marinotti⁵, Victoria K. Carpenter¹, Angus L. Dawe^{1,3}, Immo A. Hansen^{1,2,3*}

1 Department of Biology, New Mexico State University, Las Cruces, New Mexico, United States of America, **2** Institute of Applied Biosciences, New Mexico State University, Las Cruces, New Mexico, United States of America, **3** Molecular Biology Program, New Mexico State University, Las Cruces, New Mexico, United States of America, **4** Chicago Medical School, Rosalind Franklin University, North Chicago, Illinois, United States of America, **5** Department of Molecular Biology and Biochemistry, University of California Irvine, Irvine, California, United States of America

Abstract

Background: The mosquito, *Aedes aegypti*, is the principal vector of the Dengue and yellow fever viruses. During feeding, an adult female can take up more than its own body weight in vertebrate blood. After a blood meal females excrete large amounts of urine through their excretion system, the Malpighian tubules (MT). Diuresis starts within seconds after the mosquito starts feeding. Aquaporins (AQPs) are a family of membrane transporters that regulate the flow of water, glycerol and other small molecules across cellular membranes in both prokaryotic and eukaryotic cells. Our aim was to identify aquaporins that function as water channels, mediating transcellular water transport in MTs of adult female *Ae. aegypti*.

Methodology/Principal Findings: Using a bioinformatics approach we screened genome databases and identified six putative AQPs in the genome of *Ae. aegypti*. Phylogenetic analysis showed that five of the six *Ae. aegypti* AQPs have high similarity to classical water-transporting AQPs of vertebrates. Using microarray, reverse transcription and real time PCR analysis we found that all six AQPs are expressed in distinct patterns in mosquito tissues/body parts. AaAQP1, 4, and 5 are strongly expressed in the adult female MT. RNAi-mediated knockdown of the MT-expressed mosquito AQPs resulted in significantly reduced diuresis.

Conclusions/Significance: Our results support the notion that AQP1, 4, and 5 function as water transporters in the MTs of adult female *Ae. aegypti* mosquitoes. Our results demonstrate the importance of these AQPs for mosquito diuresis after blood ingestion and highlight their potential as targets for the development of novel vector control strategies.

Citation: Drake LL, Boudko DY, Marinotti O, Carpenter VK, Dawe AL, et al. (2010) The Aquaporin Gene Family of the Yellow Fever Mosquito, *Aedes aegypti*. PLoS ONE 5(12): e15578. doi:10.1371/journal.pone.0015578

Editor: Pedro Lagerblad Oliveira, Universidade Federal do Rio de Janeiro, Brazil

Received: July 29, 2010; **Accepted:** November 17, 2010; **Published:** December 29, 2010

Copyright: © 2010 Drake et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This project had two funding sources. First, start-up money given to Immo Hansen by the vice president for research of New Mexico State University. Secondly, this project was supported in part by grant #52005881 to NMSU from the Howard Hughes Medical Institute. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: immoh@nmsu.edu

Introduction

Anautogenous mosquito females need vertebrate blood for reproduction. The nutrients taken up with the blood are used to synthesize large amounts of yolk proteins that are deposited in the eggs during a process called vitellogenesis. Yolk delivers the energy and building blocks for embryogenesis. This “need for blood” in order to reproduce makes anautogenous mosquitoes effective disease vectors because they require at least one insect-host contact for every batch of eggs they develop.

During feeding, adult female *Aedes aegypti*, can take up more than their own body weight in blood [1]. This severely impairs their mobility and puts them at risk to be killed by their host or other predators. In addition, mosquitoes face the problem of high sodium content of vertebrate blood plasma and high potassium content in red blood cells. Therefore, it is essential for mosquitoes to possess an efficient system for excretion of excess water and ions while retaining the nutrients contained in the blood meal. During and after a blood meal female *Ae. aegypti* secrete large amounts of urine

through their Malpighian tubules (MT). Within the first hour after taking a blood meal (post blood meal - PBM) mosquito females can discharge more than 40% of water and sodium enclosed in the blood plasma [1]. Diuresis is under hormonal control by neuropeptide hormones secreted by the central nervous system [2]. In the current model, diuretic hormones, released seconds after start of the blood meal, stimulate the MT cells to produce the second messenger molecule cAMP which activates transcellular diuresis by increasing transepithelial cation (Na^+ , K^+) transport [2]. Another class of neuropeptide hormones, the kinins, increase intracellular calcium levels that regulate anion movement (Cl^-) into the MT lumen [3,4]. Urine produced by the MT is collected in the hind gut and subsequently forcefully ejected from the rectum in a process that involves rectal peristalsis and movement of the 7th and 8th abdominal segments. Females start expelling small urine droplets approximately 50–75 seconds after start of feeding. Urine droplets have a volume of about 10 to 12 nl and can fly up to 10 mm [1,5].

Aquaporins (AQPs) are transport channels that make cell membranes permeable to water. They are found in all plant,

animal, fungi, eubacteria, and archaea taxa studied [6,7,8]. In mammals, there are 13 AQP's and they form two subfamilies with different transport selectivity: orthodox aquaporins, which transport only water, and aquaglyceroporins, which transport glycerol, urea, small solutes, and water [9]. In the so-called hourglass model for AQP structure, the six transmembrane alpha helical domains (numbered 1 – 6) are connected by five loops termed A – E [10,11,12]. Both amino- and carboxy-terminus are located inside the cytoplasm. The transmembrane domains 2–3 and 5–6 are connected by loops B and E, both containing a highly conserved NPA (Asparagine-Proline-Alanine) motive and other conserved residues. These hydrophobic NPA loops form a ring as part of an hourglass-shaped pore within the center of the phospholipid bilayer membrane. This ring, with a diameter of 2.8 Å, is the primary filter that prevents protons from crossing through the AQP pore. Hg²⁺ ions interact with a cysteine residue close to the NPA motive in the E loop and an alanine residue in the B loop of most AQP's and efficiently obstruct water transport through the pore [13].

The activity of eukaryotic aquaporins is commonly regulated via three different mechanisms: translation, gating, or trafficking [13]. While regulation via translation is a relatively slow process, gating and trafficking can change water permeability of a membrane within seconds. Trafficking of aquaporins was first described in AQP2 in mammals where it is involved in concentrating urine in the kidneys [14]. AQP2-trafficking is controlled by a signaling cascade triggered by the neuropeptide arginine-vasopressin. The phosphorylation of conserved serine and threonine residues in the fourth loop region of AQP2 caused the redistribution of intracellular AQP storage vesicles to the plasma membrane. This resulted in a rapid increase of water permeability of the membrane. The mechanism of how AQP phosphorylation leads to the recognition of the flagged protein and subsequent vesicle movement and membrane fusion is still unknown. Gating refers to the ability of AQP's to control the flux of water by widening or constricting the channel. X-ray structures have revealed that the width of a fully opened channel can allow a single water molecule entry [13].

While vertebrate AQP's are well studied, few studies have been conducted on invertebrate AQP's (reviewed by Spring et al., 2009 [15]). DRIP (Genbank accession #: CG9023) is a partly characterized representative of *Drosophila* AQP's. It is expressed in embryonic and adult MT's of the fruit fly [16]. A mosquito AQP, a close homologue to *Drosophila* DRIP, has been cloned and characterized in the yellow fever mosquito *Ae. aegypti*. This AQP is localized in tracheolar cells associated with MT's of adult female *Ae. aegypti* mosquitoes [17]. Another mosquito aquaporin has recently been cloned and characterized from *Anopheles gambiae*. This aquaporin is a water transporter and expressed in multiple tissue and specifically in the stellate cells of MT's. It is important for water homeostasis in *An. gambiae* [18].

In order to evaluate the potential of this class of molecule as targets for vector control strategies, we have surveyed the genome of *Ae. aegypti* and identified six genes encoding putative AQP's. We show that four AQP's are expressed in the Malpighian tubules of adult females and that knockdown of three of them affects diuresis.

Results

The AQP genes of *Ae. aegypti*

BLAST searches conducted with vertebrate and *Drosophila* AQP's as query sequences allowed the identification of six loci encoding putative AQP's in the genome of *Ae. aegypti*, which were denominated *Ae. aegypti* aquaporins 1–6 (Table S1; AaAQP1–6). In

addition we identified seven putative AQP's encoded in the genome of the malaria mosquito *Anopheles gambiae* [19], eight in the *Drosophila* genome [20], six in the genomic sequence of the head louse *Pediculus humanus* [21], and seven in the genome of the red flour beetle *Tribolium castaneum* [22,23]. We also used five putative AQP of *Leishmania major*, six *Plasmodium*, and two fungal sequences of the yeast *Pichia pastoris* for our analysis.

A phylogenetic tree was constructed to indicate associations among the identified protein sequences (Figure 1). AaAQP1 forms a clade with three proteins from the other insect species and with *Drosophila* DRIP. DRIP represents a typical water transporter [16]. AaAQP2 forms a well defined orthologous clusters with other insect AQP's. AaAQP1, 2, and 3 share a clade with well characterized vertebrate AQP subfamily members that mediate water transport. Within this clade AaAQP3 forms a separate cluster together with *Drosophila* BIB (CG4722), and two *Anopheles* AQP's.

AaAQP4 and AaAQP5 belong to two separate insect-specific clades. Mosquitoes have single representatives in each of these clusters, whereas other insects have apparent gene duplications. There are three relatives of AaAQP5 in fruit fly and four in the flour beetle. AaAQP6 is closely related to *Drosophila* CG12251 and vertebrate AQP's 11 and 12.

Expression of AQP's in adult female *Ae. aegypti*

First, we analyzed AQP expression data from microarray assays performed with probes from RNAs isolated from whole mosquito females (Figure 2). As a general trend we found AQP expression down-regulated 12 and 24 h after a blood meal. At the later time points, expression returns to “non blood fed (NBF)” levels. An exception was AaAQP5 which was up regulated 12h PBM but is also down regulated at later time points.

Next, we determined AQP expression in selected organs and body parts of adult female mosquitoes before a blood meal, 3 and 24 h PBM. Using quantitative RT-PCR, we found that all six *Ae. aegypti* AQP's were expressed in adult female mosquitoes (Figure 3). Different organs/body parts varied considerably in the assortment of AQP's they expressed. The observed patterns of AQP expression were as follows:

AaAQP1. This AQP is the mosquito homologue of *Drosophila* DRIP, a highly selective water-specific channel in the fruit fly [16]. AaAQP1 was expressed in all organs and body parts examined with highest expression levels in the MT's. It is significantly up regulated in the MT's 3 h after a blood meal.

AaAQP2. We found high AaAQP2 transcript levels in all organs and body parts except the fat body. It was strongly expressed in the MT's, midgut, and ovaries at all time points. The overall expression pattern during vitellogenesis is similar to AaAQP1. It is significantly up regulated in the MT's 3 h after a blood meal.

AaAQP3. This AQP is the homologue of the *Drosophila* BIB. BIB does not function as a water channel in fruit flies but is involved in the regulation of cell adhesion [24]. AaAQP3 was weakly expressed in MT's, midgut, and ovaries and was significantly up regulated in ovaries and fat body PBM.

AaAQP4. This uncharacterized AQP was highly expressed in MT's and is significantly up regulated 3 h after a blood meal. It is down regulated in the midgut during early vitellogenesis.

AaAQP5. This uncharacterized AQP was highly expressed in all organs except the ovaries and was up regulated in most organs PBM.

AaAQP6. This uncharacterized AQP is predominantly expressed in the thorax. The highest mRNA expression levels were found in the foregut (Figure 3B).

each sequence pair. There were a total of 236 positions in the final dataset. Evolutionary analyses were conducted in MEGA4 [46]. Initial sequence alignment was completed using PROMALS3D server (PROfile Multiple Alignment with predicted Local Structures and 3D constraints) [37]. The tree is drawn using FigTree software to an approximate 3.7 Bya-long scale with relative branch lengths used to infer the tree. AQPs from different species are color-coded. Confirmed water transporters are labeled with a water molecule, confirmed aquaglyceroporins are labeled with a glycerol molecule. doi:10.1371/journal.pone.0015578.g001

Effect of AQP knockdown on diuresis after PBS injection

RNAi-mediated gene expression knockdown was determined in samples of total RNA isolated from MTs dissected three days after injection of AQP dsRNA. Knockdown was successful in the mosquito MTs with efficiencies between 95% (AaAQP5) and 60% (AaAQP2) reduction in transcript accumulation (Figure 4A).

In order to determine how fast PBS-injected mosquitoes start discharging urine, we injected female mosquitoes with 1.25 ul PBS and observed them under a stereomicroscope to determine the time period till the onset of urine discharge. Injected females started to discharge urine droplets on average at 157 s (SE = ± 16 s) after injection, thus leaving a window of two minutes to perform the first weight measurement after injection. Excretion was strongest in the first hour after injection (Figure 4B, control group, blue line).

To test the hypothesis that the AQP proteins are contributors to the function of the MT, we generated knockdown mosquitoes for each MT-expressed AQP transcript by dsRNA injection and tested for urine production using our in vivo diuresis assay (Figure 4C). Negative control mosquitoes (injected with a dsRNA directed against the unrelated jellyfish protein eGFP) excreted 39% of the injected 1.25 ul PBS within one hour. In contrast, positive control mosquitoes injected with 1.25 ul of 200 μ M HgCl₂ in PBS (known to suppress AQP activity [13]) excreted only 10% in the first hour after injection.

AaAQP1, 4, and 5 knockdown mosquitoes showed significantly reduced excretion rates compared to the eGFP dsRNA injected control (Figure 4C). Knockdown of AaAQP2 did not result in a significant effect. The strongest effect of a single gene knockdown was observed after knocking down AaAQP5 which resulted in a decline of excretion to only 22%. The combined knockdown of all four MT-expressed AQPs reduced excretion to only 18% of the injected PBS.

In order to visualize the effect of the combined knockdown on diuresis we injected control and combined knockdown mosquitoes with 2.5 ul PBS. Figure 4D shows two representatives of each group 3 h after injection.

Discussion

This report is the first comprehensive study on AQP genes and their expression in the adult yellow fever mosquito, *Aedes aegypti*. We specifically focused on AQPs expressed in the MTs, an insect organ, specialized in water and waste excretion.

An interesting result of our phylogenetic analysis (see Figure 1) is that we were unable to identify any typical aquaglyceroporin in dipteran insects. Human aquaglyceroporins, the *Homo sapiens* AQPs 3, 7, 9, and 10, form a separate clade with six *Plasmodium*, one *Leishmania*, one fungal and one louse AQP. A multiple sequence alignment of all AQPs used for our analysis (Figure S1) revealed that





Figure 2. Six *Ae. aegypti* AQPs are expressed in adult female mosquitoes. Overall expression of AaAQPs in adult female mosquitoes during vitellogenesis. Expression was determined by microarray analysis. Six replicates were analyzed, and their mean separated by Tukey-Kramer HSD ($p < 0.05$). Means which share the same letter are not significantly different. doi:10.1371/journal.pone.0015578.g002

A



B




Figure 3. The six *Ae. aegypti* AQP organ/body part expression patterns are highly diverse. A. Organ/body part expression patterns. Expression was assayed with q-RT-PCR. The results shown are representative of three separate repeats with similar results. RNA was isolated from organs/body parts of four groups of 20 mosquitoes. MT – Malpighian tubules, MG – posterior midgut, OV – ovaries, TX - thorax, FB – abdominal body wall with fat body. Three replicates were analyzed, and their mean separated by Tukey–Kramer HSD ($p < 0.05$). Means which share the same letter are not significantly different. **B.** Thorax tissue/organ expression pattern of unfed mosquitoes. Relative expression levels were determined via q-RT-PCR.

Expression values were normalized with ribosomal protein S7 expression data from the same samples. SG – salivary glands, FG – foregut, MS – wing muscle, LG – legs, WG – wings. Four replicates were analyzed, and their mean separated by Tukey–Kramer HSD ($p < 0.05$). Means which share the same letter are not significantly different. doi:10.1371/journal.pone.0015578.g003

all aquaglyceroporins share a cysteine residue at position six, C-terminal to the NPA motif in the B-loop. None of the dipteran AQPs in our analysis had a cysteine at this position. The fact that dipteran insects do not appear to possess an AQP-like glycerol transporter raises the question as to how they transport glycerol over cell membranes. Glycerol plays an important role in insect cold tolerance and diapause and is found in high concentration in the hemolymph of diapausing insects [25].

The microarray expression data was created with RNA isolated from total mosquitoes (Figure 2) and shows that AQP expression was generally down regulated after a blood meal, even at 3 h PBM. After taking a blood meal mosquitoes seek out a resting place to digest the blood and perform vitellogenesis and egg development. This resting phase extends over a period of approximately two days. During this time mosquitoes don't take up water, therefore high AQP expression levels are not necessary.

AQP expression show specific patterns in distinct organs/body parts (Figure 3). All examined organs/body parts differ in their AQP expression patterns. We were specifically interested in patterns from midgut and MTs since blood meal-derived water has to cross the midgut epithelium and subsequently the MT epithelium for excretion. AQPs 1, 2, 4, and 5 were expressed in the midgut. Interestingly, three AQPs expressed in the midgut were down regulated at the 3 h PBM time point, corresponding to the time at which the bulk of blood meal-derived water has already been excreted. Our data suggests that AQPs 1, 4, and 5 are the principal AQPs in the MTs of adult females.

We developed an *in vivo* diuresis assay for adult yellow fever mosquito females based on PBS injection. PBS is a non-toxic isotonic buffer containing sodium chloride, sodium phosphate, potassium chloride and potassium phosphate. This novel assay has several advantages over classical blood meal-based assays. Firstly,




Figure 4. Three AQPs regulate water excretion in *Ae. Aegypti*. **A.** Knockdown efficiency. AQP ds-RNA was injected in mosquitoes and transcript accumulation determined at three days post injection. eGFP ds-RNA-injected mosquitoes were used as a control and AQP expression in these mosquitoes was set as 100%. Five groups of ten knockdown mosquitoes were analyzed for each group and the data is expressed as average and standard error. **B.** Weight profiles after PBS injection in AQP knockdown and control mosquitoes. Three groups of thirty mosquitoes of a control group (blue) and AQP knockdown group (red, combined knockdown of AQP 1, 2, 4, 5) were injected with 1.25 ul PBS. Average weight was determined before injection (0 point) and in 30 min intervals after injection. **C.** Effect of AQP knockdown on excretion in PBS-injected mosquitoes. Five groups of ten mosquitoes were analyzed for each control and experimental sample. RNAi knockdown was induced three days before the experiment. Replicates were analyzed, and their mean separated by Tukey–Kramer HSD ($p < 0.05$). Bars labeled with the same letter are not significantly different. **D.** Visualization of knockdown effect. *Ae. aegypti* females; 3 h after injection of 2.5 ul PBS. The two mosquitoes on the left are representatives of 20 control mosquitoes that received eGFP dsRNA injections. The two mosquitoes on the right are representatives with combined knockdown of AQP1, 2, 4, and 5. doi:10.1371/journal.pone.0015578.g004

mosquitoes receive a standardized amount of PBS, while blood meal sizes can vary significantly. Secondly, since *Ae. aegypti* mosquitoes start secreting urine about 30 s after starting a blood meal [1,26], the determination of blood meal sizes is difficult. Applying our method, mosquitoes injected with PBS started secreting urine after 2 min, which allowed accurate measurement of their weight after injection. As mentioned above, mosquito diuresis is controlled by neuropeptide hormones that are secreted from the central nervous system in response to unknown stimuli associated with blood ingestion. PBS injection triggers this response efficiently. The rapid onset of diuresis after injection suggests that AQP activity in the MT is regulated by trafficking of AQP-containing vesicles to the plasma membrane, analogous to the processes described for the renal collection duct in human kidney [27].

RNAi is a powerful tool used to study gene function in mosquitoes. We successfully employed RNAi-mediated knockdown for the analysis of AQP function in the MT. Knockdown rates, as determined by real time PCR, were between 60 and 95% and therefore well inside the range that can be achieved by dsRNA transfection in cell culture [28]. Similar knockdown efficiencies have been found in whole mosquitoes [29].

A point of concern is that the AQP dsRNA injections likely resulted in AQP knockdown in tissues other than the MTs. However, excreted watery liquids have been shown to always pass through the MT in adult mosquitoes [1,5,30]. Therefore we expect the effects of AQP knockdown in other tissues to be negligible for the outcomes of our *in vivo* diuresis experiments.

As a positive control for successful AQP inhibition we used mercury ions (Hg^{2+}), which are capable of binding with a cysteine and an alanine within the AQP pore, thus obstructing water transport through the channel. Mosquitoes injected with 200 μM $HgCl_2$ were still able to excrete 10% of the injected fluid in one hour. This might be due to paracellular permeability, water transport through the space between the cells, which has been described in insect MTs and can be enhanced by kinins [26].

Using RNAi knockdown mosquitoes we confirmed the function of three different AQPs in the MTs of *Ae. aegypti*. While RNAi control mosquitoes were able to excrete about 40% of the injected fluid in one hour, knockdown of single AQPs (AaAQP1, 4, or 5) resulted in a significant decrease in excretion. Simultaneous knockdown of all four MT-expressed AQPs reduced excretion down to 18% of the injected fluid in one hour, indicating that a combination of AaAQPs 1, 4, and 5 performs water transport in the MTs of female *Ae. aegypti*.

The redundancy of function we have observed here has also been noted in human kidney where seven different AQP proteins are expressed [31]. One plausible explanation would be that several AQP genes with different promoters allow the fine regulation of AQP expression in stage- and tissue specific manner or after a range of stimuli.

There is a great need for the development of novel, effective insecticides to fight insect vectors, since the public health insecticides currently in use are based on only a limited number of active compounds [32]. Because of their vital importance in insect larvae and adult homeostasis, insect AQPs could become targets for the development of novel insecticides. The study presented here has identified six genes encoding putative AQP membrane transporters in *Ae. aegypti* and demonstrated the functional role of three of them in regulation of water transport. Further analysis of these AQPs and their regulation has the potential to contribute to the future development of novel anti-vector strategies.

Materials and Methods

Mosquito rearing

The *Ae. aegypti* mosquito strain UGAL was maintained in laboratory culture as has been previously described by Hays and Raikhel [33]. The strain was reared at a temperature of 28°C with 80% humidity and a photoperiod of 14 h light and 10 h dark. Larvae were fed on a diet of ground rat food, yeast and albumin (1:1:1 w/w).

Sequence Identification & Phylogenetic analysis

Predicted cDNA and deduced AA sequences of AQP family members were identified using BLAST at three databases: Ensembl [34], Genbank [35], and VectorBase [36]. A sequence alignment (supplemental online material: Figure 1) was performed using PROMALS3D software which considers structural constraints for the divergent AQPs protein sequences [37]. The reconstruction of the phylogenetic tree was performed using Mega 4 [38]. The tree was visualized using FigTree software [39].

Microarray Expression Studies

Three biological samples composed of four day old mosquitoes were collected consisting of a pool of 20 insects. Four day old females were blood fed and three biological samples (20 females each) were collected at several times after the blood meal. Total RNA was extracted with TRIzol® (Invitrogen, Carlsbad, CA) and further purified with Qiagen RNeasy columns (Qiagen, Valencia, CA) with DNase treatment according to the manufacturer's recommendations. RNA quality was assessed by capillary electrophoresis using the Agilent Bioanalyzer 2100 and spectrophotometric analysis. The RNA was reverse transcribed and amplified using WT Ovation Pico (Cat# 3300-60, NuGen, San Carlos, CA) and 2 g of this cDNA was labeled with Cy3 (Cat# 5190-1305, Agilent technologies, Santa Clara, CA) according to the Agilent Oligonucleotide Array-Based CGH For Genomic DNA Analysis Protocol (Cat# G4410-90010, Agilent technologies, Santa Clara, CA). Hybridization, washing, and scanning were processed as per the Agilent OneColor Microarray-Based Gene Expression Analysis Protocol (Cat# G4140-90040, Agilent technologies, Santa Clara, CA). Expression signals were normalized for background within chips with the Agilent spatial correction algorithm and were normalized between chips using quantile normalization. The data is available at Gene Expression Omnibus (GEO, www.ncbi.nlm.nih.gov/geo/) under series record GSE22339 [40]. Statistical analysis was performed with InStat (GraphPad Software, La Jolla, CA).

qRT-PCR Expression Studies

Gene-specific primers were developed using Primer BLAST [41]. Total RNA was obtained from different larval stages, pupae, and adult females with TRIzol® solution. Tissue-specific RNAs were isolated after dissection of samples from 30 individual mosquitoes including previtellogenic females 72 h after eclosion and female mosquitoes 3 h and 24 h post blood meal. Transcript was analyzed and quantified using quantitative RT-PCR (qPCR) using iQ Supermix (Biorad, Hercules, CA). Primers were as follows: AaAQP1f: ACC GGC ATC AGA AAG GAG AAG CG; AaAQP1 r: GCC TGC TGT TTG ATG TGT TGT GCA; AaAQP2 f: GCT CGC TCG TTT GGA ACG GC; AaAQP2 r: CAC GGT AGC GCT CTG AGG CG; AaAQP3 f: AGG TCC AGT GGG GAT GGC CC; AaAQP3 r: CAG CTG AGG TGG TGG CGG TG; AaAQP4 f: CTG CCG CCT GCA GTG TGG AA; AaAQP4 r: TGT GGA AGT TCT CGT CGG AAG ACG T; AaAQP5 f: CGG TGT TCA GGC GCG AGG TT; AaAQP5

r: ATC CCG CGT TGG TGG AAC GG; AaAQP6 f: CAG CTT GGT CGT GGC CGC AT; AaAQP6 r: GCA CAG CTC CCG CAC ACG AT.

RNAi knockdown experiments

Generation of double-stranded RNAs (dsRNA) was performed as described earlier [42] using primers with the T7 primer sequences attached. PCR product was used as template for dsRNA synthesis with the MEGAscripts T7 Kit (Ambion, Austin, Tx). Approximately 500 ng of dsRNA in 268 nl of pyrogen-free H₂O was injected into the thorax of CO₂-anesthetized female mosquitoes three days after adult emergence. The injected mosquitoes were then allowed to recover for 3 days before diuresis assay. Knockdown effectiveness was confirmed with qPCR. Primers were as follows (The T7 sequence is omitted): AaAQP1f: TAA TAC GAC TCA CTA TAG GGA GCA CTA TGG GCT GGG GCG GAG ACT; AaAQP1 r: TAA TAC GAC TCA CTA TAG GGA CGG CTG GTC CGA AAG AGC GAG CTG; AaAQP2 f: TAA TAC GAC TCA CTA TAG GGA CTG CTG GCT TAC TTG CGG CTG GCA; AaAQP2 r: TAA TAC GAC TCA CTA TAG GGG CTA CAC GGT AGC GCT CTG AGG CGG; AaAQP3 f: TAA TAC GAC TCA CTA TAG GGC CCC ATC CCC AAG CGG GTG AAC CAC; AaAQP3 r: TAA TAC GAC TCA CTA TAG GGT GGG CCA TTG GGTAGC CCC CTG GAT; AaAQP4 f: TAA TAC GAC TCA CTA TAG GGG CGG CAT CGG GTT CGG CTT CAC AGT; AaAQP4 r: TAA TAC GAC TCA CTA TAG GGT GGC CGG GTT CAT ACT CGC TCC GGT; AaAQP5 f: TAA TAC GAC TCA CTA TAG GGT ACG TTG CGG CCC AGT GCA TCG GAG; AaAQP5 r: TAA TAC GAC TCA CTA TAG GGG GAA CCT CGC GCC TGA ACA CCG TCT; AaAQP6 f: TAA TAC GAC TCA CTA TAG GGG ATC GCG GCA GTT GCT CGC CGA GTG; AaAQP6 r: TAA TAC GAC TCA CTA TAG GGA GGC CAA GCG ACA GCA CAG AGT GGC.

In vivo diuresis assay

Knockdown and control (eGFP dsRNA-injected) mosquitoes were anesthetized with CO₂ and pooled into groups of ten. Group

weight was determined on a precision weighing balance. Each mosquito was subsequently injected with 1.25 ul of PBS and the groups were immediately weighed. Diuresis was monitored by weighing mosquitoes in 30 min intervals. Control mosquitoes were injected with 1.25 ul 200 uM HgCl₂ in PBS. After the last weighing the MTs were dissected and RNA was isolated in order to determine the knockdown efficiency.

Supporting Information

Figure S1 Comprehensive alignment of 72 aquaporin family members. Sequences are from *Ae. aegypti*, *Anopheles gambiae*, *Drosophila melanogaster*, *Pediculus humanus*, *Tribolium castaneum*, *Homo sapiens*, *Leishmania major*, *Plasmodium sp.*, and *Pichia pastoris*. Darker patterns represent higher sequence similarity. (PDF)

Table S1 *Ae. aegypti* aquaporins. ¹ AAEL – VectorBase; all others – NCBI Genbank; ² [17]; ³ The BIB cluster is duplicated in the *Anopheles* genome; ⁴ This cluster is expanded in the fruit fly genome. (DOC)

Acknowledgments

The authors thank the participants of the NMSU BIOL302, Molecular Biology Techniques Laboratory, Spring 2010 class: Natalie Aarons-Cooke, Jordan Basham, Hannah Cox-Jesko, Lynda Davis, Hui Fang, Elizabeth Gamez, Anna Garliss, David Hogan, Kelsey Kalbacher, Kevin Lee, Kaitlyn Norman, Shannon Pease, Katarina Provenghi, Marisol Resendiz, Cherish Skeen, Mathew Steritz, and Mary Topalowski for their help with sequence identification and developing primers used in this project. We thank Jessica Aguirre for technical support.

Author Contributions

Conceived and designed the experiments: IAH DYB OM. Performed the experiments: LLD VKC DYB OM. Analyzed the data: IAH LLD DYB OM ALD. Contributed reagents/materials/analysis tools: DYB OM ALD IAH. Wrote the paper: LLD IAH.

References

- Clements AN, ed (1992) The Biology of Mosquitoes. London: Chapman & Hall.
- Jagge CL, Pietrantonio PV (2008) Diuretic hormone 44 receptor in Malpighian tubules of the mosquito *Aedes aegypti*: evidence for transcriptional regulation paralleling urination. *Insect Mol Biol* 17: 413–426.
- Predel R, Wegener C (2006) Biology of the CAPA peptides in insects. *Cell Mol Life Sci* 63: 2477–2490.
- Schepel SA, Fox AJ, Miyauchi JT, Sou T, Yang JD, et al. (2010) The single kinin receptor signals to separate and independent physiological pathways in Malpighian tubules of the yellow fever mosquito. *Am J Physiol Regul Integr Comp Physiol*.
- Boorman JP (1960) Observations on the feeding habits of the mosquito *Aedes (Stegomyia) aegypti* (Linnaeus): the loss of fluid after a blood-meal and the amount of blood taken during feeding. *Ann Trop Med Parasitol* 54: 8–14.
- Furukawa K, Sidoux-Walter F, Hohmann S (2009) Expression of the yeast aquaporin Aqp2 affects cell surface properties under the control of osmoregulatory and morphogenic signalling pathways. *Mol Microbiol* 74: 1272–1286.
- Nishihara E, Yokota E, Tazaki A, Orii H, Katsuhara M, et al. (2008) Presence of aquaporin and V-ATPase on the contractile vacuole of *Amoeba proteus*. *Biol Cell* 100: 179–188.
- Petterson N, Filipsson C, Becit E, Brive L, Hohmann S (2005) Aquaporins in yeasts and filamentous fungi. *Biol Cell* 97: 487–500.
- Jensen MO, Tajkhorshid E, Schulten K (2001) The mechanism of glycerol conduction in aquaglyceroporins. *Structure* 9: 1083–1093.
- Gonen T, Walz T (2006) The structure of aquaporins. *Q Rev Biophys* 39: 361–396.
- Jung JS, Preston GM, Smith BL, Guggino WB, Agre P (1994) Molecular structure of the water channel through aquaporin CHIP. The hourglass model. *J Biol Chem* 269: 14648–14654.
- Nejsum LN, Kwon TH, Jensen UB, Fumagalli O, Frokiaer J, et al. (2002) Functional requirement of aquaporin-5 in plasma membranes of sweat glands. *Proc Natl Acad Sci U S A* 99: 511–516.
- Tornroth-Horsefield S, Hedfalk K, Fischer G, Lindkvist-Petersson K, Neutze R (2010) Structural insights into eukaryotic aquaporin regulation. *FEBS Lett* 584: 2580–2588.
- Nedvetsky PI, Tamma G, Beulshausen S, Valenti G, Rosenthal W, et al. (2009) Regulation of aquaporin-2 trafficking. *Handb Exp Pharmacol*. pp 133–157.
- Spring JH, Robichaux SR, Hamlin JA (2009) The role of aquaporins in excretion in insects. *J Exp Biol* 212: 358–362.
- Kaufmann N, Mathai JC, Hill WG, Dow JA, Zeidel ML, et al. (2005) Developmental expression and biophysical characterization of a *Drosophila melanogaster* aquaporin. *Am J Physiol Cell Physiol* 289: C397–407.
- Pietrantonio PV, Jagge C, Keeley LL, Ross LS (2000) Cloning of an aquaporin-like cDNA and in situ hybridization in adults of the mosquito *Aedes aegypti* (Diptera: Culicidae). *Insect Mol Biol* 9: 407–418.
- Liu KCTH, HuangXiaoyan, ChaSung-Jac, Agre P, Rasgon J (2010) Poster Abstract LB-2046: Cloning and characterization of *Anopheles gambiae* aquaporin 1 (AgAQP1); ASTMH fifty-ninth annual meeting, Atlanta, GA. Published by the American Society of Tropical Medicine and Hygiene.
- Holt RA, Subramanian GM, Halpern A, Sutton GG, Charlab R, et al. (2002) The genome sequence of the malaria mosquito *Anopheles gambiae*. *Science* 298: 129–149.
- Adams MD, Celniker SE, Holt RA, Evans CA, Gocayne JD, et al. (2000) The genome sequence of *Drosophila melanogaster*. *Science* 287: 2185–2195.
- Pittendrigh BR, Clark JM, Johnston JS, Lee SH, Romero-Severson J, et al. (2006) Sequencing of a new target genome: the *Pediculus humanus humanus* (Phthiraptera: Pediculidae) genome project. *J Med Entomol* 43: 1103–1111.

22. Kim HS, Murphy T, Xia J, Caragea D, Park Y, et al. (2010) BeetleBase in 2010: revisions to provide comprehensive genomic information for *Tribolium castaneum*. *Nucleic Acids Res* 38: D437–442.
23. Wang L, Wang S, Li Y, Paradesi MS, Brown SJ (2007) BeetleBase: the model organism database for *Tribolium castaneum*. *Nucleic Acids Res* 35: D476–479.
24. Tatsumi K, Tsuji S, Miwa H, Morisaku T, Nuriya M, et al. (2009) *Drosophila* big brain does not act as a water channel, but mediates cell adhesion. *FEBS Lett* 583: 2077–2082.
25. Yoder JA, Benoit JB, Denlinger DL, Rivers DB (2006) Stress-induced accumulation of glycerol in the flesh fly, *Sarcophaga bullata*: evidence indicating anti-desiccant and cryoprotectant functions of this polyol and a role for the brain in coordinating the response. *J Insect Physiol* 52: 202–214.
26. Wang S, Rubenfeld A, Hayes T, Beyenbach K (1996) Leucokinin increases paracellular permeability in insect Malpighian tubules. *J Exp Biol* 199: 2537–2542.
27. Chou CL, Yip KP, Michea L, Kador K, Ferraris JD, et al. (2000) Regulation of aquaporin-2 trafficking by vasopressin in the renal collecting duct. Roles of ryanodine-sensitive Ca^{2+} stores and calmodulin. *J Biol Chem* 275: 36839–36846.
28. Shepard AR, Jacobson N, Clark AF (2005) Importance of quantitative PCR primer location for short interfering RNA efficacy determination. *Anal Biochem* 344: 287–288.
29. Souza-Neto JA, Sim S, Dimopoulos G (2009) An evolutionary conserved function of the JAK-STAT pathway in anti-dengue defense. *Proc Natl Acad Sci U S A* 106: 17841–17846.
30. Redington BC, Hockmeyer WT (1976) A method for estimating blood meal volume in *Aedes aegypti* using a radioisotope. *J Insect Physiol* 22: 961–966.
31. Ikeguchi M (2009) Water transport in aquaporins: molecular dynamics simulations. *Front Biosci* 14: 1283–1291.
32. van den Berg H (2009) Global status of DDT and its alternatives for use in vector control to prevent disease. *Environ Health Perspect* 117: 1656–1663.
33. Hays AR, Raikhel AS (1990) A novel protein produced by the vitellogenic fat body and accumulated in mosquito oocytes. *Development Genes and Evolution* 199: 114–121.
34. EMBL-EBI (2010) Ensembl *Aedes aegypti*; http://metazoa.ensembl.org/Aedes_aegypti/Info/Index. (Accessed July 2010).
35. NCBI (2010) Genbank; <http://www.ncbi.nlm.nih.gov/Genbank>. (Accessed July 2010).
36. NIAID (2010) Vectorbase; <http://aaegypti.vectorbase.org/>. (Accessed July 2010).
37. Pei J, Kim BH, Grishin NV (2008) PROMALS3D: a tool for multiple protein sequence and structure alignments. *Nucleic Acids Res* 36: 2295–2300.
38. Larkin MA, Blackshields G, Brown NP, Chenna R, McGettigan PA, et al. (2007) Clustal W and Clustal X version 2.0. *Bioinformatics* 23: 2947–2948.
39. Rambaut A FigTree; <http://tree.bio.ed.ac.uk/software/figtree/>. (Accessed June 2010).
40. Dissanayake SN, Ribeiro JM, Wang MH, Dunn WA, Yan G, et al. (2010) aeGEPUCI: a database of gene expression in the dengue vector mosquito, *Aedes aegypti*. *BMC Res Notes* 3: 248.
41. NCBI (2009) PrimerBlast. <http://www.ncbi.nlm.nih.gov/tools/primer-blast/> (Accessed July 2010).
42. Attardo GM, Hansen IA, Shiao SH, Raikhel AS (2006) Identification of two cationic amino acid transporters required for nutritional signaling during mosquito reproduction. *J Exp Biol* 209: 3071–3078.
43. Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4: 406–425.
44. Efron B, Halloran E, Holmes S (1996) Bootstrap confidence levels for phylogenetic trees. *Proc Natl Acad Sci U S A* 93: 7085–7090.
45. Zuckerkandl E, Pauling L (1965) Molecules as documents of evolutionary history. *J Theor Biol* 8: 357–366.
46. Tamura K, Dudley J, Nei M, Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* 24: 1596–1599.