

UC Berkeley

UC Berkeley Previously Published Works

Title

Draft Genome Sequence of an Escherichia coli Sequence Type 420 Isolate from a Patient with Urinary Tract Infection in Northern California

Permalink

<https://escholarship.org/uc/item/52m7109n>

Journal

Microbiology Resource Announcements, 9(23)

ISSN

2576-098X

Authors

Matsui, Yusuke
Hu, Yuan
Tarlton, Nicole J
[et al.](#)

Publication Date

2020-06-04

DOI

10.1128/mra.00251-20

Peer reviewed



Draft Genome Sequence of an *Escherichia coli* Sequence Type 420 Isolate from a Patient with Urinary Tract Infection in Northern California

 Yusuke Matsui,^a Yuan Hu,^b Nicole J. Tarlton,^a Lee W. Riley^a

^aSchool of Public Health, Division of Infectious Diseases and Vaccinology, University of California, Berkeley, Berkeley, California, USA

^bSchool of Public Health, Division of Epidemiology, University of California, Berkeley, Berkeley, California, USA

ABSTRACT The genome sequence of a uropathogenic *Escherichia coli* sequence type 420 strain isolated from a patient with urinary tract infection in northern California is described here. The draft genome sequence includes a 4.8-Mb chromosome, accompanied by a 114-kb plasmid containing IncFIB/IncFII/Col156 and a 35-kb plasmid containing IncN3.

Extraintestinal pathogenic *Escherichia coli* (ExPEC) belonging to multilocus sequence type 420 (ST420) is often reported as a low-prevalence genotype (1, 2), but it has been identified sporadically from patients with community-acquired urinary tract infection (UTI) at a college campus in northern California (3–5). Interestingly, all of those ST420 isolates were found to be susceptible to all antimicrobial agents tested (pan-susceptible) (4, 5). Here, we present the genome sequence of an ST420 ExPEC strain isolated from one of the patients (a 39-year-old female) diagnosed with UTI at the university health center in 2003 (4). In this paper and in GenBank, this ST420 isolate is designated strain IT0021. The genome sequence was determined for future use as a reference sequence.

We isolated *E. coli* from a urine sample demonstrating a viable bacterial count of $>10^4$ CFU/ml, as described previously (4, 6). *E. coli* was subcultured from a frozen glycerol stock and grown overnight in LB broth in a 37°C shaker. The next day, 2 ml of culture was pelleted in a microfuge tube, and the pellet was subjected to DNA extraction. DNA purification for whole-genome sequencing was performed with the DNeasy blood and tissue kit (Qiagen). Preparation of Illumina-compatible libraries for 300-bp paired-end reads was conducted according to a standard protocol (WaferGen Bio-systems). Libraries were sequenced on a MiSeq instrument with MiSeq reagent kit v3 (600-cycle) chemistry. Read trimming and contig assembly and analysis were performed with Geneious v.9.0.5 (Biomatters, Ltd.) (7). RAST was utilized for annotation of the draft genome (8). Plasmids, resistance genes, and putative virulence factor genes were initially identified with Web-based tools from the Center for Genomic Epidemiology (<http://www.genomicepidemiology.org>), and prophages were identified with PHASTER (9). Default parameters were used for all software unless otherwise specified.

We assembled 509,467 of 514,170 reads with the Geneious assembler. We set the threshold value of base call confidence as high quality. The *E. coli* IT0021 draft genome sequence was assembled into 133 contigs >200 bp long, with a mean contig length of 47,293 bp, a maximum contig length of 410,189 bp, and an N_{50} of 89,241 bp. The mean read coverage of the assembled contigs was approximately 25-fold. The contigs comprised a 4,695,334-bp chromosome, a 114,729-bp circular plasmid, and a 35,105-bp circular plasmid. The GC contents were 50%, 51%, and 51%, respectively.

The chromosome contained four prophages, namely, PhiV10, SfiI, mEp460, and lambda. The 114-kb plasmid contained IncFIB/IncFII/Col156 replicons. It closely resem-

Citation Matsui Y, Hu Y, Tarlton NJ, Riley LW. 2020. Draft genome sequence of an *Escherichia coli* sequence type 420 isolate from a patient with urinary tract infection in northern California. Microbiol Resour Announc 9:e00251-20. <https://doi.org/10.1128/MRA.00251-20>.

Editor David Rasko, University of Maryland School of Medicine

Copyright © 2020 Matsui et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Lee W. Riley, lwiriley@berkeley.edu.

Received 11 March 2020

Accepted 14 May 2020

Published 4 June 2020

bled pUTI89* (pSF-166-1, >99% assembled), which was found in an ST95 ExPEC isolate (10). Most ST95 strains are pansusceptible to antibiotics, and the carriage of pUTI89* is statistically significantly associated with antibiotic susceptibility (11). The plasmid carried *senB* (11), which encodes an enterotoxin, but no antibiotic resistance genes. The 35-kb plasmid contained an IncN3 replicon and closely resembled pTRE-131 (>95% of pTRE-131 assembled) (12). It contained no virulence genes or antibiotic resistance genes.

Data availability. This whole-genome shotgun project has been deposited in DDBJ/ENA/GenBank under the accession number [JAAMRS000000000](https://doi.org/10.1093/bioinformatics/bts199). The version described in this paper is version JAAMRS010000000. Raw sequence reads have been deposited in the NCBI Sequence Read Archive under BioProject number [PRJNA607787](https://doi.org/10.1093/bioinformatics/bts199) and run number [SRR1127725](https://doi.org/10.1093/bioinformatics/bts199).

ACKNOWLEDGMENTS

We thank the clinical laboratory staff members of the Tang Center at the University of California, Berkeley-affiliated health care service for their time and support on this project.

This study was supported by the Centers for Disease Control and Prevention (grant BAA200-2016-91939).

REFERENCES

- Fibke CD, Croxen MA, Geum HM, Glass M, Wong E, Avery BP, Daignault D, Mulvey MR, Reid-Smith RJ, Parmley EJ, Portt A, Boerlin P, Manges AR. 2019. Genomic epidemiology of major extraintestinal pathogenic *Escherichia coli* lineages causing urinary tract infections in young women across Canada. *Open Forum Infect Dis* 6:ofz431. <https://doi.org/10.1093/ofid/ofz431>.
- Hertz FB, Nielsen JB, Schønning K, Littauer P, Knudsen JD, Løbner-Olesen A, Frimodt-Møller N. 2016. Population structure of drug-susceptible,-resistant and ESBL-producing *Escherichia coli* from community-acquired urinary tract. *BMC Microbiol* 16:63. <https://doi.org/10.1186/s12866-016-0681-z>.
- Manges AR, Johnson JR, Foxman B, O'Bryan TT, Fullerton KE, Riley LW. 2001. Widespread distribution of urinary tract infections caused by a multidrug-resistant *Escherichia coli* clonal group. *N Engl J Med* 345:1007–1013. <https://doi.org/10.1056/NEJMoa011265>.
- Tarleton NJ, Moritz C, Adams-Sapper S, Riley LW. 2019. Genotypic analysis of uropathogenic *Escherichia coli* to understand factors that impact the prevalence of β -lactam-resistant urinary tract infections in a community. *J Glob Antimicrob Resist* 19:173–180. <https://doi.org/10.1016/j.jgar.2019.03.002>.
- Yamaji R, Rubin J, Thys E, Friedman C, Riley L. 2018. Persistent pandemic lineages of uropathogenic *Escherichia coli* in a college community from 1999 to 2017. *J Clin Microbiol* 56:e01834-17. <https://doi.org/10.1128/JCM.01834-17>.
- Yamaji R, Friedman CR, Rubin J, Suh J, Thys E, McDermott P, Hung-Fan M, Riley LW. 2018. A population-based surveillance study of shared genotypes of *Escherichia coli* isolates from retail meat and suspected cases of urinary tract infections. *mSphere* 3:e00179-18. <https://doi.org/10.1128/mSphere.00179-18>.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <https://doi.org/10.1093/bioinformatics/bts199>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a better, faster version of the PHAST phage search tool. *Nucleic Acids Res* 44:W16–W21. <https://doi.org/10.1093/nar/gkw387>.
- Stephens CM, Adams-Sapper S, Sekhon M, Johnson JR, Riley LW. 2017. Genomic analysis of factors associated with low prevalence of antibiotic resistance in extraintestinal pathogenic *Escherichia coli* sequence type 95 strains. *mSphere* 2:e00390-16. <https://doi.org/10.1128/mSphere.00390-16>.
- Cusumano CK, Hung CS, Chen SL, Hultgren SJ. 2010. Virulence plasmid harbored by uropathogenic *Escherichia coli* functions in acute stages of pathogenesis. *Infect Immun* 78:1457–1467. <https://doi.org/10.1128/IAI.01260-09>.
- Botts RT, Apffel BA, Walters CJ, Davidson KE, Echols RS, Geiger MR, Guzman VL, Haase VS, Montana MA, La Chat CA, Mielke JA, Mullen KL, Virtue CC, Brown CJ, Top EM, Cummings DE. 2017. Characterization of four multidrug resistance plasmids captured from the sediments of an urban coastal wetland. *Front Microbiol* 8:1922. <https://doi.org/10.3389/fmicb.2017.01922>.