

Lawrence Berkeley National Laboratory

Recent Work

Title

NEUTRON TRANSFER BETWEEN BCS NUCLEI

Permalink

<https://escholarship.org/uc/item/531822tj>

Authors

Kleber, M.
Holzwarth, G.

Publication Date

1972-11-01

Submitted to
Nuclear Physics

LAWRENCE
RADIATION LABORATORY

LBL-1293
Preprint c.1

FEB 9 1973

LIBRARY AND
DOCUMENTS SECTION

NEUTRON TRANSFER BETWEEN BCS NUCLEI

M. Kleber and G. Holzwarth

November 1972

Prepared for the U.S. Atomic Energy
Commission under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



LBL-1293
c.1

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

NEUTRON TRANSFER BETWEEN BCS NUCLEI*

M. Kleber[†]

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

and

G. Holzwarth[†]

Massachusetts Institute of Technology
Cambridge, Massachusetts 02139

November 1972

Abstract

Within perturbation theory we present a microscopic approach to calculate neutron transfer in the scattering of two BCS nuclei. The BCS Hamiltonian of the two coupled BCS nuclei is constructed in terms of commuting field operators. Explicit expressions for the transfer cross sections for a single neutron and for two neutrons are obtained. As an example we consider the scattering of two mercury ions below the Coulomb barrier. The limits of perturbation theory are investigated.

*Work performed under the auspices of the U. S. Atomic Energy Commission.

[†]On leave of absence from the Physik Department der Technischen Universität München, Munich, West Germany.

Neutron Transfer in Reactions Between Superfluid Nuclei

1. Introduction

In the near future scattering experiments with heavy ions will be performed in a wide energy range. A large amount of information will be obtained and has already been obtained¹⁻²⁾ by varying projectile, target, and scattering conditions. For bombarding energies below the Coulomb barrier the relative motion between a heavy projectile and a heavy target follows³⁻⁵⁾ a classical Rutherford trajectory except at backward scattering: This case excluded, the de Broglie wave length λ of the relative motion is small compared to the nuclear radii. Furthermore, the gradient of λ is small⁶⁾ compared to unity. For energies above the Coulomb barrier the second statement is not met because of strong nuclear absorption in the overlap region between projectile and target. This leads⁷⁾ to diffraction scattering and the gross structure of the angular distribution shows a diffraction pattern, predominantly of Fresnel type.

A theoretical treatment of the transition region where the incident energy is close to the Coulomb barrier should be accessible from both sides although the approach from the low energy side seems to be easier. Treating the nuclear distortion between the two heavy ions in perturbation theory one still can use the concept of a Rutherford trajectory. Nuclear distortion leads^{4,5)} to an optical potential which, of course, depends on the reactions one has in mind. The analysis⁸⁾ of the optical potential is difficult if no data on the elastic scattering are available. In this case one is forced to start from the (model) Hamiltonian and to calculate explicitly the coupling

of the two nuclei. In this paper we investigate the coupling of two superconducting nuclei where an enhanced transfer is expected because of static pair correlations.

When the nuclei approach each other their wave functions begin to overlap and the two sets of their respective field operators commute no longer. One must construct new orthogonal states^{5,9)} which depend on the relative distance and on the relative velocity. In section 2 we show how the Hamiltonian is expressed in terms of the new operators up to second order in the overlap of the wave functions. Explicit expressions for the pairing force Hamiltonian are obtained. This model Hamiltonian is used in section 3 to calculate the transfer amplitudes for a single neutron and for two neutrons. We also evaluate the elastic channel, because, if elastic scattering turns out to be very different from the Rutherford cross section then nuclear distortion can no longer be treated by perturbation theory. In section 4 we consider the various approximations for the evaluation of the transfer amplitudes. We also show the results for the scattering of two mercury ions. Section 5 contains a discussion together with some remarks on the kinematics.

An important question is the interference between Coulomb excitation and transfer. In heavy ion reactions the Coulomb field is much too strong¹⁰⁾ to allow Coulomb excitation to be treated in perturbation theory. There are, however, indications¹¹⁾ that Coulomb excitation can be rather small when the transfer takes place. This is possible because of the destructive interference between nuclear forces and the Coulomb force. Nevertheless, the transfer into excited collective states can be calculated only if Coulomb excitation is

taken into account. However, for transfer reactions into ground states or into non-collective excited states Coulomb excitation may not be so important. Having those states in mind we consider in the following the coupling of two almost identical superconducting nuclei and we neglect Coulomb excitation.

2. Orthonormal States for Neutrons

The undisturbed eigenstates $|\Phi_\mu\rangle$ of nucleus A are not orthogonal to the states $|\chi_\nu\rangle$ of nucleus B because the two sets of wave functions overlap. Having transfer reactions in mind we confine ourselves to states $|\Phi_\mu\rangle$ near the fermi levels. If the internucleus distance r (see fig. 1) is not too small we can expand the corresponding orthonormal states $|\bar{\Phi}_\mu\rangle$ and $|\bar{\chi}_\nu\rangle$ in terms of the overlap between the undisturbed states

$$|\bar{\Phi}_\mu\rangle = |\Phi_\mu\rangle - 1/2 \sum_{\pi} \epsilon_{\pi\mu}^* |\chi_\pi\rangle + 3/8 \sum_{\pi\lambda} \epsilon_{\pi\lambda} \epsilon_{\pi\mu}^* |\phi_\lambda\rangle - \dots \tag{1}$$

$$|\bar{\chi}_\nu\rangle = |\chi_\nu\rangle - 1/2 \sum_{\lambda} \epsilon_{\nu\lambda} |\phi_\lambda\rangle + 3/8 \sum_{\pi\lambda} \epsilon_{\pi\lambda}^* \epsilon_{\nu\lambda} |\chi_\pi\rangle - \dots ,$$

where

$$\epsilon_{\nu\mu} = \langle \Phi_\mu | \chi_\nu \rangle .$$

In the appendix we show that for identical nuclei the symmetric expansion (1) is equivalent to the introduction of orthonormal states which have a maximal localization. Furthermore, the symmetric expansion has the advantage of minimizing the renormalization in the elastic channel (see below). The summation indices π and λ should be restricted to the valence states in the neighborhood of the fermi levels. An unrestricted summation would lead to overcounting because the two sets $\{|\Phi\rangle\}$ and $\{|\chi\rangle\}$ are both complete sets. The nucleon annihilation operator $\psi(\underline{r})$ is expanded as

$$\psi(\underline{r}) = \sum_{\mu} a_{\mu} \bar{\Phi}_{\mu}(\underline{r}) + \sum_{\nu} b_{\nu} \bar{\chi}_{\nu}(\underline{r}) , \tag{2}$$

where the annihilation operators a_{μ} and b_{ν} now commute. We insert this expression in the Hamiltonian

$$\begin{aligned}
H = & \int d^3r \psi^\dagger(\underline{r}) (K + U_A + U_B) \psi(\underline{r}) \\
& + 1/2 \int d^3r d^3r' \psi^\dagger(\underline{r}) \psi^\dagger(\underline{r}') W(\underline{r}-\underline{r}') \psi(\underline{r}') \psi(\underline{r})
\end{aligned} \tag{3}$$

with K , U_A and U_B denoting the operator for the kinetic energy and the potential wells of the two nuclei, respectively. W is the residual nuclear interaction. Assuming that $|\phi_\mu\rangle$ and $|\chi_\nu\rangle$ are eigenfunctions of the potential wells

$$\begin{aligned}
(K + U_A) |\phi_\mu\rangle &= s_\mu |\phi_\mu\rangle, \\
(K + U_B) |\chi_\nu\rangle &= t_\nu |\chi_\nu\rangle,
\end{aligned} \tag{4}$$

one finds that up to second order in ϵ the single particle component of H is given by

$$\begin{aligned}
H_{sp} &= \int d^3r \psi^\dagger(\underline{r}) (K + U_A + U_B) \psi(\underline{r}) \\
&= \sum_{\mu\lambda} \bar{s}_{\mu\lambda} a_\mu^\dagger a_\lambda + \sum_{\nu\kappa} \bar{t}_{\nu\kappa} b_\nu^\dagger b_\kappa + \sum_{\mu\nu} (T_{\mu\nu} a_\mu^\dagger b_\nu + T_{\mu\nu}^* b_\nu^\dagger a_\mu),
\end{aligned} \tag{5}$$

where

$$\begin{aligned}
\bar{s}_{\mu\lambda} &= s_\mu \delta_{\mu\lambda} - 1/8 \sum_\nu (s_\mu + s_\lambda - 2t_\nu) \epsilon_{\nu\mu} \epsilon_{\nu\lambda}^* \\
\bar{t}_{\nu\kappa} &= t_\nu \delta_{\nu\kappa} - 1/8 \sum_\mu (t_\nu + t_\kappa - 2s_\mu) \epsilon_{\nu\mu}^* \epsilon_{\kappa\mu}
\end{aligned} \tag{5a}$$

are the renormalized single-particle energies and where

$$T_{\mu\nu} = 1/2 \langle \phi_\mu | U_A + U_B | \chi_\nu \rangle \tag{6}$$

causes single particle tunneling. It is the symmetric expansion (1) which keeps the renormalization small because the overlap $\epsilon_{\nu\mu} \epsilon_{\nu\lambda}^*$ has its maximum approximately at $s_{\mu} + s_{\lambda} - 2t_{\nu} \approx 0$.

Assuming W to be a pairing force we evaluate the two-particle component of H . Up to second order in ϵ we find

$$\begin{aligned}
 H_{\text{int}} = H - H_{\text{sp}} = & -1/4 \sum_{\pi\lambda\mu\nu} \delta_{\pi\lambda} \delta_{\mu\nu} (G_A a_{\pi}^+ a_{\lambda}^+ a_{\nu} a_{\mu} + G_B b_{\pi}^+ b_{\lambda}^+ b_{\nu} b_{\mu}) \\
 & + 1/16 \sum_{\pi\lambda\mu\nu\zeta} [(G_A \epsilon_{\zeta\pi} \epsilon_{\zeta\lambda} \delta_{\mu\nu} + G_B \epsilon_{\mu\zeta} \epsilon_{\nu\zeta} \delta_{\pi\lambda}) a_{\pi}^+ a_{\lambda}^+ b_{\nu} b_{\mu} + \text{h.c.}]
 \end{aligned}
 \tag{7}$$

As usual $\bar{\nu}$ is related to ν by time reversal. In deriving (7) we have dropped terms leading to single-particle tunneling. We further neglected the renormalization of the pairing strengths G_A and G_B . Note the repulsive character of the two-neutron tunneling part in H_{int} . One has to keep in mind that the relative phase of the operators a and b is arbitrary. The "repulsion" is changed into an "attraction" if one makes the transformation $a \rightarrow a$ and $b \rightarrow ib$. The cross section for the transfer is, of course, invariant under this transformation.

3. Transition Probabilities

The Hamiltonian H can be split into a part H_0 belonging to the unperturbed system

$$H_0 = \sum_{\mu} s_{\mu} a_{\mu}^{\dagger} a_{\mu} + \sum_{\nu} t_{\nu} b_{\nu}^{\dagger} b_{\nu} - 1/4 \sum_{\lambda \mu \nu} \delta_{\lambda \bar{\lambda}} \delta_{\mu \bar{\nu}} (G_A a_{\lambda}^{\dagger} a_{\lambda}^{\dagger} a_{\nu} a_{\mu} + G_B b_{\lambda}^{\dagger} b_{\lambda}^{\dagger} b_{\nu} b_{\mu}) \quad (8)$$

and into a part $V(t)$ containing the overlap ϵ

$$H = H_0 + V(t) \quad (9)$$

H_0 is not explicitly time dependent whereas $V(t)$ depends on time via the overlap ϵ . In the interaction representation the S-matrix is given by

$$S(t) = \hat{T} \exp \left\{ -\frac{i}{\hbar} \int_{-\infty}^t dt' V_{int}(t') \right\}, \quad (10)$$

where

$$V_{int}(t) = \exp \left(\frac{i}{\hbar} H_0 t \right) V(t) \exp \left(-\frac{i}{\hbar} H_0 t \right) \quad (11)$$

and where \hat{T} denotes the time ordering operator. The undisturbed ground state is eigenstate of H_0 . This state is a product state which we assume to be given by

$$|A; B\rangle = |BCS(A)\rangle \otimes |BCS(B)\rangle, \quad (12)$$

where BCS means the BCS-state of nucleus A and nucleus B. In the following we confine ourselves to spherical nuclei, each individual one having an even number of neutrons.

3.1. SINGLE-NEUTRON TRANSFER

The probability for the transfer of a single neutron is given by (compare ref. 12)

$$P_1 = \sum_{\mu\nu} |M_{\mu\nu}|^2, \quad (13)$$

where

$$|M_{\mu\nu}|^2 = \left| -\frac{i}{\hbar} \int_{-\infty}^{+\infty} dt' \langle A+1; B-1 | V_{\text{int}}(t') | A; B \rangle \right|^2 \quad (14)$$

reads up to second order in ϵ

$$|M_{\mu\nu}|^2 = \frac{1}{\hbar^2} \int dt_1 dt_2 u_{\mu}^2 v_{\nu}^2 T_{\mu\nu}(t_1) T_{\mu\nu}^*(t_2) \times \exp [i(\sigma_A - \sigma_B + E_{\mu} + E_{\nu})(t_1 - t_2)] \quad (15)$$

Here we have introduced BCS states of the odd system defined by their matrix elements

$$\langle A+1; B-1 | a_{\lambda}^+ b_{\mu} | A; B \rangle = u_{\mu} v_{\nu} \delta_{\lambda\mu} \delta_{\mu\nu} \quad (16)$$

The numbers u_{μ}^2 (v_{ν}^2) indicate as usual to what extent level μ (ν) of nucleus A (B) is empty (filled):

$$u_{\mu}^2 + v_{\mu}^2 = 1$$

E_{μ} and E_{ν} are the single-particle energies of nucleus A and B when pairing is taken into account, i.e.

$$E_{\mu} = [(s_{\mu} - \sigma_A)^2 + \Delta_A^2]^{1/2}, \quad (17)$$

where σ and Δ denote chemical potential and gap, respectively. The tunneling amplitudes $T_{\mu\nu}(t)$ defined in eq. (6) depend on time because they depend on the vector of relative distance \underline{r} between the two centers of nucleus A and nucleus B. Using the asymptotic expansion formulas of Buttle and Goldfarb¹³⁾ we find

$$\langle \Phi_{\mu} | U_B | \chi_{\nu} \rangle = \sqrt{4\pi} N_{\mu}^{*} \tilde{A}_{\nu} \hat{j}_{\mu} \hat{j}_{\nu} (-)^{j_{\mu}+j_{\nu}} \quad (18)$$

$$\times \sum_{\ell} (-)^{1/2(\ell_{\mu}+\ell_{\nu}-\ell)} \hat{\ell} \begin{pmatrix} \ell & j_{\mu} & j_{\nu} \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \begin{pmatrix} \ell & j_{\mu} & j_{\nu} \\ M_{\nu}-M_{\mu} & M_{\mu} & -M_{\nu} \end{pmatrix} h_{\ell}^{(1)*}(i\kappa_{\mu}r) (-)^{M_{\mu}+\frac{1}{2}} Y_{\ell M_{\nu}-M_{\mu}}(\hat{\underline{r}})$$

($\ell+\ell_{\mu}+\ell_{\nu}=\text{even}$)

We have used the notation of ref. 13

$$\hat{x} = (2x + 1)^{1/2}$$

$$\tilde{A}_{\nu} = (-)^{\ell_{\nu}} A_{\nu} = \frac{\hbar^2}{2m} N_{\nu} \kappa_{\nu}^{\ell_{\nu}} (\sigma_{\nu})^{-\ell_{\nu}-1} \quad (19)$$

N_{μ} is the normalization factor of the asymptotic solution of $|\Phi_{\mu}\rangle$ which is a spherical hankel function.

$$\Phi_{\mu}(\underline{x}) \longrightarrow N_{\mu} h_{\ell_{\mu}}^{(1)}(i\kappa_{\mu}x) Y_{\ell_{\mu} M_{\mu}}(\hat{\underline{x}}) \quad (20)$$

and $\kappa_{\mu} = (2mB_{\mu}\hbar^{-2})^{1/2}$ is defined in terms of the neutron binding energy B_{μ} .

The matrix element $\langle \Phi_{\mu} | U_A | \chi_{\nu} \rangle$ is obtained if one interchanges μ and ν in eq. (18), replaces \underline{r} by $-\underline{r}$ and takes the complex conjugate. We insert eq. (6) together with eq. (18) in eq. (15) and summing over the magnetic quantum numbers we obtain

$$P_1 = \frac{1}{\hbar^2} \sum_{\ell_1 m_1 n} u_m^2 v_n^2 \left[\hat{j}_m \hat{j}_n \hat{\ell} \begin{pmatrix} \ell & j_m & j_n \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \right]^2$$

($\ell + \ell_m + \ell_n = \text{even}$)

(21)

$$\int dt_1 dt_2 \exp \left[\frac{i}{\hbar} (\sigma_A - \sigma_B + E_m + E_n) (t_1 - t_2) \right] S_{mn}(t_1, t_2) P_\ell(r_1 r_2 / r_1 r_2)$$

with

$$S_{mn}(t_1, t_2) = \frac{1}{4} \left(N_m^* \tilde{A} \tilde{h}_\ell(\alpha_m r_1) + N_n^* \tilde{A} \tilde{h}_\ell(\alpha_n r_1) \right)$$

$$\times \left(N_m \tilde{A} \tilde{h}_\ell(\alpha_m r_2) + N_n \tilde{A} \tilde{h}_\ell(\alpha_n r_2) \right)$$
(22)

and

$$\tilde{h}_\ell(x) = i^\ell h_\ell^{(1)}(ix)$$
(23)

being real. The argument y in the Legendre polynomial $P_\ell(y)$ is the cosine of the angle between the vector $\underline{r}_1 = \underline{r}(t_1)$ and the vector $\underline{r}_2 = \underline{r}(t_2)$. The transition probability P_1 is of course independent of an arbitrary phase factor in the normalization constants N_m . In the case of spherical nuclei considered here, normalization, single-particle energies, and occupation probabilities do not depend on magnetic quantum numbers and the indices m and n stand for all single-particle quantum numbers except the magnetic quantum numbers. The integration in eq. (21) has to be taken along the classical Coulomb trajectory, thus depending on the scattering angle θ . The differential cross-section for a single-neutron transfer then reads

$$\frac{d\sigma_1}{d\Omega} = \frac{d\sigma_R}{d\Omega} P_1(\theta)$$
(24)

where $d\sigma_R/d\Omega$ is the elastic Rutherford cross section. Equation (24) only holds if the probability for elastic scattering (see below) is not very different from unity.

3.2. TWO-NEUTRON TRANSFER FROM GROUND STATE TO GROUND STATE

Up to fourth order in the overlap ϵ the probability for two neutrons being transferred into the ground state of the system (A+2; B-2) is given by

$$P_2 = |M_2^{(1)} + M_2^{(2)}|^2, \quad (25)$$

where $M_2^{(1)}$ is the contribution from the double single-particle transfer and $M_2^{(2)}$ denotes the direct pair transfer contained in H_{int} . Using the relation

$$\langle A+2; B-2 | a_{\lambda}^+ a_{\lambda}^+ b_{\nu} b_{\mu} | A; B \rangle = u_{\lambda} v_{\lambda} u_{\mu} v_{\mu} \delta_{\lambda\nu} \delta_{\lambda\mu} \quad (26)$$

we find

$$\begin{aligned} M_2^{(1)} &= \frac{-1}{\hbar^2} \int_{-\infty}^{+\infty} dt_1 dt_2 \sum_{\lambda\lambda\mu\nu} T_{\lambda\lambda}(t_1) T_{\mu\nu}(t_2) \langle A+2; B-2 | \hat{T} (a_{\lambda}^+(t_1) b_{\lambda}(t_1) a_{\mu}^+(t_2) b_{\nu}(t_2)) | A; B \rangle \\ &= \sum_{lmn}^{(l+l_m+l_n=\text{even})} u_m v_m u_n v_n \left[\hat{j}_m \hat{j}_n \hat{\ell} \begin{pmatrix} l & j_m & j_n \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \right]^2 \\ &\times \int_{-\infty}^{+\infty} dt_1 dt_2 \exp \left[\frac{i}{\hbar} (\sigma_A - \sigma_B) (t_1 + t_2) - (E_m + E_n) |t_1 - t_2| \right] S_{mn}(t_1, t_2) P_{\ell} (r_{-1-2} / r_1 r_2). \end{aligned} \quad (27)$$

\hat{T} means the time ordering operator and $a_{\lambda}^+(t)$ depends on time through

$$a_{\lambda}^+(t) = \exp \left(\frac{i}{\hbar} H_0 t \right) a_{\lambda}^+ \exp \left(- \frac{i}{\hbar} H_0 t \right). \quad (28)$$

For the direct pair transfer we obtain

$$M_2^{(2)} = -\frac{i}{8\hbar} \sum_{\ell mn} (\Delta_{A n v n}^{u v} + \Delta_{B m v m}^{u v}) \left[\hat{j}_m \hat{j}_n \hat{\ell} \begin{pmatrix} \ell & j_m & j_n \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \right]^2 \quad (29)$$

($\ell + \ell_m + \ell_n = \text{even}$)

$$\times \int_{-\infty}^{+\infty} dt \left| \frac{2m}{\hbar^2} \left[N_{n m}^* \tilde{A} \tilde{h}_\ell(r_n, r) - N_{m n} \tilde{A}^* \tilde{h}_\ell(r_m, r) \right] / (r_m^2 - r_n^2) \right|^2$$

In deriving eq. (29) we have used the relation¹³⁾

$$(r_\mu^2 - r_\nu^2) \epsilon_{\nu\mu} = \frac{2m}{\hbar^2} \left[\langle \phi_\mu | V_B | \chi_\nu \rangle - \langle \phi_\mu | V_A | \chi_\nu \rangle \right] \quad (30)$$

together with eq. (18). We also used the gap equation

$$\Delta_A = G_A \sum_{\mu > 0} u_\mu v_\mu \quad (31)$$

for nucleus A and for nucleus B.

3.3. ELASTIC CHANNEL

The elastic amplitude M_0 differs from unity because of inelastic processes. There are two second order contributions. The first arises from the renormalization of the single particle energies (eq. (5a)) and the second is due to the diagonal part of the tunneling operator $T_{mn} T_{pq}$. Writing

$$M_0 = 1 - M_0^{(1)} - M_0^{(2)}$$

we find

$$M_o^{(1)} = \frac{i}{4} \sum_{lmn} (v_m^2 - v_n^2) (s_m - t_n) \left[\hat{j}_m \hat{j}_n \hat{\ell} \begin{pmatrix} \ell & j_m & j_n \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \right]^2$$

($\ell + l_m + l_n = \text{even}$)

$$\times \int_{-\infty}^{+\infty} dt \left| \frac{2m}{\hbar^2} \left[N_n^* \tilde{A} \tilde{h}_\ell(\alpha_n r) - N_m \tilde{A}^* \tilde{h}_\ell(\alpha_m r) \right] / (\alpha_m^2 - \alpha_n^2) \right|^2$$

(32)

and

$$M_o^{(2)} = -\frac{1}{2} \sum_{lmn} \left[\hat{j}_m \hat{j}_n \hat{\ell} \begin{pmatrix} \ell & j_m & j_n \\ 0 & \frac{1}{2} & -\frac{1}{2} \end{pmatrix} \right]^2$$

($\ell + l_m + l_n = \text{even}$)

$$\times \int_{-\infty}^{+\infty} dt_1 dt_2 S_{mn}(t_1, t_2) Z_{mn}(t_1, t_2) P_\ell(r_1 r_2 / r_1 r_2)$$

(33)

where $S_{mn}(t_1, t_2)$ is defined in eq. (22) and

$$Z_{mn}(t_1, t_2) = \exp \left[i (\sigma_A - \sigma_B) (t_1 - t_2) - i (E_m + E_n) |t_1 - t_2| \right]$$

$$\times \left(v_m^2 u_n^2 \theta(t_1 - t_2) + u_m^2 v_n^2 \theta(t_2 - t_1) \right) + \{t_1 \longleftrightarrow t_2\}$$

(34)

The function $\theta(t)$ is the step function. We recall that summation index m refers to nucleus A and index n to nucleus B. The differential cross section for the elastic scattering is given by

$$\frac{d\sigma_{el}}{d\Omega} = |1 - M_o^{(1)}(\theta) - M_o^{(2)}(\theta)|^2 \frac{d\sigma_R}{d\Omega} \equiv P_o \frac{d\sigma_R}{d\Omega}$$

(35)

4. Scattering of ^{196}Hg on ^{200}Hg Below the Coulomb Barrier

For the numerical evaluation we use the following approximations:

- a) The normalization constants N_m of the asymptotic wave functions are calculated with Morinigo wave functions⁴). The effective nuclear radius R which separates the nuclear interior from the asymptotic region is chosen as $R = (1.2M^{1/3} + 1.2)$ fm, where M is the nuclear mass number.
- b) The difference quotient which appears in eqs. (29) and (32) is replaced by the symmetrized differential quotient.

$$\frac{2m}{\hbar^2} \left(N_{nm}^* \tilde{h}_\ell(\alpha_n r) - N_{mn}^* \tilde{h}_\ell(\alpha_m r) \right) / (\alpha_m^2 - \alpha_n^2) \quad (36)$$

$$\approx -\frac{1}{2} F_{mn} \left[\frac{d}{d\alpha_m} \left(\alpha_m^{\ell_m + \ell_n + 1} \tilde{h}_\ell(\alpha_m r) \right) + \frac{d}{d\alpha_n} \left(\alpha_n^{\ell_m + \ell_n + 1} \tilde{h}_\ell(\alpha_n r) \right) \right],$$

where

$$F_{mn} = N_n^* N_m / \left[\alpha_m^{\ell_m + 1} \alpha_n^{\ell_n + 1} (\alpha_m + \alpha_n) \right].$$

By numerical comparison with the exact expressions one finds that this assumption is reasonable. In order to simplify the integrals we consider only the exponential behavior of the function $\tilde{h}_\ell(\alpha r)$

$$\tilde{h}_\ell(\alpha r(t)) \approx \tilde{h}_\ell(\alpha r_0) \exp \left\{ \alpha [r_0 - r(t)] \right\}, \quad (37)$$

where r_0 is the distance of closest approach, i.e. the classical turning point. Within the same approximation we replace the arguments of the Legendre polynomials by 1, i.e. by their values at the turning point. In eq. (37) we expand the

exponent up to second order in the time t and perform all integrals analytically. In view of the rapid decrease of $h_\rho(r)$ with increasing r these approximations also seem to be justified¹⁴).

c) The single-particle energies are taken from ref. 15. The configuration space consists of 13 levels ranging from the $1h_{9/2}$ level up to the $3d_{3/2}$ level. We used the pairing rotational model¹⁶) to determine Δ and G from the two-neutron separation energies in the mercury region. For ^{200}Hg we obtain $\Delta_A = 1$ MeV and $G_A = 0.11$ MeV, in the case of ^{196}Hg we find $\Delta_B = 1.17$ MeV and $G_B = G_A$. The chemical potential for BCS nuclei is defined by

$$\sigma = -\frac{1}{4} [S(2,N) + S(2,N+2)] \quad , \quad (38)$$

where $S(2N)$ stands for the separation energy of two neutrons in a nucleus with N neutrons. From this relation the values $\sigma_A = -7.17$ MeV and $\sigma_B = -7.85$ MeV are deduced for ^{200}Hg and ^{196}Hg , respectively.

In fig. 2 we show the differential cross sections for the total one-neutron transfer in the reaction $^{200}\text{Hg}(^{196}\text{Hg}, ^{195}\text{Hg})^{201}\text{Hg}$ and for the transfer of two neutrons into the ground states of ^{194}Hg and ^{202}Hg in the reaction $^{200}\text{Hg}(^{196}\text{Hg}, ^{194}\text{Hg})^{202}\text{Hg}$ at $E_{\text{cm}} = 560$ MeV. Using the radii for nucleus A and B according to section 4a we find the Coulomb barrier at $E_{\text{cb}} = 562$ MeV. The amplitudes for the two-neutron transfer in the reaction $B(A,A+2)B-2$ are equal in BCS to those of $B(A,A-2)B+2$. The cross section for the reaction $^{200}\text{Hg}(^{196}\text{Hg}, ^{198}\text{Hg})^{198}\text{Hg}$, however, is symmetric around $\theta_{\text{cm}} = 90^\circ$ because of the two identical nuclei in the final configuration. Figure 1 also contains the function

$$F = P_0 + P_1 + P_2 + P_{-1} + P_{-2} \quad , \quad (39)$$

where P_0 is the probability for elastic scattering along the Coulomb trajectory

as defined in eq. (35). The quantities P_1 and P_2 have been introduced in section 3; they are the probabilities for the transfer of one or two neutrons to ^{200}Hg whereas P_{-1} and P_{-2} denote the transfer of one or two neutrons to ^{196}Hg . A necessary condition for perturbation theory to be valid is that F does not deviate much from unity.

In fig. 3 the energy dependence of the differential cross sections is plotted for $\theta_{\text{cm}} = 120^\circ$. Again perturbation theory is limited to $F \approx 1$.

5. Discussion

We first point out that our results should not be interpreted quantitatively. There are considerable uncertainties coming from the single-particle wave functions: If the normalization constants N_m are changed by a factor 2 then the absolute values for the cross sections for single-neutron and two-neutron transfer are changed by a factor of 16 and 256, respectively. Also uncertainties in the BCS wave functions are not irrelevant. A truncation of the configuration space would reduce the theoretical cross sections for the two-neutron transfer considerably. Therefore, our results should be compared with future experiments in terms of relative cross sections.

The two-neutron transfer shows an interesting interference pattern because of the fact that the imaginary parts of the amplitudes for double single-transfer and for pair transfer are opposite in sign. The shallow minimum in the two-neutron transfer cross section belongs to the node of the imaginary part. The break-up of a pair becomes more important for shorter coupling times because then the oscillating behavior of the exponent in eq. (27) is less relevant. Shorter coupling time means higher energies or larger scattering angles. Unfortunately, perturbation theory fails to describe the details of the second rise. The cross section for a single neutron will also deviate from the exponential law¹⁷⁾ when nuclear distortions invalidate perturbation theory. Non-perturbative theories have been discussed¹⁸⁾ in the literature although they again are restricted to the assumption that Coulomb excitation does not couple too strongly to nuclear excitation. Qualitatively, Coulomb excitation leads to a damping of the elastic channel. For quantitative results one has to treat both transfer and Coulomb excitation in a quantum mechanical way¹⁰⁾.

We briefly consider some kinematical effects. The overlap between the single-particle wave functions of the two nuclei generally depends on the velocity of the relative motion. For sub-Coulomb reactions this dependence can be neglected¹⁹). Next, it is clear that heavy-ion transfer reactions are connected with a transfer of angular momentum. There are two contributions. First, the angular momentum is changed because of the polarization of the two cores and, secondly, the transferred particle itself carries along an angular momentum in its relative motion. For bombarding energies below the Coulomb barrier it is difficult to calculate the optimum value²⁰) for the transferred angular momentum.

Finally, we mention that recoil effects²⁰) can be neglected in the scattering of two almost identical nuclei. This means that the most favourable Q-value for the neutron transfer is approximately zero, a rather well met property of the reactions we discussed in this paper.

Acknowledgments

We are indebted to Prof. H. Schmidt for his constant support and interest. One of us (M. K.) wishes to express his gratitude to Prof. N. K. Glendenning for his hospitality and to Dr. M. A. Nagarajan for useful discussions.

Appendix

We show that for identical nuclei the symmetric expansion (1) in section 2 is equivalent to the introduction of localized, orthonormal states. For simplicity we assume that there is only one state $|\phi\rangle$ in nucleus A and one state $|\chi\rangle$ in nucleus B. Since the two nuclei are identical the two eigenstates of the coupled system are

$$|\psi^{(+)}\rangle = (2(1+\epsilon))^{-1/2} (|\phi\rangle + |\chi\rangle) \quad (\text{A1})$$

and

$$|\psi^{(-)}\rangle = (2(1-\epsilon))^{-1/2} (|\phi\rangle - |\chi\rangle) \quad , \quad (\text{A2})$$

where $\epsilon = \langle \phi | \chi \rangle$ is the overlap of the two wave functions (which we assume to be real). The two eigenstates $|\psi^{(+)}\rangle$ and $|\psi^{(-)}\rangle$ are neither localized in nucleus A nor in nucleus B. The combinations

$$|\bar{\phi}\rangle = 2^{-1/2} (|\psi^{(+)}\rangle + |\psi^{(-)}\rangle) \quad (\text{A3})$$

and

$$|\bar{\chi}\rangle = 2^{-1/2} (|\psi^{(+)}\rangle - |\psi^{(-)}\rangle) \quad (\text{A4})$$

are, however, localized. Using (A1) and (A2) we can express $|\bar{\phi}\rangle$ and $|\bar{\chi}\rangle$ by the unperturbed states $|\phi\rangle$ and $|\chi\rangle$. We make a power series expansion in ϵ and get the expansion (1) of section 2

$$|\bar{\phi}\rangle = (1 + \frac{3}{8} \epsilon^2) |\phi\rangle - \frac{1}{2} \epsilon |\chi\rangle + \dots \quad (\text{A5})$$

$$|\bar{\chi}\rangle = (1 + \frac{3}{8} \epsilon^2) |\chi\rangle - \frac{1}{2} \epsilon |\phi\rangle + \dots \quad (\text{A6})$$

The generalization to identical nuclei with an arbitrary number of states is straightforward.

References

- 1) Symposium on Heavy-Ion Reactions and Many Particle Excitations, Journal de Physique, Colloque C-6 (1971)
- 2) Proceedings of the European Conference on Nuclear Physics, Aix-en-Provence, 1972
- 3) G. Breit and M. E. Ebel, Phys. Rev. 103 (1956) 679
- 4) D. Trautmann and K. Alder, Helv. Phys. Acta 43 (1970) 363
- 5) R. A. Broglia and A. Winther, Nucl. Phys. A182 (1972) 112; R. A. Broglia and A. Winther, Phys. Reports 4C (1972) 153
- 6) A. Messiah, Quantum Mechanics, Vol. 1 (North Holland, Amsterdam, 1961)
- 7) W. E. Frahn, Ann. Phys. (N. Y.) 72 (1972) 524
- 8) R. A. Broglia, S. Landowne, and A. Winther, Phys. Lett. 40B (1972) 293
A. J. Baltz and S. Kahana, Phys. Rev. Lett. 29 (1972) 1267
R. Bock and H. Yoshida, Nucl. Phys. A189 (1972) 177
P. Bonche and B. Giraud, Phys. Rev. Lett. 28 (1972) 1720
- 9) R. E. Prange, Phys. Rev. 131 (1963) 1083
- 10) R. J. Ascutto and N. K. Glendenning, Nucl. Phys. A188 (1972) 185
- 11) H. Bohn, G. Daniel, M. R. Maier, P. Kienle, J. G. Cramer, and D. Proetel, Phys. Rev. Lett. 29 (1972) 1337
- 12) V. I. Goldanskii and A. I. Larkin, Sov. Phys. JETP 26 (1968) 617
- 13) P. J. A. Buttle and L. J. B. Goldfarb, Nucl. Phys. 78 (1966) 409
- 14) L. J. B. Goldfarb and K. Wong, Nucl. Phys. A90 (1967) 361
P. J. A. Buttle and L. J. B. Goldfarb, Nucl. Phys. A176 (1971) 299
- 15) B. Sørensen, Nucl. Phys. A97 (1967) 1


- 16) J. Ginocchio and J. Weneser, Phys. Rev. 170 (1968) 859; A. Bohr, International Symposium on Nuclear Structure, Dubna, USSR, 1968 (IAEA, Vienna, 1969) p. 179; R. Beck, M. Kleber, and H. Schmidt, Z. Phys. 250 (1972) 155
- 17) K. R. Greider, Advances in Theoretical Physics, Vol. 1, ed. by L. A. Brueckner (Academic Press, New York 1965) p. 270
- 18) K. Dietrich, Phys. Lett. 32B (1970) 428; K. Dietrich, K. Hara, F. Weller, Phys. Lett. 35B (1971) 201; M. Gaudin, Nucl. Phys. A144 (1970) 191
M. Kleber and H. Schmidt, Z. Phys. 245 (1971) 68
- 19) T. Fliessbach, Z. Phys. 247 (1972) 117
- 20) D. M. Brink, Phys. Lett. 40B (1972) 37

Figure Captions

Fig. 1. Schematic picture of two nuclear potential wells A and B separated by the distance r . Horizontal lines denote the single particle levels near the fermi levels.


Fig. 2. Differential cross sections for the total single-neutron (sn) transfer $^{200}\text{Hg}(^{196}\text{Hg}, ^{195}\text{Hg})^{201}\text{Hg}$ and for the ground state two-neutron (tn) transfer $^{200}\text{Hg}(^{196}\text{Hg}, ^{194}\text{Hg})^{202}\text{Hg}$ at $E_{\text{cm}} = 560$ MeV. The function F , defined in eq. (39) depends on θ_{cm} and is a measure of the failure of perturbation theory.

Fig. 3. Differential cross sections at $\theta_{\text{cm}} = 120^\circ$ as a function of the bombarding energy E_{cm} . The symbols sn, tn are defined in fig. 2. Again F , depending now on E_{cm} , is a measure of the failure of perturbation theory.




XBL7211 - 4402

Fig. 1



XBL7211-4403

Fig. 2



XBL72II-4404

Fig. 3

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720