

Lawrence Berkeley National Laboratory

Recent Work

Title

EVIDENCE FOR A SOFT NUCLEAR-MATTER EQUATION OF STATE

Permalink

<https://escholarship.org/uc/item/53h2m6pg>

Author

Siemens, P.J.

Publication Date

1979-07-01

14



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Submitted to Physical Review Letters

EVIDENCE FOR A SOFT NUCLEAR-MATTER EQUATION OF STATE

Philip J. Siemens and Joseph I. Kapusta

July 1979

RECEIVED
LAWRENCE
BERKELEY LABORATORY

SEP 14 1979

LIBRARY AND
DOCUMENTS SECTION


TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782*

~~RECEIVED
LAWRENCE
BERKELEY LABORATORY~~

~~AUG 28 1979~~

~~LIBRARY AND
DOCUMENTS SECTION~~



14
1979

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Evidence for a Soft Nuclear-Matter Equation of State

Philip J. Siemens⁺ and Joseph I. Kapusta

Lawrence Berkeley Laboratory

University of California

Berkeley, CA 94720

ABSTRACT

The entropy of the fireball formed in central collisions of heavy nuclei at center of mass kinetic energies of a few hundred MeV per nucleon is estimated from the ratio of deuterons to protons at large transverse momentum. The observed paucity of deuterons suggests that strong attractive forces are present in hot, dense nuclear matter, or that degrees of freedom beyond the nucleon and pion may already be realized at an excitation energy of 100 MeV per baryon.

⁺On leave from the Niels Bohr Institute, University of Copenhagen, Denmark.

One of the principal motivations for accelerating heavy-ion beams to relativistic energies is the hope of producing and studying matter at baryon densities greater than are found in atomic nuclei. However, information about the properties of the dense matter thus created is obscured by the fact that the matter remains hot and dense only for a very short time, $<10^{-22}$ sec, and our observations are limited to the products emitted as it disassembles. We present arguments that the ratio R_{dp} of deuterons to protons is established during the early stages of the fireball's existence, and is little changed by the later processes of expansion and disassembly. We show that the ratio R_{dp} is proportional to the density of neutrons in phase (momentum x position) space, and thus measures the specific entropy of the fireball. Applying our arguments to recent measurements¹ of Ne + NaF and Ar + KCl at 100 to 200 MeV/nucleon c.m. kinetic energy, we find that a surprisingly large region of phase space is populated. This could be explained by strong attractive interactions in the hot dense matter, or by a proliferation of the degrees of freedom sharing the excitation energy.

It has been argued² that the time evolution of the fireball in central collisions goes through three stages: formation, explosion, and disintegration. In collisions of large nuclei, with relative velocity greater than the speed of sound, a fireball may be formed at high density. At high energies or for small nuclei, there may be partial interpenetration, in which case the initial density of the hot matter will be less than double normal density. During this stage of the reaction, most of the kinetic energy of the nuclei's relative motion is thought to be converted to thermal excitation of the dense matter. A

great deal of entropy is created in this stage of the reaction.³

During the second stage of the reaction, repeated collisions lead to a hydrodynamic expansion of the matter. This expansion is largely reversible, so that little additional entropy is generated during the expansion.^{2,3} The additional phase-space due to the expansion of matter into a larger region of position space is compensated by the shrinking of the velocity distribution into a smaller region of momentum space as the expanding matter cools. During this stage of the reaction, entropy is produced by viscous forces and by thermal conduction. Heat conduction produces entropy at a rate proportional to the square of the thermal gradients, so it should not be very large. The viscous damping of the shear-velocity gradients also produces entropy, but this merely completes the initial thermalization process.

In the third stage of the reaction, the density of particles has become so small that they seldom collide. This stage lasts until the particles reach the detectors. During this stage, the matter is no longer described hydrodynamically because the distribution of momenta at any given position and time cannot be characterized by a mean velocity and a thermal fluctuation. Instead, strong correlations develop between position and momentum, which may be used to identify the particles in a time-of-flight telescope or magnetic spectrometer. However, Liouville's theorem guarantees that the particles' density in phase space remains constant in the absence of collisions.

Due to the reaction $d + N \rightleftharpoons p + n + N$, where N is a spectator nucleon or cluster, deuterons will be constantly breaking up and reforming. If collisions are frequent enough, the deuterons will quickly

reach an equilibrium concentration determined by detailed balancing:⁴

$$e^{-\mu_d/T} d_{\tilde{d}}(R, P, S_z) = \sum_{S_z} d_{\tilde{p}}(R, P/2, s_z) d_{\tilde{n}}(R, P/2, S_z - s_z) e^{-(\mu_n + \mu_p)/T} \quad (1)$$

where $d_i(R, P, S_z)$ is the six-dimensional phase-space density (Wigner mixed density) of species i at position R with momentum P and spin projection S_z , μ_i is its chemical potential, and T is the temperature of the equilibrium thermal distribution. Since the deuteron is a very weakly-bound system we expect $\mu_d - \mu_n - \mu_p \ll T$. Summing over spins and momenta, we obtain for the ratio of deuterons to protons

$$R_{dp} = \frac{2S_d + 1}{2S_p + 1} \cdot 8 \langle \tilde{d}_{\tilde{n}} \rangle, \quad (2)$$

where $\langle \tilde{d}_{\tilde{n}} \rangle = \frac{\int \int d\tilde{p} d\tilde{r} d\tilde{p}_n d\tilde{r}_n (\tilde{p}, \tilde{r})^2}{\int \int d\tilde{p} d\tilde{r} d\tilde{p}_n d\tilde{r}_n (\tilde{p}, \tilde{r})}$ is the average phase-space density of neutrons, and assuming $d_{\tilde{n}} = d_{\tilde{p}}$.

The rate of approach to the equilibrium concentration is governed by the collision frequency, with a time constant equal to⁴

$$\tau = [n_N \langle \sigma(N+d \rightarrow n+p+N) v_{rel} \rangle]^{-1},$$

where n_N is the nucleonic density. Since $\sigma(N+d \rightarrow n+p+N)$ is similar to elastic nucleon cross-sections, the equilibration of deuteron number will be just as rapid as the thermalization of the nucleonic momentum distributions. Thus R_{dp} will be in equilibrium through the hydrodynamic expansion.⁵ In fact, since R_{dp} is just given by the mean density of neutrons in phase space $\langle \tilde{d}_{\tilde{n}} \rangle$, we can relate R_{dp} to the entropy per neutron S_n carried by the neutrons

$$S_n = 5/2 - \ln(2^{2/3} \langle d_n \rangle) = 3.95 - \ln R_{dp} . \quad (3)$$

Thus $\langle d_n \rangle$ and R_{dp} will be approximately constant throughout the hydrodynamic expansion phase.

Even after the collisions cease to maintain hydrodynamic local equilibrium, the number of deuterons does not change. If the remaining interactions are too weak to significantly alter the phase-space distributions d_n and d_p , then an alternative estimate of the number of deuterons is obtained by projecting the uncorrelated neutron and proton densities onto the deuteron wave function,⁶

$$d_d(R, P, S_z) = \sum_{s_z} \iint d\tilde{r} d\tilde{p} d_p(R + \frac{\tilde{r}}{2}, P + \frac{\tilde{p}}{2}, S_z + s_z) d_n(R - \frac{\tilde{r}}{2}, P - \frac{\tilde{p}}{2}, S_z - s_z) d_{in}(\tilde{r}, \tilde{p}, s_z). \quad (4)$$

If d_p and d_n vary slowly over the range of the deuteron's intrinsic structure $d_{in}(\tilde{r}, \tilde{p}) = (2\pi)^{-3} \int d\tilde{r}' \psi_d(\tilde{r} + \frac{\tilde{r}'}{2}) \psi_d(\tilde{r} - \frac{\tilde{r}'}{2}) \exp(ip \cdot \tilde{r}')$, we recover the relations (1) and (2) (neglecting $(\mu_d - \mu_n - \mu_p)/T$). From Liouville's theorem, $\langle d_n \rangle$ does not change after the collisions cease. We conclude that the ratio of deuterons to protons in the final state measures the entropy of the neutrons in the fireball, according to eq. 3.

The functions $d_i(\tilde{r}, \tilde{p})$ describe the distribution of particles in an individual collision. The observed distributions, on the other hand, are averages over collisions with different impact parameters and different reaction planes. Since relation (1) is non-linear, it will not in general be satisfied if average quantities are substituted. To minimize this difficulty, we follow ref. 2 and restrict our analysis to measurements of similar-sized target and projectile at large transverse

momentum. This kinematic regime is associated with central collisions of high multiplicity. The measured distributions at 90° c.m. are fit to a radially-expanding Boltzmann distribution as in ref. 2. The temperature and expansion velocity are determined from fits to the tails of pion and proton spectra; the ratio R_{dp} is then determined by the ratio of the normalizations. Sample fits are shown in Fig. 1. The resulting entropies are plotted in Fig. 2. The error bars are made large enough to include the value of R_{dp} from the total differential cross sections at 90° c.m., which include some peripherally-produced protons and deuterons of small transverse momentum. We see that the entropy decreases for larger systems or more central collisions, which would be expected since the thermalization takes place at higher mean densities in these cases. The energy per baryon is taken as the beam energy minus the energy of observed pions (assuming isospin symmetry) at 90° c.m.; the error bar indicates the size of the pion subtraction. We see that the entropy increases with energy, as expected.

To judge the significance of the result, we show the predictions for entropy versus excitation energy for a very soft equation of state at various densities in Fig. 2. The energy per baryon is given by

$$E(n,S) = E(n_o,o) + \frac{1}{9} K \left(\frac{n_o}{n} - 1 + \ln \frac{n}{n_o} \right) + E_{FG}(n,S) - E_{FG}(n,o) \quad (5)$$

where n is the baryon number density, n_o is the density of normal matter with compressibility K , and $E_{FG}(n,S)$ is the energy per baryon of a free fermi gas (evaluated in the limit of low degeneracy, corresponding to eq. 3). We see that much more entropy is created than would

be expected from even such a soft equation of state for nucleons as this. Even the inclusion of real pions in the system does not give enough entropy. This is especially surprising in light of the fact that such diverse dynamical models as fluid dynamics⁷ and intranuclear cascade⁸ predict maximum compressions of three to four.

The interesting conclusion that seems to be forced on us is that more entropy is generated during the collision than we would naively expect. It could be that there are strong attractive forces⁹ present in the hot, dense nuclear medium, thus raising the entropy. It could also be that many more mesonic and baryonic particulate degrees of freedom are excited at this energy than suspected.¹⁰ Or it could be that collective degrees of freedom, such as due to pion condensation,¹¹ are the culprits. Or perhaps the nucleons dissociate into quarks.

In any event it is important to make measurements for excitation functions, and to use heavier nuclei to reduce any unwanted surface effects. Clearly it is also important to perform more detailed numerical calculations to further support the scenario described in this letter. These could take the form of deuteron production rates in two fluid hydrodynamics⁷ or intranuclear cascade.^{8,12}

We are grateful to Marie-Claude Lemaire and Shoji Nagamiya for communication and discussion of their data, and to David Hendrie for an enlightening comment.

This work was supported by the U. S. Department of Energy, Nuclear Physics Division, under contract No. W-7405-ENG-48.


References


1. M.-C. Lemaire, S. Nagamiya, S. Schnetzer, H. Steiner and I. Tanihata, "Composite Particle Emission in Relativistic Heavy Ion Collisions," Phys. Lett., in press.
2. P.J. Siemens and J.O. Rasmussen, Phys. Rev. Lett. 42, 880 (1979).
3. M.I. Sobel, P.J. Siemens, J.P. Bondorf and H.A. Bethe, Nucl. Phys. A251, 502 (1975).
4. A. Mekjian, Phys. Rev. C17, 1051 (1978).
5. J.I. Kapusta, "Mechanisms for Deuteron Production in Relativistic Nuclear Collisions," LBL-9251 preprint.
6. R. Bond, P.J. Johansen, S.E. Koonin and S. Garpman, Phys. Lett. 71B, 43 (1977).
7. A.A. Amsden, A.S. Goldhaber, F.H. Harlow and J.R. Nix, Phys. Rev. C17, 2080 (1978).
8. R.K. Smith and M. Danos, in Proceedings of the Topical Conference on Heavy Ion Collisions, Fall Creek Falls State Park, Tennessee, 1977 (National Technical Information Service, Springfield, Va., 1977), Conf-770602, p. 363.
9. T.D. Lee and G.C. Wick, Phys. Rev. D9, 2291 (1974).
10. N.K. Glendenning and Y. Karant, Phys. Rev. Lett. 40, 374 (1978).
11. V. Ruck, M. Gyulassy and W. Greiner, Z. Phys. A277, 391 (1976).
12. E.A. Remler and A.P. Sathe, Ann. Phys. 91, 295 (1975).

FIGURE CAPTIONS

Fig. 1. Spectra at 90° c.m. for reactions $A_p + A_T \rightarrow d + X$. Data is from Lemaire et al., Ref. 1. Theoretical curves are exploding fireball model from Siemens and Rasmussen, Ref. 2, with temperature and velocity fitted to proton and pion spectra, normalization fitted to measured points.

Fig. 2. Entropy per baryon as a function of excitation energy per baryon for various reactions, including one with measured associated multiplicities greater than three. Theoretical curves are for a soft nuclear-matter equation of state at compression ratios of one, two and four (eq. 5 of the text, with $K = 200$ MeV).





This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720