

Lawrence Berkeley National Laboratory

Recent Work

Title

MESON PRODUCTION IN $p + d$ COLLISIONS AND THE 1-d-ff-ft INTERACTION. III. SPIN AND PARITY OF THE $I = 0$ ANOMALY

Permalink

<https://escholarship.org/uc/item/53j5t7v9>

Authors

Booth, Norman E.
Abashian, Alexander
Crowe, Kenneth M.

Publication Date

1963-02-24

UCRL-10409

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

MESON PRODUCTION IN $p + d$ COLLISIONS AND
THE $I=0$ $\pi-\pi$ INTERACTION
III. SPIN AND PARITY OF THE $I=0$ ANOMALY

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review

UCRL-10409

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

MESON PRODUCTION IN $p + d$ COLLISIONS AND THE $I=0$ $\pi-\pi$ INTERACTION
III. SPIN AND PARITY OF THE $I=0$ ANOMALY

Norman E. Booth, Alexander Abashian, and Kenneth M. Crowe

February 24, 1963

Meson Production in p+d Collisions
and the I=0 π - π Interaction

III. Spin and Parity of the I=0 Anomaly

Norman E. Booth, Alexander Abashian, and Kenneth M. Crowe

Lawrence Radiation Laboratory
University of California
Berkeley, California

February 24, 1963

ABSTRACT

In an auxiliary experiment an attempt was made to determine the spin and parity of the I=0 anomaly by detecting charged pions and gamma rays in coincidence with He³. The results are consistent with the interpretation that the predominant mode of decay is into two pions with a charged-to-neutral ratio of 2/1, indicating a state of spin 0 and even parity.

Meson Production in p+d Collisions
and the I=0 π - π Interaction*§

III. Spin and Parity of the I=0 Anomaly

Norman E. Booth,[†] Alexander Abashian,[‡] and Kenneth M. Crowe

Lawrence Radiation Laboratory
University of California
Berkeley, California

February 24, 1963

I. INTRODUCTION

A comparison of the momentum spectra of H^3 and of He^3 produced in p+d collisions showed the anomaly to be an I=0 isotopic spin state.¹ Since the computed resolution functions were not as wide as the anomaly, we inferred that the anomaly has an extremely short life time and that the dominant mode of decay was into two pions. In other words, the anomaly is an enhancement in the double pion production for low energies of the two-pion (2π) system. Therefore it was felt to be important to have direct information on the decay products. The experiment consisted in measuring the relative number of γ rays and charged pions in coincidence with the He^3 of several different momenta.

Table I gives the decay characteristics for particles of mass less than $3m_\pi$ for the eight simplest sets of quantum numbers. If the anomaly is $J^{PG} = 0^{++}$, then a strong decay into two pions is expected. In this case, apart from kinematical factors, the π and γ counting rates should be independent of He^3 momentum. Before the discovery² of the ω^0 , there was some speculation³ that the anomaly might be the 1^{--} particle postulated by Nambu.⁴ For this assignment the anomaly decays primarily into $\pi^0 + \gamma$, and one would expect an increase in the γ - He^3 coincidence rate in the region of the anomaly. For most of the other decay modes listed in Table I, there should also be a difference in the π - He^3 and γ - He^3 rates in the region of the anomaly.

II. EXPERIMENT

The experiment was performed in a two-week period at the end of the second run at the 184-inch cyclotron.⁵ For this auxiliary experiment the D_2 gas target was replaced by a modified version of the Y-shaped target used in the first run.⁶ The modification consisted of a long thin stainless steel "bubble" wall 0.028-in. thick on the side of the target opposite the He^3 or H^3 momentum-selected beam.


For He^3 produced at a laboratory angle of 11.8 deg and with momenta corresponding to the peak of the anomaly, i. e. $w \approx 310$ MeV, the "w system", which we denote by w, is produced at a laboratory angle of 84 deg. At this angle on the opposite side of the proton beam from the He^3 spectrometer a well-shielded scintillation counter telescope was set up, as shown in Fig. 1. The half-angle subtended by the telescope was about 8 deg. Charged pions were detected by the coincidence 123, and γ rays by the coincidence $\bar{1}234$ with 1/4 in. of Pb placed between 1 and 2.

The experiment was made possible by the installation in the cyclotron of the auxiliary dee system,⁷ which increased the duty cycle by a factor of 50 to 100. Although the proton beam was stopped in the back wall of the same cave containing the target and counter telescope, this caused very little trouble. The important source of accidentals was other interactions in the D_2 target in the same rf pulse as a He^3 production.

Let us consider the kinematics of the pions and γ rays from the reactions



and



for a fixed He^3 laboratory angle. Just at threshold, i. e., $w \approx 2m_\pi$ the pions move out in a narrow cone at roughly 90 deg to the incident beam direction.

As w increases and the pions have more relative energy, the angle of the cone rapidly increases and the axis of the cone rotates forward. For charged pions with $w = 310$ MeV, the pion telescope is on the axis of the cone at 84 deg in the lab system and the half-angle of the cone is 30 deg. Figure 2 shows some of the lab angular distributions of the charged pions for the case of isotropic emission in the w -system center of mass. The dashed part of each curve corresponds to pions going backward in this center-of-mass (c.m.) system which, however, are confined to the forward cone in the lab system. Some of these pions cannot get through the counter telescope because of their low range. We therefore took pion data only with the 1/4 in. of Pb in, so that none of these backward pions were detected. A range curve taken by placing Cu before counter 4 gave results consistent with the expected range for the high-energy group of pions.

The decay of neutral pions from reaction (1b) into γ rays destroys the cone, and although the angular distribution of the γ rays is still peaked along the p_w direction, it falls off so slowly that the detection efficiency of the telescope for γ rays varies slowly with w .

The fast coincidences $\pi = 123$ and $\gamma = \overline{1234}$ were made in Wenzel coincidence circuits.⁸ The counters were delayed and plateaued and the circuits matched by placing the telescope in the 11.8-deg momentum-selected beam and setting the time of flight for pions. The outputs of the Wenzel circuits were put into coincidence with a pulse signifying an identified momentum-selected He^3 , as well as a pulse from the first time-of-flight counter (S_1 in Fig. 6 of paper I). Three of these second circuits were used--one for the γ - He^3 coincidence and two for the π - He^3 coincidence. Of the latter, one was used to measure genuine coincidences and one to measure π - He^3 accidentals--the roles of the two π - He^3 circuits being frequently interchanged.

In order to increase the counting rate, the momentum-selecting grids were all connected together in the coincidence requirement. We took measurements at nine different values of the He^3 momentum, concentrating mainly on the anomaly but obtaining enough data on either side for comparison. Gamma- He^3 coincidences were measured with Pb in and Pb out. One π - He^3 circuit measured events with normal delay, while the other measured accidentals. Then another run was taken with the roles of the two π - He^3 circuits interchanged. Since the two π - He^3 circuits could have different thresholds for accidentals and because the π - He^3 accidentals were large (about 30%), the following criteria were used:

- (a) If in two consecutive runs (roles of circuits interchanged between runs) there was no noticeable change of beam intensity or other conditions, we used the data from each circuit independently.
- (b) If the beam intensity did change, we used the data from each run independently. Several runs were taken for each He^3 momentum. A mean π - He^3 rate for each momentum was obtained by weighting the results of the different runs according to the statistical errors. The ratio of deviation from the mean to statistical error was then computed for each measurement. A plot of the ratios for all the measurements is shown in Fig. 3. The shape of the plot is in good agreement with the normal distribution, shown as the dashed curve, but it is too wide. To take account of the indicated systematic error, we have increased the error of each measurement by 25%.

A similar plot for the γ - He^3 data indicated that the errors should be about 15% larger. Tables II and III respectively show typical sets of π and γ measurements. The results of all measurements are summarized in Table IV.

III. ANALYSIS

As the first step in the analysis we choose to divide out the kinematical factors in the π and γ rates. The relative efficiency for π and γ detection as a function of w must be computed. First, the He^3 momentum scale was translated to a scale of w . Then assuming that in 2π production the pions are emitted isotropically in the w system, the laboratory distributions of π 's and γ 's were weighted by the experimental angular resolution and resolution in w , the latter being determined by the momentum resolution. This gives the relative efficiency, which is compared with the experimental points in Fig. 4. Here we plot the observed π and γ rates vs w . The dashed curve is the prediction for the pion rate when the resolution is neglected. The effect of the resolution is easily calculated for $w > 300$ MeV. However, it is very difficult to estimate for w near $2m_\pi$ because the pion cone is changing so rapidly. In the absence of an exact calculation we have chosen to draw a smooth curve through the first three points and to join it to our resolution calculation at $w = 300$ MeV. The solid curve through the γ points also has the resolution folded in. The only factor neglected here is the dependence of the detection efficiency upon γ -ray energy. This has been neglected because it is a smooth slowly varying function of w . The solid curves are close to that expected for $I=0$ production of two pions only. We see from Fig. 4 that there is no marked deviation of the data from the solid curves or the hypothesis of 2π production in the region of $w=310$ MeV.

To facilitate comparison with the results to be expected from some of the other decay modes of Table I we have divided the values of experimental points by values of the corresponding points from the solid curves of Fig. 4. The results are shown in Fig. 5

Before computing the expected results for some other decay mode, we must consider two factors. Firstly, there is a more or less constant background of charged pions due to $J=1$ production. This can be determined from our measurements of $p+d \rightarrow H^3 + \pi^+ + \pi^0$, and is easily taken into account. Secondly, the other decay modes of Table I may have rather complicated angular distributions. We shall neglect this fact since we have no idea what the angular distribution might be and consider each decay mode only on the basis of the numbers of charged pions and γ -rays.

With these considerations we have plotted in Fig. 5 the expected results for each of several of the decay modes listed in Table I. Note that in Fig. 5 the pion data excludes any mode consisting only of neutral pions and γ -rays. The 2π , $2\pi\gamma$, $\pi^+\pi^-\gamma$, and $\pi^+\pi^-2\gamma$ modes fit the pion data reasonably well. Of these, the 2π and $\pi^+\pi^-\gamma$ modes fit the γ -ray data well. The $2\pi\gamma$ and $\pi^+\pi^-2\gamma$ modes fit the γ -ray data rather poorly (with χ^2 probabilities of 1 and 6% respectively). We consider them to be possible but not very likely candidates.⁹

IV. RESULTS

Referring to Table I, we conclude the following regarding possible J^{PG} assignments:

0^{++} : The dominant mode of decay is into two pions, and this assignment is allowed on the basis of our data.

0^{+-} : The $J=2$ barrier in the $2\pi\gamma$ mode tends to favor the $\pi^+\pi^-2\gamma$ mode. On the basis of the data, neither mode seems very likely. We conclude that this assignment is possible but not so likely.

0^{-+} : This is the same assignment as the η and it seems very probable that the 2γ mode dominates over $\pi^+\pi^-\gamma$ for a mass of 310 MeV. Therefore this assignment appears very unlikely.

0^{--} : Here we come to a conclusion similar to that for 0^{+-} not so likely.

1^{++} : Jacobsohn and Henley calculated the ratio $\pi^+\pi^-\gamma/\pi^02\gamma$ to be 25/1 for a mass of 550 MeV.¹⁰ We would expect this ratio to be much less at 310 MeV for two reasons. The $J=1$ angular-momentum barrier in $\pi^+\pi^-\gamma$ is much more important so close to threshold. Also, the relative phase space is decreased. Therefore, we might expect the $\pi^02\gamma$ mode to be dominant. However, this type of consideration is subject to criticism, and therefore the 1^{++} assignment is not excluded.

1^{+-} : The $\pi^0\gamma$ mode appears very unlikely on the basis of our data.

1^{-+} : We have the same situation as for 1^{++} , i. e., not excluded.

1^{--} : We have the same situation as for 1^{+-} , i. e., very unlikely.

V. CONCLUSIONS

The measurement of charged pions and γ -rays in coincidence with He^3 in the region of the ABC anomaly gives no indication that the anomaly is anything other than an enhancement in the $I=0$ two-pion production. However, the possible assignments 0^{+-} , 0^{--} , 1^{++} , and 1^{-+} , are not definitely excluded. The first two of these appear not so likely on the basis of the data. The last two appear doubtful on the basis of an extrapolation of the calculations of Henley and Jacobsohn.¹⁰ However these considerations are not sufficient to exclude them as possible assignments.

In the following paper (IV) the width of the anomaly is compared with the experimental resolution. The apparent lifetime of the order of 10^{-23} sec. makes 0^{++} the only reasonable assignment. All other modes are expected to have lifetimes 10^3 to 10^4 times this.

ACKNOWLEDGMENTS

We wish to acknowledge the generous assistance of Messers R. B. Bacastow, R. L. Beck, P. B. Beilin, G. M. Bingham, J. B. Czirr, H. W. Kruger, T. Maung, and R. E. Shafer who helped in the experiment. We are grateful to Messers Edwin F. McLaughlin and William L. Pope and the hydrogen target crew for design, testing, and installation of the high-pressure gas targets. Finally, we wish to thank Mr. James Vale and the entire cyclotron crew for capable and reliable operation of the cyclotron.

FOOTNOTES AND REFERENCES

- * This work was performed under the auspices of the U. S. Atomic Energy Commission.
- § Paper III in a series comprising UCRL-10407, 10408, 10409, and 10410.
- † Present address: The Enrico Fermi Institute for Nuclear Studies, The University of Chicago, Chicago, Illinois.
- ‡ Present address: Department of Physics, University of Illinois, Urbana, Illinois.
1. See previous papers I and II; see N. E. Booth, A. Abashian, and K. M. Crowe, Phys. Rev. Letters 7, 35 (1961), and A. Abashian, N. E. Booth, and K. M. Crowe, Phys. Rev. Letters 5, 258 (1960).
 2. B. C. Maglic, L. W. Alvarez, A. H. Rosenfeld, and M. L. Stevenson, Phys. Rev. Letters 7, 178 (1961).
 3. See for example R. G. Sachs and B. Sakita, Phys. Rev. Letters 6, 306 (1961); see also, the discussion following N. E. Booth, A. Abashian, and K. M. Crowe, Revs. Modern Phys. 33, 393 (1961).
 4. Y. Nambu, Phys. Rev. 106, 1366 (1957).
 5. N. E. Booth, A. Abashian, and K. M. Crowe, Phys. Rev. Letters 7, 35 (1961).
 6. A. Abashian, N. E. Booth, and K. M. Crowe, Phys. Rev. Letters 5, 258 (1960).
 7. Quentin A. Kerns, Lawrence Radiation Laboratory Electronic Engineering Note EET-743, 1961 (unpublished).
 8. William A. Wenzel, Lawrence Radiation Laboratory Report UCRL-8000, October 1957 (unpublished).
 9. There is also the suggestion of Akimov et al. that the anomaly may be associated with threshold effects due to the endothermic process

$\pi^0 + \pi^0 \rightarrow \pi^+ + \pi^-$ [see Yu. K. Akimov, V. I. Komarov, K. S. Marish, O. V. Savchenko, and L. M. Soroko, Nucl. Phys. 30, 258 (1962)].

This would give effectively a $2\pi^0$ decay scheme for the anomaly, which fits our data poorly. However we cannot exclude this effect as being partially responsible for the anomaly.

10. E. M. Henley and B. A. Jacobsohn, Phys. Rev. 127, 1829 (1962).

Table I. Some possible neutral mesons with $I=0$.^a

J^{PG} ^b	Dominant decay mode			Comments
	Strong	Order e	Order e^2	
0^{++}	$\pi\pi$			
0^{+-}		$2\pi\gamma$	$\pi^+\pi^-2\gamma$	$J_{\pi\pi}=2$ in $2\pi\gamma$ and $\pi^+\pi^-/\pi^0\pi^0=2/1$.
0^{-+}		$\pi^+\pi^-\gamma$	2γ	$J_{\pi\pi}=1$ in $\pi^+\pi^-\gamma$.
0^{--}		$2\pi\gamma$	$\pi^+\pi^-2\gamma$ $\pi^0e^+e^-$	$J_{\pi\pi}=2$ in $2\pi\gamma$ and $\pi^+\pi^-/\pi^0\pi^0=2/1$.
1^{++}		$\pi^+\pi^-\gamma$	$\pi^02\gamma$	$J_{\pi\pi}=1$ in $\pi^+\pi^-\gamma$.
1^{+-}		$\pi^0\gamma$		
1^{-+}		$\pi^+\pi^-\gamma$	$\pi^02\gamma$	$J_{\pi\pi}=1$ in $\pi^+\pi^-\gamma$.
1^{--}		$\pi^0\gamma$		

^a This table is a summary of the more complete table given by E. M. Henley and B. A. Jacobsohn, Phys. Rev. 128, 1394 (1962).

^b For neutral non-strange mesons of $I=0$, we have $G=C$.

Table II. Typical set of π data.^a

Normal delay	Criterion for accidentals	Accidentals	Net rate
4.87 ± 0.60	(b)	2.23 ± 0.43	2.64 ± 0.74
6.77 ± 0.51	(b)	3.13 ± 0.38	3.64 ± 0.63
4.60 ± 0.47	(a)	1.74 ± 0.20	2.96 ± 0.51
2.81 ± 0.31	(b)	1.11 ± 0.20	1.70 ± 0.37
5.28 ± 0.54	(b)	2.26 ± 0.35	3.02 ± 0.64
5.92 ± 0.54	(b)	3.56 ± 0.41	2.36 ± 0.68
4.94 ± 0.33	(a)	2.01 ± 0.31	2.93 ± 0.45
4.03 ± 0.37	(b)	1.35 ± 0.21	2.68 ± 0.42
Weighted mean =			2.58 ± 0.18

^a Magnets set for $p/z = 610$ MeV/c. Mean value of w is 342 MeV. All rates are per 100 He³ counts.

Table III. Typical set of γ data.^a

Pb in	Pb out	Net
1.10 ± 0.16	0.33 ± 0.10	0.77 ± 0.19
0.84 ± 0.14	0.25 ± 0.11	0.59 ± 0.18
0.76 ± 0.11	0.18 ± 0.09	0.58 ± 0.14
2.09 ± 0.34	0.89 ± 0.34	1.20 ± 0.48
Weighted mean = 0.65 ± 0.09		

^a See footnote a, Table II.

Table IV. Summary of π and γ coincidence data.

$(p/z)_s$ Observed He ³ momentum (MeV/c)	Mean value of w (MeV)	π			γ		
		Coinci- dences per 100 He ³	Errors		Coinci- dences per 100 He ³	Errors	
			Statis- tical	Total		Statis- tical	Total
680	251	3.59	0.48	0.60	0.91	0.35	0.40
670	268	5.91	0.44	0.55	1.04	0.47	0.54
660	284.5	7.20	0.22	0.27	0.83	0.13	0.15
650	298	5.62	0.18	0.22	0.40	0.09	0.10
640	312	5.27	0.20	0.25	0.62	0.12	0.14
630	324	3.64	0.18	0.22	0.57	0.10	0.12
610	342	2.58	0.18	0.22	0.65	0.09	0.10
590	355	2.22	0.27	0.34	0.43	0.16	0.18
550	372	1.47	0.30	0.37	0.38	0.14	0.16

Table V. Goodness of fit and its probability for four possible decay modes.

Assumed decay of anomaly	Spin and parity	χ^2	$P(\chi^2)$ (%)
2π	0^+	8.7	40
$\pi^+\pi^-\gamma$	1^-	7.2	50
2γ	0^-	18.2	2
$\pi^0\gamma$	1^+ or 1^-	21.8	1
Expected $\chi^2 = 9 - 1 = 8$			

FIGURE LEGENDS

- Fig. 1. Experimental arrangement showing D_2 target and π and γ telescope. The vertical and horizontal dimensions, respectively, of the scintillators were (1) 13 by 20 cm, (2) 10 by 18 cm, (3) 10 by 15 cm, and (4) 20 by 20 cm.
- Fig. 2. Kinematics of charged pions from $p+d \rightarrow He^3 + \pi^+ + \pi^-$ for $T_p = 743$ MeV and a He^3 laboratory angle of 11.8 deg. The angle θ_w is the laboratory angle of the vector sum of the momenta of the two pions, θ_π is the pion laboratory angle, and $dN_\pi/d\Omega$ is the pion laboratory cross section.
- Fig. 3. Test of consistency of the measurements of the π - He^3 coincidence rates. The histogram shows the deviations of 76 separate measurements in terms of their statistical errors. The dashed curve is the normal distribution. The solid curve results if the errors on the measurements are increased by 25%.
- Fig. 4. Plot of the observed π - He^3 and γ - He^3 coincidence rates as a function of w , the total energy in the 2π barycentric system. The curve at the top of the figure shows the observed He^3 rate for comparison.
- Fig. 5. Values of the pion and gamma rates divided by the corresponding points on the solid curves of Fig. 4. If the anomaly is an enhancement in the 2π production, then a horizontal straight line is expected in the two cases. The curves correspond to various assumptions about the decay mode of the anomaly.




Fig. 1






Fig. 2




MU-27802

Fig. 3



MU-27818

Fig. 4



MU-29787

Fig. 5

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

