

UCSF

UC San Francisco Previously Published Works

Title

Prenatal tricuspid valve size as a predictor of postnatal outcome in patients with severe pulmonary stenosis or pulmonary atresia with intact ventricular septum.

Permalink

<https://escholarship.org/uc/item/55j4707m>

Journal

Fetal diagnosis and therapy, 35(2)

ISSN

1015-3837

Authors

Lowenthal, Alexander
Lemley, Breniel
Kipps, Alaina K
et al.

Publication Date

2014

DOI

10.1159/000357429

Peer reviewed

Prenatal Tricuspid Valve Size as a Predictor of Postnatal Outcome in Patients with Severe Pulmonary Stenosis or Pulmonary Atresia with Intact Ventricular Septum

Alexander Lowenthal^a Breniel Lemley^a Alaina K. Kipps^a Michael M. Brook^a
Anita J. Moon-Grady^{a, b}

Departments of ^aPediatrics and ^bFetal Treatment Center, University of California, San Francisco, Calif., USA

Key Words

Pulmonary atresia with intact ventricular septum · Prenatal predictors · Postnatal outcome · Tricuspid regurgitation

Abstract

Introduction: Tricuspid valve (TV) size at birth correlates with intervention strategy in patients with severe pulmonary stenosis (SPS) or pulmonary atresia/intact ventricular septum (PA/IVS). Prenatal features that might predict postnatal TV size have not been well studied. We hypothesized that prenatal echocardiographic measurements predict the postnatal TV Z-score in fetuses with SPS and PA/IVS. **Materials and Methods:** We identified 16 neonates (gestational age 28 ± 4.8 weeks) with a fetal diagnosis of SPS or PA/IVS from 2001 to 2010. Measurements were performed offline. ROC (receiver operating characteristic) analysis was used to generate AUC (areas under the curve) for each of the variables. **Results:** AUC was 0.94 for tricuspid to mitral valve (TV/MV) ratio, 0.88 for TV Z-score, and 0.85 for TV inflow duration. A cut-off value of >0.63 for TV/MV yielded a sensitivity of 78%, specificity of 100% for predicting postnatal TV Z-score >-3 . Neonates with TV Z-score ≥ -3 and all fetuses with antegrade flow across the pulmonary valve or more than moderate tricuspid regurgitation had biventric-

ular circulation in follow-up. **Conclusion:** Fetal TV/MV >0.63 predicts favorable TV Z-score at birth in patients with SPS and PA/IVS. Antegrade pulmonary valve flow and more than moderate tricuspid regurgitation also conferred a favorable outcome.

© 2014 S. Karger AG, Basel

Introduction

Severe pulmonary stenosis (SPS) and pulmonary atresia with intact ventricular septum (PA/IVS) are cardiac anomalies in which the right ventricular (RV) outflow is obstructed with varying degrees of RV hypoplasia in those with PA/IVS. The majority of both PA/IVS and SPS have no known underlying genetic/chromosomal cause or affiliation with any other major congenital malformations [1]. Depending on in utero RV growth in the face of this obstruction, SPS and PA/IVS may exhibit a spectrum of severity, requiring postnatal interventions including balloon pulmonary valvuloplasty alone, surgical biventricular repair, initial palliation with shunt and bidirectional Glenn with RV outflow tract reconstruction (so-called ‘1.5 ventricle repair’), or staged single-ventricle palliation.

PA/IVS can be diagnosed in mid-gestation due to an abnormal four-chamber view and sometimes is made in the first trimester as either atresia or severe stenosis [2, 3]. Albeit it is not uncommon for the RV to be of a relatively good size at 18–22 weeks despite even severe obstruction, such examples may be overlooked in screening programs. This is of importance as RV growth may be subsequently impaired leading to a much smaller ventricular size at birth [4, 5].

Thus, both postnatally and, more importantly prenatally, at diagnosis the counseling health practitioner has to be able to predict the likeliness of a univentricular (including the possibility of 1.5 ventricle) (UV) or biventricular (BV) repair, and if UV repairs appears the imminent outcome, whether fetal pulmonary valve balloon valvuloplasty with its associated risks should be offered [6–8].

In a landmark paper reviewing the outcome of PA/IVS, a tricuspid valve (TV) size at birth with a Z-score <-3 was associated with single-ventricle outcome [9]. There have been previous articles that attempted to predict BV vs. UV outcome in this unique patient population utilizing fetal morphologic and hemodynamic parameters [6–8, 10, 11]. Prenatal echocardiographic features that might specifically predict postnatal TV size Z-score have not been as well studied.

Our objective was to determine whether prenatal TV size or other fetal echo-derived cardiac variables can predict favorable postnatal TV diameter in fetuses with SPS and PA/IVS.

Material and Methods

Review of medical records was approved by the Committee for Human Research at the University of California, San Francisco (UCSF). We searched the UCSF Benioff Children's Hospital Fetal Cardiovascular Program echocardiography database for all fetal echocardiograms performed and interpreted as SPS or PA/IVS between 2001 and 2010. We identified 16 neonates (gestational age 28 ± 4.8 weeks) with a fetal diagnosis of SPS or PA/IVS. PA/IVS was diagnosed by the absence of flow across the pulmonary valve and SPS by the presence of thickened and domed pulmonary valve cusps with a pinhole jet of flow, and in both, with Doppler evidence of reversal of flow in the ductus arteriosus [12]. The main predictors that we evaluated were: TV dimensions, TV/MV ratio, TV inflow duration, RV/LV length ratio, degree of TR, flow patterns in TV, PV, systemic veins and DA.

All echocardiograms were reviewed independently by two echocardiographers unaware of postnatal outcomes (A.M.G., A.K.K.) for confirmation of anatomy and measurements of TV, pulmonary valve, mitral valve (MV), and right heart dimensions. Z-scores were generated to normalize for differences in size and gestational age (based on previously published normative data)

[13]. Subsequently, postnatal echocardiograms and medical records from these patients were reviewed. Interventions including single-ventricle palliation and BV repair were recorded. The presence of retrograde blood flow in the ductus arteriosus, presence/degree of tricuspid regurgitation (color Doppler assessment, as follows: small/narrow jet extending only partly into the receiving chamber graded as mild, larger/wider jet extending midway or further into the atrium as moderate, and a wider/broad jet with associated right atrial enlargement), and sinusoids were assessed. Venous Doppler waveforms of the ductus venosus and the inferior vena cava were classified as normal or abnormal (fig. 1).

PA/IVS and SPS were defined as obstruction to the RV outflow necessitating ductal derived pulmonary blood flow via reversal of flow across the ductus arteriosus during gestation and maintenance of ductal patency postnatally.

Statistical Analysis

T test was used for comparisons in the case of continuous variables and Fisher's exact testing for categorical variables. A p value <0.05 was considered statistically significant. Receiver operator curves (ROC) were generated for continuous variables.

Results

During the study period, 22 fetuses met inclusion criteria; of these, there were 7 pregnancy terminations (all prior to 24 weeks' gestation). From the continuing pregnancies, 9 newborns had TV Z-scores >-3 at birth and 6 had Z-scores <-3 . All of the patients with an initial TV Z-score >-3 underwent balloon valvuloplasty alone or in combination with surgical relief of obstruction and were considered BV (including 1 with superior cavopulmonary anastomosis maintained, so-called 1.5 ventricle circulation) at the time of last follow-up. All of the patients with Z-scores <-3 underwent surgically staged procedures resulting in single-ventricle palliation (fig. 2).

Data derived from the initial echocardiograms are listed in table 1. There was no significant difference in gestational age at diagnosis in either group. Patients who had PV forward flow, moderate to severe tricuspid regurgitation, or normal venous Doppler patterns all had a postnatal Z-score >-3 . Patients who had sinusoids all had a postnatal Z-score of <-3 . A statistically significant difference was found between those with a neonatal TV Z-score >-3 versus those with a Z-score <-3 in the following variables: fetal TV/MV ratio (0.74 ± 0.2 vs. 0.4 ± 0.2 ; $p = 0.002$), fetal TV Z-score (-2.28 ± 1.8 vs. -6.45 ± 2.8 ; $p = 0.003$), fetal TV inflow duration (0.37 ± 0.06 vs. 0.27 ± 0.09 ; $p = 0.046$) and fetal RV/LV length ratio (0.72 ± 0.2 vs. 0.38 ± 0.1 ; $p = 0.004$) (table 1).

Using these continuous variables as diagnostic tests for an outcome of a Z-score >-3 , ROC curves (fig. 3)




Fig. 1. Illustration of select fetal echocardiographic data collected. For 2D measurements, markers demonstrate the method of measurement; valve annulus was measured in diastole at hinge points of the respective (tricuspid, TV, and mitral, MV) valves and right and left ventricular (RV, LV) length measured at end-diastole when chamber size was at its largest. Venous spectral Doppler patterns were scored as normal or abnormal as illustrated. DV = Ductus venosus; IVC = inferior vena cava.

yielded the following: for TV/MV ratio areas under the curve (AUC) was 0.94, with a cut-off of 0.63 yielding a specificity of 100% and sensitivity of 78%. For RV/LV length ratio, AUC was 0.90, with a cut-off of 0.54 yielding a specificity of 83% and a sensitivity of 89%. For TV Z-score, AUC was 0.88, with a cut-off of -4 yielding a specificity of 83% and sensitivity of 90%. For TV inflow duration as a percentage of the total cardiac cycle, AUC was 0.85, with a cut-off of 0.31 yielding a specificity of 80% and sensitivity of 83%. More than mild tricuspid regurgitation and presence of antegrade flow across the pulmonary valve in fetal life were never present in the patients with an unfavorable TV Z-score at birth, suggesting that these findings confer a favorable advantage.

Discussion

This retrospective study identified several fetal ultrasound markers that are strong predictors of postnatal TV Z-score >-3 , a strong known predictor for BV outcome. Fetal TV/MV ratio demonstrated the best trade-off of sensitivity and specificity for predicting a favorable postnatal TV Z-score of >-3 . Using fetal TV/MV ratio, patients with a value >0.63 consistently had a postnatal Z-score >-3 , which in our series was secondarily always correlated with a BV outcome. Utilizing this ratio avoids the need for gestational age-specific normal data. In addition, none of the fetuses in this small series who presented with demonstrable forward flow across the pulmonary valve or with more than moderate TR required UV palliation.




Fig. 2. Flow diagram showing outcomes for the 22 fetuses identified from our database.

Table 1. Data derived from the initial echocardiograms

Fetal echocardiographic data	Neonatal TV Z >-3 (n = 9)	Neonatal TV Z <-3 (n = 6)	p value
GA at Dx, weeks (±SD)	27.1±4.8	29.5±3.8	NS
TV/MV ratio (±SD)	0.74±0.2	0.4±0.2	0.002
TV Z score (±SD)	-2.28±1.8	-6.45±2.8	0.003
TV inflow duration (±SD)	0.37±0.06	0.27±0.09	0.046
RV/LV length ratio (±SD)	0.72±0.2	0.38±0/1	0.004
None/mild TR	6	6	NS
Moderate/severe TR	4	0	see Discussion
PV forward flow	5	0	see Discussion
TV monophasic	1	4	NS
Abnormal systemic venous Dopplers	5	5	NS
Retrograde DA flow	7*	6	NS

DA = Ductus arteriosus; Dx = diagnosis; GA = gestational age; LV = left ventricle; MV = mitral valve annulus; NS = not significant; RV = right ventricle; TR = tricuspid regurgitation; TV = tricuspid valve annulus.

* Two fetuses had equivocal imaging on the initial study but on follow-up fetal examination and in our database had flow reversal in the ductus arteriosus.

As prenatal diagnosis of PA/IVS has been associated with a high rate of terminations of pregnancy ranging between 61 and 82% due to concerns regarding RV growth [14, 15], early and accurate prediction of postnatal outcome is of great importance as many parents decide to continue the pregnancy when BV repair is likely. Several studies have looked at prenatal determinants of postnatal

outcome in fetuses with PA/IVS (table 2), many of which have had findings similar to ours.

In keeping with our study, Peterson et al. [10] showed that fetal TV measurements had similar predictive characteristics to postnatal TV measurements, with a fetal TV Z-score of ≤ -4 beyond 23 weeks of gestation or absolute TV annulus of ≤ 5 mm beyond 30 weeks of gestation pre-




Fig. 3. ROC curves generated for each of the continuous variables reported. Individual values for AUC are shown. LV = Left ventricle; MV = mitral valve; RV = right ventricle; TV = tricuspid valve.

dicted an eventual single-ventricle repair or death. Salvin et al. [11] showed that in 13 fetuses with a mid-gestation fetal TV Z-score <-3 , 1 achieved BV repair compared with all of those with a TV Z-score >-3 achieving a BV repair. They found that a fetal TV Z-score >-3 correlated with a BV outcome in all but 1 of their patients. The same results were obtained after applying the cut-off value of the TV Z-score proposed by Gardiner et al. [8] in fetuses before 26 weeks of gestation. Our results suggest that a smaller TV Z-score of -4 may still be compatible as a good outcome.

The first published multiparameter scoring system studied 27 patients and included four variables: TV/MV ratio, RV/LV length ratio, indexed tricuspid inflow duration and the presence of ventriculocoronary connections [6]. Looking at UV palliation as an outcome, Roman et al. [6] found that a cut-off value of <0.7 for TV/MV ratio yielded a sensitivity of 100% and a specificity of 75%, which is similar to our findings. A slightly higher cut-off for TV/MV ratio of >0.83 was utilized in the 4-point scoring system proposed by Gomez-Montes et al. [7]. Utilizing statistical modeling, Gardiner et al. [8] devised the right atrial pressure (RAP) score to predict single versus BV outcome in the fetus with PA/IVS. The RAP score is a combination of severity of tricuspid regurgitation, waveform characteristics of the ductus venosus and restriction

of the interatrial septum. They concluded that the RAP score is a good independent predictor of early outcome with a RAP score >3 predicting a BV outcome with an AUC of 0.833. When they combined the RAP score with morphological measurements (specifically Z-scores of the tricuspid and pulmonary valve and tricuspid to mitral valve ratio), they were able to predict a BV circulation with a sensitivity of 92% and specificity of 100% before 26 weeks. In their study the detection of fetal coronary fistulae did not necessarily preclude a BV circulation.

In our study, moderate to severe tricuspid regurgitation during fetal life actually served as a marker for post-natal tricuspid Z-scores of >-3 , suggesting that tricuspid regurgitation might be a beneficial factor in TV size. This is in agreement with Iacobelli et al. [16] who found that lack of tricuspid regurgitation in PA/IVS confers a higher risk of ventriculocoronary connections and subsequently a higher risk of needing single-ventricle palliation post-natally. Conversely, the degree of TV regurgitation did not differ between the two outcome groups in the study performed by Roman et al. [6].

Venous Doppler abnormality was not found to be predictive of outcome in our series. This is not surprising, given the not infrequent observation of abnormal venous Doppler pattern consistent with 'a' wave transmission as reported by Berg et al. [17] in patients with

Table 2. Published studies of predictors of outcome in the prenatal diagnosis of PA/IVS

Reference (first author)	Number of subjects studied: PA/IVS or SPS	Mean age of gestation when first studied, weeks	Percentage that achieved biventricular circulation	Predictors of outcome		Comments
				predictors of good/BV outcome	predictors of poor/UV outcome	
Peterson (2006) [10]	28	16–37	42		TV Z-score ≤ -4 after 23 weeks; TV annulus ≤ 5 mm after 30 weeks; RV/LV length or width < 0.5 and/or no tricuspid regurgitation	
Salvin (2006) [11]	25	18–29	30	TV Z-score > 3 ; rate of growth of TV		More likely to achieve a BV repair
Roman (2007) [6]	27	19–31	70		RV/LV length < 0.6 ; TV/MV maximal diameter < 0.7 ; TV inflow $< 31.5\%$ cardiac cycle length; presence of coronary fistulae	Before 31 weeks, if 3/4 criteria were met: 100% sensitivity and 75% specificity for a non-BV repair
Gardiner (2008) [8]	21 (3 of the patients underwent fetal valvuloplasty)	16–34	52	PV Z-score > -1 or TV Z-score > -3.4 before 23 weeks; median TV Z-score > -3.95 before 26 weeks gestational; median PV Z-score > -2.8 and medium TV/MV ratio > 0.7 at 26–31 weeks; median TV Z-score > 3.9 and medium TV/MV ratio > 0.59 after 31 weeks		TV Z-score was a good predictor at all ages
Iacobelli (2008) [16]	17	24–32			Absence of tricuspid regurgitation TV/MV ratio $< 0-56$	
Gomez-Montes (2011) [7]	16	20–28	75		Scoring system for non-BV outcome: TV/MV ratio ≤ 0.83 ; PV/AV ratio ≤ 0.75 ; TV inflow duration/cardiac cycle length $\leq 36.5\%$; RV/LV length ratio ≤ 0.64	Before 28 weeks, if 3/4 markers are present, this predicts a non-BV outcome with sensitivity of 100% and specificity of 92% (both 100% if all four criteria are fulfilled)
Lowenthal [current paper]	15	22–33	60	TV/MV > 0.63 antegrade PV flow $>$ moderate TR		Predicts favorable TV Z-score at birth leading to BV repair

LV = Left ventricle; MV = mitral valve; RV = right ventricle; TV = tricuspid valve; PA/IVS = pulmonary atresia intact ventricular septum; PV = pulmonary valve.

right heart obstructive lesions. This may be due to alterations in atrial and/or ventricular compliance in this population, rather than a harbinger of impending hydrops/death as it may be in fetuses without structural abnormality.

The most significant limitations to this study, though not different from all other studies published on this subject, include the retrospective study design and the small number of patients due in part to elective termination of a large number of identified pregnancies. We did

not have longitudinal data for analysis during these pregnancies; TV annulus growth may be impaired through gestation (crossing Z-score lines) as is true in all forms of left and right outflow obstruction [4]. RV length as well may progressively shorten in critical outflow lesions with little ventricular inflow [6–8]. Unfortunately, due to the high pregnancy termination rates in our study and in others, there are not sufficient available data to prognosticate well-based earlier observations and subsequent growth velocities, therefore extrapola-

tion of our results should be limited to the gestational ages reported. However, it is clear that with the addition of our small series that a definitive scoring system for this complex and intriguing question will only stem from a concerted effort by a multitude of centers pooling their data. An additional limitation of our study that is also inherent in other studies is the inconsistency that may be introduced in using different published or internally derived Z-score determination algorithms. Certainly larger series combining institutional experience will need to give attention to the method by which Z-score formulas are chosen and adhere to consistency in this regard.

We conclude that a fetal TV/MV ratio >0.63 is a strong prenatal predictor of a favorable postnatal TV Z-score at birth in patients with SPS and PA/IVS. In addition, antegrade pulmonary valve flow and more than moderate tri-

cuspid regurgitation also conferred a favorable outcome. Though postnatal Z-scores in the range of -3 are usually considered borderline for BV outcome, our results suggest that a smaller TV Z-score of as small as -4 may still be compatible as a good outcome in some patients. This information will be helpful in counseling patients regarding both diagnosis and prognosis for their fetus.

Acknowledgements

This publication was supported by NIH/NCRR UCSF-CTSI Grant No. UL1 RR024131. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the NIH. B. Lemley was supported by the Biomedical and Health Sciences Internship Program at the University of California San Francisco and by the National Step-Up program funded by NIH/NIDDK, #5R25DK078382.

References

- Shinebourne EA, Rigby ML, Carvalho JS: Pulmonary atresia with intact ventricular septum: from fetus to adult: congenital heart disease. *Heart* 2008;94:1350–1357.
- Allan LD, Crawford DC, Tynan MJ: Pulmonary atresia in prenatal life. *J Am Coll Cardiol* 1986;8:1131–1136.
- Maeno YV, Boutin C, Hornberger LK, McCrindle BW, Cavalle-Garrido T, Gladman G, Smallhorn JF: Prenatal diagnosis of right ventricular outflow tract obstruction with intact ventricular septum, and detection of ventriculocoronary connections. *Heart* 1999;81:661–668.
- Hornberger LK, Need L, Benacerraf BR: Development of significant left and right ventricular hypoplasia in the second and third trimester fetus. *J Ultrasound Med* 1996;15:655–659.
- Hornberger LK, Sanders SP, Sahn DJ, Rice MJ, Spevak PJ, Benacerraf BR, McDonald RW, Colan SD: In utero pulmonary artery and aortic growth and potential for progression of pulmonary outflow tract obstruction in tetralogy of Fallot. *J Am Coll Cardiol* 1995;25:739–745.
- Roman KS, Fouron JC, Nii M, Smallhorn JF, Chaturvedi R, Jaeggi ET: Determinants of outcome in fetal pulmonary valve stenosis or atresia with intact ventricular septum. *Am J Cardiol* 2007;99:699–703.
- Gomez-Montes E, Herraiz I, Mendoza A, Albert L, Hernandez-Garcia JM, Galindo A: Pulmonary atresia/critical stenosis with intact ventricular septum: prediction of outcome in the second trimester of pregnancy. *Prenat Diagn* 2011;31:372–379.
- Gardiner HM, Belmar C, Tulzer G, Barlow A, Pasquini L, Carvalho JS, Daubeney PE, Rigby ML, Gordon F, Kulinskaya E, Franklin RC: Morphologic and functional predictors of eventual circulation in the fetus with pulmonary atresia or critical pulmonary stenosis with intact septum. *J Am Coll Cardiol* 2008;51:1299–1308.
- Hanley FL, Sade RM, Blackstone EH, Kirklin JW, Freedom RM, Nanda NC: Outcomes in neonatal pulmonary atresia with intact ventricular septum. A multi-institutional study. *J Thorac Cardiovasc Surg* 1993;105:406–423.
- Peterson RE, Levi DS, Williams RJ, Lai WW, Sklansky MS, Drant S: Echocardiographic predictors of outcome in fetuses with pulmonary atresia with intact ventricular septum. *J Am Soc Echocardiogr* 2006;19:1393–1400.
- Salvin JW, McElhinney DB, Colan SD, Gauvreau K, del Nido PJ, Jenkins KJ, Lock JE, Tworetzky W: Fetal tricuspid valve size and growth as predictors of outcome in pulmonary atresia with intact ventricular septum. *Pediatrics* 2006;118:e415–e420.
- Berning RA, Silverman NH, Villegas M, Sahn DJ, Martin GR, Rice MJ: Reversed shunting across the ductus arteriosus or atrial septum in utero heralds severe congenital heart disease. *J Am Coll Cardiol* 1996;27:481–486.
- Schneider C, McCrindle BW, Carvalho JS, Hornberger LK, McCarthy KP, Daubeney PE: Development of Z-scores for fetal cardiac dimensions from echocardiography. *Ultrasound Obstet Gynecol* 2005;26:599–605.
- Daubeney PE, Sharland GK, Cook AC, Keeton BR, Anderson RH, Webber SA: Pulmonary atresia with intact ventricular septum: impact of fetal echocardiography on incidence at birth and postnatal outcome. UK and Eire Collaborative Study of Pulmonary Atresia with Intact Ventricular Septum. *Circulation* 1998;98:562–566.
- Todros T, Paladini D, Chiappa E, Russo MG, Gaglioti P, Pacileo G, Cau MA, Martinelli P: Pulmonary stenosis and atresia with intact ventricular septum during prenatal life. *Ultrasound Obstet Gynecol* 2003;21:228–233.
- Iacobelli R, Pasquini L, Toscano A, Raimondi F, Michielon G, Tozzi AE, Sanders SP: Role of tricuspid regurgitation in fetal echocardiographic diagnosis of pulmonary atresia with intact ventricular septum. *Ultrasound Obstet Gynecol* 2008;32:31–35.
- Berg C, Kremer C, Geipel A, Kohl T, Germer U, Gembruch U: Ductus venosus blood flow alterations in fetuses with obstructive lesions of the right heart. *Ultrasound Obstet Gynecol* 2006;28:137–142.