

eScholarship

Combinatorial Theory

Title

RSK tableaux and box-ball systems

Permalink

<https://escholarship.org/uc/item/5700d4s0>

Journal

Combinatorial Theory, 3(2)

ISSN

2766-1334

Authors

Drucker, Ben
Garcia, Eli
Gunawan, Emily
[et al.](#)

Publication Date

2023

DOI

10.5070/C63261978

Supplemental Material

<https://escholarship.org/uc/item/5700d4s0#supplemental>

Copyright Information

Copyright 2023 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RSK TABLEAUX AND BOX-BALL SYSTEMS

Ben Drucker^{*1}, Eli Garcia^{†2}, Emily Gunawan^{‡3}, Aubrey Rumbolt^{§4}, and Rose Silver^{¶5}

¹*Pacific Northwest National Laboratory, Richland, WA, U.S.A*
ben.drucker@icloud.com

²*Massachusetts Institute of Technology, Cambridge, MA, U.S.A*
tbone@mit.edu

³*David and Judi Proctor Department of Mathematics, University of Oklahoma, Norman, OK, U.S.A*
egunawan@ou.edu

⁴*Charles H. McCann Technical School, North Adams, MA, U.S.A*
ar366599@wne.edu

⁵*Khoury College of Computer Sciences, Northeastern University, Boston, MA, U.S.A*
silver.r@northeastern.edu

Submitted: Mar 1, 2022; Accepted: Jan 19, 2023; Published: Sep, 15 2023

© The authors. Released under the CC BY license (International 4.0).

Abstract. A box-ball system is a discrete dynamical system whose dynamics come from the balls jumping according to certain rules. A permutation on n objects gives a box-ball system state by assigning its one-line notation to n consecutive boxes. After a finite number of steps, a box-ball system will reach a steady state. From any steady state, we can construct a tableau called the soliton decomposition of the box-ball system. We prove that if the soliton decomposition of a permutation w is a standard tableau or if its shape coincides with the Robinson–Schensted (RS) partition of w , then the soliton decomposition of w and the RS insertion tableau of w are equal. We also use row reading words, Knuth moves, RS recording tableaux, and a localized version of Greene’s theorem (proven recently by Lewis, Lyu, Pylyavskyy, and Sen) to study various properties of a box-ball system.

Keywords. Permutations, box-ball systems, soliton cellular automata, Young tableaux, Robinson–Schensted–Knuth correspondence, Greene’s theorem, Knuth equivalence

Mathematics Subject Classifications. 05A05, 05A17, 37B15

*Supported by the University of Connecticut Mathematics REU and NSF (DMS-1950543).

†Supported by the University of Connecticut Mathematics REU and NSF (DMS-1950543).

‡Supported by the University of Connecticut Mathematics REU and NSF (DMS-1950543) and by the Isaac Newton Institute for Mathematical Sciences (funded by EPSRC Grant Number EP/R014604/1) during the programme *Cluster algebras and representation theory*.

§Supported by the University of Connecticut Mathematics REU and NSF (DMS-1950543).

¶Supported by the University of Connecticut Mathematics REU and NSF (DMS-1950543).

1. Introduction

A *box-ball system (BBS)* is a collection of discrete time states. At each state, we have an injective map from n balls (labeled by the integers from 1 to n) to boxes (labeled by the natural numbers); each box can fit at most one ball. The dynamics come from the balls jumping according to certain rules. Let S_n denote the set of permutations on $[n] = \{1, 2, \dots, n\}$. A permutation w in S_n gives a box-ball system state by assigning the one-line notation of the permutation to n consecutive boxes. Given a BBS state at time t , we compute the BBS state at time $t + 1$ by applying one BBS move, which is the process of moving each integer to the nearest empty box to its right, beginning with the smallest. See Figure 1.1. This version of the box-ball system was introduced in [Tak93] and is an extension of the box-ball system first invented by Takahashi and Satsuma in [TS90].




Figure 1.1: Performing a BBS move on $w = 452361$.

A *soliton* is a maximal consecutive increasing sequence of balls which is preserved by all subsequent BBS moves. After a finite number of BBS moves, a box-ball system containing a configuration w will reach a *steady state*, decomposing into solitons whose sizes are weakly decreasing from right to left, that is, forming an integer partition shape. From such a state, we can construct the *soliton decomposition of the box-ball system*, denoted SD , by stacking solitons so that the rightmost soliton is placed on the first row, the soliton to its left is placed on the second row, and so on. We obtain a tableau where each row is increasing but which may or may not be standard. The *soliton decomposition of a permutation w* is the soliton decomposition of the box-ball system containing w .

Figure 1.2 shows the state of the box-ball system containing $w = 452361$ from $t = 0$ to $t = 4$. Note that steady state is first reached at $t = 3$. The soliton decomposition of $w = 452361$ is the tableau

$$SD(w) = \begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & 5 & \\ \hline 4 & & \\ \hline \end{array}.$$

In this example, the soliton decomposition is a standard tableau, but most permutations have soliton decompositions which are not standard. The tableau $SD(w)$ has shape $(3, 2, 1)$. We will refer to the shape of the soliton decomposition as the *BBS soliton partition*.

Theorem A (Theorem 4.2). *Suppose w is a permutation. Then the following are equivalent:*

1. $SD(w) = P(w)$.
2. $SD(w)$ is a standard tableau.
3. The shape of $SD(w)$ equals the shape of $P(w)$.

The key ingredients of our proof are Greene’s theorem (Theorem 2.2) and a result of Fukuda which says that the RS insertion tableau is an invariant of a box-ball system (Theorem 3.3). The proof that part (3) implies part (2) was suggested to us by Darij Grinberg.

1.2. Tableau reading words

We study the connection between steady-state configurations and row reading words.

Proposition B (Proposition 5.1). *A permutation r is in steady state if and only if r is the row reading word of a standard tableau.*

Next, we represent a box-ball system state as an array containing integers from 1 to n called the *configuration array*. This array has increasing rows but not necessarily increasing columns; it also may not have a valid skew shape and it may be disconnected. Proposition B turns out to be a special case of the following.

Proposition C (Proposition 5.2). *A BBS configuration w is in steady state if and only if the configuration array of w is a standard skew tableau whose rows are weakly decreasing in length.*

We will prove Proposition C in Section 5 using the carrier algorithm. Note that Proposition C is a corollary of a characterization for steady state given by Lewis, Lyu, Pylyavskyy, and Sen in [LLPS19, Proof of Lemma 2.1 and 2.3].

1.3. Recording tableaux and time to steady state

We also study the relationship between the RS recording tableau of a permutation and the behavior of its box-ball system. The number of BBS moves required for a permutation w to reach steady state is called the *steady-state time* of w . For example, as illustrated in Figure 1.2, the steady-state time of the permutation 452361 is 3.

Theorem D (Theorem 6.7). *If $n \geq 5$, let*

$$\widehat{Q} := \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline n & \\ \hline \end{array} \dots \begin{array}{|c|c|} \hline n-2 & n-1 \\ \hline \end{array}.$$

If $Q(w) = \widehat{Q}$, then w first reaches steady state at time $n - 3$.

This particular recording tableau is special; we conjecture that all other permutations in S_n have steady-state time smaller than $n - 3$.

Conjecture 1.1. A permutation in S_n whose recording tableau is not equal to \widehat{Q} has steady-state time smaller than $n - 3$.

Furthermore, we conjecture that Theorem D is a special case of the following general phenomenon.

Conjecture 1.2. If two permutations v and w are such that $Q(v) = Q(w)$, then v and w have the same steady-state time.

Conjecture 1.2 is proven in a sequel to this paper [CFG⁺22].

Remark 1.3. Conjecture 1.2 would simplify the proof of Theorem D; it would simply require demonstrating that one single permutation whose recording tableau is \widehat{Q} has steady-state time $n - 3$.

1.4. Types of Knuth moves

The RS insertion tableau is preserved under any Knuth move [Knu70]. In contrast, the soliton decomposition is only preserved under certain types of Knuth moves.

Definition 1.4 (Knuth Moves). Suppose $v, w \in S_n$ and $x < y < z$.

1. We say that v and w differ by a Knuth relation of the *first kind* (K_1) if

$$v = v_1 \dots yxz \dots v_n \text{ and } w = v_1 \dots yzx \dots v_n \text{ or vice versa.}$$

2. We say that v and w differ by a Knuth relation of the *second kind* (K_2) if

$$v = v_1 \dots xzy \dots v_n \text{ and } w = v_1 \dots zxy \dots v_n \text{ or vice versa.}$$

In addition, We say that v and w differ by a Knuth relation of *both kinds* (K_B) if they differ by a Knuth relation of the first kind (K_1) and of the second kind (K_2), that is,

$$v = v_1 \dots y_1 x z y_2 \dots v_n \text{ and } w = v_1 \dots y_1 z x y_2 \dots v_n \text{ or vice versa}$$

where $x < y_1 < z$ and $x < y_2 < z$.

Note that, when we apply a K_1 move (respectively, a K_2 move), the move may or may not be a K_B move. If we apply a K_B move, then it is both a K_1 move and a K_2 move.

A *proper K_1 move* is a K_1 move which is not K_B , and a *proper K_2 move* is a K_2 move which is not K_B .

When performing a Knuth move, if we replace an “ xz ” pattern with a “ zx ” pattern, we denote this with a superscript “+.” Otherwise, if we replace a “ zx ” pattern with an “ xz ” pattern, we denote this with a superscript “-.” For example, if $x < y_1 < z$ and $x < y_2 < z$, the move $y_1 x z y_2 \mapsto y_1 z x y_2$ is denoted K_B^+ .

We say that v and w are *Knuth equivalent* if they differ by a finite sequence of Knuth relations.

Using the localized version of Greene’s Theorem given in Section 2.2, we prove a partial characterization of the BBS soliton partition in terms of types of Knuth moves.

Theorem E (Theorem 7.1). *If v and w are related by a sequence of Knuth moves containing an odd number of K_B moves, then $\text{SD}(v) \neq \text{SD}(w)$. If v and w are related by a sequence of non- K_B Knuth moves, then $\text{sh SD}(v) = \text{sh SD}(w)$.*

We also use non- K_B Knuth moves to give a family of permutations which have steady-state time 1.

Theorem F (Theorem 7.4). *Let r be the row reading word of a standard tableau. If w is a permutation which is related to r by one proper K_1 move or one proper K_2 move, then the steady-state time of w is 1.*

The paper is organized as follows. In the next two sections, we review materials in the literature that we will use to prove our results. First, we review Greene’s theorem in Section 2.1 and Lewis, Lyu, Pylyavskyy, and Sen’s localized Greene’s theorem in Section 2.2. Next, we review Fukuda’s carrier algorithm and its connection to the RS insertion tableaux in Section 3. In Section 4, we prove Theorem A. In Section 5, we define the configuration array and use the carrier algorithm to prove Proposition C. Section 6 is devoted to the proof of Theorem D. We prove the two results involving types of Knuth moves (Theorem E and Theorem F) in Section 7.

2. Greene’s theorem and a localized version of Greene’s theorem

In the 1970s, Greene showed that the RS partition of a permutation and its conjugate record the numbers of disjoint unions of increasing and decreasing sequences of the permutation, which we explain in Section 2.1. Lewis, Lyu, Pylyavskyy, and Sen recently showed that the BBS soliton partition of a permutation and its conjugate record a localized version of Greene’s theorem statistics. They studied an alternate version of the box-ball system, so in Section 2.2 we reframe their result to match our box-ball convention.

2.1. Greene’s theorem and RS partition

In this section, we review Greene’s theorem [Gre74, Theorem 3.1], which states that the RS partition of a permutation and its conjugate record the numbers of disjoint unions of increasing and decreasing sequences of the permutation. For more details, see for example Chapter 3 of the textbook [Sag01].

Definition 2.1 (longest k -increasing and k -decreasing subsequences). A subsequence σ of w is called k -increasing if, as a set, it can be written as a disjoint union

$$\sigma = \sigma_1 \sqcup \sigma_2 \sqcup \cdots \sqcup \sigma_k$$

where each σ_i is an increasing subsequence of w . If each σ_i is a decreasing subsequence of w , we say that σ is k -decreasing. Let

$i_k(w)$ denote the length of a longest k -increasing subsequence of w
and $d_k(w)$ denote the length of a longest k -decreasing subsequence of w .

Theorem 2.2 ([Gre74, Theorem 3.1]). *Suppose $w \in S_n$. Let $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ denote the RS partition of w , that is, let $\lambda = \text{sh } P(w)$. Let $\mu = (\mu_1, \mu_2, \mu_3, \dots)$ denote the conjugate of λ . Then, for any k ,*

$$i_k(w) = \lambda_1 + \lambda_2 + \dots + \lambda_k,$$

$$d_k(w) = \mu_1 + \mu_2 + \dots + \mu_k.$$

Example 2.3. Let $w = 5623714$. For short, we write $i_k := i_k(w)$ and $d_k := d_k(w)$. The longest 1-increasing subsequences are

$$567, \quad 237, \quad \text{and} \quad 234.$$

The longest 2-increasing subsequence is given by

$$562374 = 567 \sqcup 234.$$

A longest 3-increasing subsequence (among others) is given by

$$5623714 = 56 \sqcup 237 \sqcup 14.$$

Thus,

$$i_1 = 3, \quad i_2 = 6, \quad \text{and} \quad i_k = 7 \text{ if } k \geq 3.$$

Similarly, the longest 1-decreasing subsequences are

$$521, \quad 621, \quad 531, \quad \text{and} \quad 631.$$

A longest 2-decreasing subsequence (among others) is given by

$$52714 = 521 \sqcup 74.$$

A longest 3-decreasing subsequence (among others) is given by

$$5623714 = 52 \sqcup 631 \sqcup 74.$$

Thus,

$$d_1 = 3, \quad d_2 = 5, \quad \text{and} \quad d_k = 7 \text{ if } k \geq 3.$$

By Theorem 2.2, the RS partition is equal to $\lambda = (i_1, i_2 - i_1, i_3 - i_2) = (3, 3, 1)$ and the conjugate of the RS partition is $\mu = (d_1, d_2 - d_1, d_3 - d_2) = (3, 2, 2)$. We can verify this by computing the RS tableaux

$$P(w) = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 6 & 7 \\ \hline 5 & & \\ \hline \end{array}, \quad Q(w) = \begin{array}{|c|c|c|} \hline 1 & 2 & 5 \\ \hline 3 & 4 & 7 \\ \hline 6 & & \\ \hline \end{array}.$$

2.2. Localized Greene's theorem and BBS soliton partition

In [LLPS19, Lemma 2.1] and the blog post [Lew19], Lewis, Lyu, Pylyavskyy, and Sen presented a localized version of Greene's theorem. They studied an alternate version of the box-ball system, and in this section we reframe their result to match our box-ball convention.

Definition 2.4 (A localized version of longest k -increasing subsequences). If u is a sequence, let $i(u)$ denote the length of a longest increasing subsequence of u .

For $w \in S_n$ and $k \geq 1$, we define

$$I_k(w) = \max_{w=u_1|\dots|u_k} \sum_{j=1}^k i(u_j),$$

where the maximum is taken over ways of writing w as a concatenation $u_1 | \dots | u_k$ of consecutive subsequences. That is, we consider all ways to break w into k consecutive subsequences, sum the $i(u_j)$ values for each way, and let $I_k(w)$ be the maximum sum.

If u is a sequence of ℓ elements, an integer $m \in [\ell - 1]$ is called a *descent* of u if $u_m > u_{m+1}$.

Definition 2.5 (A localized version of longest k -decreasing subsequences). Let

$$D(u) := \begin{cases} 0 & \text{if } u \text{ is empty} \\ 1 + |\{\text{descents of } u\}| & \text{otherwise.} \end{cases}$$

For $w \in S_n$ and $k \geq 1$, we define

$$D_k(w) = \max_{w=u_1 \sqcup \dots \sqcup u_k} \sum_{j=1}^k D(u_j),$$

where the maximum is taken over ways to write w as the union of disjoint subsequences u_j of w . Notice that we only require u_1, \dots, u_k to be disjoint, *not* consecutive, in contrast to the procedure for calculating $I_k(w)$.

The following lemma is a corollary of [LLPS19, Lemma 2.1].

Lemma 2.6 (A localized version of Greene's theorem). *Suppose that $w \in S_n$. Let $\Lambda = (\Lambda_1, \Lambda_2, \Lambda_3, \dots)$ denote the BBS soliton partition of w , that is, let $\Lambda = \text{sh SD}(w)$. Let $M = (M_1, M_2, M_3, \dots)$ denote the conjugate of Λ . Then, for any k ,*

$$\begin{aligned} I_k(w) &= \Lambda_1 + \Lambda_2 + \dots + \Lambda_k, \\ D_k(w) &= M_1 + M_2 + \dots + M_k. \end{aligned}$$

Example 2.7. Let $w = 5623714$, the permutation used in Example 2.3. For short, we write $I_k := I_k(w)$ and $D_k := D_k(w)$. Then

$$\begin{aligned} I_1 &= i(w) = 3 \text{ (since the longest increasing subsequences are } 567, 237, \text{ and } 234), \\ I_2 &= 5 \text{ (witnessed by } 56|23714 \text{ or } 56237|14), \\ I_3 &= 7 \text{ (witnessed uniquely by } 56|237|14), \text{ and} \\ I_k &= 7 \text{ for all } k \geq 3. \end{aligned}$$

We have

$$\begin{aligned}
 D_1 &= D(w) = 1 + |\text{descents of } 5623714| = 1 + |\{2, 5\}| = 3, \\
 D_2 &= 6 \text{ (one can take subsequences } 531 \text{ and } 6274, \text{ among other partitions),} \\
 D_3 &= 7 \text{ (one can take subsequences } 52, 631, \text{ and } 74, \text{ among other partitions), and} \\
 D_k &= 7 \text{ for all } k \geq 3.
 \end{aligned}$$

By Lemma 2.6, we have that $\text{sh SD}(w) = (I_1, I_2 - I_1, I_3 - I_2) = (3, 2, 2)$ and its conjugate is $(D_1, D_2 - D_1, D_3 - D_2) = (3, 3, 1)$. We can verify this by computing the soliton decomposition $\text{SD}(w)$, which turns out to be the nonstandard tableau

$$\begin{array}{|c|c|c|}
 \hline
 1 & 3 & 4 \\
 \hline
 2 & 7 & \\
 \hline
 5 & 6 & \\
 \hline
 \end{array} .$$

Note that, in this example, $\text{SD}(w) \neq P(w)$, demonstrating Theorem A. Also, in this example, $\text{sh SD}(w) = (3, 2, 2)$ is smaller than $\text{sh } P(w) = (3, 3, 1)$ in the dominance partial order.

Corollary 2.8. *If $w \in S_n$, then the BBS soliton partition of w is smaller or equal to the RS partition of w in the dominance partial order.*

Proof. Let $\Lambda = (\Lambda_1, \Lambda_2, \Lambda_3, \dots)$ denote $\text{sh SD}(w)$ and let $\lambda = (\lambda_1, \lambda_2, \lambda_3, \dots)$ denote $\text{sh } P(w)$. Then, for all $k = 1, 2, \dots$, we have

$$\begin{aligned}
 \Lambda_1 + \Lambda_2 + \dots + \Lambda_k &= I_k(w) \text{ by localized Greene's theorem (Lemma 2.6)} \\
 &\leq i_k(w) \text{ since } I_k(w) \text{ gives the length of a } k\text{-increasing subsequence of } w \\
 &= \lambda_1 + \lambda_2 + \dots + \lambda_k \text{ by Greene's theorem (Theorem 2.2).} \quad \square
 \end{aligned}$$

3. Fukuda's carrier algorithm


In this section, we review the carrier algorithm and the fact that the RS insertion tableau is an invariant of a box-ball system (BBS).

3.1. Carrier algorithm

The carrier algorithm is a way to describe a BBS move as a sequence of local operations of inserting and bumping numbers in and out of a carrier filled with a weakly increasing string. A version of the carrier algorithm was first introduced in [TM97], and the version of the carrier algorithm we use in this paper comes from [Fuk04, Section 3.3]. Given a BBS state at time t , the carrier algorithm is used to calculate the state at time $t + 1$. We describe the process in Algorithm 1. Note that, after each insertion and ejection step, the sequence in the carrier is weakly increasing.

Algorithm 1 The carrier algorithm [Fuk04].

```

1: begin carrier algorithm
2: |   Set  $e := n + 1$ , so that  $e$  is considered to be larger than any ball
3: |   Set  $B :=$  the configuration of the BBS at time  $t$ , where each empty box is replaced with
   |   an  $e$  and the first (leftmost) element of  $B$  is the integer in the first (leftmost) nonempty
   |   box in the configuration and the last (rightmost) element of  $B$  is the integer in the last
   |   (rightmost) nonempty box of the configuration
4: |   Let  $\ell$  denote the number of elements (including the  $e$ 's) of  $B$ 
5: |   Fill the "carrier"  $\mathcal{C}$  —depicted —with  $n$  copies of  $e$ 
6: |   Write  $B$  to the right of  $\mathcal{C}$ 
7: |   begin Process 1: insertion process
8: |   |   for all  $i$  in  $\{1, 2, \dots, \ell\}$  do
9: |   |   |   Set  $p$  to be the  $i^{\text{th}}$  leftmost element of  $B$ 
10: |   |   |   begin element ejection process
11: |   |   |   |   if an element in  $\mathcal{C}$  is larger than  $p$  then
12: |   |   |   |   |   Set  $s :=$  smallest element in  $\mathcal{C}$  larger than  $p$ . If  $s = e$ , pick the leftmost  $e$ 
13: |   |   |   |   |   Eject  $s$  from  $\mathcal{C}$  and put it immediately to the left of  $\mathcal{C}$ 
14: |   |   |   |   |   insert  $p$  in the place of  $s$ 
15: |   |   |   |   else
16: |   |   |   |   |   Set  $s :=$  the leftmost (smallest) element in  $\mathcal{C}$ 
17: |   |   |   |   |   Eject  $s$  from  $\mathcal{C}$  and put it immediately to the left of  $\mathcal{C}$ 
18: |   |   |   |   |   ► Note: There are now  $n - 1$  elements in  $\mathcal{C}$ 
19: |   |   |   |   |   Shift each element of  $\mathcal{C}$  to the left by one
20: |   |   |   |   |   Place  $p$  in the rightmost location in  $\mathcal{C}$ 
21: |   |   |   |   |   ► Note: There are now  $n$  elements in  $\mathcal{C}$ 
22: |   |   |   |   end if
23: |   |   |   end element ejection process
24: |   |   end for
25: |   end Process 1: insertion process
26: |   begin Process 2: flushing process
27: |   |   while there are non- $e$  elements in  $\mathcal{C}$  do
28: |   |   |   Set  $p := e$ 
29: |   |   |   Perform the element ejection process (see line 16)
30: |   |   end while
31: |   end Process 2: flushing process
32: |   ► Note: The current elements to the left of  $\mathcal{C}$  correspond to the  $t + 1$  state of the BBS
33: end carrier algorithm

```

Example 3.1. We compute the configuration at time $t = 3$ of the box-ball system from Figure 1.2 by applying the carrier algorithm to the configuration at time $t = 2$. Following Algorithm 1, we set $B := 452ee136$. The carrier algorithm then proceeds as follows.

```

begin Process 1: insertion process
    eeeeee 452ee136
    e 4eeee 52ee136
    ee 45eeee 2ee136
    ee4 25eeee ee136
    ee42 5eeee e136
    ee425 eeeeee 136
    ee425e 1eeee 36
    ee425ee 13eeee 6
    ee425eee 136eee
end insertion process

begin Process 2: flushing process
    ee425eee 136eee ← e
    ee425eee1 36eeee ← e
    ee425eee13 6eeee ← e
    ee425eee136 eeeeee
end flushing process
    
```

The elements $ee425eee136$ to the left of \mathcal{C} correspond to the configuration at time $t = 3$ given in Figure 1.2.

3.2. The RS insertion tableau is an invariant of a box-ball system

Remark 3.2 ([Fuk04, Remark 4]). The carrier algorithm can be viewed as a sequence of Knuth moves. Consider the insertion of \mathbf{p} into the carrier. Note that, since our carrier can carry n elements, if $\mathbf{p} \neq e$, then the carrier must contain a number (possibly e) greater than \mathbf{p} . If $\mathbf{p} = e$, then no number in the carrier is greater than \mathbf{p} .

First, suppose $\mathbf{p} \neq e$, and let C_p denote the smallest element in the carrier which is greater than \mathbf{p} .

- (i) If C_p is the smallest element in the carrier, then the insertion process is equivalent to applying a sequence of K_1^- moves

$$\underbrace{C_p z_1 z_2 \cdots z_{\ell-1} z_\ell}_{\mathcal{C}} \mathbf{p}$$

$$\begin{array}{c}
C_p z_1 z_2 \cdots z_{\ell-1} \mathbf{p} z_\ell \\
\vdots \\
C_p z_1 \mathbf{p} z_2 \cdots z_{\ell-1} z_\ell \\
C_p \mathbf{p} z_1 z_2 \cdots z_\ell.
\end{array}$$

- (ii) If C_p is the largest element in the carrier, then the insertion process is equivalent to applying a sequence of K_2^+ moves

$$\begin{array}{c}
\underbrace{x_1 x_2 \cdots x_{m-1} x_m C_p}_{\mathbf{p}} \\
x_1 x_2 \cdots x_{m-1} C_p x_m \mathbf{p} \\
\vdots \\
x_1 C_p x_2 \cdots x_{m-1} x_m \mathbf{p} \\
C_p \underbrace{x_1 x_2 \cdots x_{m-1} x_m}_{\mathbf{p}}.
\end{array}$$

- (iii) If C_p is neither the smallest nor the largest element in the carrier, then the insertion process is equivalent to applying a sequence of K_1^- moves

$$\begin{array}{c}
\underbrace{x_1 x_2 \cdots x_{m-1} x_m C_p z_1 z_2 \cdots z_{\ell-1} z_\ell}_{\mathbf{p}} \\
x_1 x_2 \cdots x_{m-1} x_m C_p z_1 z_2 \cdots z_{\ell-1} \mathbf{p} z_\ell \\
\vdots \\
x_1 x_2 \cdots x_{m-1} x_m C_p z_1 \mathbf{p} z_2 \cdots z_{\ell-1} z_\ell \\
x_1 x_2 \cdots x_{m-1} x_m C_p \mathbf{p} z_1 z_2 \cdots z_{\ell-1} z_\ell
\end{array}$$

followed by a sequence of K_2^+ moves

$$\begin{array}{c}
x_1 x_2 \cdots x_{m-1} x_m C_p \mathbf{p} z_1 z_2 \cdots z_{\ell-1} z_\ell \\
x_1 x_2 \cdots x_{m-1} C_p x_m \mathbf{p} z_1 z_2 \cdots z_{\ell-1} z_\ell \\
\vdots \\
x_1 C_p x_2 \cdots x_{m-1} x_m \mathbf{p} z_1 z_2 \cdots z_{\ell-1} z_\ell \\
C_p \underbrace{x_1 x_2 \cdots x_{m-1} x_m}_{\mathbf{p}} z_1 z_2 \cdots z_{\ell-1} z_\ell.
\end{array}$$

Next, suppose $\mathbf{p} = e$. Then \mathbf{p} is greater than or equal to every element in the carrier, and the insertion process is equivalent to applying the trivial transformation

$$\begin{array}{c}
\underbrace{x_1 x_2 \cdots x_n}_{\mathbf{p}} \\
x_1 \underbrace{x_2 \cdots x_n}_{\mathbf{p}}.
\end{array}$$

Theorem 3.3 ([Fuk04, Theorem 3.1]). *The RS insertion tableau is a conserved quantity under the time evolution of the BBS, i.e., the RS insertion tableau is preserved under each BBS move. More precisely, let B_t be the state of a box-ball system at time t . Let $B_t^!$ be the permutation created from B_t by removing all e 's. Then $\mathbb{P}(B_t^!)$ is identical for all t .*

Example 3.4. As shown in Figure 1.2, the configurations 452361, ee45e2136, eeee452ee136, and eeeee425eee136 are in the same box-ball system. As Theorem 3.3 tells us, the permutations 452361, 452136, and 425136 have the same RS insertion tableau

$$P(452361) = P(452136) = P(425136) = \begin{array}{|c|c|c|} \hline 1 & 3 & 6 \\ \hline 2 & 5 & \\ \hline 4 & & \\ \hline \end{array}.$$

Corollary 3.5. Let w be a permutation. If r is the row reading word of $SD(w)$, then $P(w) = P(r)$.

Proof. Let r be the row reading word of $SD(w)$. By definition of the soliton decomposition tableau, we know that r is the order in which the balls of w are configured once we reach a steady state. Therefore, r is a state in the box-ball system containing w . Theorem 3.3 tells us that the RS insertion tableau is preserved under a sequence of box-ball moves, so $P(w) = P(r)$. \square

Example 3.6. Let $w = 5623714$, the permutation from Section 2, and let r be the row reading word of $SD(w)$. We have

$$SD(w) = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 7 & \\ \hline 5 & 6 & \\ \hline \end{array}, \quad r = 5627134, \quad \text{and} \quad P(w) = \begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 2 & 6 & 7 \\ \hline 5 & & \\ \hline \end{array} = P(r).$$

In Example 3.4, the soliton decomposition coincides with the RS insertion tableau of the box-ball system, but in Example 3.6 these two tableaux do not coincide. In the next section we discuss when $SD(w) = P(w)$.

4. When the soliton decomposition and the RS insertion tableau coincide

In this section, we will prove Theorem 4.2. One direction of our proof uses the following lemma, which was communicated to us by Darij Grinberg.

Lemma 4.1. Suppose S is a row-strict tableau, that is, every row is increasing (with no restrictions on the columns). Let r be the row reading word of S . If $\text{sh } S = \text{sh } P(r)$, then S is standard, that is, every column of S is increasing.

Proof. Suppose S is not standard. Then S has two adjacent entries in a column which are out of order. Indexing our rows from top to bottom and our columns from left to right, this means there is a column (say, column c) for which the entry in some row k is bigger than the entry immediately below it. Let y be the entry in the k -th row, c -th column of S , and let x be the entry immediately below it (in the $k + 1$ -th row, c -th column of S).

Since r is the row reading word of S and since each row of S is increasing, we can construct a list of k disjoint increasing subsequences of r : The first $k - 1$ increasing subsequences of r are the first $k - 1$ rows of S . The k -th increasing subsequence starts in row $k + 1$, column 1 of S , moving along the same row until we get to column c (with entry x), then going up to row k above (which has entry y), then continuing to the end of row k .

The length of the k -th increasing subsequence is larger (by 1) than the length of the k -th row of S . So the total number of letters in our list of k disjoint increasing subsequences of r is larger by 1 than the total length of the first k rows of S . Thus, Greene's theorem (Theorem 2.2) says that the total length of the first k rows of the RS insertion tableau $P(r)$ of r is larger (at least by 1) than the total length of the first k rows of S . Therefore, the shape of S is not equal to the shape of $P(r)$. \square

The following theorem gives a characterization of permutations whose soliton decompositions are equal to their RS insertion tableaux.

Theorem 4.2. *Let w be a permutation. Then the following are equivalent:*

1. $\text{SD}(w) = P(w)$.
2. $\text{SD}(w)$ is a standard tableau.
3. The shape of $\text{SD}(w)$ equals the shape of $P(w)$.

Proof. Certainly (1) implies (2) and (3). We will show that (2) implies (1) and (3) implies (2).

Let r be the row reading word of $\text{SD}(w)$. By Corollary 3.5, we have

$$P(w) = P(r). \quad (4.1)$$

First, we show that (2) implies (1). Suppose that $\text{SD}(w)$ is a standard tableau T . Since r is the row reading word of T , we have $P(r) = T$ by (1.1). Combining this equality with (4.1), we get $P(w) = P(r) = T = \text{SD}(w)$.

Next, we show that (3) implies (2). Let S denote $\text{SD}(w)$, and note that $\text{SD}(w)$ is a row-strict tableau by construction. Suppose $\text{sh } S = \text{sh } P(w)$. Since $P(w) = P(r)$ by (4.1), we have $\text{sh } P(w) = \text{sh } P(r)$, so $\text{sh } S = \text{sh } P(w) = \text{sh } P(r)$. Because S is a row-strict tableau, the permutation r is the row reading word of S , and $\text{sh } S = \text{sh } P(r)$, Lemma 4.1 tells us that S is standard. \square

Corollary 4.3. *Let w be a permutation. Then the following five statements are equivalent:*

1. $\text{SD}(w) = P(w)$.
2. $\text{SD}(w)$ is a standard tableau.
3. The shape of $\text{SD}(w)$ equals the shape of $P(w)$.
4. For all $k \geq 1$, we have

$$I_k(w) = i_k(w).$$

5. For all $k \geq 1$, we have

$$D_k(w) = d_k(w).$$

The symbols I_k and D_k are the statistics from localized Greene's theorem (Section 2.2) and i_k and d_k are the statistics from Greene's theorem (Section 2.1).

Proof. For short, we write $i_k := i_k(w)$, $I_k := I_k(w)$, $d_k := d_k(w)$, and $D_k := D_k(w)$. By localized Greene’s theorem (Lemma 2.6),

the shape of $SD(w)$ is $(I_1, I_2 - I_1, I_3 - I_2, \dots)$ and
the shape of the conjugate of $SD(w)$ is $(D_1, D_2 - D_1, D_3 - D_2, \dots)$.

By Greene’s theorem (Theorem 2.2),

the shape of $P(w)$ is $(i_1, i_2 - i_1, i_3 - i_2, \dots)$ and
the shape of the conjugate of $P(w)$ is $(d_1, d_2 - d_1, d_3 - d_2, \dots)$.

Combining these facts, we conclude that $\text{sh } SD(w) = \text{sh } P(w)$ if and only if $I_k = i_k$ for all $k \geq 1$ if and only if $D_k = d_k$ for all $k \geq 1$. □

Example 4.4. Let $w = 5623714$. From Examples 2.3 and 2.7, we have $I_2(w) = 5 < 6 = i_2(w)$. So all the other items of Corollary 4.3 must also be false.

5. Reading words and steady states

We study the steady-state configurations of a box-ball system. The main result of this section (Proposition 5.2) is a corollary of [LLPS19, Proof of Lemma 2.1 and 2.3].

5.1. Reading words of standard tableaux

The permutations which reach their steady state at time 0 are precisely the row reading words of standard tableaux.

Proposition 5.1. *A permutation r has steady-state time 0 if and only if r is the row reading word of a standard tableau.*

In particular, if r is the row reading word of a standard tableau T , then $T = SD(r)$. In the next section, the standard tableau in Proposition 5.1 is generalized to standard skew tableaux whose rows are weakly decreasing in length.

5.2. Reading words of standard skew tableaux

A BBS state can be represented as a *configuration array* containing the integers from 1 to n as follows: scanning the boxes from right to left, each *increasing run* (maximal consecutive increasing string of balls) becomes a row in the array. A string of g empty boxes indicates that the next row below should be shifted g spaces to the left. Note that this array has increasing rows but not necessarily increasing columns; it may be disconnected and it may not have a valid skew shape.

Proposition 5.2. *A BBS configuration is in steady state if and only if its configuration array is a standard (possibly disconnected) skew tableau whose rows are weakly decreasing in length.*

We will give a proof in Section 5.3.

Example 5.3. Let $w = 5623714$, the example we use in Section 2. The following are the box-ball system states from time $t = 0$ to $t = 4$ and their configuration arrays.

$t = 0$	$5623714e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>1</td><td>4</td></tr> <tr><td>2</td><td>3</td><td>7</td></tr> <tr><td>5</td><td>6</td></tr> </table>	1	4	2	3	7	5	6
1	4								
2	3	7							
5	6								
$t = 1$	$ee56e27134e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>1</td><td>3</td><td>4</td></tr> <tr><td>2</td><td>7</td></tr> <tr><td>5</td><td>6</td></tr> </table>	1	3	4	2	7	5	6
1	3	4							
2	7								
5	6								
$t = 2$	$eeee56e27e134e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>1</td><td>3</td><td>4</td></tr> <tr><td>2</td><td>7</td></tr> <tr><td>5</td><td>6</td></tr> </table>	1	3	4	2	7	5	6
1	3	4							
2	7								
5	6								
$t = 3$	$eeeeee56e27ee134e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>1</td><td>3</td><td>4</td></tr> <tr><td>2</td><td>7</td></tr> <tr><td>5</td><td>6</td></tr> </table>	1	3	4	2	7	5	6
1	3	4							
2	7								
5	6								
$t = 4$	$eeeeeeee56e27eee134e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>1</td><td>3</td><td>4</td></tr> <tr><td>2</td><td>7</td></tr> <tr><td>5</td><td>6</td></tr> </table>	1	3	4	2	7	5	6
1	3	4							
2	7								
5	6								

In this box-ball system, all configurations at time $t \geq 1$ are in steady state.

Example 5.4. The following is an example of a non-steady-state BBS configuration and its configuration array. Note that the configuration array is a standard skew tableau but its rows are not weakly decreasing in length.

$\dots e137e2469ee58e\dots$	<table border="1" style="border-collapse: collapse; text-align: center;"> <tr><td>5</td><td>8</td></tr> <tr><td>2</td><td>4</td><td>6</td><td>9</td></tr> <tr><td>1</td><td>3</td><td>7</td></tr> </table>	5	8	2	4	6	9	1	3	7
5	8									
2	4	6	9							
1	3	7								

5.3. Separation condition

A ‘separation condition’ for steady state is given in statement (43) in [LLPS19]. In Lemmas 5.5 and 5.6, we reframe this characterization for steady state in terms of our version of the box-ball system. Proposition 5.2 follows directly from these two lemmas.

Lemma 5.5 (Separation condition). *Let a BBS configuration be in steady state. Suppose two adjacent solitons L (the left soliton with length ℓ) and R (the right soliton) are separated by g empty boxes, where $g < \ell$. Then, for $i = 1, 2, \dots, \ell - g$,*

*the i -th smallest ball of the right soliton R is smaller than
the $(i + g)$ -th smallest ball of the left soliton L .*

Proof. We apply one BBS move to the configuration via the carrier algorithm. Suppose $L = L_1L_2\dots L_\ell$ and $R = R_1R_2\dots R_r$ are the two leftmost solitons.

Our initial setup with n copies of e in the carrier is

$$\underbrace{ee \cdots e}_{\text{carrier}} L_1 \cdots L_\ell \overbrace{e \cdots e}^{g \text{ copies}} R_1 \cdots R_r \cdots$$

First, we simply insert L_1, \dots, L_ℓ into the carrier. Since L is increasing, each time we insert a ball of L , we eject a copy of e . We get

$$\overbrace{e \cdots e}^{\ell \text{ copies}} \underbrace{L_1 \cdots L_\ell ee \cdots e}_{\text{carrier}} \overbrace{e \cdots e}^{g \text{ copies}} R_1 \cdots R_r \cdots \tag{5.1}$$

Next, we insert the g copies of e into the carrier and eject L_1, \dots, L_g :

$$e \cdots e \underbrace{L_1 \cdots L_g}_{\text{first } g \text{ balls}} \overbrace{L_{g+1} \cdots L_\ell}^{\ell - g \text{ balls}} ee \cdots e R_1 \cdots R_r \cdots$$

Since we started with a steady-state configuration, the left soliton L must stay intact at the end of the carrier algorithm. So, for each $i = 1, \dots, \ell - g$, as we insert R_i , we must eject L_{g+i} , and get

$$e \cdots e L_1 \cdots L_g \underbrace{L_{g+1} \cdots L_\ell}_{\ell - g \text{ balls}} \overbrace{R_1 \cdots R_{\ell - g}} \underbrace{ee \cdots e}_{\text{carrier}} R_{\ell - g + 1} \cdots R_r \cdots$$

So we must have $R_i < L_{g+i}$ for $i = 1, 2, \dots, \ell - g$, as needed.

After we insert the rest of the elements of R into the carrier, we have

$$e \cdots e L_1 \cdots L_\ell \overbrace{ee \cdots e}^{r - \ell + g \text{ copies}} \underbrace{R_1 \cdots R_r}_{\text{carrier}} ee \cdots e \cdots$$

If we have a third soliton located to the right of R , we would be in the same situation as (5.1). We then repeat the same process for the rest of the solitons and arrive at the same conclusion. \square

Lemma 5.6 (Sufficient condition for steady state). *Suppose a BBS configuration w satisfies the following.*

1. *The configuration array of w has rows of weakly decreasing length.*
2. *The configuration array of w is standard; that is, if two adjacent maximal consecutive increasing blocks L (the left block with length ℓ) and R (the right block) of w are separated by g empty boxes such that $g < \ell$, then, for $i = 1, 2, \dots, \ell - g$,*

*the i -th ball of the right block R is smaller than
the $(i + g)$ -th ball of the left block L .*

Then w is in steady state.

Proof. Suppose w is the configuration at time t . We apply the carrier algorithm to get the configuration at time $t + 1$. Suppose $L = L_1 L_2 \dots L_\ell$ and $R = R_1 R_2 \dots R_r$ are the two leftmost increasing runs (maximal consecutive increasing blocks of balls).

Prior to applying the carrier algorithm, we have

$$\underbrace{ee \dots e}_{\text{carrier}} \underbrace{L_1 \dots L_\ell}_{\text{first run}} \overbrace{e \dots e}^{g \text{ copies}} \underbrace{R_1 \dots R_r}_{\text{second run}} \dots$$

First, we insert each of L_1, \dots, L_ℓ into the carrier and eject an e each time. We get

$$\overbrace{e \dots e}^{\ell \text{ copies}} \underbrace{L_1 \dots L_\ell}_{\text{first run}} ee \dots e \overbrace{e \dots e}^{g \text{ copies}} R_1 \dots R_r \quad (5.2)$$

Next, we insert the g copies of e into the carrier and eject L_1, \dots, L_g . There are two cases: either (a) $g \geq \ell$ or (b) $g < \ell$.

(a) First, suppose that $g \geq \ell$. Then all of L_1, \dots, L_ℓ are ejected and the carrier is now empty:

$$e \dots e \underbrace{L_1 \dots L_\ell}_{\text{first run}} \overbrace{e \dots e}^{g-\ell} \underbrace{ee \dots e}_{\text{second run}} R_1 \dots R_r \dots$$

We proceed by inserting R_1, \dots, R_r into the carrier. Since R is increasing, we eject r copies of e 's:

$$e \dots e L_1 \dots L_\ell \overbrace{e \dots e}^{g-\ell} \overbrace{e \dots e}^r \underbrace{R_1 \dots R_r}_{\text{second run}} ee \dots e \dots$$

(b) Second, suppose $g < \ell$. After L_1, \dots, L_g are ejected, we have

$$e \dots e \underbrace{L_1 \dots L_g}_{\text{first } g \text{ balls}} \overbrace{L_{g+1} \dots L_\ell}^{\ell-g \text{ balls}} ee \dots e \underbrace{R_1 \dots R_r}_{\text{second run}} \dots$$

We proceed by inserting R_1, \dots, R_r into the carrier. We have $\ell \leq r$ by assumption part (1) and $R_i < L_{g+i}$ for $i = 1, 2, \dots, \ell - g$ by assumption part (2). Therefore, as we insert $R_1, \dots, R_{\ell-g}$, we must eject L_{g+1}, \dots, L_ℓ , and we get

$$e \dots e L_1 \dots L_g \underbrace{L_{g+1} \dots L_\ell}_{\ell-g \text{ balls}} \underbrace{R_1 \dots R_{\ell-g}}_{\text{second run}} ee \dots e R_{\ell-g+1} \dots R_r \dots$$

After we insert the rest of the elements of R into the carrier, we have

$$e \dots e L_1 \dots L_\ell \overbrace{ee \dots e}^{r-\ell+g} \underbrace{R_1 \dots R_r}_{\text{second run}} ee \dots e \dots$$

In both cases, at time $t+1$ there are at least $r-\ell+g$ empty boxes to the right of L . Since $\ell \leq r$, we have $g \leq r - \ell + g$, so there are at least as many empty boxes to the right of L as at time t . Furthermore, the increasing run L stays together.

If we have a third increasing run $S = S_1 \dots S_s$ to the right of R (with a gap of g' empty boxes), we would be in the same situation as (5.2). After inserting the elements of S into the carrier, we would have

$$e \dots e L_1 \dots L_\ell \overbrace{ee \dots e}^{r-\ell+g} R_1 \dots R_r \overbrace{ee \dots e}^{s-r+g'} \underbrace{S_1 \dots S_s} ee \dots e \dots$$

Again, there are at least as many empty boxes to the right of R at time $t+1$ than at time t , and R stays together.

At the end of the carrier algorithm, the increasing runs stay together, their order stays the same, and the gap of empty boxes between each pair of adjacent sequences is at least as large as at time t . The new configuration satisfies both part (1) and (2) of the assumption. By induction, subsequent carrier algorithm applications leave the order of the increasing runs unchanged, so these increasing runs are in fact solitons. \square

By the two lemmas above, we have Proposition 5.2: a box-ball configuration is in steady state if and only if (1) its configuration array has rows of weakly decreasing length and (2) each column of the configuration array is increasing.

6. A recording tableau giving $n-3$ steady-state time

In this section, we prove Theorem 6.7, which states that all permutations in S_n with a certain recording tableau have box-ball steady-state time $n-3$. We conjecture that all other permutations in S_n have steady-state time smaller than $n-3$ (Conjecture 1.1).

Theorem 6.7 turns out to be a special case of a general phenomenon, which is proven in a sequel to this paper [CFG⁺22]: if two permutations have the same recording tableau, then they have the same BBS steady-state time (Conjecture 1.2).

6.1. A recording tableau giving $n-3$ steady-state time

Definition 6.1. If $n \geq 5$, let \widehat{Q} denote the tableau

$$\begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline n & \\ \hline \end{array} \dots \begin{array}{|c|c|} \hline n-2 & n-1 \\ \hline \end{array}$$

Let $S_n(\widehat{Q})$ be the set of permutations $w \in S_n$ such that its recording tableau $Q(w)$ is equal to \widehat{Q} .

Example 6.2. For $n = 5$, the five permutations of $S_n(\widehat{Q})$ are the following.

45132 25143 35142 45231 35241

For $n = 6$, the sixteen permutations of $S_n(\widehat{Q})$ are as follows.

451362 251463 351462 452361 352461 561243 261354 361254
461253 561342 261453 361452 461352 562341 362451 462351

Note that one of our running examples, 452361, is in $S_6(\widehat{Q})$. As illustrated in Figure 1.2, its steady-state time is $3 = 6 - 3$.

Remark 6.3. It follows from Definition 6.1 that the RS algorithm induces a bijection from $S_n(\widehat{Q})$ onto the set of standard tableaux of shape $(n-3, 2, 1)$, so $S_n(\widehat{Q})$ is counted by the sequence [OEI, A077415].

The rest of this section is devoted to proving Theorem 6.7, which states that every permutation in $S_n(\widehat{Q})$ has steady-state time $n - 3$.

6.2. Lemmas for Theorem 6.7

Lemma 6.4. *Let $n \geq 5$, and suppose $w \in S_n(\widehat{Q})$. Then w is not the union of two increasing subsequences.*

Proof. The recording tableau of w is equal to \widehat{Q} , which has height 3. Therefore, the RS partition of w has three parts. By Greene's theorem (Theorem 2.2), w is not the union of two increasing subsequences. \square

Lemma 6.5. *Let $n \geq 5$, and suppose $w = w_1 w_2 \dots w_n \in S_n(\widehat{Q})$. Then w satisfies the following.*

1. $w_3 < w_4 < \dots < w_{n-1}$
2. $w_n < w_2$
3. $w_1 < w_2$
4. $w_3 < w_1$
5. $w_3 < w_2$
6. $w_4 < w_2$

Proof. Since $w \in S_n(\widehat{Q})$, the recording tableau of w is equal to \widehat{Q} . We will use the inverse RS algorithm¹ to construct w . Let $P = P(w)$ and $Q = Q(w)$. Denote the entries in the top row

¹For definition of the inverse RS algorithm, see, for example, the textbook [Sag01, Section 3.1].

of P by a_1, a_2, \dots, a_q (where $q = n - 3$), the second row of P by b_1 and b_2 , and the entry in the third row of P by c_1 . Hence, the starting pair P and Q is

$$P = \begin{array}{cccc|c|c} a_1 & a_2 & a_3 & a_4 & \dots & a_{q-1} & a_q \\ b_1 & b_2 & & & & & \\ c_1 & & & & & & \end{array} \qquad Q = \begin{array}{cccc|c|c} 1 & 2 & 5 & 6 & \dots & n-2 & n-1 \\ 3 & 4 & & & & & \\ n & & & & & & \end{array}$$

Since P is standard, we know that $b_1 < c_1$. The other entry b_2 in the second row is larger than b_1 . If $b_2 < c_1$, let b_y equal b_2 . Otherwise, let b_y be b_1 . In other words, we let b_y denote the largest element in the second row which is smaller than c_1 . Similarly, let a_x denote the largest element in the first row which is smaller than b_y . The first step of the inverse RS algorithm tells us that $w_n = a_x$.

After the first step in the inverse RS algorithm, we get the pair of tableaux

$$P_{n-1} = \begin{array}{cccc|c|c} \alpha_1 & \alpha_2 & \alpha_3 & \alpha_4 & \dots & \alpha_{q-1} & \alpha_q \\ \beta_1 & \beta_2 & & & & & \end{array} \qquad Q_{n-1} = \begin{array}{cccc|c|c} 1 & 2 & 5 & 6 & \dots & n-2 & n-1 \\ 3 & 4 & & & & & \end{array}$$

We now pause to observe two facts that will be referenced at the end of this proof. First, note that P_{n-1} is standard by definition of the inverse RS algorithm. Thus,

$$\alpha_1, \alpha_2, \dots, \alpha_q \text{ is increasing.} \tag{6.1}$$

Second, we note that

$$a_x < \beta_2, \tag{6.2}$$

as we now explain. Recall that $w_n = a_x$, so, using the original RS algorithm, we insert a_x into P_{n-1} to get P . Since row 1 of P_{n-1} and row 1 of P have the same size, we know that a_x bumps a number in row 1 of P_{n-1} to row 2. Let

a_i denote the smallest entry in row 1 of P_{n-1} which is greater than a_x .

The RS algorithm replaces a_i with a_x and bumps a_i to row 2. Since row 2 of P_{n-1} and row 2 of P have the same size, we know that a_i bumps a number in row 2 of P_{n-1} . So a_i must be smaller than β_2 . Since $a_x < a_i$, we have $a_x < \beta_2$. This concludes our explanation for (6.2).

We also note that

$$\beta_1 < \beta_2, \tag{6.3}$$

$$\alpha_1 < \beta_1, \text{ and} \tag{6.4}$$

$$\alpha_2 < \beta_2, \tag{6.5}$$

since P_{n-1} is standard. We will reference these inequalities at the end of this proof.

If $n > 5$, the numbers $n - 1, n - 2, \dots, 6, 5$ are in the first row of Q , so the next steps in the inverse RS algorithm are to remove elements $\alpha_q, \alpha_{q-1}, \dots, \alpha_4, \alpha_3$ from P_{n-1} , in that order. Hence, the last $n - 4$ letters of w are $\alpha_3, \alpha_4, \dots, \alpha_{q-1}, \alpha_q, a_x$.

The new pair of tableaux is

$$P_4 = \begin{array}{|c|c|} \hline \alpha_1 & \alpha_2 \\ \hline \beta_1 & \beta_2 \\ \hline \end{array} \quad Q_4 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & 4 \\ \hline \end{array}.$$

Note that 4 is the bottom right corner of Q_4 . Since $\alpha_2 < \beta_2$ by (6.5), we know that α_2 is the largest element in row 1 of P which is smaller than β_2 . So $w_4 = \alpha_2$, and the last $n - 3$ letters of w are $\alpha_2, \alpha_3, \alpha_4, \dots, \alpha_q, \alpha_{q-1}, a_x$.

The new pair of tableaux is

$$P_3 = \begin{array}{|c|c|} \hline \alpha_1 & \beta_2 \\ \hline \beta_1 & \\ \hline \end{array} \quad Q_3 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline 3 & \\ \hline \end{array}.$$

Note that 3 is in the second row of Q_3 . We know from (6.3) that β_2 is larger than β_1 , so α_1 is the largest element in row 1 smaller than β_1 . Thus, $w_3 = \alpha_1$. So the last $n - 2$ letters of w are $\alpha_1, \alpha_2, \alpha_3, \alpha_4, \dots, \alpha_q, \alpha_{q-1}, a_x$. The new pair of tableaux is

$$P_2 = \begin{array}{|c|c|} \hline \beta_1 & \beta_2 \\ \hline \end{array} \quad Q_2 = \begin{array}{|c|c|} \hline 1 & 2 \\ \hline \end{array}.$$

We then remove β_2 and β_1 from P_2 , in that order.

Therefore,

$$w = \underbrace{\beta_1 \beta_2}_{\text{increasing}} \underbrace{\alpha_1 \alpha_2 \alpha_3 \alpha_4 \dots \alpha_{q-1} \alpha_q}_{\text{increasing}} a_x.$$

We now have all the necessary information to prove all parts of the lemma.

1. The subsequence w_3, w_4, \dots, w_{n-1} is increasing because it is equal to the sequence $\alpha_1, \alpha_2, \dots, \alpha_q$, which is increasing due to (6.1). This proves part (1).
2. We have $w_n < w_2$ from (6.2), since $w_n = a_x$ and $w_2 = \beta_2$. This proves part (2).
3. We have $w_1 < w_2$ from (6.3), since $w_1 = \beta_1$ and $w_2 = \beta_2$. This proves part (3).
4. We have $w_3 < w_1$ from (6.4), since $w_3 = \alpha_1$ and $w_1 = \beta_1$. This proves part (4).
5. We have $w_3 < w_2$ since $w_1 < w_2$ and $w_3 < w_1$. This proves part (5).
6. We have $w_4 < w_2$ from (6.5), since $w_4 = \alpha_2$ and $w_2 = \beta_2$. This proves part (6). □

Lemma 6.6. Suppose $w = w_1 \dots w_n \in S_n(\widehat{Q})$.

1. Either $w_n = 1$ or $w_3 = 1$.
2. If $w_3 = 1$, then $w_1 = 2, w_4 = 2$, or $w_n = 2$.
3. If $w_3 = 1$ and $w_1 = 2$, then $w_4 = 3$ or $w_n = 3$.

Proof. 1. Suppose $w_n \neq 1$. Since both w_1, w_2 and w_3, \dots, w_{n-1} are increasing subsequences by Lemma 6.5(3),(1), either $w_1 = 1$ or $w_3 = 1$. Since $w_3 < w_1$ by Lemma 6.5(4), we must have $w_3 = 1$.

2. Assume $w_3 = 1$. We will show that $w_2 \neq 2$ and that none of w_5, \dots, w_{n-1} is equal to 2 (hence $w_1 = 2, w_4 = 2$, or $w_n = 2$). Since $w_3 < w_2$ and $w_4 < w_2$ by Lemma 6.5(5),(6) and since w is a permutation, we must have $2 < w_2$. Similarly, since $w_3 < w_4 < w_5 < \dots < w_{n-1}$ by Lemma 6.5(1) and since w is a permutation, each of w_5, \dots, w_{n-1} must be larger than 2.

3. Suppose $w_3 = 1$ and $w_1 = 2$. We will prove that $w_2 \neq 3$ and none of w_5, \dots, w_{n-1} is equal to 3 (hence $w_4 = 3$ or $w_n = 3$). Since $w_n \notin \{1, 2\}$, we have $2 < w_n$. By Lemma 6.5(2), we have $w_n < w_2$. So $2 < w_n < w_2$, which implies that w_2 is larger than 3 (since w is a permutation). Similarly, since $w_3 < w_4$ by Lemma 6.5(1) and $w_1 = 2$, we must have $2 < w_4 < w_5 < \dots < w_{n-1}$ by Lemma 6.5(1). So each of w_5, \dots, w_{n-1} is larger than 3 (since w is a permutation). \square

6.3. Proof of Theorem 6.7

Theorem 6.7. *If $n \geq 5$, every permutation in $S_n(\widehat{Q})$ has steady-state time $n - 3$.*

Proof. Suppose $w = w_1 \dots w_n \in S_n(\widehat{Q})$ is the box-ball configuration at time 0. We will show that w first reaches steady state at time $t = n - 3$.

Let j be the smallest number in $\{3, 4, \dots, n - 1\}$ such that $w_n < w_j$. We claim that the box-ball configuration at time $t = 1$ is

$$\begin{array}{ccccccc}
 & & & & \overset{n-5}{\text{copies}} & & \\
 & & & & \underbrace{\hspace{2cm}} & & \\
 e e & \underbrace{w_1 w_2}_{\text{increasing block}} & e e e \dots e & x & \underbrace{1 y_1 y_2 \dots y_{n-4}}_{\text{increasing block}} & & \\
 & & & & & &
 \end{array} \tag{6.6}$$

where $x = w_j$, there are $(n - 5)$ copies of e between w_2 and x , and $y_1 < y_2 < \dots < y_{n-4}$.

To prove this claim, consider the following cases. Due to Lemma 6.6, these five cases cover all possibilities.

1. $w_n = 1$
2. $w_3 = 1$ and $w_n = 2$
3. $w_3 = 1, w_1 = 2$, and $w_n = 3$
4. $w_3 = 1, w_1 = 2$, and $w_4 = 3$
5. $w_3 = 1$ and $w_4 = 2$

First, suppose $w_n = 1$. Lemma 6.5 tells us that w_3 is smaller than each w_i except for $w_n = 1$, so we must have $w_3 = 2$ and $j = 3$:

$$w_1 w_2 \underbrace{w_3}_2 w_4 w_5 \dots w_{n-1} \underbrace{w_n}_1 .$$

Since $w_1 < w_2$ and $w_4 < w_5 < \dots < w_{n-1}$ and since $w_4 < w_2$, applying one box-ball move to w results in the configuration

$$e e w_1 w_2 \overbrace{e e e \dots e}^{n-5} \underbrace{w_3}_x 1 w_4 w_5 \dots w_{n-1}$$

where there are $(n - 5)$ copies of e between w_2 and $x = w_3 = 2$.

Second, suppose $w_3 = 1$ and $w_n = 2$:

$$w_1 w_2 \underbrace{w_3}_1 w_4 w_5 \dots w_{n-1} \underbrace{w_n}_2 .$$

Since $w_1 < w_2$ and $w_4 < w_5 < \dots < w_{n-1}$ and since $w_4 < w_2$, applying one box-ball move to w results in the configuration

$$e e w_1 w_2 \overbrace{e e e \dots e}^{n-5 \text{ copies}} \underbrace{w_4}_x 1 2 w_5 w_6 \dots w_{n-1}$$

where there are $(n - 5)$ copies of e between w_2 and $x = w_4$. In this case, $w_3 = 1$ is not bigger than $w_n = 2$, but w_4 must be bigger than $w_n = 2$ since $w_4 \notin \{1, 2\}$, so $j = 4$.

Third, suppose $w_3 = 1$ and $w_1 = 2$ and $w_n = 3$. Lemma 6.5 tells us that w_4 is smaller than each of the w_i (except for $w_3 = 1$, $w_1 = 2$, and $w_n = 3$), so w_4 must be 4:

$$\underbrace{w_1}_2 w_2 \underbrace{w_3}_1 \underbrace{w_4}_4 w_5 \dots w_{n-1} \underbrace{w_n}_3 .$$

Using the same reasoning as in the previous two cases, applying one box-ball move to w results in the configuration

$$e e \underbrace{w_1}_2 w_2 \overbrace{e e e \dots e}^{n-5 \text{ copies}} \underbrace{w_4}_x 1 w_n w_5 w_6 \dots w_{n-1}$$

where there are $(n - 5)$ copies of e between w_2 and $x = w_4$. In this case, $j = 4$ since $w_3 = 1$ is not larger than $w_n = 3$ but $w_4 = 4$ is.

Finally, suppose we have one of the last two cases, so $w_3 = 1$ and $w_4 < w_n$:

$$w_1 w_2 \underbrace{w_3}_1 w_4 w_5 \dots w_{n-1} \underbrace{w_n}_{\text{larger than } w_4}$$

Since $w_1 < w_2$ and $w_4 < w_5 < \dots < w_{j-1} < w_n < w_j < \dots < w_{n-1}$ and since $w_4 < w_2$, applying one box-ball move to w results in the configuration

$$e e w_1 w_2 \overbrace{e e e \dots e}^{n-5 \text{ copies}} \underbrace{w_j}_x 1 w_4 w_5 \dots w_{j-1} w_n w_{j+1} \dots w_{n-1}$$

where there are $(n - 5)$ e 's between w_2 and $x = w_j$. In this case, $j \geq 5$ since w_4 is smaller than w_n . This concludes the proof of our claim that the box-ball configuration at time $t = 1$ is as given in (6.6).

Now we perform another box-ball move to reach the configuration at $t = 2$. If $n > 5$, in the configuration at $t = 2$, there are $(n - 6)$ e 's between w_2 and x :

$$e e e e w_1 w_2 \overbrace{e e \dots e}^{n-6 \text{ copies}} x \overbrace{e e \dots e}^{n-4 \text{ copies}} \underbrace{1 y_1 y_2 \dots y_{n-4}}_{\text{increasing block}}.$$

In fact, at each BBS move, the increasing sequence w_1, w_2 moves together two spaces to the right, the singleton x moves one space to the right, and the increasing sequence $1, y_1, y_2, \dots, y_{n-4}$ moves $n - 3$ spaces to the right. So the number of e 's between w_2 and x decreases by 1 after each BBS move. The configuration at $t = n - 4$ is

$$\dots e e e w_1 w_2 x e e e \dots e e e \underbrace{1 y_1 y_2 \dots y_{n-4}}_{\text{increasing block}}.$$

We claim that

$$x < w_2,$$

which we now prove. Recall that $x = w_j$, where j is the smallest number in $\{3, 4, \dots, n - 1\}$ such that $w_n < w_j$. If $w_2 < w_j$, then $w_1 < w_2 < w_j < w_{j+1} < \dots < w_{n-1}$ and the remaining w_i 's form two increasing subsequences of w whose union is w . This contradicts Lemma 6.4, so indeed $x < w_2$.

Since $x < w_2$, we have either $x < w_1 < w_2$ or $w_1 < x < w_2$. If $x < w_1 < w_2$, then the configuration at $t = n - 3$ is

$$\dots e e w_1 \underbrace{x w_2}_{\text{increasing block}} e e \dots e e \underbrace{1 y_1 y_2 \dots y_{n-4}}_{\text{increasing block}}.$$

If $w_1 < x < w_2$, then the configuration at $t = n - 3$ is

$$\dots e e w_2 \underbrace{w_1 x}_{\text{increasing block}} e e \dots e e \underbrace{1 y_1 y_2 \dots y_{n-4}}_{\text{increasing block}}.$$

Either way, the configuration array at $t = n - 3$ is a standard skew tableau whose rows have length $n - 3, 2$, and 1 . By Proposition 5.2, the configuration at $t = n - 3$ is in steady state.

The configuration at $t = n - 4$ is not yet in steady-state, as the relative positions of w_1, w_2 , and x in the configuration at $t = n - 4$ differ from the configuration at $t = n - 3$. Therefore, $t = n - 3$ is the minimum steady-state time of w . □

7. Knuth moves

We study how types of Knuth moves (Definition 1.4) play a role in a box-ball system. In Section 7.1, we prove that a non- K_B Knuth move preserves the shape of a soliton decomposition and that a K_B move changes it (Theorem 7.1). In Section 7.2, we prove that every permutation which is one non- K_B Knuth move from a row reading word has steady-state time 1 (Theorem 7.4).

7.1. Soliton decompositions are preserved by certain Knuth moves

Using the localized version of Greene's Theorem given in Section 2.2, we prove a partial characterization of the shape of SD in terms of types of Knuth moves.

Theorem 7.1. *Suppose v and w are two permutations in the same Knuth equivalence class.*

1. *If v and w are related by a sequence of Knuth moves containing an odd number of K_B moves, then $\text{SD}(v) \neq \text{SD}(w)$.*
2. *If v and w are related by a sequence of non- K_B Knuth moves, then $\text{sh SD}(v) = \text{sh SD}(w)$.*

Proof. To prove part (1), we observe that a K_B^+ move decreases the number of descents by 1, and a K_B^- move increases the number of descents by 1. Since the height the partition $\text{sh SD}(w)$ is equal to

$$D_1(w) = 1 + |\{\text{descents of } w\}|$$

by Lemma 2.6, it follows that applying an odd number of K_B moves to w changes $\text{sh SD}(w)$.

To prove part (2), suppose $x, y \in S_n$ are related by a proper K_1 or proper K_2 move. Due to Lemma 2.6, it suffices to prove that $D_k(x) = D_k(y)$ for all k . This breaks down into two main cases: case (i), where $y = K_1^+(x)$, and case (ii), where $y = K_2^+(x)$. These further divide into the following subcases, where $a < b < c$ in all cases:

- | | | | | |
|-----|-----|------------------------------------|----|--|
| i. | (a) | $y = \dots bca$
$x = \dots bac$ | or | $y = \dots bcad \dots$ with $c < d$
$x = \dots bacd \dots$ |
| | (b) | $y = \dots bca$
$x = \dots bac$ | or | $y = \dots bca a'$ with $a' < a$
$x = \dots bac a'$ |
| ii. | (a) | $y = cab \dots$
$x = acb \dots$ | or | $y = \dots dcab \dots$ with $c < d$
$x = \dots dacb \dots$ |
| | (b) | $y = cab \dots$
$x = acb \dots$ | or | $y = \dots a' cab \dots$ with $a' < a$
$x = \dots a' acb \dots$ |

The proofs are similar for each case. We include a partial proof of case (ia). Suppose

$$\begin{aligned} y &= \dots bca \\ x &= \dots bac \end{aligned}$$

or

$$\begin{aligned} y &= \dots bcad \dots \\ x &= \dots bacd \dots \end{aligned}$$

where $a < b < c < d$. The idea is to show that $D_k(y) \leq D_k(x)$ and $D_k(x) \leq D_k(y)$ for all k , from which the result follows.

Let $k \geq 1$. To show $D_k(y) \leq D_k(x)$, suppose that u_1, \dots, u_k are disjoint subsequences of y such that

$$D_k(y) = D(u_1) + \dots + D(u_k).$$

We will produce disjoint subsequences u'_1, \dots, u'_k of x where

$$D(u_1) + \dots + D(u_k) \leq D(u'_1) + \dots + D(u'_k).$$

First, suppose that c and a are in different subsequences. Then set $u'_i := u_i$ for each $1 \leq i \leq k$. Since $D(u_1) + \dots + D(u_k) = D(u'_1) + \dots + D(u'_k)$, we have $D_k(y) \leq D_k(x)$.

Next, suppose that b, c , and a are in the same subsequence u_j of y . Define u'_j to be the subsequence of x which is obtained from u_j by swapping c, a with a, c . Define $u'_i := u_i$ for all $i \neq j$. Then, since $a < b < c$, we have

$$D(u_j) = D(\dots, b, c, a, \dots) \leq D(\dots, b, a, c, \dots) = D(u'_j),$$

so $D_k(y) \leq D_k(x)$.

Lastly, suppose that c and a are in the same subsequence, say u_1 , and b is in a different subsequence, say u_2 . Write u_1 as a concatenation

$$u_1 = \underbrace{(\dots, c)}_{u_1^1} \sqcup \underbrace{(a, \dots)}_{u_1^2}$$

of two subsequences u_1^1 and u_1^2 , respectively. Write u_2 as a concatenation

$$u_2 = \underbrace{(\dots, b)}_{u_2^1} \sqcup \underbrace{(\dots)}_{u_2^2}$$

of two subsequences u_2^1 and u_2^2 , respectively. Define

$$\begin{aligned} u'_1 &:= u_2^1 \sqcup u_1^2 = (\dots, b) \sqcup (a, \dots), \\ u'_2 &:= u_1^1 \sqcup u_2^2 = (\dots, c) \sqcup (\dots), \end{aligned}$$

and $u'_i := u_i$ for all $i \notin \{1, 2\}$. Then, since $a < b < c$,

$$D(u_1) + D(u_2) \leq D(u'_1) + D(u'_2),$$

so $D_k(y) \leq D_k(x)$. The proof of the reverse inequality $D_k(x) \leq D_k(y)$ is similar. □

Theorem 7.1 allow us to use Knuth moves to find a subset of permutations whose soliton decomposition and RS insertion tableau coincide.

Corollary 7.2 (Corollary of Theorem 4.2 and Theorem 7.1). *Let $w \in S_n$, let $T = P(w)$, and let r be the row reading word of T .*

1. *If w is related to r by a sequence of Knuth moves containing an odd number of K_B moves, then $SD(w) \neq P(w) = T$.*
2. *If w is related to r by a sequence of non- K_B moves, then $SD(w) = P(w) = T$.*

Example 7.3. The permutations 362514 and 632514 are the reading words of the tableaux

$$\begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & 6 \\ \hline \end{array}
 \quad \text{and} \quad
 \begin{array}{|c|c|} \hline 1 & 4 \\ \hline 2 & 5 \\ \hline 3 & \\ \hline 6 & \\ \hline \end{array}
 \quad \text{respectively.}$$

Figure 7.1 (respectively, Figure 7.2) shows all permutations in the Knuth equivalence class of $r = 362514$ (respectively, $r = 632514$). The corresponding soliton decomposition is drawn next to each permutation. An edge with label K_1 (respectively, K_2) indicates that the Knuth move is a proper K_1 (respectively, K_2) move. An edge with label K_B indicates that the Knuth move is both K_1 and K_2 . The permutations are arranged such that they form a subdiagram of the Hasse diagram of the right weak order² on the symmetric group S_6 .

7.2. Permutations one proper Knuth move from a row reading word

Theorem 7.4. *Let r be the row reading word of a standard tableau. Suppose w is a permutation which is related to r by one proper K_1 move or one proper K_2 move. Then w has steady-state time 1.*

If w is one K_B move from the row reading word of a standard tableau, then the steady-state time of w may be 1 or greater than 1. See Example 7.5.

Example 7.5. The corresponding steady-state times are given next to each permutation in the two Knuth equivalence classes of Figures 7.1 and 7.2.

In Figure 7.1, the permutation 362154 is one K_B^- move from r , and its steady-state time is $t = 2$. Another permutation, 326514, is also one K_B^- move from r , and its steady-state time is $t = 1$.

In Figure 7.2, we can perform a K_B^- move and also a proper K_1^+ move on r (see Lemma 7.6). The permutation 635214 is one proper K_1^+ move from r , and its steady-state time is $t = 1$, illustrating Lemma 7.9.

²For definition of the right weak order, see, for example, the textbook [BB05, Section 3.1].




Figure 7.1: The Knuth equivalence class of $r = 362514$, with their soliton decompositions and steady-state times.




Figure 7.2: The Knuth equivalence class of $r = 632514$, with their soliton decompositions and steady-state times.

7.2.1 Proof of Theorem 7.4

Theorem 7.4 follows from the following four lemmas.

Lemma 7.6. *Let $r = r_1 r_2 \dots r_n$ be the row reading word of a standard tableau P .*

1. *If one performs a K_1^- move on r , the move is K_B .*
2. *Suppose we are able to perform a K_1^+ move $yxz \mapsto yzx$ (where $x < y < z$) on r . If $r_1 \neq y$, we must have*

$$r = r_1 \dots \underbrace{r_\ell y x z}_{\text{decreasing}} \dots r_{n-1} r_n \quad (7.1)$$

where $r_1 > r_2 > \dots > r_\ell > y > x$. The tableau P must be of the form given in Figure 7.3, where the entry y is in its own row, and the row immediately above y starts with entries x, z .

3. *If one performs a K_1^+ move on r , the move is not K_B .*

$$\begin{array}{c}
 \vdots \dots \\
 \begin{array}{|c|c|c|c|}
 \hline
 a_1 & a_2 & a_3 & \dots \\
 \hline
 x & z & b_1 & b_2 \dots \\
 \hline
 y & & & \text{(possibly with no } b_i\text{'s)} \\
 \hline
 r_\ell & & & \\
 \hline
 \vdots & & & \\
 \hline
 r_1 & & & \\
 \hline
 \end{array}
 \end{array}$$

Figure 7.3: General form of a standard tableau P whose row reading word can undergo a K_1^+ move.

Proof. First, we prove part (1) of the lemma. Suppose we perform a K_1^- move $yzx \mapsto yxz$ (where $x < y < z$) on r . Since r is the row reading word of P , the tableau P must contain a subtableau

$$\begin{array}{|c|c|}
 \hline
 x & b \\
 \hline
 y & z \\
 \hline
 \end{array}
 \quad \text{or} \quad
 \begin{array}{|c|}
 \hline
 x \\
 \hline
 \end{array}
 \dots
 \begin{array}{|c|c|}
 \hline
 \dots & b \\
 \hline
 y & z \\
 \hline
 \end{array}.$$


Since the rows and columns of P are increasing, we must have $x < b < z$. Thus, r must contain a consecutive subsequence $yzxb'$ where $x < b' \leq b < z$, so the K_1^- move $yzx \mapsto yxz$ is K_B^- .

Now suppose we perform a K_1^+ move $yxz \mapsto yzx$ on r . First, we prove part (2). Since $x < y < z$ and P is standard, the entry y must be the only element in its row in P , that is, the rows of P containing x, y, z are of the form

$$\begin{array}{|c|c|}
 \hline
 x & z \\
 \hline
 y & \\
 \hline
 \end{array}
 \dots$$

If $r_1 = y$, then we are done. Suppose $r_1 \neq y$, and write $r = r_1 r_2 \dots r_\ell y x z \dots r_n$. Since the rows of P are weakly decreasing in length, the rows of P below y are of size 1. Since P is standard, we have $r_1 > r_2 > \dots > r_\ell > y$. So r is of the form given in (7.1) and P is of the form given in Figure 7.3.

Finally, to prove part (3) of the lemma, we prove that this K_1^+ move is not a K_B move. If $r_n = z$, then we know this K_1^+ move is not K_B . Suppose $r_n \neq z$, so $r = r_1 \dots y x z b \dots r_n$ for some b . Since r is the row reading word of P , either the entry b is immediately above x in P or the entry b is immediately to the right of z in P :



Since P is standard, either $b < x$ or $z < b$. Either way, this K_1^+ move is not K_B . □

Lemma 7.7. *Let $r = r_1 r_2 \dots r_n$ be the row reading word of a standard tableau P .*

1. *It is impossible to perform a K_2^+ move on r .*
2. *Suppose we are able to perform a K_2^- move $zxy \mapsto xzy$ (where $x < y < z$) which is not a K_B move on r . If $r_1 \neq z$, we have*

$$r = r_1 \dots \underbrace{r_\ell z x}_{\text{decreasing}} y \dots r_{n-1} r_n \tag{7.2}$$

where $r_1 > r_2 > \dots > r_\ell > z$. The tableau P must be of the form given in Figure 7.4, where the entry z is in its own row, and the row immediately above z starts with entries x, y .




Figure 7.4: General form of a standard tableau P whose row reading word can undergo a K_2^- move which is not K_B .

Proof. First, we prove part (1) of the lemma. Assume (for the sake of contradiction) that one could perform a K_2^+ move on r . Then r must contain a xzy pattern. Hence, since r is the row reading word of P , the tableau P must contain the following subtableau:

$$\begin{array}{c} \boxed{y} \cdots \\ \boxed{x} \quad \boxed{z} \end{array}$$

Notice that y is north or northwest of x but $x < y$. This is a contradiction to the fact that P is a standard tableau. Therefore, we cannot perform a K_2^+ move on r .

Next, we prove part (2) of the lemma. Suppose we perform a K_2^- move $zxy \mapsto xzy$ on r which is not a K_B move. If $r_1 = z$, then the last two rows of P are of the form

$$\begin{array}{cc} \boxed{x} & \boxed{y} & \cdots \\ \boxed{z} & & \end{array},$$

so P is of the form given in Figure 7.4.

Suppose $r_1 \neq z$, and write $r = r_1 \dots r_\ell z x y \dots r_{n-1} r_n$. Since our K_2^- move is not K_B , we must have either $r_\ell < x$ or $z < r_\ell$. Since P is standard and x is in the first column, we cannot have $r_\ell < x$. So $z < r_\ell$. Therefore z is in its own row in P . Since the rows of P are weakly decreasing in length, the rows of P below z are of size 1. Since P is standard, we have $r_1 > r_2 > \dots > r_\ell$. So r is of the form given in (7.2) and P is of the form given in Figure 7.4. \square

Remark 7.8. In general, a K_2^- move on the row reading word of a standard tableau may (or may not) be K_B .

The proofs of the next two lemmas, Lemmas 7.9 and 7.10, are similar.

Lemma 7.9. *Suppose $r = r_1 r_2 \dots r_n \in S_n$ is the row reading word of a standard tableau P . Let w be a permutation which differs from r by one proper K_1 move. Then w first reaches its steady state at $t = 1$.*

Proof. By Lemma 7.6, applying a K_1 move that is not K_B to r must be a K_1^+ move $yxz \mapsto yzx$ such that

$$\begin{aligned} r &= \underbrace{r_1 r_2 \dots r_\ell y x z \dots r_{n-1} r_n}_{\text{decreasing}} \\ w = K_1^+(r) &= \underbrace{r_1 r_2 \dots r_\ell y z x \dots r_{n-1} r_n}_{\text{decreasing}} \end{aligned}$$

where $r_1 > r_2 > \dots > r_\ell > y$ (if $r_1 \neq y$) and $x < y < z$.

We apply the carrier algorithm to w . First, we insert $r_1, r_2, \dots, r_\ell, y$ into the carrier. Since these are decreasing, we eject $e, r_1, r_2, \dots, r_\ell$ from the carrier in consecutive order:

$$\begin{array}{c}
 \underbrace{ee \cdots e}_{\text{carrier}} \underbrace{r_1 r_2 r_3 \cdots r_{\ell-1} r_\ell y}_{\text{decreasing}} z x \cdots r_n \\
 e \underbrace{r_1 ee \cdots e}_{\text{decreasing}} r_2 r_3 \cdots r_{\ell-1} r_\ell y z x \cdots r_n \\
 e r_1 \underbrace{r_2 ee \cdots e}_{\text{decreasing}} r_3 \cdots r_{\ell-1} r_\ell y z x \cdots r_n \\
 \vdots \\
 e r_1 r_2 \cdots r_\ell \underbrace{y ee \cdots e}_{\text{decreasing}} z x \cdots r_n
 \end{array}$$

Next, we insert z into the carrier. Since the only non- e entry in the carrier, y , is smaller than z , we eject an e :

$$e e r_1 r_2 \cdots r_\ell \underbrace{y z ee \cdots e}_{\text{decreasing}} x r_{\ell+4} \cdots r_n$$

Next, we insert x into the carrier. Since $x < y < z$, we eject y and get

$$e e r_1 r_2 \cdots r_\ell e y \underbrace{x z ee \cdots e}_{\text{decreasing}} r_{\ell+4} \cdots r_n$$

Note that the string

$$x z r_{\ell+4} \cdots r_{n-1} r_n$$

is equal to the consecutive subsequence $r_{\ell+2} \cdots r_{n-1} r_n$ of r . This string is the row reading word of the subtableau (possibly with no b_i 's)

$$\begin{array}{c}
 \vdots \quad \dots \\
 \begin{array}{|c|c|c|c|}
 \hline
 a_1 & a_2 & a_3 & \cdots \\
 \hline
 x & z & b_1 & b_2 \cdots \\
 \hline
 \end{array}
 \end{array}$$

of P , where P is given in Figure 7.3. Since this subtableau has the shape of a partition and has increasing rows and columns, completing the carrier algorithm yields the configuration at time $t = 1$:

$$e e r_1 r_2 \cdots r_\ell e y \overbrace{e e \cdots e}^{\text{0 or more copies}} x z b_1 b_2 b_3 \cdots a_1 a_2 \cdots r_{n-1} r_n \underbrace{e e \cdots e}_{\text{decreasing}}.$$

The configuration array at $t = 1$ is the skew tableau created by taking P and shifting some of the rows to the right. Since P is standard tableau with partition shape to begin with, the configuration array is a standard skew tableau with weakly increasing rows. By Proposition 5.2, the configuration at $t = 1$ is in steady state. \square

Lemma 7.10. *Suppose $r = r_1 r_2 \dots r_n \in S_n$ is the row reading word of a standard tableau P . Let w be a permutation which differs from r by one proper K_2 move. Then w first reaches its steady state at $t = 1$.*

Proof. By Lemma 7.7, applying a K_2 move that is not K_B to r must be a K_2^- move $zxy \mapsto xzy$ to r such that

$$\begin{aligned} r &= r_1 \dots r_\ell z x y \dots r_{n-1} r_n \\ w &= K_2^-(r) = r_1 \dots r_\ell x z y \dots r_{n-1} r_n \end{aligned}$$

where $r_1 > r_2 > \dots > r_\ell > z$ (if $r_1 \neq z$) and $x < y < z$.

As in the proof of Lemma 7.9, we apply the carrier algorithm to w . We insert the decreasing sequence $r_1, r_2, \dots, r_\ell, x$ into the carrier and eject $e, r_1, r_2, \dots, r_\ell$, in that order. As we insert z and y , we eject e and z , in that order:

$$\begin{array}{c} \underbrace{e \dots e}_{\text{carrier}} \underbrace{r_1 r_2 r_3 \dots r_\ell x}_{\text{decreasing}} z y \dots r_{n-1} r_n \\ e \underbrace{r_1 e e \dots e}_{\text{decreasing}} \underbrace{r_2 r_3 \dots r_\ell x}_{\text{decreasing}} z y \dots r_{n-1} r_n \\ e r_1 \underbrace{r_2 e e \dots e}_{\text{decreasing}} \underbrace{r_3 \dots r_\ell x}_{\text{decreasing}} z y \dots r_{n-1} r_n \\ \vdots \\ e r_1 r_2 \dots r_\ell \underbrace{x e \dots e}_{\text{decreasing}} z y \dots r_{n-1} r_n \\ e r_1 r_2 \dots r_\ell e \underbrace{x z e \dots e}_{\text{decreasing}} y r_{\ell+4} \dots r_n \\ e r_1 r_2 \dots r_\ell e z \underbrace{x y e \dots e}_{\text{decreasing}} r_{\ell+4} \dots r_n \end{array}$$

Note that the string

$$x y r_{\ell+4} \dots r_{n-1} r_n$$

is equal to the consecutive subsequence $r_{\ell+2} \dots r_{n-1} r_n$ of r . This string is the row reading word of the subtableau (possibly with no b_i 's)

$$\begin{array}{c} \vdots \dots \\ \begin{array}{|c|c|c|c|} \hline a_1 & a_2 & a_3 & \dots \\ \hline x & y & b_1 & b_2 \dots \\ \hline \end{array} \end{array}$$

of P , where P is given in Figure 7.4. Since this subtableau has the shape of a partition and has increasing rows and columns, completing the carrier algorithm yields the configuration at time $t = 1$:

$$e e r_1 r_2 \dots r_\ell e z \overbrace{e \dots e}^{\text{0 or more copies}} x y b_1 b_2 b_3 \dots a_1 a_2 \dots r_{n-1} r_n \underbrace{e e \dots e}_{\text{decreasing}}.$$

The configuration array at $t = 1$ is the skew tableau created by taking P and shifting some of the rows to the right. Since P is standard tableau with partition shape to begin with, the configuration array is a standard skew tableau with weakly increasing rows. By Proposition 5.2, the configuration at $t = 1$ is in steady state. \square

Acknowledgements

This research project started during the University of Connecticut 2020 Mathematics REU. Our project was inspired by a blog post [Lew19] for the University of Minnesota's Open Problems in Algebraic Combinatorics (OPAC) conference and conversations with Joel B. Lewis. We thank Ian Whitehead for serving as a faculty mentor to B. Drucker's research course in Fall 2020 and for helpful suggestions. We also thank Pavlo Pylyavskyy and Rei Inoue for useful comments and Marisa Cofie, Olivia Fugikawa, Madelyn Stewart, and David Zeng for many discussions during SUMRY 2021. Special thanks to Darij Grinberg for proving one of our conjectures and for helpful feedback. This work also benefited from computation using SAGEMATH [Dev20] and the High Performance Computing facility at the University of Connecticut. Finally, we thank the anonymous referees whose suggestions helped improve and clarify this paper.

References

- [BB05] Anders Björner and Francesco Brenti. *Combinatorics of Coxeter groups*, volume 231 of *Graduate Texts in Mathematics*. Springer, New York, 2005. doi:10.1007/3-540-27596-7.
- [CFG⁺22] Marisa Cofie, Olivia Fugikawa, Emily Gunawan, Madelyn Stewart, and David Zeng. Box-ball systems and RSK recording tableaux. 2022. arXiv:2209.09277v1.
- [Dev20] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 9.1)*, 2020. URL: <https://www.sagemath.org>.
- [Fuk04] Kaori Fukuda. Box-ball systems and Robinson-Schensted-Knuth correspondence. *J. Algebraic Combin.*, 19(1):67–89, 2004. doi:10.1023/B:JACO.0000022567.30060.3a.
- [Gre74] Curtis Greene. An extension of Schensted's theorem. *Advances in Math.*, 14:254–265, 1974. doi:10.1016/0001-8708(74)90031-0.
- [Knu70] Donald E. Knuth. Permutations, matrices, and generalized Young tableaux. *Pacific J. Math.*, 34:709–727, 1970. doi:10.2140/pjm.1970.34.709.
- [Lew19] Joel B. Lewis. A localized version of Greene's theorem. <https://realopacblog.wordpress.com/2019/11/24/a-localized-version-of-greenes-theorem>, 2019.
- [LLPS19] Joel Lewis, Hanbaek Lyu, Pavlo Pylyavskyy, and Arnab Sen. Scaling limit of soliton lengths in a multicolor box-ball system. 2019. arXiv:1911.04458v2.
- [OEI] OEIS Foundation Inc. (2022). The On-Line Encyclopedia of Integer Sequences. <http://oeis.org>.

- [Sag01] Bruce E. Sagan. *The symmetric group*, volume 203 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 2001. Representations, combinatorial algorithms, and symmetric functions. doi:10.1007/978-1-4757-6804-6.
- [Sag20] Bruce E. Sagan. *Combinatorics: the art of counting*, volume 210 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, [2020] ©2020. doi:10.1090/gsm/210.
- [Sch61] C. Schensted. Longest increasing and decreasing subsequences. *Canadian J. Math.*, 13:179–191, 1961. doi:10.4153/CJM-1961-015-3.
- [Tak93] Daisuke Takahashi. On some soliton systems defined by using boxes and balls. In *Proceedings of the international symposium on nonlinear theory and its applications (NOLTA'93)*, pages 555–558, 1993. URL: <https://hakotama.jp/laboratory/works/public/93t-nolta.pdf>.
- [TM97] Daisuke Takahashi and Junta Matsukidaira. Box and ball system with a carrier and ultradiscrete modified KdV equation. *J. Phys. A*, 30(21):L733–L739, 1997. doi:10.1088/0305-4470/30/21/005.
- [TS90] Daisuke Takahashi and Junkichi Satsuma. A soliton cellular automaton. *J. Phys. Soc. Japan*, 59(10):3514–3519, 1990. doi:10.1143/JPSJ.59.3514.