

UCLA

UCLA Previously Published Works

Title

The Role of Family in Diabetes Management for Mexican American Adults.

Permalink

<https://escholarship.org/uc/item/5761q5km>

Journal

Hispanic health care international : the official journal of the National Association of Hispanic Nurses, 22(2)

Authors

Jordan, Olivia

Benitez, Amanda

Burnet, Deborah

et al.

Publication Date

2024-06-01

DOI



10.1177/15404153231206086

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

The Role of Family in Diabetes Management for Mexican American Adults

Olivia J. Jordan, MD¹ , Amanda Benitez, MPH² ,
Deborah L. Burnet, MD, MA³, Michael T. Quinn, PhD³
and Arshiya A. Baig, MD, MPH³

Hispanic Health Care International
2024, Vol. 22(2) 109–118
© The Author(s) 2023



Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/15404153231206086
journals.sagepub.com/home/hci



Abstract

Introduction: The purpose of this study was to characterize how family influences diabetes self-management in Mexican American adults. **Methods:** Data were analyzed from previously collected data that included 34 semi-structured interviews with Hispanic adults with diabetes and six focus groups with 37 adults with diabetes and family members. Themes related to family and diabetes management were identified and analyzed using a modified template approach. **Results:** Family-related facilitators to T2DM self-management were (1) provides support, (2) provides motivation, and (3) desire to protect family from diabetes. Family-related challenges were (1) lack of support, (2) family responsibilities, and (3) stress related to family. Diabetes education was shared with family members. Family member perspectives on T2DM included (1) not knowing how to help, (2) effect on emotional wellbeing, (3) diabetes affects the whole family, and (4) family provides support. **Conclusion:** Most participants with T2DM felt supported by family, but many desired more social support and support surrounding dietary changes from family. Many felt family did not understand what living with diabetes meant for them. Most family members wished to learn more about how to help. Future interventions should include family members and teach them supportive strategies to support beneficial diabetes self-management behaviors.

Keywords

Mexican American, type 2 diabetes, family, diabetes self-management

Mexican Americans have the highest prevalence of type 2 diabetes (T2DM) within the U.S. Hispanic community with 14.4% affected, and are disproportionately affected by diabetes-related comorbidities (“Center for Disease Control and Prevention” [CDC], 2023; Haw et al., 2021; National Center for Health Statistics, 2022). Engaging in diabetes self-care behaviors, including exercise, monitoring blood glucose levels, and following a healthy diet, is vital for adequate diabetes management (“American Diabetes Association” [ADA], 2022). However, Hispanics encounter barriers to engage in adequate diabetes self-care behaviors which contribute to worse health outcomes (Aguayo-Mazzucato et al., 2019). Implemented diabetes self-education programs have been found to increase knowledge of diabetes self-care behaviors and improve glycosylated hemoglobin (A1C) (ADA, 2022).

Diabetes self-management education can be significantly impacted by family support (McEwen et al., 2019). Often, family members improve a patient’s well-being, ability to follow up with medical treatments, and adherence to positive lifestyle changes by way of their expressed thoughts and attitudes, similar to findings by Fort et al. (2020). However, at times, family’s nonsupportive behaviors can intensify the stressors that come with managing a disease (Mayberry et al., 2019), and can

be associated with being less adherent to one’s diabetes medication regimen and having poorer glucose control (Mayberry et al., 2014).

Culturally tailored interventions that incorporate family (Ingram et al., 2007) have led to statistically significant increases in diabetes self-care behaviors in Mexican Americans in the U.S. border region (Teufel-Shone et al., 2005). However, in the Mexican American population, few studies have qualitatively explored how family supports and hinders diabetes self-management, which is concerning given worse complications seen in this subgroup compared to other Hispanic American adults (CDC, 2023). Some recent studies have gathered perspectives on the relationship between Mexican Americans with T2DM and their children and in rural populations, but few

¹ Department of Medicine, Internal Medicine Resident, University of California, Los Angeles, Los Angeles, CA, USA

² Enlace Chicago, Chicago, IL, USA

³ University of Chicago, Section of General Internal Medicine, Chicago, IL, USA

Corresponding Author:

Olivia J. Jordan, University of California, Los Angeles, 757 Westwood Plaza, Suite 7501, Los Angeles, CA 90095, USA.
Email: olivajordan@gmail.com

have been done within the family unit or in other geographic areas with a high concentration of Mexican Americans (McEwen & Murdaugh, 2014; McEwen et al., 2017; Rascón et al., 2022).

Chicago, Illinois' Hispanic population is largely Mexican American ("2020 Census Hispanic", 2020), with this population experiencing higher rates of T2DM (16.7%) than Chicago's general Hispanic population (13.2%) ("Adult Diabetes Rate", 2021). Despite its reported benefits, the effects of family on diabetes self-management behaviors have not been qualitatively described in this urban setting that is home to a significant proportion of Mexican Americans. Through qualitative analysis, the purpose of this study was to identify how family influences diabetes self-care management in Mexican Americans, and how family without T2DM is impacted by the disease.

Methods

Study design

This qualitative study used data from two separate research studies and one additional dataset (Baig et al., 2012, 2014, 2015, 2016). The current study design uses the qualitative research method of modified template analysis to explore diabetes self-care management behaviors in the Mexican American community using previously transcribed data. Complete details regarding study design, data collection, and data analyses of the prior datasets can be found in Table 1.

In the first study, participants were recruited to elicit solutions of how to address T2DM disparities in the Hispanic community (Baig et al., 2012). This study was conducted in partnership with community members at two predominantly Mexican American churches and used community-based participatory research (CBPR) methods, a research methodology that focuses on collaborating with community members throughout the research process (Wells & Jones, 2009), to elicit ideas of effective diabetes interventions (group vs. one-on-one) that might happen in a church setting. The second study (Baig et al., 2015) was a randomized, controlled pilot study to improve diabetes self-management in the Hispanic population. The intervention arm included eight weekly 90-min group classes that were conducted

by lay members who encouraged using intrinsic motivators to elicit behavioral change; in the control arm, participants were invited to a 90-min lecture on diabetes self-management by a bilingual community health educator. In this study, all participants could opt to take pictures that represented or were related to their T2DM and management; semi-structured interviews were conducted to inform subsequent research in creating a culturally-tailored diabetes intervention (Baig et al., 2014, 2016).

The final dataset used randomly selected participants from the second study described. These data were used in a culturally-competent training for healthcare providers that included direct patient audio narratives describing the importance of chosen photographs (Baig et al., 2014, 2016). An objective of the interviews was to explore beliefs, behaviors, and challenges to diabetes self-management.

Setting

All study subjects were recruited in a predominantly low-income, Mexican American immigrant neighborhood in Chicago, Illinois in which many risk factors exist that lead to poor health and higher diabetes-related mortality (Community Snapshot: South Lawndale, 2016). The research team recruited participants through posted flyers at churches, verbal referral at health fairs and church events, and word-of-mouth in the community.

Data collection

The first study was conducted between February and April 2009. Six focus groups were conducted with 37 adults diagnosed with T2DM or who had a family member with T2DM. These groups were led by a trained bilingual community member. Interviewers followed a semi-structured questioning guide that was intended to elicit opinions on proposed interventions and to elicit perceived community needs and barriers to diabetes self-management (Appendix A). Five groups were conducted in English, and one in Spanish that was translated into English by a professional translation service. The sessions lasted approximately 90 min each and were audio recorded and transcribed verbatim using professional software.

Table 1. Description of Previously Collected Study Data.

	[Deidentified Study 1] (2012)	[Deidentified Study 2] (2015)	[Deidentified Dataset] (2014, 2016)
Study design	CBPR guided focus groups	Randomized, controlled, pilot study	Qualitative research study
Study setting	South Lawndale; a predominantly low-income, Mexican American immigrant neighborhood in Chicago, Illinois		
Study sample	37	27	14
Data collection	Six focus groups led by trained members; five in English, one in Spanish	Postintervention interviews using semistructured interview guide	Postintervention interviews using semistructured interview guide
Data analysis	Modified template approach, HyperRESEARCH	Modified template approach, HyperRESEARCH	Modified template approach, HyperRESEARCH

CBPR; community based participatory research.

For the randomized pilot study, 100 participants were recruited from May 2011 to June 2012. Of this study's 100 participants, 30 were contacted to complete post-study interviews between July 2012 and February 2013 to gather personal experiences and social support as it relates to diabetes self-management and study participation. Interviewers followed a question guide that included two family-focused questions that asked if the program helped to involve family in diabetes management and if they would recommend the program to a family member (Appendix B). These interviews were conducted by trained research assistants in a private room at a local health center, lasted 45–60 min each, and were led by a trained bilingual interviewer. Interviews were transcribed verbatim using professional software; three were not transcribed due to administrative error.

For the final dataset, an additional fourteen interviews were conducted with participants from the randomized study; seven of these participants had not previously been interviewed. Interviews were completed between June 2013 and August 2013, lasted approximately 30 min each, and were audio-recorded and transcribed in a similar fashion to the prior studies described. Trained research staff led participant interviews that discussed the rationale for taking the photographs, and worries, challenges, motivators, and successes that the photo represented. The potential role of family in diabetes management for each participant was discussed in the interview (Appendix C).

Data analysis

One researcher grouped representative quotations into themes using inductive and deductive analysis and developed an initial codebook. Data were then analyzed using a modified template approach and code reports were generated in HyperRESEARCH. This researcher further used hierarchical coding to identify sub-themes and representative quotes. A second researcher independently reviewed code hierarchies and reviewed subthemes in collaboration with the first researcher. The first researcher subsequently refined subthemes based on the discussion and coded representative quotations into a list that was finalized in discussion with the second researcher (Morse, 1995).

Ethical considerations

All studies from which the current study's data were derived were approved by the Institutional Review Board at The University of Chicago. All participants provided written informed consent.

Results

Data were collected from a total of 71 participants (37 from focus groups, 34 from one-on-one interviews). Participants had a mean (SD) age of 55 (11) years and 80% were female. Eighty-nine percent self-identified as Hispanic, and the majority

of participants reported being of Mexican descent (86%) and born in Mexico (72%). Fifty-eight participants (82%) reported a diagnosis of T2DM and 13 (18%) noted having an immediate family member with T2DM. Seventy-seven percent of participants were born outside of the United States. Spanish was the primary language spoken at home for 82% of participants.

Quotations were grouped into four categories. The first three categories pertained to participants with T2DM, and the fourth included perspectives from family members of those with the disease.

Family-Related Facilitators to Participants' Diabetes Self-Management (Table 2)

Provides support. Family as a source of support for diabetes management was expressed by 39 (55%) of 71 participants. Twenty-two participants mentioned that family provided instrumental support in providing healthy food, medication adherence, and transportation to doctor's appointments. Several mentioned families were accommodating to their needs, and a couple stated family was involved in day-to-day management of their diabetes. Some participants also discussed how family members provided informational support by giving advice, recommendations, and reminders to assist with diabetes management as well as emotional and companionship support. One noted:

"My daughter bought some little boxes like this ... she arranges everything for me so that ... I don't forget [to take my medication]."

Provides motivation. Family as a motivator for participating in diabetes self-care behaviors was expressed by 24 (34%) of 71 participants (Table 2). Being able to support family members in the future and enjoy life with grandchildren were frequently mentioned as motivation to improve self-care. Many of these participants said family members who had already suffered from diabetes complications provided motivation to maintain their own health. Many (20%) of the participants mentioned having brothers, sisters, parents, and grandparents with diabetes, whereas only one participant spoke of having no history of diabetes in her family. One participant shared:

"... If I didn't do what I was supposed to have done, I'd probably be in a hospital, somewhere laid up, sick ... No, I don't want that. I've seen my mother already there and um, that was too much for me."

Desire to protect family from diabetes. Twenty-one (30%) of all participants reported wanting to be more informed about diabetes in order to prevent diabetes in family members and protect them from its complications. One participant stated:

"[Diabetes] has affected me primarily in that I'm thinking about not wanting my children to inherit it from me ... I am now making sure that my kids exercise and eat well so that they won't end up like me."

Table 2. Family-Related Facilitators to Participant's Diabetes Self-Management.

Theme	Subtheme	Example Quote
Support	Family is a source of support	Yes, ... <i>family plays an important role. The family ... is a foundation as well.</i>
	Family is involved in day-to-day management of diabetes	<i>[My blood sugar is] almost always high. My daughter tells me, "You haven't checked yourself?" I tell her, "I forgot." I like to check it before I eat breakfast and she reminds me ...</i>
	Family is a source of emotional support	<i>Everybody worried about me. And they told me that, they saw that, when they saw me and said that I was going to attend a program, they said that "this was good, how did I feel?"</i>
	Family is a source of instrumental support	<i>[My youngest son] ... bought me a little scale to weigh the food. And the first days, he was cooking for me, he was weighing and boiling, and he froze [the food] for me. ... I did have a lot of support, a lot, a lot of support from my family. "My daughter bought some little boxes like this ... Yeah, she arranges everything for me so that ... I don't forget [to take my medication]."</i>
	Family is a source of informational support	<i>[My daughter's] always forcing me to eat healthier. "Don't have white sugar, have brown sugar. Eat more vegetables."</i>
	Family is a source of companionship support	<i>I go [exercise] with my husband, yeah, and it makes him happy.</i>
Motivation	Family is a source of motivation for diabetes management	<i>I was motivated to try and not be so ill so that I could help my parents. I was motivated by the child I take care of. [I attended classes] to inform myself about diabetes ... but more because I know that my granddaughters might have [diabetes]. And [I'm] making the changes for myself and also for them. I'm going to take care of myself because I want to see more generations of my family and that was my motivation to keep going. Because I was getting like depressed, and it was when I said to myself, hey – positive. I had to keep going and that was when I got out of depression, because I wanted to see more generations of my family.</i>
	Suffering or death related to diabetes in the family as motivation for self-care	<i>I was gonna die if I didn't [change]. If I, if I didn't do what I was supposed to have done, I'd probably be in a hospital, somewhere laid up, sick ... and ... No, I don't want that. I've seen my mother already there and um, that was too much for me.</i>
Desire to Protect Family from Diabetes	<i>[Diabetes] has affected me primarily in that I'm thinking about not wanting my children to inherit it from me ... I am now making sure that my kids exercise and eat well so that they won't end up like me.</i>	

Family-Related Challenges to Participants' Diabetes Self-Care (Table 3)

Lack of support. Lack of social support from family was expressed by 15 (21%) of 71 participants. Some participants mentioned they did not have family members around or who cared about their diabetes management with one stating:

"I am here by myself. In other words, I am here with my four kids, no one else. So, I don't have any relatives and my children, well they want to be taken care of, not the other way around."

Nine participants noted that family members did not understand what is it like to live with diabetes.

Fifteen (21%) of 71 participants stated they had difficulty engaging family in a healthy lifestyle with respect to exercise

and food. Some participants explained that they had difficulty maintaining healthy lifestyle behaviors when with family, such as resisting unhealthy foods, and at times felt wearied by their family members' unhealthy behaviors:

"[I consume] a lot of tortillas. When you're eating with a lot of other people, and they heat them up and say 'Here are the tortillas!' As my aunt says, 'take them and eat up!' Yeah, more than anything, it's [hard to resist] tortillas."

However, one participant mentioned she had little difficulty changing their own eating habits with regards to appropriate food portions. Another participant did not live with any family, which made it easier to adhere to a healthy diet.

Seven participants (10%) expressed that diabetes had negatively affected their relationships with family members.

Table 3. Family-Related Challenges to Participant’s Diabetes Self-Management.

Theme	Subtheme	Example Quote
Lack of Support	Not knowing what it is like to live with diabetes	... I watch them eat it and serve a ton and everything. And oh my God. How do they not understand that I am not well and to see those attitudes and instead of helping me they make me worse because I am going to want to serve myself like them, eat the same that they’re is eating ...
	Lack of social support	I am here by myself. In other words, I am here with my four kids, no one else. So, I don’t have any relatives and my children, well they want to be taken care of, not the other way around.
	Diabetes has negative effect on family relationship	Well in my house, one gets depressed a lot and one has less control over one’s nerves. In other words, I used to be very tranquil before and I could control myself. With the diabetes, I feel that there’s no control, so then I ... for example, with my children, in order to not get dramatic when something happens, well the only thing I do is lock myself in my room and that’s where I sleep, but I don’t sleep.
	Difficulty maintaining changes when with family	[I consume] a lot of tortillas. [W]hen you’re eating with a lot of other people, and they heat them up and say “Here are the tortillas!” As my aunt says, “take them and eat up!” Yeah, more than anything, it’s [hard to resist] tortillas.
	Difficulty motivating family to engage in a healthy lifestyle with participant	I want to encourage [my kids] to come with me walking or to exercise, but they don’t want to. They are active in school but they don’t come. They don’t want to go through that. [My family is] used to eating differently and so, some of them want to eat what I eat and others do not. For example, if I eat vegetables, they do not want vegetables, uh-huh. If I eat, for example, fish, they do not want fish.
Family Responsibilities	I am in charge of taking [my mother] to the doctor, and my father-in-law when it is my turn, my husband, two daughters, three single daughters, three dogs that are like children also, so I am very busy I would like to have a bit more time to do a bit more [exercise].	
Stress Related to Family	Well, imagine. Not being able to be with my mom affects me. Not being able to go see her. Yes, it does affect me but I try to stay calm because I know that it is harmful to my health. My husband is continuously sick, and I think that is what makes my pressure a bit [higher], because there is a lot more pressure ... in my body. Because it is like a stress that you carry, a very strong burden inside.	

Family responsibilities. Of all participants, 16 (23%) were routinely busy taking children to and from school or taking family members to appointments, which competed with their ability to provide self-care. Two participants had difficult family environments that interfered with management behaviors. One participant noted:

I am in charge of taking [my mother] to the doctor, and my father-in-law when it is my turn, my husband, two daughters, three single daughters, three dogs that are like children also, so I am very busy. I would like to have a bit more time to do a bit more [exercise].”

Stress related to family. Seven participants (10%) with diabetes mentioned stress from family as negatively affecting their self-care, citing sick family members, issues coping with family stress, and the stress of living separately from family members.

Impact of Diabetes Education Information on Family (Table 4)

Sharing information. Of all participants, 16 (23%) reported they disseminated information to family members, with many

sharing healthy eating tips and materials from the study intervention:

“My 18-year old nephew who I took care of when he was young, tells me “Godmother, you know, ever since you have been giving my mom the handouts, she has had everyone on a diet.” And he has lost a lot of weight. He says, “My mom only wants to feed us lettuce like rabbits.”

Some participants encountered family tension when sharing diabetes information, but most (23%) said that the knowledge they shared with family led to a positive impact on family behavior, especially regarding diet.

Multiple participants mentioned the desire to include family members, especially children, in diabetes education programs to prevent the disease in future generations.

Perceived improvement in family relationships. Four participants noted that sharing diabetes knowledge with family helped involve them in their diabetes management, improved communication within the family, and allowed them to share their feelings about diabetes:

Table 4. Impact of Diabetes Education Information on Family.

Theme	Subtheme	Example quote
Sharing Information	Disseminating information to family members	<i>My father-in-law who is diabetic ... is an old man with 87 years ... everything that I know, I pass on to him ... So, in other words what I learn, I pay it forward.</i>
	Shared information led to positive changes in family behavior	<i>My 18-year old nephew who I took care of when he was young, tells me "Godmother, you know, ever since you have been giving my mom the handouts, she has had everyone on a diet." And he has lost a lot of weight. He says, "My mom only wants to feed us lettuce like rabbits." My sister's husband has been able to control his sugar levels somewhat because of the handouts I have shared with them. He used to buy lots of packs of flour tortillas because he liked to eat burritos. He says, "Now I only eat a burrito weekly, before it was every day; instead of sliced red meat, they are filled with fish."</i>
Perceived Improvement in Family Relationships	<i>Now I am very happy with my family because we all get together and talk about this program. We talk about diabetes and about how we can take care of ourselves, how we have to eat, and what exercises we can do I explain to my family how my attitude has improved ... We communicate much better.</i>	

"Now I am very happy with my family because we all get together and talk about ... diabetes and about how we can take care of ourselves, how we have to eat, and what exercises we can do ... We communicate much better."

Family Member Perspectives on Diabetes Management (Table 5)

Family does not know how to help. Of 13 family members, 11 (85%) also expressed a desire to be more educated about diabetes or have more education provided to family members. One family member stated family responsibilities interfered with providing more support. One participant expressed her issues providing support in the context of their cultural upbringing:

"You need to deal with customs and traditions. You need to find out the best way to educate yourself ..."

Some expressed feeling helpless, had difficulty changing the family environment to support healthy behaviors, and lacked knowledge of how to cook according to a diabetic diet:

"That's the main problem, the food, the meals, and you have to cook for yourself and you have to cook different for somebody else."

One participant noted getting support from other community members:

"I think that parents in the Hispanic community are very closed, like 'this is my problem', we're gonna deal with it as a family we don't need anybody else's help. But as a teacher I have parents who'll come and ask me 'What do you think about this?'"

Effect on emotional wellbeing. Six (46%) of all family members of someone with diabetes reported that diabetes had affected

their emotions, inciting worry, stress, and anger. One participant stated cultural traditions of respecting elders contributed to stress when helping manage their loved one's diabetes:

"You know, he's your dad and you're trying to at the same time please him but also in our culture ... you respect your elders and you can't always tell them no ... it's just a lot of stress and you try to figure out how are you gonna balance everything."

Diabetes affects the whole family. Four family members of participants with diabetes reported that diabetes affected the entire family.

"Whether you want it or not, it (diabetes) affects the whole family, because the family has to be there for support."

Family provides support. Five family members expressed they did provide support for those with diabetes.

Discussion

This study outlines various key themes regarding the family role in diabetes self-management among Mexican American adults, including perspectives from family members. This relationship has been investigated, but previous qualitative studies have focused on Mexican Americans in different geographic areas or pairs (Hu et al., 2013; McEwen et al., 2017) and, to our knowledge, this is the only study in recent years that categorizes this relationship in an urban city with a majority Mexican American Hispanic population.

These results emphasize that family provides varying types of support for diabetes self-care behaviors, and often is a motivator to improving self-management behaviors. Similar studies have noted grandchildren, spouses, and other family members as motivation (Hu et al., 2013; Perez-Brescia et al., 2022;

Table 5. Family Member Perspectives on Diabetes Management.

Theme	Subtheme	Example Quote
Family Does Not Know How to Help	Feeling helpless	<i>I think that parents in the Hispanic community are very closed, like “this is my problem”, we’re gonna deal with it as a family we don’t need anybody else’s help. But as a teacher I have parents who’ll come and ask me “What do you think about this?”, “What do you think about that?” [I]t’s like the not, not, not knowing how to help, because we don’t spend all day with that person ... and sometimes, one can see that that person isn’t doing what they are supposed to be doing, right ... So then [one feels] like impotence, right? To not be able to do more ...</i>
	Issues changing the family environment	<i>That’s the main problem, the food, the meals, and you have to cook for yourself and you have to cook different for somebody else. They’re trying to work, they’re trying to commute, number one, to the area they need to go to and then they come back, they take care of their kids and sometimes they’re just tired. What do you do? You know to go pick up some hotdogs, go pick up some tacos, go pick up whatever.</i>
	Lack of knowledge of how to cook according to a diabetic diet	<i>... I just don’t know what to cook to bring change in his diet, change cakes, different recipes, and I, everyday I’m trying to do something different but this is, I need to know what to cook and what I could use in substitute for ... things ... like that.</i>
	Family wants to learn more about diabetes	<i>You need to deal with customs and traditions. You need to find out the best way to educate yourself ... to try to understand the subject and how to understand how you will feel with the problem and send the right message. I think it’s good for [family] to come just to be a support to you, I mean, this is why I am here, it’s just to kind of understand my father a little. I’m also concerned about my daughters. Their grandmother was also diabetic. I don’t know how alcohol also affected the diabetes ... if it really progressed to even more, maybe not taking care of themselves as much. Though my daughters aren’t alcoholics but it’s just the fact that I don’t know how it is, how I can educate them ... I wanna see how I can help them.</i>
Effect on Emotional Wellbeing	Cultural barriers influence wellbeing	<i>I think with my father [who has diabetes] ... you know, we don’t want it to control our lives but at the same time we, I know that if he doesn’t eat right it’s gonna make him sick ... so it just kind of concerns me. You know, he’s your dad and you’re trying to at the same time please him but also in our culture, you know, you respect your elders and you can’t always tell them no ... it’s just a lot of stress and you try to figure out how are you gonna balance everything.</i>
Diabetes Affects the Whole Family	<i>The disease of ... a diabetic isn’t of ... exactly one person. Whether you want it or not, it affects the whole family, because the family has to be there for support. So, like, if we were here one family, and only one of us was diabetic, we would have this whole family supporting us. ... it’s like any other disease, like an alcoholic. An alcoholic isn’t just the only affected. The entire family is affected.</i>	
Family Provides Support	<i>Well, I would help by cooking. By trying to cook something for everyone, except that since the person is diabetic I would try to follow the same diet. I would not cook things that would be harmful for that person. That’s how I would help.</i>	

Rascón et al., 2022; Schmittdiel et al., 2018). Despite this knowledge, the proportion of Mexican Americans with poorly controlled T2DM (Hemoglobin A1C >9%) remains almost double that of non-Hispanic whites (CDC, 2023). Our study provides detailed evidence on the struggles those with diabetes face and how we can ameliorate them in Mexican American families.

Our research enforces that diet and T2DM are inextricably linked. Dietary changes hold a complex role in self-management behaviors. Often, conflict within families centered around lack of support for healthy eating and poor food choices by family

members, which is similar to qualitative research results by Toney et al. (2023). Our study highlights the complexities of this theme within the Mexican American community, and future directions should encourage dietary recommendations that are culturally adaptable in Mexican American families.

Participants often shared diabetes education information within the family unit which led to positive behavioral changes in family members. There are increased odds of unrecognized prediabetes in the Hispanic population (Cowie et al., 2018). Research on non-Hispanic adults has shown that family members of those newly diagnosed with T2DM often

implement significant changes in health-related behaviors both consciously and subconsciously (Schmittiel et al., 2018). In research performed by Reininger et al. (2020), involving family in a diabetes intervention led to improved A1C at a 12-month interval, which suggests a longer intervention period may be more beneficial in populations who experience more social or cultural barriers to health. To address shortcomings in preventative health, diabetes interventions could be more effective by including family, perhaps with a focus on shareable resources and hands-on demonstrations to encourage the exponential spread of health benefits to at-risk family members.

The interference of family responsibilities and family-related stress with diabetes self-management noted amongst our study participants has also been found in other studies including Mexican Americans, (Hu et al., 2013) and reflects how the existing healthcare system structure is intertwined with social determinants of health. Revising policies to expand transportation access and utilization may unload that burden off of caregivers, and also may address limited transportation options to healthy foods if they are further from grocery stores than convenience stores. We corroborate with existing literature that policies should focus on improving systematic support to ensure that interventions shown to improve diabetes outcomes can be translated effectively into the real world (Marquez et al., 2019).

Many family members in this study felt helpless due their lack of knowledge of how to best be a support person for one with diabetes. Prior studies have noted similar findings among family members of Mexican American patients (Hu et al., 2013; McEwen et al., 2017) with one intervention that involved teaching family how to best provide positive social and emotional support leading to clinically significant weight loss (Rosas et al., 2020). There is evidence that inclusion of a family member in a group support intervention can lead to statistically significant improvements in fasting blood sugar levels and diabetes knowledge in a rural Mexican American population and improvements in diabetes outcomes (Brown & Hanis, 1995; Mendez-Luck et al., 2019). However, family-based interventions remain to be adopted as mainstream despite this knowledge and the continued disparities in T2DM prevalence and its complications in Mexican Americans today. Our study adds qualitative evidence to the shortcomings and necessities of diabetes education in this Hispanic subgroup. It is worthwhile to prioritize efforts to include family in interventions to improve T2DM outcomes, especially in urban, Mexican American populations where the data are lacking.

These results show that family members often sought other trusted community members outside of the family unit, such as teachers, to guide behaviors. Using community health workers, or promotoras (lay community-based health outreach partner) has led to improved lifestyle management in interventional studies (Flores-Luevano et al., 2020). While not directly family, extending the familial relationship to trustworthy citizens and forming interventions that utilize these members could help close the disparities gap.

Generally, our findings are similar to those found in other ethnic minorities. In certain East Asian groups, family is a motivator but also presents dietary challenges, (Liu et al., 2022; Nguyen & Jones, 2021), with an added pressure has been noted to improve self-care to avoid becoming a burden to family as they age (Yoon et al., 2022). Park et al. (2022) found that patients with T2DM trusted health information when explained by family members more than providers, which may be why *promotoras* previously described have improved diabetes self-management. Similar thoughts on barriers and motivators regarding family and diabetes self-management have been described in qualitative literature focusing on black and South Asian communities (Deol et al., 2022; Lee et al., 2019; Rao et al., 2021). This suggests the desires expressed in our study are generalizable to other ethnic minorities facing health disparities.

Limitations of this study included that it was completed in an urban, Midwestern city with a large Mexican American population, and our findings may not be generalizable to all Hispanic populations. Furthermore, the study population was chosen by convenience sampling and collected on a volunteer basis, which may not match a general population, but our results are consistent with other studies with Mexican Americans with diabetes (Hu et al., 2013; McEwen & Murdaugh, 2014). Our study population also consisted of mostly women, so results may not be applicable for Mexican American men. Since our study design analyzed interviews retrospectively and was exploratory in nature, our data may not capture the entire breadth of opinions and beliefs of the family role for Mexican American adults with T2DM, although it is similar to findings in other ethnic groups.

Conclusion

We found that while lack of social support, family responsibilities, and stress from family can undermine successful diabetes management, many participants with diabetes had family available to support their diabetes management but wanted them to provide more social support. Seeing family members suffer with diabetes and preventing diabetes in future generations were noted as motivators. Diabetes knowledge was shared within the family unit, and family members had a desire to learn more about the disease, but didn't know how to best support at times due to conflict from cultural family norms. Further research should use our findings to inform culturally-tailored diabetes education efforts in order to improve health-care disparities in the Hispanic community.

Acknowledgements

This project was funded by the 2017 Summer Research Program at The University of Chicago Pritzker School of Medicine. This research was supported by grants from the University of Chicago Clinical and Translational Science Award (UL1RR024999), the National Institute of Diabetes and Digestive and Kidney Diseases Diabetes Research and Training Center (P60 DK20595) and the Chicago Center for

Diabetes Translation Research (P30 DK092949). Dr. Baig was supported by a NIDDK Mentored Patient-Oriented Career Development Award (K23 DK087903-01A1). Acknowledgements are made to the Little Village Community Advisory Board for their feedback and collaboration and Erin Staab, MPH for their help with this study. We have no conflict of interest to disclose.



Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Olivia J. Jordan  <https://orcid.org/0000-0002-1161-955X>
Amanda Benitez  <https://orcid.org/0000-0002-6177-4611>

Supplemental material

Supplemental material for this article is available online.

References

- 2020 Census Hispanic or Latino Population Data (2020). Great Cities Institute, University of Illinois at Chicago. <https://greatcities.uic.edu/2021/11/24/2020-census-hispanic-or-latino-population-data/>
- Adult Diabetes Rate (2021). Chicago Health Atlas. <https://www.chicagohealthatlas.org/indicators/diabetes>
- Aguayo-Mazzucato, C., Diaque, P., Hernandez, S., Rosas, S., Kostic, A., & Caballero, A. E. (2019). Understanding the growing epidemic of type 2 diabetes in the Hispanic population living in the United States. *Diabetes/Metabolism Research and Reviews*, 35(2), e3097. <https://doi.org/10.1002/dmrr.3097>
- American Diabetes Association Professional Practice Committee (2022). 5. Facilitating behavior change and well-being to improve health outcomes: *Standards of Medical Care in Diabetes—2022*. *Diabetes Care*, 45(Supplement_1), S60–S82. <https://doi.org/10.2337/dc22-S005>
- Baig, A. A., Locklin, C. A., Wilkes, A. E., Oborski, D. D., Acevedo, J. C., Gorawara-Bhat, R., Quinn, M. T., Burnet, D. L., & Chin, M. H. (2012). “One can learn from other people’s experiences”: Latino adults’ preferences for peer-based diabetes interventions. *The Diabetes Educator*, 38(5), 733–741. <https://doi.org/10.1177/0145721712455700>
- Baig, A. A., Benitez, A., Paredes, A. Z., Gao, Y., Monnot, L., Brazda, K., & Quinn, M. T. (2014). 37th annual meeting of the society of general internal medicine. <https://doi.org/10.1007/s11606-014-2834-9>
- Baig, A. A., Benitez, A., Locklin, C. A., Gao, Y., Lee, S. M., Quinn, M. T., Solomon, M. C., Sánchez-Johnsen, L., Burnet, D. L., & Chin, M. H., & Little Village Community Advisory Board (2015). Picture good health: A church-based self-management intervention among Latino adults with diabetes. *Journal of General Internal Medicine*, 30(10), 1481–1490. <https://doi.org/10.1007/s11606-015-3339-x2>
- Baig, A. A., Benitez, A., & Acevedo, M., (2016). *Local patients, local stories*. American Public Health Association Annual Meeting.
- Brown, S. A., & Hanis, C. L. (1995). A community-based, culturally sensitive education and group-support intervention for Mexican Americans with NIDDM: A pilot study of efficacy. *The Diabetes Educator*, 21(3), 203–210. <https://doi.org/10.1177/014572179502100307>
- Centers for Disease Control and Prevention. (2023). *National Diabetes Statistics Report*. <https://www.cdc.gov/diabetes/data/statistics-report/index.html>
- Community Snapshot: South Lawndale. (2016). Rush University Medical Center. <https://www.rush.edu/sites/default/files/2020-10/chna-south-lawndale.pdf>
- Cowie, C. C., Casagrande, S. S., & Geiss, L. S. (2018). Prevalence and incidence of type 2 diabetes and prediabetes. In Cowie, C. C., Casagrande, S. S., Geiss, L. S., Menke, A., Cissell, M. A., Eberhardt, M. S., Meigs, J. B., Gregg, E. W., Knowler, W. C., Barrett-Connor, E., Becker, D. J., Brancati, F. L., Boyko, E. J., Herman, W. H., Howard, B. V., Narayan, K. M. V., Rewers, M., & Fradkin, J. E., (Eds.), *Diabetes in America* (3rd ed.). National Institute of Diabetes and Digestive and Kidney Diseases (US).
- Deol, R. M., Thompson, L. M., Chun, K. M., & Chesla, C. (2022). Managing type 2 diabetes: Beliefs and daily practices in first generation Asian Indians in the United States. *SAGE Open Nursing*, 8, 23779608211054814. <https://doi.org/10.1177/23779608211054814>
- Flores-Luevano, S., Pacheco, M., Shokar, G. S., Dwivedi, A. K., & Shokar, N. K. (2020). Impact of a culturally tailored diabetes education and empowerment program in a Mexican American population along the US/Mexico border: A pragmatic study. *Journal of Clinical Medicine Research*, 12(8), 517–529. <https://doi.org/10.14740/jocmr4273>
- Fort, M. P., Steiner, J. F., Santos, C., Moore, K. R., Villaverde, M., Nease, D. E., Ortega, D., & Manson, S. M. (2020). Opportunities, challenges, and strategies for engaging family in diabetes and hypertension management: A qualitative study. *Journal of Health Care for the Poor and Underserved*, 31(2), 827–844. <https://doi.org/10.1353/hpu.2020.0063>
- Haw, J. S., Shah, M., Turbow, S., Egeolu, M., & Umpierrez, G. (2021). Diabetes complications in racial and ethnic minority populations in the USA. *Current Diabetes Reports*, 21(1), 2. <https://doi.org/10.1007/s11892-020-01369-x>
- Hu, J., Amirehsani, K., Wallace, D. C., & Letvak, S. (2013). Perceptions of barriers in managing diabetes: Perspectives of Hispanic immigrant patients and family members. *The Diabetes Educator*, 39(4), 494–503. <https://doi.org/10.1177/0145721713486200>
- Ingram, M., Torres, E., Redondo, F., Bradford, G., Wang, C., & O’Toole, M. L. (2007). The impact of promotoras on social support and glycemic control among members of a farmworker community on the US-Mexico border. *The Diabetes Educator*, 33(Suppl 6), 172S–178S. <https://doi.org/10.1177/0145721707304170>
- Lee, L. T., Jung, S. E., Bowen, P., Clay, O. J., Locher, J. L., & Cherrington, A. L. (2019). Understanding the dietary habits of black men with diabetes. *The Journal for Nurse Practitioners : JNP*, 15(5), 365–369. <https://doi.org/10.1016/j.nurpra.2018.12.023>
- Liu, Y., Jiang, J., You, W., Gong, D., Ma, X., Wu, M., & Li, F. (2022). Exploring facilitators and barriers to self-management engagement of Chinese people with type 2 diabetes mellitus and

- poor blood glucose control: A descriptive qualitative study. *BMC Endocrine Disorders*, 22(1), 294. <https://doi.org/10.1186/s12902-022-01214-0>
- Marquez, I., Calman, N., & Crump, C. (2019). A framework for addressing diabetes-related disparities in US Latino populations. *Journal of Community Health*, 44(2), 412–422. <https://doi.org/10.1007/s10900-018-0574-1>
- Mayberry, L. S., Berg, C. A., Greevy, R. A. Jr., & Wallston, K. A. (2019). Assessing helpful and harmful family and friend involvement in adults' type 2 diabetes self-management. *Patient Education and Counseling*, 102(7), 1380–1388. <https://doi.org/10.1016/j.pec.2019.02.027>
- Mayberry, L. S., Rothman, R. L., & Osborn, C. Y. (2014). Family members' obstructive behaviors appear to be more harmful among adults with type 2 diabetes and limited health literacy. *Journal of Health Communication*, 19(2, Suppl 2), 132–143. <https://doi.org/10.1080/10810730.2014.938840>
- McEwen, M. M., & Murdaugh, C. (2014). Partnering with families to refine and expand a diabetes intervention for Mexican Americans. *The Diabetes Educator*, 40(4), 488–495. <https://doi.org/10.1177/0145721714528996>
- McEwen, M. M., Pasvogel, A., & Murdaugh, C. (2019). Effects of a family-based diabetes intervention on family social capital outcomes for Mexican American adults. *The Diabetes Educator*, 45(3), 272–286. <https://doi.org/10.1177/0145721719837899>
- McEwen, M. M., Pasvogel, A., Murdaugh, C., & Hepworth, J. (2017). Effects of a family-based diabetes intervention on behavioral and biological outcomes for Mexican American adults. *The Diabetes Educator*, 43(3), 272–285. <https://doi.org/10.1177/0145721717706031>
- Mendez-Luck, C. A., Miranda, J., Mangione, C. M., Yoon, J., & VanGarde, A. (2019). The *Juntos* pilot study: A diabetes management intervention for Latino caregiving dyads. *The Diabetes Educator*, 45(5), 507–519. <https://doi.org/10.1177/0145721719866619>
- Morse, J. M. (1995). The significance of saturation. In Morse, J. (Ed.), *Qualitative health research* (pp. 147–149). Sage Publications.
- National Center for Health Statistics (2022). *Health, United States, 2020-2021: Table 14*. <https://www.cdc.gov/nchs/hus/data-finder.htm>
- Nguyen, A. T., & Jones, E. J. (2021). Diabetes beliefs, culturally influenced self-management practices, and interventions among Vietnamese adults: A systematic review. *Journal of Transcultural Nursing : Official Journal of the Transcultural Nursing Society*, 32(3), 266–275. <https://doi.org/10.1177/1043659620988317>
- Park, L., Vang, A., Yang, B., & Quanbeck, A. (2022). Barriers to type 2 diabetes mellitus management for older Hmong patients with minimal English language skills: Accounts from caregivers, case managers, and clinicians. *Journal of Racial and Ethnic Health Disparities*, 1–8. Advance online publication. <https://doi.org/10.1007/s40615-022-01480-7>
- Perez-Brescia, M., Beck, C. T., Alicea Planas, J., Newlin-Lew, K. H., Whittemore, R., & Juarez, A. (2022). *Familismo Primero* and *Puerta Cerrada* in self-managing diabetes among Hispanics: A qualitative meta-synthesis. *Journal of Transcultural Nursing : Official Journal of the Transcultural Nursing Society*, 33(6), 666–674. <https://doi.org/10.1177/10436596221109834>
- Rao, D., Meyer, J., Maurer, M., & Shyanbola, O. O. (2021). Perceptions of psychosocial and interpersonal factors affecting self-management behaviors among African Americans with diabetes. *Exploratory Research in Clinical and Social Pharmacy*, 3, 100057. <https://doi.org/10.1016/j.rcsop.2022.100196>
- Rascón, A. M., McEwen, M. M., & Crist, J. D. (2022). “My problems aren't their fault”: Mexican American women managing diabetes while caring for grandchildren. *Journal of Women & Aging*, 34(6), 745–756. <https://doi.org/10.1080/08952841.2021.1969861>
- Reininger, B. M., Lee, M., Hessabi, M., Mitchell-Bennett, L. A., Sifuentes, M. R., Guerra, J. A., Ayala, C. D., Xu, T., Polletta, V., Flynn, A., & Rahbar, M. H. (2020). Improved diabetes control among low-income Mexican Americans through community-clinical interventions: Results of an RCT. *BMJ Open Diabetes Research & Care*, 8(1), e000867. <https://doi.org/10.1136/bmjdr-2019-000867>
- Rosas, L. G., Lv, N., Xiao, L., Lewis, M. A., Venditti, E. M. J., Zavella, P., Azar, K., & Ma, J. (2020). Effect of a culturally adapted behavioral intervention for Latino adults on weight loss over 2 years: A randomized clinical trial. *JAMA Network Open*, 3(12), e2027744. <https://doi.org/10.1001/jamanetworkopen.2020.27744>
- Schmittdiel, J. A., Cunningham, S. A., Adams, S. R., Nielsen, J., & Ali, M. K. (2018). Influence of a new diabetes diagnosis on the health behaviors of the patient's partner. *Annals of Family Medicine*, 16(4), 290–295. <https://doi.org/10.1370/afm.2259>
- Teufel-Shone, N. I., Drummond, R., & Rawiel, U. (2005). Developing and adapting a family-based diabetes program at the U.S.-Mexico border. *Preventing Chronic Disease*, 2(1), A20. <https://pubmed.ncbi.nlm.nih.gov/15670473/>
- Toney, A. M., Pinerros-Leano, M., Pérez-Flores, N. J., Gomez, D., & Aguayo, L. (2023). ‘It is in our hands-why wait until you are sick?’: Perceptions about diabetes prevention of Latina mothers in Mexico and the United States. *Diabetic Medicine : A Journal of the British Diabetic Association*, e15060. Advance online publication. <https://doi.org/10.1111/dme.15060>
- Wells, K., & Jones, L. (2009). Research” in community-partnered, participatory research. *JAMA*, 302(3), 320–321. <https://doi.org/10.1001/jama.2009.1033>
- Yoon, S., Kwan, Y. H., Phang, J. K., Tan, W. B., & Low, L. L. (2022). Personal goals, barriers to self-management and desired mHealth application features to improve self-care in multi-ethnic Asian patients with type 2 diabetes: A qualitative study. *International Journal of Environmental Research and Public Health*, 19(22), 15415. <https://doi.org/10.3390/ijerph192215415>