

UC Merced

UC Merced Previously Published Works

Title

Genome of the Root-Associated Plant Growth-Promoting Bacterium *Variovorax paradoxus* Strain EPS

Permalink

<https://escholarship.org/uc/item/57n0w4tp>

Journal

Microbiology Resource Announcements, 1(5)

ISSN

2169-8287

Authors

Han, Jong-In
Spain, Jim C
Leadbetter, Jared R
[et al.](#)

Publication Date

2013-10-31

DOI

10.1128/genomea.00843-13

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Genome of the Root-Associated Plant Growth-Promoting Bacterium *Variovorax paradoxus* Strain EPS

Jong-In Han,^a Jim C. Spain,^b Jared R. Leadbetter,^c Galina Ovchinnikova,^d Lynne A. Goodwin,^e Cliff S. Han,^e Tanja Woyke,^e Karen W. Davenport,^d Paul M. Orwin^f

Department of Civil and Environmental Engineering, KAIST, Daejeon, Republic of Korea^a; School of Civil and Environmental Engineering, Georgia Institute of Technology, Atlanta, Georgia, USA^b; Divisions of Biology and Environmental Science and Engineering, California Institute of Technology, Pasadena, California, USA^c; DOE Joint Genome Institute, Walnut Creek, California, USA^d; Los Alamos National Laboratory, Los Alamos, New Mexico, USA^e; Department of Biology, California State University at San Bernardino, San Bernardino, California, USA^f

***Variovorax paradoxus* is a ubiquitous betaproteobacterium involved in plant growth promotion, the degradation of xenobiotics, and quorum-quenching activity. The genome of *V. paradoxus* strain EPS consists of a single circular chromosome of 6,550,056 bp, with a 66.48% G+C content.**

Received 18 September 2013 Accepted 27 September 2013 Published 24 October 2013

Citation Han J-I, Spain J-C, Leadbetter J-R, Ovchinnikova G, Goodwin L-A, Han C-S, Woyke T, Davenport K-W, Orwin P-M. 2013. Genome of the root-associated plant growth-promoting bacterium *Variovorax paradoxus* strain EPS. *Genome Announc.* 1(5):e00843-13. doi:10.1128/genomeA.00843-13.

Copyright © 2013 Han et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](http://creativecommons.org/licenses/by/3.0/).

Address correspondence to Paul M. Orwin, porwin@csusb.edu.

Here, we report the finished genome sequence of *Variovorax paradoxus* strain EPS. *V. paradoxus* EPS was cultured on low-nutrient agar (5 mg/liter yeast extract [YE]) from the rhizosphere community of the sunflower (*Helianthus annuus*) on the campus of California State University, San Bernardino, CA. Other bacterial strains from this species have been identified in many different environments, including diverse soils, endophytic growth (1), and the human oral microbiota (2). Strains within this species have been shown to degrade acyl-homoserine lactone signaling compounds (3), pesticides (4), and nitrotyrosines (5), and they promote plant growth in several crops (6, 7). The genome of the potato endophyte strain *V. paradoxus* S110 was previously reported, consisting of two chromosomes with a total genome size of 6.7 Mb (1). The surface attachment and motility characteristics of *V. paradoxus* EPS have also been evaluated previously (8, 9).

The genome of *V. paradoxus* EPS was sequenced using high-throughput sequencing approaches (Illumina and 454, 30× genome coverage) from pure culture-derived genomic DNA (Promega). Sequencing and annotation were performed at the Joint Genome Institute Oak Ridge National Laboratory (JGI-ORNL) and JGI Production Genomics Facility (JGI-PGF), and finishing was done by the JGI Los Alamos National Laboratory (JGI-LANL). The finished genome was assembled using Newbler version 2.3. The genome is 6,550,056 bases in length, is organized into a single circular chromosome, and has a G+C content of 66.48%. A total of 6,088 genes were annotated, including 6,020 putative protein-coding sequences (91.25% of total bases). Of these, functions have been predicted for 73.92% (4,499 loci). A gene for 1-aminocyclopropane-1-carboxylate (ACC) deaminase was identified (Varpa_5820), consistent with a role as a plant-growth promoting rhizobacterium, along with a gene for acyl-homoserine lactone acylase activity (Varpa_4314) described previously in other *V. paradoxus* strains (1, 3). Although this bacterium is motile using a single polar flagellum, the flagellar locus was not clearly identified in this chromosomal sequence, as it was in *V. paradoxus*

strain S110. Putative pilus loci were identified, as were a number of secretion systems. Two potential prophage elements (one potentially complete prophage) were also identified with the PHAST tool (<http://phast.wishartlab.com> [10]) in positions 1534102 to 1541586 and 2169456 to 2239232. An additional potential prophage region starting at tRNA-Lys (position 2933124, with several phage-related genes between open reading frames [ORFs] Varpa_2722 to Varpa_2770) was identified by inspection and subsequent BLAST analysis. The annotated genome data are available in IMG (see <http://img.jgi.doe.gov>, taxon ID 649633106), as well as NCBI (taxon ID595537) and the Genomes OnLine Database (GOLD card no. Gc01580).

Nucleotide sequence accession number. The finished and annotated genome sequence has been deposited in GenBank under the accession no. CP002417.

ACKNOWLEDGMENTS

This work was funded by CSP_788249 from the Department of Energy Joint Genome Institute, Walnut Creek, CA. The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under contract no. DE-AC02-05CH11231.

REFERENCES

1. Han JI, Choi HK, Lee SW, Orwin PM, Kim J, LaRoe SL, Kim TG, O'Neil J, Leadbetter JR, Lee SY, Hur CG, Spain JC, Ovchinnikova G, Goodwin L, Han C. 2011. Complete genome sequence of the metabolically versatile plant growth-promoting endophyte *Variovorax paradoxus* S110. *J. Bacteriol.* 193:1183–1190.
2. Anesti V, McDonald IR, Ramaswamy M, Wade WG, Kelly DP, Wood AP. 2005. Isolation and molecular detection of methylotrophic bacteria occurring in the human mouth. *Environ. Microbiol.* 7:1227–1238.
3. Leadbetter JR, Greenberg EP. 2000. Metabolism of acyl-homoserine lactone quorum-sensing signals by *Variovorax paradoxus*. *J. Bacteriol.* 182:6921–6926.
4. Dejonghe W, Berteloot E, Goris J, Boon N, Crul K, Maertens S, Höfte M, De Vos P, Verstraete W, Top EM. 2003. Synergistic degradation of

- linuron by a bacterial consortium and isolation of a single linuron-degrading *Variovorax* strain. *Appl. Environ. Microbiol.* **69**:1532–1541.
5. Nishino SF, Spain JC. 2006. Biodegradation of 3-nitrotyrosine by *Burkholderia* sp. strain JS165 and *Variovorax paradoxus* JS171. *Appl. Environ. Microbiol.* **72**:1040–1044.
 6. Belimov AA, Dodd IC, Hontzeas N, Theobald JC, Safronova VI, Davies WJ. 2009. Rhizosphere bacteria containing 1-aminocyclopropane-1-carboxylate deaminase increase yield of plants grown in drying soil via both local and systemic hormone signalling. *New Phytol.* **181**:413–423.
 7. Jiang F, Chen L, Belimov AA, Shaposhnikov AI, Gong F, Meng X, Hartung W, Jeschke DW, Davies WJ, Dodd IC. 2012. Multiple impacts of the plant growth-promoting rhizobacterium *Variovorax paradoxus* 5C-2 on nutrient and ABA relations of *Pisum sativum*. *J. Exp. Bot.* **63**: 6421–6430.
 8. Jamieson WD, Pehl MJ, Gregory GA, Orwin PM. 2009. Coordinated surface activities in *Variovorax paradoxus* EPS. *BMC Microbiol.* **9**:124.
 9. Pehl MJ, Jamieson WD, Kong K, Forbester JL, Fredendall RJ, Gregory GA, McFarland JE, Healy JM, Orwin PM. 2012. Genes that influence swarming motility and biofilm formation in *Variovorax paradoxus* EPS. *PLoS One* **7**:e31832. doi:[10.1371/journal.pone.0031832](https://doi.org/10.1371/journal.pone.0031832).
 10. Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool. *Nucleic Acids Res.* **39**:W347–W352.