

UC San Diego

UC San Diego Previously Published Works

Title

Society of Anesthesia and Sleep Medicine Guidelines on Preoperative Screening and Assessment of Adult Patients With Obstructive Sleep Apnea

Permalink

<https://escholarship.org/uc/item/58h4r790>

Journal

Anesthesia & Analgesia, 123(2)

ISSN

0003-2999

Authors

Chung, Frances

Memtsoudis, Stavros G

Ramachandran, Satya Krishna

et al.

Publication Date

2016-08-01

DOI

10.1213/ane.0000000000001416

Peer reviewed

OPEN

Society of Anesthesia and Sleep Medicine Guidelines on Preoperative Screening and Assessment of Patients With Obstructive Sleep Apnea

Frances Chung, MBBS, FRCPC,* Stavros Memtsoudis, MD, PhD,† Satya Krishna Ramachandran, MD,‡ Mahesh Nagappa, MD,§ Mathias Opperer, MD,|| Crispiana Cozowicz, MD,¶# Sara Patrawala, MD,**†† David Lam, BSc,* Anjana Kumar, BSc,‡ Girish P. Joshi, MD,‡‡ John Fleetham, MD,§§ Najib Ayas, MD,||| Nancy Collop, MD,¶¶ Anthony Doufas, MD, PhD,## Matthias Eikermann, MD, PhD,*** Marina Englesakis, HBA, MLIS,††† Bhargavi Gali, MD,‡‡‡ Peter Gay, MD,§§§ Adrian Hernandez, MD, MSc, PhD, FACC,||||| Roop Kaw, MD,¶¶¶ Eric Kezirian, MD,### Atul Malhotra, MD,**** Babak Mokhlesi, MD,†††† Sairam Parthasarathy, MD,‡‡‡‡ Tracey Stierer, MD,§§§§ Frank J. Pöpl, MD,|||||||¶¶¶¶ David R. Hillman, MD,#### and Dennis Auckley, MD*****

The purpose of the Society of Anesthesia and Sleep Medicine guideline on preoperative screening and assessment of patients with obstructive sleep apnea (OSA) is to present recommendations based on the available clinical evidence on the topic where possible. As very few well-performed randomized studies in this field of perioperative care are available, most of the recommendations were developed by experts in the field through consensus processes involving utilization of evidence grading to indicate the level of evidence upon which recommendations were based. This guideline may not be appropriate for all clinical situations and all patients. The decision whether to follow these recommendations must be made by a responsible physician on an individual basis. Protocols should be developed by individual institutions taking into account the patients' conditions, extent of interventions and available resources. This practice guideline is not intended to define standards of care or represent absolute requirements for patient care. The adherence to these guidelines cannot in any way guarantee successful outcomes and is rather meant to help individuals and institutions formulate plans to better deal with the challenges posed by perioperative patients with OSA. These recommendations reflect the current state of knowledge and its interpretation by a group of experts in the field at the time of publication. While these guidelines will be periodically updated, new information that becomes available between updates should be taken into account. Deviations in practice from guidelines may be justifiable and such deviations should not be interpreted as a basis for claims of negligence. (Anesth Analg 2016;XXX:00–00)

Obstructive sleep apnea (OSA) is a serious condition characterized by repeated episodes of complete or partial obstruction of the upper airway. These

episodes are accompanied by varying degrees of arterial oxygen desaturation and sympathetic activation. They are usually terminated by brief cortical arousals or, occasionally,

From the *Department of Anesthesiology, Toronto Western Hospital, University Health Network, University of Toronto, Toronto, Ontario, Canada; †Department of Anesthesiology, Weill Cornell Medical College and Hospital for Special Surgery, New York, New York; ‡Department of Anesthesiology, University of Michigan, Ann Arbor, Michigan; §Department of Anesthesiology and Perioperative Medicine, University Hospital, St. Joseph's Hospital and Victoria Hospital, London Health Sciences Centre and St. Joseph's Health care, Western University, London, Ontario, Canada; ||Paracelsus Medical University, Department of Anesthesiology, Perioperative Medicine and Intensive Care, Salzburg, Austria; ¶Department of Anesthesiology, Hospital for Special Surgery, Weill Cornell Medical College New York, New York; #Perioperative Medicine and Intensive Care Medicine, Paracelsus Medical University, Salzburg, Austria; **Department of Medicine, University of California San Diego, San Diego, California; ††Sparrow Hospital, Lansing, Michigan; ‡‡Department of Anesthesiology and Pain Management, University of Texas Southwestern Medical School, Texas; §§Department of Medicine, Division of Respiratory Medicine, The University of British Columbia, Vancouver, BC, Canada; |||University of British Columbia, Vancouver, BC, Canada; ¶¶Department of Medicine, Emory University, Atlanta, Georgia; ##Department of Anesthesiology, Perioperative and Pain Medicine, Stanford University Medical Center, Palo Alto, California; ***Department of Anesthesia, Critical Care and Pain Medicine, Harvard University, Cambridge, Massachusetts; †††Library and Information Services, University Health Network, University of Toronto, Toronto, Ontario, Canada; ‡‡‡Department of Anesthesiology, Mayo Clinic, Rochester, Minnesota;

§§§Department of Pulmonary, Critical Care and Sleep Medicine, Mayo Clinic, Rochester, Minnesota; |||||School of Medicine, Universidad Peruana de Ciencias Aplicadas (UPC), Lima, Peru; ¶¶¶Departments of Hospital Medicine and Outcomes Research (Anesthesiology Institute), Cleveland Clinic Lerner College of Medicine, Cleveland, Ohio; ###Department of Otolaryngology—Head and Neck Surgery, Keck School of Medicine of USC, Los Angeles, California; ****Division of Pulmonary and Critical Care Medicine, University of California San Diego, San Diego, California; ††††Department of Pulmonary and Critical Care, University of Chicago, Chicago, Illinois; #####Department of Pulmonary, Allergy, Critical Care, and Sleep Medicine, University of Arizona, Tucson, Arizona; §§§§Department of Anesthesiology and Critical Care Medicine, Johns Hopkins University, Baltimore, Maryland; |||||Department of Anaesthesiology and Intensive Care Medicine, University Witten/Herdecke, Cologne, Germany; ¶¶¶¶Department of Paediatric Anaesthesia, Cologne Medical Centre, Cologne, Germany; ####Department of Pulmonary Physiology and Sleep Medicine, Sir Charles Gairdner Hospital, University of Western Australia, Nedlands, Australia; and *****Division of Pulmonary, Critical Care and Sleep Medicine, Case Western Reserve University, School of Medicine, Cleveland, Ohio.

Accepted for publication April 23, 2016.

Funding: None.

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal's website (www.anesthesia-analgesia.org).

Reprints will not be available from the authors.

Address correspondence to Frances Chung, MBBS, FRCPC, Department of Anesthesiology, Toronto Western Hospital, University Health Network, University of Toronto, 399 Bathurst Street, McL 2-405 Toronto, ON, Canada M5T 2S8. Address e-mail to frances.chung@uhn.ca.

Copyright © 2016 International Anesthesia Research Society. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially. DOI: 10.1213/ANE.0000000000001416

awakenings. Resultant sleep disruption is responsible for the commonly associated symptom of excessive daytime sleepiness. Habitual snoring usually coexists. Apart from these symptoms, which can be very intrusive, OSA can lead to adverse health outcomes, including cerebrovascular disease, cardiovascular disorders (eg, hypertension, ischemic heart disease, arrhythmias, pulmonary hypertension, and congestive heart failure), metabolic syndrome, depression, and increased risk of accidents.¹⁻⁸ Numerous factors including alcohol consumption, smoking, obesity, increased neck circumference, male sex, advanced age, tonsillar and adenoidal hypertrophy, macroglossia, nasal obstruction, and craniofacial abnormalities increase the risk of OSA.⁹⁻¹²

Age and sex are important influences. Twenty-seven percent of women and 43% of men ages 50 to 70 years old are estimated to have OSA versus 9% of women and 26% of men in the 30- to 49-years category.^{13,14} Up to 80% of individuals with moderate-to-severe OSA may remain undiagnosed.^{15,16} Importantly, OSA is common in patients who present for surgery, with estimates ranging from the prevalence of the general population to as high as 70% in select populations (eg, bariatric surgical patients).^{17,18} Similar to the general population, most patients with OSA remain undiagnosed at the time of surgery.^{19,20} The diagnosis of OSA is confirmed by observing the number of apnea and/or hypopnea episodes per hour (the apnea-hypopnea index [AHI]) on an overnight polysomnography. Cutoffs for AHI have been used to define the severity of OSA. The American Academy of Sleep Medicine defines mild OSA as AHI 5 to 15 events/h, moderate OSA as AHI 15 to 30 events/h, and severe OSA as AHI >30 events/h.²¹ The severity of OSA as measured by the AHI may worsen in OSA patients during the postoperative period.^{22,23} Studies on several large databases, including millions of patients and 2 meta-analyses of 13 and 17 studies, respectively, have found that patients with OSA are at increased risk of postoperative complications.²⁴⁻²⁹

With an aging population and increasing rates of obesity, the prevalence of OSA is likely to increase.³⁰ Coupled with an overall growth in the number of surgical procedures being performed, this suggests that the number of patients presenting for surgery with OSA will grow substantially. Whether their OSA is diagnosed or undiagnosed, and whether or not those who are diagnosed are on effective treatment (commonly continuous positive airway pressure [CPAP] therapy),³¹ appropriate perioperative planning is challenging in these patients. Absence of clear recommendations is problematic, and clinicians often are faced with difficult ad hoc decisions regarding preoperative preparation, intraoperative management, and postoperative monitoring and implementation of treatment.

Perioperative complications related to OSA have increasingly resulted in malpractice lawsuits.³²⁻³⁹ The Society of Anesthesia and Sleep Medicine and the Anesthesia Closed Claims Project have established a registry in North America to accurately record adverse anesthetic events related to OSA.³⁶ Accumulation of these data will help to identify those aspects of OSA that have greatest influence on postoperative complications, allowing better risk stratification and risk prevention.³⁶ These uncertainties and the present lack of clear recommendations underline why guidelines based on

the best-available evidence (or expert opinion where such evidence is lacking) are needed to inform perioperative management of patients with diagnosed or suspected OSA. Because of the large volume of medical literature relating to perioperative management, this first Society of Anesthesia and Sleep Medicine (SASM) guideline concentrates on the initial aspect of this subject: preoperative evaluation. Intraoperative and postoperative management of OSA are not considered in this guideline and await future development, based on a similarly thorough literature reviews.

What Other Guidelines and Reviews Are Available?

Previous practice guidelines⁴⁰⁻⁴³ and reviews^{12,44} have been published by the American Society of Anesthesiologists (ASA),^{40,41} the Society for Ambulatory Anesthesia,⁴² the American Academy of Sleep Medicine,⁴³ in *Chest*,⁴⁴ and the *Canadian Journal of Anesthesia*.¹²

Why Was This Guideline Developed?

While worthy contributions, these previously published practice guidelines, and related reviews, have some obvious deficiencies. Most have failed to sufficiently examine and objectively establish OSA as a perioperative risk factor. Moreover, few have focused on an objective assessment of how to identify patients with OSA. An example is provided by the ASA Practice Guideline for the Perioperative Management of Patients with Obstructive Sleep Apnea, which was published in 2014,⁴¹ updating their 2006 guideline.⁴⁰ It recommends preoperative screening to identify undiagnosed OSA, preoperative initiation of CPAP therapy for OSA where possible, and postoperative monitoring of OSA patients.⁴¹ However, despite best intentions, a significant dilemma occurs because these broadly based recommendations are factored on limited evidence, and the cost of implementation is high. Given this problem, a more focused and practical approach is needed to balance the desire to reduce postoperative adverse events and improve long-term health benefits with appropriate allocation of health care resources.

How Does This Guideline Differ From Existing Guidelines?

This guideline was developed to provide a more effective and effective approach to preoperative workup of patients with diagnosed or suspected OSA. In doing so, it carefully examines the existing evidence-base for preoperative screening and preparation of patients with OSA, as well as the perioperative use of CPAP in patients with diagnosed or undiagnosed OSA. The Task Force recognizes the difficulties posed by late recognition of OSA during preoperative workup, and thus, the recommendations provide a practical balance between the desire to minimize postoperative complications and the requirement for efficient use of health care resources.

To enhance the value of these guidelines for treating physicians, we have categorized patients into 3 clinically relevant groups: diagnosed OSA, treated; diagnosed OSA, partially treated/untreated; and suspected OSA. The ASA 2014 guideline does not make these distinctions, thus limiting its utility in the clinical setting.⁴¹ It indicated "CPAP

should be initiated if patients have severe OSA," with no support for this statement.⁴¹ Our recommendations are based on a thorough evaluation of the literature, including a meta-analysis of preoperative CPAP use. In addition, as discussed later in this guideline, we recommend against delaying or canceling surgery for further workup except in particular groups of patients, with additional problems suggesting disturbed ventilation or gas exchange, such as evidence of hypoventilation, severe pulmonary hypertension, or resting hypoxemia in the absence of other diagnosed cardiopulmonary disease. An executive summary is attached as an addendum.

Aims

This guideline focuses on the preoperative screening and preparation of adult surgical patients scheduled for elective surgery, including implementation of CPAP therapy in the preoperative and postoperative period. The specific aims were to (1) summarize the current state of knowledge regarding the impact of OSA on perioperative outcomes; (2) determine the appropriate preoperative assessments for patients with suspected or diagnosed OSA scheduled for elective surgery; and (3) evaluate the current evidence informing best practices in preoperative management of diagnosed OSA, treated; diagnosed OSA, partially treated/untreated; and suspected OSA. In areas lacking sufficient published evidence, the task force sought to establish expert consensus. Patients affected by sleep-disordered breathing unrelated to OSA were not considered in this guideline, because there is insufficient evidence on which to base recommendations regarding preoperative management for these disorders. These include hypoventilation syndromes, periodic breathing, and central sleep apnea unrelated to OSA.

Guideline Task Force

The task force was composed of 28 members of SASM, an international society devoted to advancing standards of care for clinical problems shared by anesthesiology and sleep medicine. It included 12 anesthesiologists, 9 sleep medicine specialists, 2 hospitalists, 1 otolaryngologist, 2 research assistants, a research librarian, and a clinical epidemiologist. Members of the Task Force share expertise in the topics of sleep-disordered breathing and anesthesia and practice both in academic and in nonacademic settings in various parts of the United States, Canada, Europe, Australia, and South America.

METHODS

Research Questions

A systematic review of the literature addressing the preoperative assessment and preparation of patients with OSA in different areas was conducted. Research questions were divided into 3 groups (Table 1). The first group assessed the literature to determine how OSA affects perioperative outcomes (group 1). The second group investigated methods of screening preoperative patients to identify those at high risk for OSA in the preoperative period (group 2). The third group examined the literature to guide recommendations for best practices regarding preoperative management

Table 1. Selection Criteria and Study Questions

Population	Adults with suspected or diagnosed OSA undergoing elective surgery
Language	English
Questions	<ol style="list-style-type: none"> Does a diagnosis of OSA change postoperative outcomes? Preoperative assessment for OSA <ol style="list-style-type: none"> Should adult patients at risk for OSA be identified before surgery? Which tools can be used to identify surgical patients with suspected OSA in the preoperative period? What is the clinical value of performing additional preoperative tests? What are the best preoperative practices in patients who were diagnosed with OSA, nonadherent with CPAP therapy or have a high pretest probability for OSA? <ol style="list-style-type: none"> What is the evidence of efficacy of CPAP in surgical patients in the perioperative period? What are the best preoperative practices aimed at improving outcomes for surgical patients with known OSA who are adherent to PAP therapy? What are the best preoperative practices aimed at improving outcomes for surgical patients with known OSA but decline or are poorly adherent to therapy? What are the best preoperative practices aimed at improving outcomes for surgical patients with a high probability of OSA?
Study designs	Randomized controlled trials, prospective observational trials, and retrospective trials, systematic reviews, meta-analysis, and clinical guidelines.

Abbreviations: CPAP, continuous positive airway pressure; OSA, obstructive sleep apnea; PAP, positive airway pressure.

of patients with diagnosed OSA on therapy, patients with diagnosed OSA who decline or are nonadherent to therapy, and those with a high pretest probability for OSA (group 3). Leaders and members of each group are listed at the end of the manuscript.

Literature Search Strategy

With the help of a research librarian, depending on the topic, literature search was performed to include publications from 1946 to June 2014. Databases searched included PubMed-Medline, Medline in-process, and other nonindexed citations, Embase, Cochrane Central Register of Controlled Trials, Cochrane Database of Systematic Reviews (2005 to June 2014), and Health Technology Assessment (fourth Quarter 2013). Continued literature surveillance was done through November 2015. Our search was restricted to studies in adults (18 years of age and older) and published in the English language only. All duplicate records were removed.

In group 1, the search used the controlled vocabulary terms and key words: "sleep apnea, obstructive," "postoperative period," "complications" or "outcome," "perioperative care," "intraoperative monitoring," "postoperative monitoring," "perioperative complications," "intraoperative complications," "postoperative complications," "outcome," "risk," "morbidity," "mortality" and "death," and also "obstructive sleep apnea," "obstructive sleep apnea syndrome," "sleep disordered breathing," "obesity hypoventilation syndrome," "apnoea or apnea," "hypopnea or hypopnea."

In group 2, the search used the controlled vocabulary terms and key words: "mass screening," "polysomnography," "questionnaire," "sleep apnea," "sleep disordered breathing," "obesity hypoventilation syndrome," "Pickwick syndrome," "Cheyne Stokes respiration," "obesity hypoventilation syndrome," "central apnea," "diagnostic test," "probability," "sensitivity and specificity," "accuracy," and "diagnosis."

In group 3, the search used the controlled vocabulary terms and key words: "preoperative care," "preoperative evaluation," "preoperative assessment," "patient selection," "identified high-risk patient," "physician's practice patterns," "referral," "consult," and "gatekeeping." To determine the efficacy of CPAP in the surgical patients, the search used the MESH key words: "obstructive," "sleep apnea," "continuous positive airway pressure," "obstructive sleep apnea," "obstructive sleep apnea syndrome," "obstructive sleep apnoea syndrome," "sleep disordered breathing," "obesity hypoventilation syndrome," "apnoea or apnea," "hypopnoea or hypopnea," "auto-titrating positive airway pressure," "positive airway pressure," "bilevel positive airway pressure," "postoperative" and one of "complications" or "outcome," "perioperative care," "intraoperative care," "postoperative care," "intraoperative monitoring," "postoperative monitoring," "perioperative complications," "intraoperative complications," "postoperative complications," "outcome," "risk," "morbidity" or "mortality," and "death." The full search strategies used in Medline for the different questions are shown in the Supplemental Digital Content 1, Tables S1 to S7 (<http://links.lww.com/AA/B442>).

Study Selection and Grading of Strength of Evidence

In their respective groups, 2 or more reviewers assessed titles and abstracts, to identify whether studies met inclusion and exclusion criteria and graded the level of evidence. The reviewers for each respective group are listed at the end of the manuscript. The search results were assessed in a stepwise manner. In the first step of the review, relevant studies were selected after we reviewed the title of the search results. In the second step, the abstracts were screened to decide whether the eligibility criteria were met. In the third step, the selected articles were reviewed and relevant data were extracted. A citation search by manual review of references from primary or review articles was performed, and all relevant results were compiled. The number of studies excluded and the reason for exclusion of studies were recorded. Any disagreements were resolved by consensus or by consulting with the main SASM groups via face-to-face meetings, teleconferences, or e-mail communication. Study designs included randomized controlled trials, prospective observational studies, and retrospective studies, case series, systematic reviews, meta-analyses, and guidelines.

Data extracted from these studies included the type of study, level of evidence, demographic data, associated comorbid conditions, methods of OSA diagnoses, screening tools, types of OSA therapy, types of procedure, intraoperative and postoperative adverse events, hospital readmission

rates, mortality after surgery in OSA patients, preparation for surgery, sleep medicine referral, and CPAP therapy.

Exclusion Criteria

For group 1, regarding the evaluation of postoperative outcomes in OSA, studies not reporting on a control group or at least 1 postoperative adverse outcome and/or mortality were excluded. For group 2, regarding preoperative screening and assessment of patients with OSA, studies were excluded if there was no information on screening performed for OSA, for defining the standards of accuracy, or for best clinical assessment tests when screening and classifying the risk of OSA. For group 3, regarding preparation of OSA patients for surgery, studies were excluded if there was no information concerning strategies or treatment interventions in patients at high risk for OSA or for patients with OSA but nonadherent with treatment. For use of CPAP in surgical patients, studies were excluded if they involved upper airway surgery, provided no outcome data, or if patients did not use either preoperative or postoperative CPAP.

The included studies were graded by the reviewers for strength of evidence according to the Grading of Recommendations, Assessment, Development, and Evaluation (GRADE) system^{45,46} (Table 2). The SASM Guideline Task Force met at special sessions at the 2014 and 2015 SASM annual meeting, as well as during multiple teleconferences. The preliminary results were presented and discussions regarding expert member recommendations took place to consider the implications of their findings.

RECOMMENDATIONS

As discussed previously, recommendations were graded according to the GRADE Working Group^{45,46} (Table 2). The process of building recommendations included not only quality of evidence (confidence in effect estimates) but also the balance between benefit and harm of interventions, the values and preferences of patients, and the use of resources.^{47,48} The recommendations were based on data obtained from the application of general principles of safe

Table 2. Significance of the 4 Levels of Evidence

Quality Level	Definitions
High	Very confident that the true effect lies close to that of the estimate of the effect (eg, RCT with consistent results or observational studies with very large effect size)
Moderate	Moderately confident in the effect estimate: the true effect is likely to be close to the estimate of the effect, but there is a possibility that it is substantially different (eg, RCT with methodological limitations or observational studies with large effect size)
Low	Confidence in the effect estimate is limited: The true effect may be substantially different from the estimate of the effect (eg, RCT with very serious limitation or observational studies without exceptional strength)
Very low	Little confidence in the effect estimate: The true effects are likely to be substantially different from the estimate of effect (eg, case reports and case series)

Abbreviation: RCT, randomized controlled trial.

Modified with permission from Balshem et al.⁴⁶

perioperative care. The benefits and harms of interventions and clinical practice information were considered to ensure that the recommendations preserved patient safety, clinical validity, and usefulness.

The recommendation strength was classified as either “strong” or “weak.”^{47,48} Recommendations were made to administer or not administer an intervention based on trade-offs between benefits on the one hand and downsides (harms, burden, and cost) on the other.⁴⁹ A strong recommendation was given to administer an intervention or offer a treatment if the desirable effects of an outcome or an intervention clearly outweigh the undesirable effects. Conversely, if downsides outweigh benefits, the guideline will recommend against the implementation of such a treatment.⁴⁹ If the overall effects were uncertain, a weak recommendation, also known as conditional, discretionary or qualified recommendation was given. Essentially, a weak recommendation means that the desirable effects of adherence probably outweigh the undesirable effects. In addition, the weak assessment rating was assigned if the evidence was of low quality or the evidence indicated that both desirable and undesirable effects were evenly present. Both strong and weak recommendations have a direction: either for or against. In some instances, the evidence is low because of an absence of published literature, but there is strong support for the statement by consensus of the experts. By use of the GRADE approach, this potential dichotomy is presented for the reader to review and interpret.

Recommendations may be conditional on patient values and preferences, the resources available, or the setting in which the interventions will be implemented.^{47,48} Because resources are valued differently in the different parts of the world, it is not feasible to make judgments about the appropriateness of resource use in this guideline.

Recommendations may be used at the discretion of the patient and clinician or qualified health care professional with an explanation about the issues that would lead decisions to vary. The evidence categories were established upon the level of evidence and agreement among the members of the SASM guideline group (Table 3).^{47–49} In case of disagreement on the recommendations among the members of the SASM task force, the Delphi method was used to collect comments from individuals on their recommendations. Final decisions were made by SASM task force or the results of Delphi methods on those specific questions were discussed.⁵⁰

1. PREOPERATIVE DIAGNOSIS OF OSA

1.1. Question: Does a Diagnosis of OSA Change Postoperative Outcomes?

Recommendation 1.1.1: Patients With a Diagnosis of OSA Should Be Considered to Be at Increased Risk for Perioperative Complications (Level of Evidence: Moderate. Grade of Recommendation: Strong for)

The notion that OSA negatively affects postoperative outcomes is held widely among perioperative physicians; however, prospective, controlled studies with sufficient power to support this assumption are rare, and randomization may not be feasible. In recent years, 2 independent meta-analyses have evaluated studies comparing outcomes in patients with and without a diagnosis of OSA and concluded that

Table 3. Going From Evidence to Recommendations

Strong recommendation	All or almost all informed people would make the recommended choice for or against an intervention. <ul style="list-style-type: none"> • Uniformity of choice • Clinicians spend less time on decision-making process • Focus efforts on overcoming barriers to implementation or adherence • For patients: Most individuals in this situation would want the recommended course of action and only a small proportion would not. • For clinicians: Most patients with the disease should receive the intervention. • For policy makers: The recommendations can be adapted as policy in most situations.
Weak recommendation (discretionary)	Most informed people would choose the recommended course of action, but a substantial number would not. <ul style="list-style-type: none"> • Variability of choice • Clinicians need to devote time to the decision-making process • Ensure patients understand the implications of their choices • For patients: The majority of individuals in this situation would want the suggested course of action, but many would not. • For clinicians: Decision aids may be useful in helping individuals make decisions consistent with their value and preference • For policy makers: Policy making will require substantial debates and involvement of many stakeholders.

Modified with permission from Andrews et al^{47,48} and Schünemann et al.⁴⁹

such a diagnosis is associated with increased risk of postoperative complications.^{24,25} The cardiopulmonary adverse events were increased by 2- to 3-fold.^{24,25} Nevertheless, these analyses used fairly strict inclusion criteria, resulting in the inclusion of 13 and 17 studies only, respectively. Additional literature available on the topic was not considered in these meta-analyses, including recent analyses generated from data collected in large national databases on millions of patients.^{26–29,51} Although the accuracy of studies that use *International Classification of Diseases, Ninth Revision (ICD-9)* coding to identify patients with OSA in Canada has been recently criticized, the point of the critique might not be applicable to other databases, and separate validation would be necessary to verify sensitivity and specificity.⁵² Moreover, potential deficiencies of coding systems in OSA identification would label a mixture of OSA and non-OSA patients with the diagnosis of OSA, and potential misclassification would bias results to the null. Considering the fact that patients with ICD-9 code of OSA are likely to have the disease and many who remain undiagnosed are included in the control group as is likely the case in database studies, the associations found in these analyses could actually underestimate the true effects.

The SASM Task Force on preoperative screening and assessment of patients with OSA sought to perform a comprehensive evaluation of the available information to

Table 4. Studies Evaluating the Association of OSA and Postoperative Complications, Resource Utilization, and Mortality in Cases Utilizing General or Neuraxial Anesthesia

Impact of OSA on Outcomes	No. of Studies	OSA (n)	Non-OSA (n)	Detrimental Impact of OSA (Studies)	Beneficial Impact of OSA (Studies)	No Significant Impact of OSA (Studies)	GRADE Quality of Evidence
Pulmonary complications	17	274 832	7 765 190	11 ^{26,27,29,57-64}	0	6 ^{28,65-69}	Low
Desaturation	14	1993	4898	8 ^{22,23,59,63,70-73}	0	6 ^{58,74-78}	Moderate
Difficult intubation	6	29779	202 698	4 ⁷⁹⁻⁸²	1 ⁸³	1 ⁸⁴	Moderate
Cardiac complications	11	68 543	770 121	2 ^{27,64}	0	9 ^{57-61,63,65,67,72}	Low
Atrial fibrillation	6	88 636	1 065 165	5 ^{28,29,85-87}	0	1 ⁷²	Low
Combined complications	11	52 561	565 842	9 [*]	0	2 ^{68,90}	Low
Resource utilization	19	118 597	934 986	11 [†]	2 ^{28,65}	6 ^{67,72,82,94-96}	Moderate
Other outcomes	4	36 339	192 896	1 ⁷⁴	2 ^{97,98}	1 ⁹⁹	Low
Mortality	13	202 447	2 084 951	1 ⁵⁷	3 ^{28,29,56}	9 ^{9,27,60,65-68,72,94}	Very low

Abbreviations: GRADE, Grading of Recommendations, Assessment, Development, and Evaluation; OSA, obstructive sleep apnea.

formulate an opinion on the question: "Does the diagnosis of OSA change postoperative outcomes?" This is of utmost importance because it forms the basis for clinical decision making regarding OSA in the perioperative setting and allows for the allocation of appropriate resources toward the prevention of adverse outcomes. Studies were reviewed irrespective of OSA definition and included those basing identification using ICD-9 coding, polysomnography, and chart diagnosis. Analyses that used data from population-based administrative databases also were included because these reported on large numbers of patients and are powered to investigate rare outcomes. No meta-analysis was performed, however, as the pooling of data from database studies has the possibility of case-overlap with other such investigations. The systematic review on the impact of OSA on postoperative outcomes was published by the group separately.⁵³ It should be mentioned that additional studies have been published on the topic after the literature search was performed, further supporting the concept of OSA as a perioperative risk factor for complications.^{54,55}

The literature on postoperative complications consists primarily of observational studies, and the majority of publications support the notion that the presence of OSA negatively influences perioperative outcomes. OSA may negatively affect respiratory outcomes and may be linked to postoperative cardiovascular events, such as the development of atrial fibrillation.²⁹ Although speculative, commonly found comorbid conditions such as pulmonary hypertension and heart disease may contribute to these findings, suggesting that cardiac in addition to respiratory monitoring for selected patients with OSA may be considered.

The literature on the association between OSA and perioperative mortality is less clear. Current data suggest an association between OSA and lower 30-day mortality. The studies reporting a reduced association of the perioperative mortality with OSA are based on population-based database analyses with residual confounding, dependence on ICD-9 diagnosis of OSA, and an inability to determine causality.^{28,29,56} It is therefore plausible that increased vigilance afforded to patients with OSA by care providers may reduce fatal outcomes as a result of complications, without influencing the latter per se. In view of ethical considerations, it is now difficult to perform randomized controlled trials in patients with OSA to determine its associations with postoperative complications, particularly in patients with severe OSA. Because of the large number of patients

and the large number of trials, the overall quality of the body of evidence was considered to be moderate using the GRADE approach.⁴⁶

Summary of the Literature Review

Implementing the described search strategy, the initial search yielded 5617 references (S2). After title screening, 510 abstracts were identified and reviewed, and 449 were excluded. Finally, 63 pertinent studies that addressed the issue of perioperative complications in patients with OSA were identified. A summary of the effects of OSA on various outcome categories can be found in Tables 4 and 5.

A total of 52 studies were identified that reported on the association of OSA with select perioperative outcomes for surgeries under general or neuraxial anesthesia. Eleven studies based the diagnosis of OSA on ICD-9 coding,[‡] 18 used polysomnography,[§] 6 used chart diagnoses,^{80,81,83,89,92,98} 16 used screening questionnaires,^{||} and 1 study⁷³ referred to "clinical diagnosis."

In total, the included studies reported on 413 576 OSA patients (diagnosed by ICD-9 coding, polysomnography, chart or clinical diagnoses, and screening questionnaires) and 8 557 044 control (non-OSA) patients. In summary, 17 studies reported on pulmonary complications,^{26-29,57-69} 14 on oxygen desaturation events,[¶] 6 on difficult intubation,⁷⁹⁻⁸⁴ 11 on cardiac complications,[#] 6 on atrial fibrillation,^{28,29,72,85-87} 11 on varying composite outcomes of minor/major complications,^{**} 19 on the utilization of resources,^{††} 4 on various other outcomes,^{74,97-99} and 13 on the outcome of mortality.^{‡‡} In only 7 studies was OSA associated with a beneficial effect for one of the outcomes studied.^{§§} The level of evidence to judge the impact of OSA on various outcomes ranged from very low to moderate.⁴⁶

*References 27, 51, 58-60, 62, 70, 88, 89.

†References 27, 51, 57-59, 70, 78, 89, 91-93.

‡References 26-29, 51, 56-58, 65, 91, 97.

§References 22, 23, 59-62, 66-68, 74, 75, 79, 84, 85, 88, 94-96.

||References 63, 64, 69-72, 76-78, 82, 86, 87, 90, 93, 99, 111.

¶References 22, 23, 58, 59, 63, 70-78.

#References 27, 57-61, 63-65, 67, 72.

**References 27, 51, 58-60, 62, 68, 70, 88-90.

††References 27, 28, 51, 57-59, 65, 67, 70, 72, 78, 82, 89, 91-96.

‡‡References 27-29, 56, 57, 60, 65-68, 72, 94, 111.

§§References 28, 29, 56, 65, 83, 97, 98.

Table 5. Studies Evaluating the Association of OSA and Postoperative Complications, Resource Utilization, and Mortality in Cases Utilizing Sedation Only

Impact of OSA on Outcomes	No. of Studies	OSA (n)	Non-OSA (n)	Detrimental Impact of OSA	Beneficial Impact of OSA	No Significant Impact of OSA	GRADE Quality of Evidence
Oxygen desaturation	7	610	713	3 studies ¹⁰⁰⁻¹⁰²	0 studies	4 studies ¹⁰³⁻¹⁰⁶	Moderate
Airway maneuvers	3	263	297	1 study ¹⁰⁰	0 studies	2 studies ^{107,108}	Very low
Combined complications	3	978	808	0 studies	0 studies	3 studies ¹⁰⁸⁻¹¹⁰	Low

Abbreviations: GRADE, Grading of Recommendations, Assessment, Development, and Evaluation; OSA, obstructive sleep apnea.

The majority of the studies reported worse outcomes among patients with OSA compared with the control group, 11 of 17 reporting on pulmonary complications,^{26,27,29,57-64} 8 of 14 reporting on oxygen desaturation events,^{22,23,59,63,70-73} 4 of 6 reporting on difficult intubation,⁷⁹⁻⁸² 2 of 11 reporting on cardiac complications,^{27,64} 5 of 6 reporting on atrial fibrillation,^{28,29,85-87} 9 of 11 on the composite outcome of various complications,¹¹¹ and 11 of 19 reporting on resource utilization.¹¹¹ Regarding mortality, 9 of 13 studies reported no association,^{##} 3 reported lower mortality in the OSA group,^{28,29,56} and 1 study reported an increase in mortality among OSA patients.⁵⁷ The only study reporting increased mortality was a population-based database study, which found an association between diagnosis of OSA by ICD-9 code and increased mortality in patients undergoing revision knee or hip arthroplasties.⁵⁷ Detailed distribution of the number of studies with adverse, beneficial, or no association with OSA and various perioperative outcomes is summarized in Table 4.

Eleven studies reported on outcomes in patients with OSA undergoing procedures requiring sedation.¹⁰⁰⁻¹¹⁰ Three defined the presence of OSA based on polysomnography,^{104,105,109} 2 on chart diagnosis,^{101,103} and 6 on screening questionnaires.^{100,102,106-108,110} The level of evidence of surveyed studies for various outcomes using the GRADE approach was low to moderate.⁴⁶

To summarize the studies on patients undergoing sedation, 7 studies reported on oxygen desaturation as an outcome,¹⁰⁰⁻¹⁰⁶ 3 on composite minor/major complications,¹⁰⁸⁻¹¹⁰ and 3 on airway maneuvers such as chin lifts, head repositioning, increased oxygen flow by nasopharyngeal or oropharyngeal cannula, or mask airway^{100,107,108} (Table 5). Three studies reported an adverse association of OSA with oxygen desaturation,¹⁰⁰⁻¹⁰² and 1 study reported an adverse association of OSA with necessary airway maneuvers.¹⁰⁰ No study reported better outcomes among OSA patients. Overall, the presence of OSA negatively influences perioperative outcomes in patients undergoing either general or neuraxial anesthesia or sedation.

2. PREOPERATIVE ASSESSMENT FOR OSA

2.1. Question: Should Adult Patients at Risk for OSA Be Identified Before Surgery?

Recommendation 2.1.1: Adult Patients at Risk for OSA Should Be Identified Before Surgery (Level of Evidence: Low, Grade of Recommendation: Weak for)

Routine preoperative screening for OSA in patients presenting for surgery may identify the majority of the patients

with OSA and may provide opportunities for heightened awareness and potential risk reduction by implementing appropriate preoperative,^{61,112-114} intraoperative, and postoperative interventions.^{12,40-44,63,115} The prevalence of patients at risk of OSA (STOP-Bang ≥ 3) in the surgical population is high.^{64,100,108,116,117} When compared with patients with STOP-Bang scores of 0 to 2 (low-risk OSA), surgical patients with STOP-Bang scores of ≥ 3 had an increased rate of perioperative complications.^{***} may justify the use of a screening tool to risk stratify the high-risk OSA patients. A screening tool may help the perioperative team to establish a strategy to reduce the risk of perioperative complications.¹²¹ Practice groups should consider making OSA screening part of standard preanesthetic evaluation.

Although patients identified to be at high risk for OSA have been shown to have increased perioperative complications,^{54,55,62,64,90,93} only a small proportion of them will be affected, because the modest specificity of current screening tools means many patients identified as "at risk" for OSA may not have the disease. Second, not all that do will be at increased risk beyond emergence from anesthesia because of the low-risk nature of their procedure. Third, the OSA phenotype and the severity of OSA may play an additional role. Although the ultimate goal is to minimize risk of postoperative complication as much as feasible by ensuring that every patient with suspected OSA is identified, it is clear that such an approach will result in a challenging logistical, economic, and clinical burden for healthcare providers. Hence, a balance has to be struck between the desire to minimize postoperative complications and the responsible use of resources. The realistic goal is to risk stratify those at particular risk and suggest methods to prevent or treat problems without creating undue economic burden on the health care system.

There is some supporting evidence to suggest that identifying OSA in certain patient populations, such as morbidly obese patients undergoing bariatric surgery or obstetric patients with high-risk pregnancies, may be useful.

Given the adverse maternal,¹²²⁻¹²⁴ fetal,¹²⁵ and perioperative complications associated with OSA, prompt diagnosis and treatment of OSA is critical in the obstetric population.¹²⁶ Although not all obstetric patients are at high risk for OSA, several prospective studies suggest that pregnant women with preeclampsia,^{127,128} gestational hypertension,¹²⁹ gestational diabetes,¹³⁰ asthma,¹³¹ body mass index (BMI) ≥ 35 kg/m², greater prepregnancy BMI, and excessive weight gain during pregnancy place obstetric patients at high-risk for OSA.¹³²⁻¹³⁵ Unfortunately, screening tools appear less accurate and have not been validated in the obstetric

|||References 27, 51, 58-60, 62, 70, 88, 89.

¶¶References 27, 51, 57-59, 70, 78, 89, 91-93.

##References 27, 60, 66-68, 72, 91, 94, 111.

***References 54, 55, 62, 64, 78, 90, 93, 100, 102, 108, 111, 118-120.

population,¹³⁶ and currently, there are no pregnancy-specific guidelines for the diagnosis and treatment of OSA. With the lack of evidence, the best option is to extrapolate the perioperative evaluation and management approach recommended for other patients with diagnosed or suspected OSA to the obstetric population.

Fifty percent of the patients with severe obesity (BMI ≥ 40 kg/m²) have been found to have OSA, and 10% to 20% of these have obesity hypoventilation syndrome.¹³⁷ Patients with a BMI ≥ 40 kg/m² have increased postoperative mortality and morbidity, especially from thromboembolic, infectious, and surgical complications.¹³⁸ The risk further increases with the presence of OSA.¹³⁹ Those at such substantial risk may be first identified by preoperative screening, using a questionnaire such as STOP-Bang.¹¹⁶ The STOP-Bang score has been validated in both obese and morbidly obese surgical patients. A STOP-Bang score of 4 has a high sensitivity of 88% for identifying patients with severe OSA, whereas increasing the cutoff threshold to 6 increases the specificity for severe OSA (although with a lower sensitivity).¹⁴⁰ Steps to identify those likely to have OSA among morbidly obese patients are becoming accepted practice before bariatric surgery.

It is important to note that, at present, there is limited evidence supporting the use of preoperative screening tools for OSA as a practice to reduce patient complications.¹¹¹ However, the expert recommendation for preoperative screening reflects a growing consensus that identifying patients at high risk for OSA before surgery for targeted perioperative precautions and interventions may help to reduce patient complications. For example, there is evidence supporting the avoidance of general anesthesia in patients with OSA who are undergoing specific procedures such as joint arthroplasty.¹⁴¹ An important public health implication of a preoperative OSA screening program is that it could improve the long-term health of patients through better OSA treatment and a reduction of the associated consequences. It is plausible that preoperative diagnosis and optimization of patients with OSA may lead to similar benefits as those achieved by perioperative diagnosis and treatment of other chronic diseases such as coronary artery disease and type 2 diabetes. Undoubtedly, further research in this area is needed, including the development of equally sensitive but more specific screening tools than those that currently exist.

2.2. Question: Which Tools Can Be Used to Identify Surgical Patients With Suspected OSA in the Preoperative Period?

Recommendation 2.2.1: Screening Tools Such As STOP-Bang,¹¹⁶ P-SAP,¹⁴² Berlin,⁶² and ASA Check List⁶² Can Be Used as Preoperative Screening Tools to Identify Patients With Suspected OSA (Level of Evidence: Moderate. Grade of Recommendation: Strong for)

The majority of OSA patients presenting for surgery are undiagnosed and lack sufficient time before surgery to undergo formal diagnostic preoperative polysomnography testing, making screening relevant for perioperative clinical care. Previous systematic reviews, meta-analyses,^{143,144} and guidelines^{12,40-44} have recommended preoperative OSA screening, but with a low level of evidence and a low level of certainty for Ambulatory



Anesthesia guideline,⁴² do not provide recommendations for specific screening tests based on diagnostic accuracy. The choice of screening test is further complicated by the fact that there are many different screening approaches including questionnaires and clinical models with additional screening techniques such as upper airway imaging and overnight oximetry.^{143,144} Furthermore, the majority of studies investigating screening for OSA are designed around higher risk populations (eg, sleep clinic populations), and many are not validated in surgical populations.

The comparative accuracy of different clinical tests for OSA has been reviewed extensively.^{143,144} Specific screening tools are associated with significantly greater accuracy than others, but there is a large heterogeneity within each individual screening test. This means that any given screening tool will have varying accuracy across different populations and may not have the same accuracy when implemented in clinical practice.

The 2 major considerations for choosing screening tests for OSA are feasibility and reliability. Because many patients presenting for surgery may be screened on the day of or within 1 to 2 days of surgery, and some within a few weeks before surgery, the most feasible tests are questionnaires or simple clinical models. Questionnaires are the most commonly used method and have modest accuracy, whereas clinical models that incorporate symptom review with simple clinical measurements are superior to questionnaires.

Implementing the search strategy on screening, the initial electronic search yielded 9565 references (S3). After screening of titles, abstracts, and manuscripts, 9562 articles were excluded. Three pertinent publications addressing the issue of screening of surgical patients for OSA with validation by laboratory polysomnography were identified.^{62,116,142} The STOP-Bang tool,¹¹⁶ P-SAP score,¹⁴² Berlin questionnaire,⁶² and ASA checklist⁶² are screening tests that were evaluated in the surgical population and found to have comparable accuracy (Table 6).

The STOP-Bang tool is the most validated screening tool in surgical patients^{116,117,140,145-149} and also has been validated in sleep clinic patients¹⁵⁰⁻¹⁵⁹ and the general population¹⁶⁰ to detect patients at high risk of OSA. In the surgical patients, a greater STOP-Bang score is associated with a greater probability of moderate-to-severe OSA.^{116,117,145,146} A combination of a STOP score ≥ 2 + BMI > 35 kg/m² or male sex is associated with a greater risk of OSA.^{145,147} The inverse relationship between sensitivity and specificity at increasing STOP-Bang diagnostic thresholds influences the relative rates of missed diagnoses and wasted resource utilization in diagnosing OSA. It is also important to recognize that the predictive values of the tool will be lower in many preoperative settings that have lower prevalence of OSA. Despite these limitations, the updated STOP-Bang tool adds clinical value to the preoperative assessment as it provides a relatively easy method of dichotomous risk stratification of high or low risk of OSA.^{145,147}

Other screening tools such as the P-SAP score also perform like the STOP-Bang tool when increasing the test threshold, with similar implications for missed diagnoses and resource utilization.¹⁴²

Table 6. Comparison of OSA Screening Tools in Surgical Patients

	STOP-Bang Questionnaire ¹¹⁶ (n = 177)	Berlin Questionnaire ⁶² (n = 177)	ASA Checklist ⁶² (n = 177)	P-SAP Score ¹⁴² (n = 511)
Sensitivity	83.6 (75.8–89.7)	68.9 (59.8–76.9)	72.1 (63.3–79.9)	93.9 (91.8–96.6)
Specificity	56.3 (42.3–69.6)	56.4 (42.3–69.7)	38.2 (25.4–52.3)	32.3 (23.2–46.7)
PPV ^a	81.0 (73.0–87.4)	77.9 (68.8–85.2)	72.1 (63.3–79.9)	10.0 (9.0–24.0)
NPV ^a	60.7 (46.1–74.1)	44.9 (32.9–57.4)	38.2 (25.4–52.3)	99.0 (98.0–99.0)
LR+	1.9 (1.40–2.61)	1.57 (1.17–2.36)	1.16 (0.94–1.51)	1.38 (1.37–1.39)
LR–	0.29 (0.18–0.46)	0.55 (0.39–0.79)	0.73 (0.47–1.13)	0.18 (0.16–0.21)
DOR	6.58 (3.03–14.36)	2.85 (1.48–5.50)	1.59 (0.81–3.13)	7.40 (6.48–8.45)
ROC	0.80	0.69	0.78	0.82

Abbreviations: ASA, American Society of Anesthesiologists; DOR, diagnostic odds ratio; LR+, positive likelihood ratio; LR–, negative likelihood ratio; NPV, negative predictive value; OSA, obstructive sleep apnea; PPV, positive predictive value; ROC, area under receiver operating characteristic curve.

^aPredictive values are highly dependent on the prevalence of OSA, which was 69% in the evaluation of STOP-Bang, Berlin, and ASA checklist, and 7.1% for the P-SAP score.

Local practices should decide about the threshold of screening tests, after considering the implications for missed diagnoses and cost of care. The trade-off between sensitivity and specificity means that at lower thresholds, groups should expect improved sensitivity with potentially increased resource utilization, whereas increasing the threshold will result in loss of sensitivity and increased false-negative rates but improved resource utilization. A higher threshold should be adopted in the population with a lower prevalence of OSA.

Because no single test has perfect sensitivity and specificity, there will be varying proportions of false positives and false negatives with any screening program. It is perhaps relevant to consider that a preoperative 12-lead electrocardiogram has <50% sensitivity for the detection of left ventricular hypertrophy and acute myocardial infarction.^{161,162} More advanced screening tests for OSA with greater prediction accuracy have been validated in the literature,¹⁴³ but their complexity and applicability in the surgical population are yet to be evaluated.

Currently, there is inadequate evidence in the literature to recommend the use of sleep testing such as polysomnography in the preoperative period. A future area in need of clarification regarding preoperative screening for OSA is its timing in context of the surgical date. Although experts acknowledged the challenges involved, there was agreement that the greatest opportunity for preoperative diagnosis and treatment would come only with a more deliberate push toward earlier screening. Primary care providers and surgeons should share responsibility for early identification of OSA in patients who are scheduled for surgery as this would allow more time to optimize preoperative preparation. This could involve clinic-based screening at the initial surgical visit.

2.3. Question: What Is the Clinical Value of Performing Additional Preoperative Tests?

Recommendation 2.3.1: There Is Insufficient Evidence to Support Canceling or Delaying Surgery to Perform More Advanced Screening Techniques or Sleep Testing to Diagnose OSA in Those Patients Identified as Being at High Risk of OSA Preoperatively, Unless There Is Evidence of an Associated Significant or Uncontrolled Systemic Disease or Additional Problems With Ventilation or Gas Exchange (Level of Evidence: Low. Grade of Recommendation: Weak for)

Because screening instruments perform better in patients with greater OSA severity, screen-positive patients with greater threshold values should be assumed to have

moderate-to-severe OSA in the absence of diagnostic polysomnography.^{116,117,145,146} The inclusion of preoperative serum bicarbonate level may improve the predictive accuracy of the screening instrument.¹⁶³ Although AHI is the most commonly used metric of OSA severity,²¹ other parameters such as oxygen desaturation index, or cumulated duration of oxygen desaturation <90% may improve prediction of postoperative complications.¹⁶⁴ Further research in this area will help to better define which patients with diagnosed or suspected OSA are at most risk of postoperative complications.

The aforementioned recommendation does not relate to procedures in which polysomnography is performed as part of the accepted preoperative management, usually because of a high prevalence of OSA (eg, bariatric surgery, tonsillectomy, or upper airway surgery for OSA). Summaries of the different recommendations for screening of surgical patients for OSA are listed in Table 7.

3. BEST PREOPERATIVE PRACTICES IN PATIENTS WHO ARE DIAGNOSED WITH OSA, NONADHERENT WITH PRESCRIBED THERAPY, OR HAVE A HIGH PRETEST PROBABILITY FOR OSA

Because OSA remains undiagnosed in the majority of surgical patients, many patients will be identified as having a high probability for OSA for the first time during the preoperative screening process.^{19,20} In addition, many patients with an established diagnosis of OSA either refuse to use or are poorly adherent with the prescribed therapy.¹⁶⁵ As discussed herein, there are limited data to suggest that preoperative positive airway pressure (PAP) therapy, in the form of CPAP, autotitrated positive airway pressure (APAP), or bilevel positive airway pressure may improve perioperative outcomes.^{61,112–114} Alternative therapies used to treat OSA, such as surgery, oral appliances, negative pressure devices, hypoglossal nerve stimulation, positional therapies, nasal resistive valves, and other treatments have not been studied systematically in the perioperative setting, although some patients may be using these as primary therapy for their OSA.

Ideally, identification of patients with OSA, whether adherent or nonadherent to therapy, and those with suspected OSA should take place well in advance of elective surgery to allow time for potential evaluation and management of OSA preoperatively. In clinical practice, many patients, however, are identified close to the operative time, often just days before or even immediately before surgery. Because of a lack

Table 7. Summary of Recommendations for Screening to Identify Patients With Suspected OSA

Recommendations	Level of Evidence	Grade of Recommendation
1.1.1 Patients with a diagnosis of OSA should be considered to be at increased risk for perioperative complications	Moderate	Strong
2.1.1 Adult patients at risk for OSA should be identified before surgery	Low	Weak
2.2.1 Screening tools such as STOP-Bang, P-SAP, Berlin, and ASA checklist can be used as preoperative screening tools to identify patients with suspected OSA	Moderate	Strong
2.3.1 Insufficient evidence exists to support canceling or delaying surgery to formally diagnose OSA in those patients identified as being at high risk of OSA preoperatively, unless there is evidence of uncontrolled systemic disease or additional problems with ventilation or gas exchange	Low	Weak

Abbreviations: ASA, American Society of Anesthesiologists; OSA, obstructive sleep apnea.

of evidence to guide clinical decision making, considerable uncertainty exists regarding the need for further preoperative interventions. The objective in this section is to provide recommendations for the optimal preoperative practices for the care of diagnosed or suspected OSA patients to reduce perioperative complications. We first examined the evidence of efficacy of CPAP for surgical patients in the perioperative period. The recommendations are divided into 3 parts: (i) surgical patients with diagnosed OSA who are adherent to PAP therapy; (ii) surgical patients with diagnosed OSA who are nonadherent to PAP therapy; and (iii) surgical patients with a high probability for having OSA. It should be noted that because there is limited literature on the best preoperative practices in patients with diagnosed or suspected OSA, some of the recommendations are based on data or evidence that is extrapolated from a nonsurgical setting.

Evidence of Efficacy of CPAP for Surgical Patient in the Perioperative Period

Previous guidelines have suggested that CPAP therapy should be considered preoperatively when OSA is severe intraoperatively; in patients under sedation if they have used CPAP previously; and postoperatively, when feasible, particularly if frequent or severe airway obstruction is evident during postoperative monitoring.^{12,41-44} A trade-off occurs when one considers the balance between potentially improved adverse events versus the increased utilization of health care resources resulting from additional interventions and monitoring.¹⁶⁶ At present, <20% of patients with a diagnosis of OSA receive CPAP therapy, are observed in advanced care settings, or both in actual clinical practice.²⁷

Implementing our search strategy on the use of perioperative CPAP, the initial electronic search identified 1970 articles (S4-7). After screening of the titles, abstracts, and manuscripts, 1964 articles were excluded. Six pertinent studies addressing the issue of surgical patients with OSA with or without CPAP were identified.^{58,66,77,89,167,168} Further citation search yielded 3 more relevant articles.^{55,61,112} The types of studies and the level of evidence are shown in Table 8.^{†††} In the 9 studies analyzed, the total number of OSA patients with either preoperative/postoperative CPAP therapy was 4380, and the total number of diagnosed or suspected OSA patients without preoperative/postoperative CPAP therapy was 3649.^{†††}

In patients with OSA, 9 studies demonstrated that CPAP applied preoperatively and/or postoperatively may have

some beneficial effect on postoperative complications. These studies are mainly of low to moderate quality with 2 RCTs, 2 prospective cohorts, 4 retrospective matched cohorts, and 1 case series (Table 9).^{†††} In 2 studies, CPAP significantly reduced the postoperative AHI compared with the preoperative baseline (preoperative AHI versus postoperative AHI: 37 ± 19 vs 12 ± 16 events/h).¹¹⁴ A recent meta-analysis showed a trend toward significance for reduction in the length of hospital stay of 0.4 days favoring the CPAP group (CPAP versus no CPAP: 4.0 ± 4 vs 4.4 ± 8 days, $P = .05$).¹¹⁴

In addition to the studies noted previously, 2 recent large retrospective studies suggest potential efficacy of CPAP in patients with diagnosed OSA.^{61,112} In the first study, those patients diagnosed with OSA and provided with a prescription for CPAP therapy before their surgery had a significantly reduced risk for cardiovascular adverse events compared with those undiagnosed OSA patients (odds ratio [OR], 0.34; 95% confidence interval, 0.15–0.77; $P = .009$).⁶¹ Patients with severe OSA or undiagnosed OSA were found to have significantly increased risk for cardiorespiratory complications.⁶¹

The second study found that the untreated OSA patients had greater cardiopulmonary complication rates compared with those prescribed CPAP therapy (risk-adjusted rates 6.7% vs 4%; adjusted OR = 1.8, $P = .001$).¹¹² Both myocardial infarction (adjusted OR = 2.6, $P = .031$) and unplanned reintubations (adjusted OR = 2.5, $P = .003$) were significantly greater in untreated OSA patients.¹¹² These 2 large database studies provide preliminary evidence supporting preoperative diagnosis of OSA and treatment with CPAP therapy.^{61,112}

There are 3 studies, 2 prospective and 1 retrospective, on empiric APAP therapy in the perioperative care of patients with newly diagnosed untreated OSA.^{77,165,168} Each study reported a different primary outcome: perioperative AHI versus length of stay versus APAP adherence. None of the studies was sufficiently powered to address the impact of APAP on postoperative complications; and in one study, patients with suspected OSA who were at >10% risk for postoperative cardiac complications as estimated by American College of Physicians criteria were excluded if they had an abnormal overnight oximetry study.⁷⁷ The adherence to APAP therapy in these studies was uniformly low. Approximately 45% of patients with newly diagnosed OSA were adherent to APAP therapy in the perioperative period.^{165,168}

†††References 55, 58, 61, 66, 77, 89, 112, 167, 168.

Table 8. Effect of CPAP on Postoperative Outcomes in Surgical Patients With OSA

Author	Study Type	Level of Evidence	N	Results
Rennotte et al ¹⁶⁷	Case series report Postoperative CPAP	Very Low	16	No CPAP (2) vs CPAP (14) 1st patient died; 2nd patient: serious postoperative Cx 14 patients CPAP Rx, no Cx
Gupta et al ⁸⁹	Retrospective case control study Home CPAP	Low	101	No CPAP (68) vs CPAP group (33) Serious Cx: 31% vs 9% ($P = .02$) Total ICU: 32.3% vs 3% ($P = .001$) Unplanned ICU: 27.9% vs 3% ($P = .003$) Hospital stay: 7 ± 3 vs 6 ± 2 days ($P = .03$)
Jensen et al ⁶⁶	Prospective cohort study Home CPAP	Low	284	No CPAP/BPAP (140) vs CPAP/BPAP (144) Pneumonia: 2.1% vs 0.7% (NS) Hospital stay: 2.2 vs 2.7 days ($P = .24$)
Liao et al ⁵⁸	Retrospective matched cohort study Home CPAP	Low	240	No CPAP (90) vs CPAP (150) Postoperative Cx: 46.6% vs 40.6% ($P = .36$)
Liao et al ¹⁶⁸	Randomized controlled study Preoperative CPAP	Moderate	177	No APAP (90) vs APAP (87) Postoperative Cx: 48.3% vs 48.3% (NS) Preoperative AHI vs postoperative AHI (N3) APAP: 30 to 3.0 ($P < .001$) No APAP: 30 to 32 ($P = .3$) Hospital stay: 4.3 ± 6 vs 3.5 ± 6 days ($P = .36$)
O'Gorman et al ¹⁷⁷	Randomized controlled study Postoperative CPAP	Low	138	No APAP (43) vs APAP (43) Any Cx: 20.9% vs 23.3% ($P = 1.0$) No significant difference between LOS
Mutter et al ⁶¹	Retrospective Matched cohort study CPAP prescription	Moderate	20,488	Respiratory Cx OSA + CPAP (2629) vs undiagnosed OSA (1569) OR 0.68 (95% CI, 0.27–1.71, $P = .41$) Cardiovascular Cx OSA + CPAP (2496) vs undiagnosed OSA (1489) OR 0.34 (95% CI, 0.15–0.77, $P = .009$)
Abdelsattar et al ¹¹²	Prospective cohort study Home CPAP	Moderate	2646	Untreated OSA (1465) vs treated OSA (1181) Cardiopulmonary Cx 6.7% vs 4.0%; aOR = 1.8 ($P = .001$) Unplanned reintubations aOR = 2.5 ($P = .003$) Myocardial infarction aOR = 2.6 ($P = .03$)
Proczko et al ⁵⁵	Retrospective cohort Home CPAP	Low	693	OSA + home CPAP (99) vs STOP-Bang ≥ 3 (182) Death: 0% vs 1% ($P = .5$) Hospital stay: 3.2 vs 4.1 days ($P < .0001$) Pneumonia: 2% vs 9.3% ($P < .04$)

Abbreviations: AHI, apnea-hypopnea index; aOR, adjusted odds ratio; APAP, autotitrated positive airway pressure; BPAP, bilevel PAP; CI, confidence interval; CPAP, continuous positive airway pressure; Cx, complications; ICU, intensive care unit; LOS, length of hospital stay; N-CPAP, nasal continuous positive airway pressure; NS, not significant; OSA, obstructive sleep apnea; Rx, treated.

From Chung et al.¹¹³

3.1. Question: What Are the Best Preoperative Practices Aimed at Improving Outcomes for Surgical Patients With Diagnosed OSA Who Are Adherent to PAP Therapy?

Recommendation 3.1.1: The Patient, Surgeon, Anesthesiologist, and the Health Care Team Should Be Aware Before the Procedure That the Patient Carries a Diagnosis of OSA, Which May Increase Morbidity Associated With Surgery (Level of Evidence: Low. Grade of Recommendation: Strong for)

The consensus of the SASM guideline committee^{24–29,58} that all perioperative providers, and the patient, should be aware that the patient has a known diagnosis of OSA, which could potentially adversely affect his or her clinical course. It should be noted if the patient is on PAP or alternative therapy at home. Patients with OSA who have had surgery as a treatment for OSA, as well as those who have been treated with an oral appliance or nasal resistive valves, but have not had follow-up sleep testing to confirm resolution or control of the OSA, should be regarded as having a high

probability of having untreated OSA. Identifying these patients preoperatively will allow for the implementation of specific practices and protocols aimed at minimizing perioperative risk.^{12,41–44}

Recommendation 3.1.2: Suggest That Consideration Be Given to Obtaining the Results of the Sleep Study and the Recommended PAP Setting Before Surgery (Level of Evidence: Low. Grade of Recommendation: Weak for)

The utility of knowing the results of the patient's sleep study and their correct PAP setting before surgery has not been studied. The consensus of the committee^{24–29,58} at obtaining results of past sleep studies, as well as documenting the recommended PAP setting, is meaningful for patient care. Reviewing results of the sleep studies can confirm the presence, type, and severity of sleep apnea. This may influence clinical decision making regarding postoperative monitoring and the use of opioids in these patients. Knowing the correct PAP setting guides adjustment of therapy to provide patients with effective upper airway support during sleep

Table 9. Best Preoperative Practices for Surgical Patients With Known OSA, Adherent or Nonadherent to PAP Therapy, or a High Probability of OSA

Recommendations	Level of Evidence	Grade of Recommendation
3.1 Surgical patients with OSA who are adherent to PAP therapy		
3.1.1 The patient and the health care team should be aware that a diagnosis of OSA may be associated with increased postoperative morbidity	Low	Strong
3.1.2 Consideration should be given in obtaining the results of the sleep study and the recommended PAP setting before surgery	Low	Weak
3.1.3 Facilities should consider having PAP equipment available for perioperative use or have the patient bring their own PAP equipment to the surgical facility	Low	Strong
3.1.4 Patients should continue to wear their PAP device at appropriate times during their stay in the hospital, both preoperatively and postoperatively	Moderate	Strong
3.2 Surgical patients with OSA but decline or are poorly adherent to PAP therapy		
3.2.1 The patient and the health care team should be aware that untreated OSA may be associated with increased postoperative morbidity	Low	Strong
3.2.2 Consideration should be given to obtaining the results of the sleep study and the recommended PAP setting before surgery	Low	Weak
3.2.3 Facilities should have PAP equipment for perioperative use or have the patient bring their own PAP equipment with them to the surgical facility	Low	Strong
3.2.4 Additional evaluation for preoperative cardiopulmonary optimization should be considered in patients with known OSA who are nonadherent or poorly adherent to PAP therapy and have uncontrolled systemic conditions or additional problems with ventilation or gas exchange such as: (i) hypoventilation syndromes, (ii) severe pulmonary hypertension, and (iii) resting hypoxemia in the absence of other cardiopulmonary disease	Low	Weak
3.2.5 Untreated OSA patients with optimized comorbid conditions may proceed to surgery if provided strategies for mitigation of postoperative complications are implemented. The risks and benefits of the decision should include consultation with the surgeon and the patient	Low	Weak
3.2.6 Patients should be encouraged to wear their PAP device at appropriate times during their stay in the hospital, both preoperatively and postoperatively	Moderate	Strong
3.3 Surgical patients who have a high probability for OSA		
3.3.1 The patient and the health care team should be aware that a high probability of OSA may increase postoperative morbidity	Low	Strong
3.3.2 Additional evaluation for preoperative cardiopulmonary optimization should be considered in patients who have a high probability of having OSA and have uncontrolled systemic conditions or additional problems with ventilation or gas exchange such as: (i) hypoventilation syndromes, (ii) severe pulmonary hypertension, and (iii) resting hypoxemia in the absence of other cardiopulmonary disease	Low	Weak
3.3.3 Patients who have a high probability of having OSA may proceed to surgery in the same manner as those with a confirmed diagnosis, provided strategies for mitigation of postoperative complications are implemented. Alternatively, they may be referred for further evaluation and treatment. The risks and benefits of the decision should include consultation with the surgeon and the patient	Low	Weak
3.3.4 Patients should be advised to notify their primary medical provider that they were found to have a high probability of having OSA, thus allowing for appropriate referral for further evaluation	Low	Weak

Abbreviations: OSA, obstructive sleep apnea; PAP, positive airway pressure.

while recovering from surgery. In addition, data obtained from the patient's PAP compliance card may help provide objective evidence of the use and efficacy of their therapy.

Recommendation 3.1.3: We Suggest That Facilities Should Consider Having PAP Equipment for Perioperative Use Available or Have the Patient Bring Their Own PAP Equipment to the Surgical Facility (Level of Evidence: Low. Grade of Recommendation: Strong for)

To provide appropriate PAP therapy, either the facility at which the surgery is performed or the patients themselves will need to provide the proper equipment. Local institutional policies may determine how this matter is to be handled as logistical capabilities and expertise may vary.

Recommendation 3.1.4: Patients Should Continue to Wear Their PAP Device at Appropriate Times During Their Stay in the Hospital, Both Preoperatively and Postoperatively (Level of Evidence: Moderate. Grade of Recommendation: Strong for)

There is a limited-but-growing body of literature suggesting that PAP therapy for OSA patients may play a role in

attenuating postoperative complications.^{61,112-114} In addition, the acute withdrawal of PAP therapy in patients adherent to treatment has been shown to result in recurrence of OSA and OSA-related symptoms within 1 to 3 days and physiologic derangements within 2 weeks.^{31,169} We recommend the continued use of PAP therapy at previously prescribed settings during periods of sleep while hospitalized. Adjustments may need to be made to the settings to account for perioperative changes, such as facial swelling, upper airway edema, fluid shifts,¹⁷⁰⁻¹⁷² pharmacotherapy, and respiratory function. It should be noted that CPAP therapy is not an alternative to appropriate monitoring.

Patients using alternative therapies for OSA, such as oral appliances, body positioners, nasal resistive valves, oral negative pressure devices, and hypoglossal nerve stimulators, should be encouraged to continue use of their therapy in the perioperative setting. Additional consultation with the patient's sleep specialist may be considered if there are questions regarding the effectiveness, settings, or use of these devices while in the postoperative care setting.

3.2. Question: What Are the Best Preoperative Practices Aimed at Improving Outcomes for Surgical Patients With Diagnosed OSA but Decline or Are Poorly Adherent to PAP Therapy?

Recommendations 3.2.1 to 3.2.3 Are Identical to Recommendations 3.1.1 to 3.1.3

Recommendation 3.2.4: We Suggest That Additional Evaluation for Preoperative Cardiopulmonary Optimization Be Considered in Patients Who Have a Known Diagnosis of OSA and Are Nonadherent or Poorly Adherent to PAP Therapy and Where There Is Indication of Uncontrolled Systemic Conditions or Additional Problems With Ventilation or Gas Exchange. These Conditions Include, But May Not Be Limited to (i) Hypoventilation Syndromes, (ii) Severe Pulmonary Hypertension, and (iii) Resting Hypoxemia Not Attributable to Other Cardiopulmonary Disease (Level of Evidence: Low. Grade of Recommendation: Weak for):

The committee recognizes that there is a paucity of data to support delaying surgery for patients with untreated OSA. However, patients with certain comorbidities, such as clinically significant hypoventilation¹⁷³ or pulmonary hypertension,^{174,175} may be more likely to develop postoperative complications, and these conditions commonly coexist with OSA. Resting hypoxemia in the absence of other known cardiopulmonary disease should also warrant consideration for further preoperative evaluation.¹⁶⁴ A combination of OSA and acute or chronic cardiopulmonary diseases needs to be considered in such a scenario. When untreated OSA is present, benefit may exist to initiating and optimizing PAP therapy to appropriately treat OSA while simultaneously enhancing postoperative respiratory function.^{61,112–114,176}

Recommendation 3.2.5: We Suggest That Untreated OSA Patients With Optimized Comorbid Conditions May Proceed to Surgery Provided Strategies for Mitigation of Postoperative Complications Are Implemented. The Risks and Benefits of the Decision Should Include Consultation and Discussion With the Surgeon and the Patient (Level of Evidence: Low. Grade of Recommendation: Weak for):

There are multiple factors that need to be considered in this decision-making process, and generalizations lacking supportive evidence are not prudent. Specific comorbidities (as noted in 3.2.4), the urgency of the surgery, the type of surgical procedure, the anticipated need for high-dose postoperative opioids, and the availability of postoperative monitoring for opioid-related adverse events may influence the decision whether to proceed with surgery or to delay for further preoperative evaluation.^{177–180} Ultimately, a decision to proceed or delay surgery needs to be made on an individual basis and should involve relevant perioperative providers, including the surgeon and the patient. Patients should be fully informed of the increased risk of postoperative complications if they have untreated OSA. For example, postoperative respiratory-related adverse events, including reintubation and respiratory failure, may be the result of an imbalance between enhanced pain processes and increased sensitivity to anesthetics and/or opioids in patients with some specific OSA phenotypes.^{180–182}

Evaluation and management of OSA by a practitioner with expertise in sleep medicine has been shown to reduce discontinuation of PAP therapy¹⁸³ and improve CPAP

adherence.^{183–185} Sleep medicine specialists can investigate reasons for refusal of therapy or poor adherence to therapy in patients diagnosed with OSA and implement strategies to improve acceptance and/or adherence to CPAP therapy.

Alternative management strategies, including oral appliances, positional therapy (eg, body positioners, elevating the head of the bed), upper airway surgery, and negative pressure devices be considered in patients unable to use PAP therapy. However, data supporting their use in the perioperative period are lacking, and their implementation in the postoperative setting may be limited.

Recommendation 3.2.6: Patients Should Be Encouraged to Wear Their PAP Device at Appropriate Times During Their Stay in the Hospital, Both Preoperatively and Postoperatively (Level of Evidence: Moderate. Grade of Recommendation: Strong for):

As noted earlier, accumulating evidence suggests that PAP therapy for OSA patients may play a role in reducing postoperative complications.^{61,112–114} We recommend patients be encouraged to use their PAP therapy at their previously prescribed setting during sleep while in the hospital. Simple maneuvers such as refitting a mask, the addition of heated humidification, or the control of nasal congestion with nasal corticosteroid sprays may help patients to adhere to therapy, and consideration should be given to inquiring about these issues in poorly adherent patients.¹⁸⁶ Consultation with the patient's sleep specialist may be considered if there are additional questions regarding optimizing PAP usage in the postoperative care setting.

3.3. Question: What Are the Best Preoperative Practices Aimed at Improving Outcomes for Surgical Patients Who Have a High Probability for OSA?

Recommendation 3.3.1: The Patient, Surgeon, Anesthesiologist, and the Health Care Team Should Be Aware Before the Procedure That the Patient Has a High Probability of Having OSA, Which May Increase the Morbidity Associated With Surgery (Level of Evidence: Low. Grade of Recommendation: Strong for)

The consensus of the SASM guideline committee that all perioperative providers, as well as the patient, should be aware that the patient has a high probability of having OSA, and this could potentially adversely affect their clinical course.^{24–29} Identifying these patients preoperatively will allow for the implementation of specific practices and protocols aimed at minimizing perioperative risk.^{12,41–44}

Recommendation 3.3.2: We Suggest That Additional Evaluation for Preoperative Cardiopulmonary Optimization Be Considered in Patients Who Have a High Probability of Having OSA and Where There Is Indication of Uncontrolled Systemic Conditions or Additional Problems With Ventilation or Gas Exchange. These Conditions Include, but May Not Be Limited to (i) Hypoventilation Syndromes, (ii) Severe Pulmonary Hypertension, and (iii) Resting Hypoxemia Not Attributable to Other Cardiopulmonary Disease (Level of Evidence: Low. Grade of Recommendation: Weak for)

The committee recognizes that there is a paucity of data to support delaying surgery for patients with a high

probability of having OSA. However, as noted in 3.2.3, patients with certain comorbidities, such as clinically significant hypoventilation¹⁷³ or pulmonary hypertension,^{174,175} may be more likely to develop postoperative complications, and these conditions commonly coexist with OSA. Unexplained preoperative hypoxemia should also warrant consideration for further preoperative evaluation.¹⁶⁴

Recommendation 3.3.3: We Suggest That Patients Who Have a High Probability of Having OSA May Proceed to Surgery in the Same Manner as Those With a Confirmed Diagnosis, Provided Strategies for Mitigation of Postoperative Complications Are Implemented. Alternatively, They May Be Referred for Further Evaluation and Treatment. The Risks and Benefits of the Decision Should Include Consultation and Discussion With the Surgeon and the Patient (Level of Evidence: Low. Grade of Recommendation: Weak for)

The preoperative decision-making process can be difficult when considering patients who have a high probability of having OSA as there is little evidence to guide recommendations. Specific comorbidities (as noted in 3.3.2), the urgency of the surgery, the type of surgical procedure, the anticipated need for high-dose postoperative opioids, and the availability of postoperative monitoring for opioid-related adverse events all may influence the decision of whether to proceed with surgery or to delay for further preoperative evaluation.^{177–180} In a retrospective study, using a risk-stratified management protocol, patients with a high probability of OSA were managed safely without formal polysomnography confirmation.¹⁸⁷ Ultimately, the decision to proceed or delay surgery needs to be made on an individual basis and should involve relevant perioperative providers, including the surgeon and the patient. Patients should be fully informed of the increased risk of postoperative complications if they have untreated OSA. There should be a low threshold for the use of postoperative monitoring of oxygenation and ventilation in these patients.^{177–182} Judgment regarding whether to do so may be made in the course of a prolonged postanesthetic care unit stay.⁶³

If additional preoperative sleep testing is pursued, the primary goals should include diagnosing the presence and categorizing the severity of OSA, as well as implementing appropriate therapy. While this can be most comprehensively accomplished by detailed polysomnography, limited availability and cost may limit the feasibility and home sleep monitoring could be a reasonable option for high probability patients.¹⁸³

Recommendation 3.3.4: Patients Should Be Advised to Notify Their Primary Medical Provider That They Were Found to Have a High Probability of Having OSA, Thus Allowing for Appropriate Referral for Further Evaluation (Level of Evidence: Low. Grade of Recommendation: Weak for)

As OSA is associated with numerous poor health outcomes including risk of death, cardiovascular events, and average number of days hospitalized,^{188–190} and treatment of OSA appears to improve these outcomes,^{191,192} it is clinically important to diagnose OSA in those patients screened at high risk and advise these patients to have ongoing, long-term management. Relevant to identifying patients in preoperative screening, a recent 2-year follow-up survey study found

that patients diagnosed with OSA in the preoperative setting who were adherent to CPAP therapy experienced long-term health benefits and a reduction in medication usage.¹⁹³

It should be recognized that questionnaires or other screening processes are imperfect diagnostic tools with low to moderate specificity.^{62,116,142,145} As such they tend to overdiagnose, with a significant proportion of patients labeled as at risk of OSA when they may not have the condition. It is important to differentiate between patients identified as at increased risk of having OSA and those who have been accurately diagnosed with OSA. If a patient is overdiagnosed or misdiagnosed as having OSA, this could have negative consequences in terms of any future surgical procedure, health insurance costs, and the ability to continue to drive. Accurate diagnosis is a necessary precursor to appropriate treatment. Conversely, ruling out OSA in patients who screened as at risk for OSA eliminates many of the negative consequences from overdiagnosis. The committee recommends that patients identified as being at high risk of OSA during preoperative screening be advised to notify their primary care provider of this status so that referral for further evaluation can be discussed. Summaries of the recommendations for optimal preoperative practices in patients with diagnosed or suspected OSA are listed in Table 9.

RESEARCH QUESTIONS

Our careful review of the literature presents the best evidence available regarding preoperative evaluation of patients with diagnosed or suspected OSA. However, there are many clinically relevant questions that remain unanswered. Further work is needed to answer questions such as: (1) How do we better stratify risk among these patients? Are there phenotypic characteristics among OSA patients that identify those at particular risk? Are there better sleep study metrics than AHI to identify perioperative risk? (2) Which surgical patients with diagnosed OSA, partially treated/untreated, or suspected OSA should be referred for specialist preoperative evaluation and management? Under what circumstances should surgery be delayed to allow this? (3) Which patients are likely to require continuous monitoring beyond the immediate recovery period? Can they reliably be identified preoperatively? (4) When is the optimal time to implement PAP therapy perioperatively? (5) What are the best modalities of PAP therapy for perioperative use? (6) What are the obstacles to implementing a perioperative PAP protocol? (7) What are the cost implications and cost-effectiveness of preoperative screening for OSA risk?

SUMMARY

Worldwide, 234 million major surgical procedures are undertaken annually.¹⁹⁴ In view of the increased risk of perioperative complications associated with it, surgical safety for patients with OSA is a substantial concern on a global level. As of July 2015, the Joint Commission, Division of Health Care Improvement, had received 61 sentinel event reports in which a patient was diagnosed with or suspected of having OSA.¹⁹⁵ The Joint Commission cited the following concerns regarding the care of patients in the setting of OSA: (1) lack of training for health care professionals to screen for and recognize OSA; (2) failure to assess patients

for OSA; (3) lack of guidelines for the care and treatment of individuals at risk for and those diagnosed with OSA; (4) failure to implement appropriate monitoring of patients with risk factors associated with OSA; (5) lack of communication among health care providers regarding patients with OSA or potential risk factors associated with OSA; and (6) lack of postoperative evaluation and treatment for OSA.¹⁹⁶ These are all areas waiting for further improvement.

The primary goal of this SASM guideline is to ensure optimal preoperative evaluation of patients with diagnosed or suspected OSA to improve patient safety. There is now firm evidence that these patients are at increased risk of postoperative complications. Preoperative evaluation, best done in an anesthesia preoperative clinic, enhances anesthetic management of chronic medical conditions such as OSA through documentation of health status and, where appropriate, initiation of investigations to pursue diagnostic possibilities raised in the screening process including referral to appropriate subspecialists.¹⁹³ Primary care providers and surgeons require education regarding the increased risk of postoperative complications present in diagnosed or undiagnosed OSA patients. They should be alert to the possibility of OSA in their patients and attempt to identify it early in the surgical planning process to allow optimal preoperative preparation and risk stratification.

OSA is a heterogeneous disorder with multiple causes and different phenotypes.¹⁹⁶ The clinical practice of sleep medicine is changing rapidly with wider availability of simple home-based diagnostics and more readily acceptable treatments.¹⁹⁷ There is some evidence supporting the desirability of preoperative diagnosis of OSA and treatment with CPAP therapy.^{61,112–114} The time between the decision to perform surgery and the date of the surgery should allow for optimization of patient health, including identification and management of OSA where it exists. With proper training in sleep medicine, anesthesiologists may be able to diagnose and treat OSA in the perioperative setting.¹⁹⁸ The overall costs of health care could be reduced by treating sleep apnea as the number of hospital admissions and their costs have been shown to be lower in OSA patients on CPAP versus no therapy.^{199,200}

At present, we believe that there is insufficient evidence to support canceling or delaying surgery to perform sleep studies and to initiate PAP therapy in those patients identified as being at high risk of OSA preoperatively, unless there is evidence of significant or uncontrolled systemic disease. Additional subspecialty evaluation for preoperative cardiopulmonary optimization should be considered in patients who have a known diagnosis of OSA, are nonadherent to PAP therapy, and have uncontrolled systemic conditions such as hypoventilation,¹⁷³ severe pulmonary hypertension,^{174,175} or resting hypoxemia in the absence of other known cardiopulmonary disease.¹⁶⁴ These concerns may also apply to patients with suspected OSA who have these problems. These considerations make a case for fast track consultation and PAP therapy services, facilitating access to diagnosis and treatment for such patients.

Regardless of whether the diagnosis of OSA was established before surgery or not, anesthesiologists, surgeons, and institutions should develop an institutional protocol

for patients with known or suspected OSA, including type of anesthesia,²⁰¹ choice of medications,^{202,203} postoperative analgesic regimens,^{179–182} monitoring,^{177,178} and appropriate preoperative/postoperative referral to reduce complications and ensure the best possible patient outcome.^{204,205}

RECOMMENDATIONS: EXECUTIVE SUMMARY

- Patients with obstructive sleep apnea (OSA) undergoing procedures under anesthesia are at increased risk for perioperative complications compared with patients without the disease diagnosis. Identifying patients at high risk for OSA before surgery for targeted perioperative precautions and interventions may help to reduce perioperative patient complications.
- Screening tools help to risk stratify patients with suspected OSA with reasonable accuracy. Practice groups should consider making OSA screening part of standard preanesthetic evaluation.
- There is insufficient evidence in the current literature to support canceling or delaying surgery for a formal diagnosis (laboratory or home polysomnography) in patients with suspected OSA, unless there is evidence of an associated significant or uncontrolled systemic disease or additional problems with ventilation or gas exchange.
- The patient and the health care team should be aware that both diagnosed OSA (whether treated, partially treated, or untreated) and suspected OSA may be associated with increased postoperative morbidity.
- If available, consideration should be given to obtaining the results of the sleep study and, where applicable, the patient's recommended positive airway pressure (PAP) setting before surgery.
- If resources allow, facilities should consider having PAP equipment for perioperative use or have patients bring their own PAP equipment with them to the surgical facility.
- Additional evaluation to allow preoperative cardiopulmonary optimization should be considered in patients with diagnosed, partially treated/untreated, and suspected OSA where there is indication of an associated significant or uncontrolled systemic disease or additional problems with ventilation or gas exchange such as: (i) hypoventilation syndromes, (ii) severe pulmonary hypertension, and (iii) resting hypoxemia in the absence of other cardiopulmonary disease.
- Where management of comorbid conditions has been optimized, patients with diagnosed, partially treated, untreated OSA, or suspected OSA may proceed to surgery provided strategies for mitigation of postoperative complications are implemented.
- The risks and benefits of the decision should include consultation and discussion with the surgeon and the patient.
- The use of PAP therapy in previously undiagnosed but suspected OSA patients should be considered case by

case. Because of the lack of evidence from randomized controlled trials, we cannot recommend its routine use.

- Continued use of PAP therapy at previously prescribed settings is recommended during periods of sleep while hospitalized, both preoperatively and postoperatively. Adjustments may need to be made to the settings to account for perioperative changes such as facial swelling, upper airway edema, fluid shifts, pharmacotherapy, and respiratory function.

ACKNOWLEDGMENT

The SASM task force is divided into 3 groups to answer the key questions.

Does a diagnosis of OSA change postoperative outcomes?
Dennis Auckley, Crispiana Cozowicz, Frances Chung, Roop Kaw, Stavros Memtsoudis (leader), Babak Mokhlesi, and Mathias Opperer.

Preoperative assessment: How do we screen for high risk of OSA in the preoperative period?

Dennis Auckley, Frances Chung (co-leader), Anthony G. Doufas, Matthias Eikermann, Bhargavi Gali, Girish Joshi, Atul Malhotra, Mahesh Nagappa, Satya Krishna Ramachandran (leader), Sara Patrawala, and Tracey Stierer.

What are the best practices regarding preoperative management of patients with diagnosed OSA on therapy, patients with diagnosed OSA who decline or are nonadherent to therapy, and those with a high pretest probability for OSA?

Dennis Auckley (leader), Najib Ayas, Frances Chung (co-leader), Nancy Collop, Anthony G. Doufas, John Fleetham, Peter Gay, Girish P. Joshi, David R. Hillman, Roop Kaw, Eric Kezirian, Babak Mokhlesi, Mahesh Nagappa, Sai Parthasarathy, and Frank Wappler.

Reviewers of literature

Does a diagnosis of obstructive sleep apnea change outcome?
Crispiana Cozowicz, Stavros Memtsoudis, and Mathias Opperer.

Should all patients at risk for OSA be identified before surgery?

Frances Chung, Anjana Kumar, Mahesh Nagappa, Sara Patrawala, and Satya Krishna Ramachandran.

Preoperative preparation and efficacy of CPAP for surgical patient in the perioperative period

Dennis Auckley, Frances Chung, David Lam, and Mahesh Nagappa.

DISCLOSURES

Name: Frances Chung, MBBS, FRCPC.

Contribution: This author helped design the study, conduct the study, analyze the data, write the manuscript, and attests to the integrity of the original data and has approved the final manuscript. Dr. Chung is the archival author.

Attestation: Frances Chung approved the final manuscript.

Conflicts of interest: STOP-Bang tool is proprietary to University Health Network, Royalties from UpToDate, research grants from ResMed Foundation and Pfizer.

Name: Stavros Memtsoudis, MD, PhD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Stavros Memtsoudis approved the final manuscript.

Conflicts of interest: Stavros Memtsoudis reports no conflicts of interest.

Name: Satya Krishna Ramachandran, MD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Satya Krishna Ramachandran approved the final manuscript.

Conflicts of interest: Satya Krishna Ramachandran funded research from Merck, Sharp and Dohme, NJ.

Name: Mahesh Nagappa, MD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Mahesh Nagappa approved the final manuscript.

Conflicts of interest: Mahesh Nagappa reports no conflicts of interest.

Name: Mathias Opperer, MD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Mathias Opperer approved the final manuscript.

Conflicts of interest: Mathias Opperer reports no conflicts of interest.

Name: Crispiana Cozowicz, MD.

Contribution: This author helped conduct the study, analyze the data, and write the manuscript.

Attestation: Crispiana Cozowicz approved the final manuscript.

Conflicts of interest: Crispiana Cozowicz reports no conflicts of interest.

Name: Sara Patrawala, MD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Sara Patrawala approved the final manuscript.

Conflicts of interest: Sara Patrawala reports no conflicts of interest.

Name: David Lam, BSc.

Contribution: This author helped conduct the study, analyze the data, and write the manuscript.

Attestation: David Lam approved the final manuscript.

Conflicts of interest: David Lam reports no conflicts of interest.

Name: Anjana Kumar, BSc.

Contribution: This author helped conduct the study and analyze the data.

Attestation: Anjana Kumar approved the final manuscript.

Conflicts of interest: Anjana Kumar reports no conflicts of interest.

Name: Girish P. Joshi, MD.

Contribution: This author helped write the manuscript.

Conflicts of interest: Girish P. Joshi received an honorarium from Baxter Pharmaceuticals, Pacira Pharmaceuticals, and Mallinckrodt Pharmaceuticals.

Attestation: Girish P. Joshi approved the final manuscript.

Name: John Fleetham, MD.

Contribution: This author helped write the manuscript.

Attestation: John Fleetham approved the final manuscript.

Conflicts of interest: John Fleetham reports no conflicts of interest.

Name: Najib Ayas, MD.

Contribution: This author helped write the manuscript.

Attestation: Najib Ayas approved the final manuscript.

Conflicts of interest: Najib Ayas reports no conflicts of interest.

Name: Nancy Collop, MD.

Contribution: This author helped write the manuscript.

Attestation: Nancy Collop approved the final manuscript.

Conflicts of interest: Nancy Collop received royalties from UpToDate, and honorariums from Best Doctors and Jazz Pharmaceuticals.

Name: Anthony Doufas, MD, PhD.

Contribution: This author helped write the manuscript.

Attestation: Anthony Doufas approved the final manuscript.

Conflicts of interest: Anthony Doufas reports no conflicts of interest.

Name: Matthias Eikermann, MD, PhD.

Contribution: This author helped write the manuscript.

Attestation: Matthias Eikermann approved the final manuscript.

Conflicts of interest: ResMed Foundation.

Name: Marina Englesakis, HBA, MLIS.

Contribution: This author helped literature search and helped write the manuscript.

Attestation: Marina Englesakis approved the final manuscript.

Conflicts of interest: Marina Englesakis reports no conflicts of interest.

Name: Bhargavi Gali, MD.

Contribution: This author helped write the manuscript.

Attestation: Bhargavi Gali approved the final manuscript.

Conflicts of interest: Bhargavi Gali reports no conflicts of interest.

Name: Peter Gay, MD.

Contribution: This author helped write the manuscript.

Attestation: Peter Gay approved the final manuscript.

Conflicts of interest: Peter Gay reports no conflicts of interest.

Name: Adrian Hernandez, MD, MSc, PhD, FACC.

Contribution: This author helped write the manuscript.

Attestation: Adrian Hernandez approved the final manuscript.

Conflicts of interest: Adrian Hernandez reported no conflicts of interest.

Name: Roop Kaw, MD.

Contribution: This author helped write the manuscript.

Attestation: Roop Kaw approved the final manuscript.

Conflicts of interest: Roop Kaw reports no conflicts of interest.

Name: Eric Kezirian, MD.

Contribution: This author helped write the manuscript.

Attestation: Eric Kezirian approved the final manuscript.

Conflicts of interest: Nyxoah: Medical Advisory Board, Consultant, Inspire Medical Systems: Research Support, Consultant (previously), Pillar Palatal: Medical Advisory Board, Consultant, ReVENT Medical: Medical Advisory Board, Consultant, Split Rock Scientific: Consultant, Berendo Scientific: Consultant, Gerard Scientific: Consultant

Name: Atul Malhotra, MD.

Contribution: This author helped write the manuscript.

Attestation: Atul Malhotra approved the final manuscript.

Conflicts of interest: Atul Malhotra reports no conflicts of interest.

Name: Babak Mokhlesi, MD.

Contribution: This author helped write the manuscript.

Attestation: Babak Mokhlesi approved the final manuscript

Conflicts of interest: Babak Mokhlesi reports no conflicts of interest.

Name: Sairam Parthasarathy, MD.

Contribution: This author helped write the manuscript.

Attestation: Sairam Parthasarathy approved the final manuscript.

Conflicts of interest: Grants from NIH/NHLBI, Patient Centered Outcomes Research Institute, US Department of Defense, NIH (National Cancer Institute) NCI, US Department of Army, Johrei Institute, personal fees from American Academy of Sleep Medicine, American College of Chest Physicians, Philips-Respironics, Inc., UpToDate Inc., and

Vaoptherm, Inc. Non-financial support from National Center for Sleep Disorders Research of the NIH (NHLBI), grants from Younes Sleep Technologies, Ltd., Niveus Medical Inc., Philips-Respironics, Inc., a patent UA 14-018 U.S.S.N. 61/884,654; PTAS 502570970 (Home breathing device) pending.

Name: Tracey Stierer, MD.

Contribution: This author helped write the manuscript.

Attestation: Tracey Stierer approved the final manuscript.

Conflicts of interest: Tracey Stierer reports no conflicts of interest.

Name: Frank Wappler, MD.

Contribution: This author helped write the manuscript.

Attestation: Frank Wappler approved the final manuscript.

Conflicts of interest: Frank Wappler reported no conflicts of interest.

Name: David R. Hillman, MD.

Contribution: This author helped write the manuscript.

Attestation: David Hillman approved the final manuscript.

Conflicts of interest: David Hillman reports no conflicts of interest.

Name: Dennis Auckley, MD.

Contribution: This author helped design the study, conduct the study, analyze the data, and write the manuscript.

Attestation: Dennis Auckley approved the final manuscript.

Conflicts of interest: Dennis Auckley reports no conflicts of interest.

REFERENCES

1. Marin JM, Carrizo SJ, Vicente E, Agusti AG. Long-term cardiovascular outcomes in men with obstructive sleep apnoea-hypopnoea with or without treatment with continuous positive airway pressure: an observational study. *Lancet*. 2005;365:1046–1053.
2. Peker Y, Kraiczi H, Hedner J, Löth S, Johansson A, Bende M. An independent association between obstructive sleep apnoea and coronary artery disease. *Eur Respir J*. 1999;14:179–184.
3. Mehra R, Benjamin EJ, Shahar E, et al; Sleep Heart Health Study. Association of nocturnal arrhythmias with sleep-disordered breathing: The Sleep Heart Health Study. *Am J Respir Crit Care Med*. 2006;173:910–916.
4. Arzt M, Young T, Finn L, Skatrud JB, Bradley TD. Association of sleep-disordered breathing and the occurrence of stroke. *Am J Respir Crit Care Med*. 2005;172:1447–1451.
5. Yaggi HK, Concato J, Kernan WN, Lichtman JH, Brass LM, Mohsenin V. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med*. 2005;353:2034–2041.
6. Ip MS, Lam B, Ng MM, Lam WK, Tsang KW, Lam KS. Obstructive sleep apnea is independently associated with insulin resistance. *Am J Respir Crit Care Med*. 2002;165:670–676.
7. Coughlin SR, Mawdsley L, Mugarza JA, Calverley PM, Wilding JP. Obstructive sleep apnoea is independently associated with an increased prevalence of metabolic syndrome. *Eur Heart J*. 2004;25:735–741.
8. Yaffe K, Laffan AM, Harrison SL, et al. Sleep-disordered breathing, hypoxia, and risk of mild cognitive impairment and dementia in older women. *JAMA*. 2011;306:613–619.
9. Punjabi NM. The epidemiology of adult obstructive sleep apnea. *Proc Am Thorac Soc*. 2008;5:136–143.
10. Isono S. Obstructive sleep apnea of obese adults: pathophysiology and perioperative airway management. *Anesthesiology*. 2009;110:908–921.
11. Chung SA, Yuan H, Chung F. A systemic review of obstructive sleep apnea and its implications for anesthesiologists. *Anesth Analg*. 2008;107:1543–1563.
12. Seet E, Chung F. Management of sleep apnea in adults—functional algorithms for the perioperative period: continuing professional development. *Can J Anaesth*. 2010;57:849–864.
13. Peppard PE, Young T, Barnett JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol*. 2013;177:1006–1014.

14. Heinzer R, Vat S, Marques-Vidal P, et al. Prevalence of sleep-disordered breathing in the general population: the HypnoLaus study. *Lancet Respir Med*. 2015;3:310–318.
15. Young T, Evans L, Finn L, Palta M. Estimation of the clinically diagnosed proportion of sleep apnea syndrome in middle-aged men and women. *Sleep*. 1997;20:705–706.
16. Chen X, Wang R, Zee P, et al. Racial/ethnic differences in sleep disturbances: The Multi-Ethnic Study of Atherosclerosis (MESA). *Sleep*. 2015;38:877–888.
17. Hollowell PT, Stellato TA, Schuster M, et al. Potentially life-threatening sleep apnea is unrecognized without aggressive evaluation. *Am J Surg*. 2007;193:364–367.
18. Frey WC, Pilcher J. Obstructive sleep-related breathing disorders in patients evaluated for bariatric surgery. *Obes Surg*. 2003;13:676–683.
19. Singh M, Liao P, Kobah S, Wijeyesundera DN, Shapiro C, Chung F. Proportion of surgical patients with undiagnosed obstructive sleep apnoea. *Br J Anaesth*. 2013;110:629–636.
20. Finkel KJ, Searleman AC, Tymkew H, et al. Prevalence of undiagnosed obstructive sleep apnea among adult surgical patients in an academic medical center. *Sleep Med*. 2009;10:558.
21. Berry RB, Budhiraja R, Gottlieb DJ, et al; American Academy of Sleep Medicine. Rules for scoring respiratory events in sleep: update of the 2007 AASM Manual for the Scoring of Sleep and Associated Events. Deliberations of the Sleep Apnea Definitions Task Force of the American Academy of Sleep Medicine. *J Clin Sleep Med*. 2012;8:597–619.
22. Chung F, Liao P, Elsaid H, Shapiro CM, Kang W. Factors associated with postoperative exacerbation of sleep-disordered breathing. *Anesthesiology*. 2014;120:299–311.
23. Chung F, Liao P, Yegneswaran B, Shapiro CM, Kang W. Postoperative changes in sleep-disordered breathing and sleep architecture in patients with obstructive sleep apnea. *Anesthesiology*. 2014;120:287–298.
24. Kaw R, Chung F, Pasupuleti V, Mehta J, Gay PC, Hernandez AV. Meta-analysis of the association between obstructive sleep apnoea and postoperative outcome. *Br J Anaesth*. 2012;109:897–906.
25. Hai F, Porhomayon J, Vermont L, Frydrych L, Jaoude P, El-Solh AA. Postoperative complications in patients with obstructive sleep apnea: a meta-analysis. *J Clin Anesth*. 2014;26:591–600.
26. Memtsoudis S, Liu SS, Ma Y, et al. Perioperative pulmonary outcomes in patients with sleep apnea after noncardiac surgery. *Anesth Analg*. 2011;112:113–121.
27. Memtsoudis SG, Stundner O, Rasul R, et al. The impact of sleep apnea on postoperative utilization of resources and adverse outcomes. *Anesth Analg*. 2014;118:407–418.
28. Mokhlesi B, Hovda MD, Vekhter B, Arora VM, Chung F, Meltzer DO. Sleep-disordered breathing and postoperative outcomes after bariatric surgery: analysis of the nationwide inpatient sample. *Obes Surg*. 2013;23:1842–1851.
29. Mokhlesi B, Hovda MD, Vekhter B, Arora VM, Chung F, Meltzer DO. Sleep-disordered breathing and postoperative outcomes after elective surgery: analysis of the nationwide inpatient sample. *Chest*. 2013;144:903–914.
30. Kezirian EJ, Maselli J, Vittinghoff E, Goldberg AN, Auerbach AD. Obstructive sleep apnea surgery practice patterns in the United States: 2000 to 2006. *Otolaryngol Head Neck Surg*. 2010;143:441–447.
31. Young LR, Taxin ZH, Norman RG, Walsleben JA, Rapoport DM, Ayappa I. Response to CPAP withdrawal in patients with mild versus severe obstructive sleep apnea/hypopnea syndrome. *Sleep*. 2013;36:405–412.
32. Fouladpour N, Jesudoss R, Bolden N, Shaman Z, Auckley D. Perioperative Complications in Obstructive Sleep Apnea Patients Undergoing Surgery: A Review of the Legal Literature. *Anesth Analg*. 2016;122:145–151.
33. Svider PE, Pashkova AA, Folbe AJ, et al. Obstructive sleep apnea: strategies for minimizing liability and enhancing patient safety. *Otolaryngol Head Neck Surg*. 2013;149:947–953.
34. Lee LA, Caplan RA, Stephens LS, et al. Postoperative opioid-induced respiratory depression: a closed claims analysis. *Anesthesiology*. 2015;122:659–665.
35. Benumof JL. Mismanagement of obstructive sleep apnea may result in finding these patients dead in bed. *Can J Anaesth*. 2016;63:3–7.
36. Society of Anesthesia and Sleep Medicine Obstructive Sleep Apnea Death and Near Miss Registry. Available at: <http://depts.washington.edu/asaccp/projects/obstructive-sleep-apnea-osa-death-near-miss-registry>. Accessed January 12, 2016.
37. Cottam D, Lord J, Dallal RM, et al. Medicolegal analysis of 100 malpractice claims against bariatric surgeons. *Surg Obes Relat Dis*. 2007;3:60–66.
38. Coté CJ, Posner KL, Domino KB. Death or neurologic injury after tonsillectomy in children with a focus on obstructive sleep apnea: Houston, we have a problem! *Anesth Analg*. 2014;118:1276–1283.
39. Subramanyam R, Chidambaram V, Ding L, Myer CM III, Sadhasivam S. Anesthesia and opioids-related malpractice claims following tonsillectomy in USA: LexisNexis claims database 1984–2012. *Paediatr Anaesth*. 2014;24:412–420.
40. Gross JB, Bachenberg KL, Benumof JL, et al; American Society of Anesthesiologists Task Force on Perioperative Management. Practice guidelines for the perioperative management of patients with obstructive sleep apnea: a report by the American Society of Anesthesiologists Task Force on Perioperative Management of patients with obstructive sleep apnea. *Anesthesiology*. 2006;104:1081–1093.
41. Gross JB, Apfelbaum JL, Caplan RA, et al. Practice guidelines for the perioperative management of patients with obstructive sleep apnea. An updated report by the American Society of Anesthesiologists Task Force on perioperative management of patients with obstructive sleep apnea. *Anesthesiology*. 2014;120:268–286.
42. Joshi GP, Ankichetty SP, Gan TJ, Chung F. Society for Ambulatory Anesthesia consensus statement on preoperative selection of adult patients with obstructive sleep apnea scheduled for ambulatory surgery. *Anesth Analg*. 2012;115:1060–1068.
43. Meoli AL, Rosen CL, Kristo D, et al; Clinical Practice Review Committee; American Academy of Sleep Medicine. Upper airway management of the adult patient with obstructive sleep apnea in the perioperative period—avoiding complications. *Sleep*. 2003;26:1060–1065.
44. Adesanya AO, Lee W, Greilich NB, Joshi GP. Perioperative management of obstructive sleep apnea. *Chest*. 2010;138:1489–1498.
45. Guyatt GH, Oxman AD, Vist GE, et al; GRADE Working Group. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ*. 2008;336:924–926.
46. Balshem H, Helfand M, Schünemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol*. 2011;64:401–406.
47. Andrews JC, Schünemann HJ, Oxman AD, et al. GRADE guidelines: 15. Going from evidence to recommendation—determinants of a recommendation’s direction and strength. *J Clin Epidemiol*. 2013;66:726–735.
48. Andrews J, Guyatt G, Oxman AD, et al. GRADE guidelines: 14. Going from evidence to recommendations: the significance and presentation of recommendations. *J Clin Epidemiol*. 2013;66:719–725.
49. Schünemann HJ, Jaeschke R, Cook DJ, et al; ATSS Statements Development and Implementation Committee. An official ATS statement: grading the quality of evidence and strength of recommendations in ATS guidelines and recommendations. *Am J Respir Crit Care Med*. 2006;174:605–614.
50. Dalkey N, Helmer O. An experimental application of the Delphi method to the use of experts. *Management Science*. 1963;9:458–467.
51. Stundner O, Chiu YL, Sun X, et al. Sleep apnoea adversely affects the outcome in patients who undergo posterior lumbar fusion: a population-based study. *Bone Joint J*. 2014;96-B:242–248.
52. McIsaac DI, Gershon A, Wijeyesundera D, Bryson GL, Badner N, van Walraven C. Identifying obstructive sleep apnea in administrative data: a study of diagnostic accuracy. *Anesthesiology*. 2015;123:253–263.
53. Opperer M, Cozowicz C, Bugada D, et al. Does obstructive sleep apnea influence perioperative outcome? A qualitative systematic review for the society of anesthesia and sleep medicine.

- task force on preoperative preparation of patients with sleep-disordered breathing. *Anesth Analg*. 2016;122:1334.
54. Seet E, Chua M, Liaw CM. High STOP-BANG questionnaire scores predict intraoperative and early postoperative adverse events. *Singapore Med J*. 2015;56:212–216.
 55. Proczko MA, Stepaniak PS, de Quelerij M, et al. STOP-Bang and the effect on patient outcome and length of hospital stay when patients are not using continuous positive airway pressure. *J Anesth*. 2014;28:891–897.
 56. Nguyen NT, Masoomi H, Laugouner K, et al. Predictive factors of mortality in bariatric surgery: data from the Nationwide Inpatient Sample. *Surgery*. 2011;150:347–351.
 57. D'Apuzzo MR, Browne JA. Obstructive sleep apnea as a risk factor for postoperative complications after revision joint arthroplasty. *J Arthroplasty*. 2012;27:95–98.
 58. Liao P, Yegneswaran B, Vairavanathan S, Zilberman P, Chung F. Postoperative complications in patients with obstructive sleep apnea: a retrospective matched cohort study. *Can J Anaesth*. 2009;56:819–828.
 59. Kaw R, Pasupuleti V, Walker E, Ramaswamy A, Foldvary-Schafer N. Postoperative complications in patients with obstructive sleep apnea. *Chest*. 2012;141:436–441.
 60. Mador MJ, Goplani S, Gottumukkala VA, et al. Postoperative complications in obstructive sleep apnea. *Sleep Breath*. 2013;17:727–734.
 61. Mutter TC, Chateau D, Moffatt M, Ramsey C, Roos LL, Kryger M. A matched cohort study of postoperative outcomes in obstructive sleep apnea: could preoperative diagnosis and treatment prevent complications? *Anesthesiology*. 2014;121:707–718.
 62. Chung F, Yegneswaran B, Liao P, et al. Validation of the Berlin questionnaire and American Society of Anesthesiologists checklist as screening tools for obstructive sleep apnea in surgical patients. *Anesthesiology*. 2008;108:822–830.
 63. Gali B, Whalen FX, Schroeder DR, Gay PC, Plevak DJ. Identification of patients at risk for postoperative respiratory complications using a preoperative obstructive sleep apnea screening tool and postanesthesia care assessment. *Anesthesiology*. 2009;110:869–877.
 64. Corso RM, Petrini F, Buccioli M, et al. Clinical utility of preoperative screening with STOP-Bang questionnaire in elective surgery. *Minerva Anestesiol*. 2014;80:877–884.
 65. Griffin JW, Novicoff WM, Browne JA, Brockmeier SF. Obstructive sleep apnea as a risk factor after shoulder arthroplasty. *J Shoulder Elbow Surg*. 2013;22:e6–e9.
 66. Jensen C, Tejirian T, Lewis C, Yadegar J, Dutton E, Mehran A. Postoperative CPAP and BiPAP use can be safely omitted after laparoscopic Roux-en-Y gastric bypass. *Surg Obes Relat Dis*. 2008;4:512–514.
 67. Kaw R, Golish J, Ghamande S, Burgess R, Foldvary N, Walker E. Incremental risk of obstructive sleep apnea on cardiac surgical outcomes. *J Cardiovasc Surg (Torino)*. 2006;47:683–689.
 68. Weingarten TN, Flores AS, McKenzie JA, et al. Obstructive sleep apnoea and perioperative complications in bariatric patients. *Br J Anaesth*. 2011;106:131–139.
 69. Ursavaş A, Güven T, Coskun F, Ege E, Yilmazlar A. Association between self reported snoring, STOP questionnaire and postoperative pulmonary complications in patients scheduled for orthopaedic surgery. *Multidiscip Respir Med*. 2013;8:3.
 70. Berend KR, Ajluni AF, Núñez-García LA, Lombardi AV, Adams JB. Prevalence and management of obstructive sleep apnea in patients undergoing total joint arthroplasty. *J Arthroplasty*. 2010;25:54–57.
 71. Gali B, Whalen FX Jr, Gay PC, et al. Management plan to reduce risks in perioperative care of patients with presumed obstructive sleep apnea syndrome. *J Clin Sleep Med*. 2007;3:582–588.
 72. Munish M, Sharma V, Yarussi KM, Sifain A, Porhomonayon J, Nader N. The use of practice guidelines by the American Society of Anesthesiologists for the identification of surgical patients at high risk of sleep apnea. *Chron Respir Dis*. 2012;9:221–230.
 73. Blake DW, Chia PH, Donnan G, Williams DL. Preoperative assessment for obstructive sleep apnoea and the prediction of postoperative respiratory obstruction and hypoxaemia. *Anaesth Intensive Care*. 2008;36:379–384.
 74. Flink BJ, Rivelli SK, Cox EA, et al. Obstructive sleep apnea and incidence of postoperative delirium after elective knee replacement in the nondemented elderly. *Anesthesiology*. 2012;116:788–796.
 75. Ahmad S, Nagle A, McCarthy RJ, Fitzgerald PC, Sullivan JT, Prystowsky J. Postoperative hypoxemia in morbidly obese patients with and without obstructive sleep apnea undergoing laparoscopic bariatric surgery. *Anesth Analg*. 2008;107:138–143.
 76. Eikermann M, Garzon-Serrano J, Kwo J, Grosse-Sundrup M, Schmidt U, Bigatello L. Do patients with obstructive sleep apnea have an increased risk of desaturation during induction of anesthesia for weight loss surgery? *Open Respir Med J*. 2010;4:58–62.
 77. O'Gorman SM, Gay PC, Morgenthaler TI. Does autotitrating positive airway pressure therapy improve postoperative outcome in patients at risk for obstructive sleep apnea syndrome? A randomized controlled clinical trial. *Chest*. 2013;144:72–78.
 78. Pereira H, Xará D, Mendonça J, Santos A, Abelha FJ. Patients with a high risk for obstructive sleep apnea syndrome: postoperative respiratory complications. *Rev Port Pneumol*. 2013;19:144–151.
 79. Iyer US, Koh KF, Chia NC, Macachor J, Cheng A. Perioperative risk factors in obese patients for bariatric surgery: a Singapore experience. *Singapore Med J*. 2011;52:94–99.
 80. Kheterpal S, Healy D, Aziz MF, et al; Multicenter Perioperative Outcomes Group (MPOG) Perioperative Clinical Research Committee. Incidence, predictors, and outcome of difficult mask ventilation combined with difficult laryngoscopy: a report from the multicenter perioperative outcomes group. *Anesthesiology*. 2013;119:1360–1369.
 81. Kheterpal S, Martin L, Shanks AM, Tremper KK. Prediction and outcomes of impossible mask ventilation: a review of 50,000 anesthetics. *Anesthesiology*. 2009;110:891–897.
 82. Stierer TL, Wright C, George A, Thompson RE, Wu CL, Collop N. Risk assessment of obstructive sleep apnea in a population of patients undergoing ambulatory surgery. *J Clin Sleep Med*. 2010;6:467–472.
 83. Vest D, Lee D, Newcome K, Stamper H. A retrospective review of difficult intubations: is obstructive sleep apnea a predictor? *Clin Nurse Spec*. 2013;27:128–131.
 84. Neligan PJ, Porter S, Max B, Malhotra G, Greenblatt EP, Ochroch EA. Obstructive sleep apnea is not a risk factor for difficult intubation in morbidly obese patients. *Anesth Analg*. 2009;109:1182–1186.
 85. Moe T, Gullsbj S, Rabben T, Eriksson P. Sleep-disordered breathing: a novel predictor of atrial fibrillation after coronary artery bypass surgery. *Coron Artery Dis*. 1996;7:475–478.
 86. Mungan U, Ozeke O, Mavioglu L, et al. The role of the preoperative screening of sleep apnoea by Berlin Questionnaire and Epworth Sleepiness Scale for postoperative atrial fibrillation. *Heart Lung Circ*. 2013;22:38–42.
 87. van Oosten EM, Hamilton A, Petsikas D, et al. Effect of preoperative obstructive sleep apnea on the frequency of atrial fibrillation after coronary artery bypass grafting. *Am J Cardiol*. 2014;113:919–923.
 88. Hwang D, Shakir N, Limann B, et al. Association of sleep-disordered breathing with postoperative complications. *Chest*. 2008;133:1128–1134.
 89. Gupta RM, Parvizi J, Hanssen AD, Gay PC. Postoperative complications in patients with obstructive sleep apnea syndrome undergoing hip or knee replacement: a case-control study. *Mayo Clin Proc*. 2001;76:897–905.
 90. Vasu TS, Doghramji K, Cavallazzi R, et al. Obstructive sleep apnea syndrome and postoperative complications: clinical use of the STOP-BANG questionnaire. *Arch Otolaryngol Head Neck Surg*. 2010;136:1020–1024.
 91. Kamath AF, McAuliffe CL, Baldwin KD, Lucas JB, Kosseim LM, Israelite CL. Unplanned admission to the intensive care unit after total hip arthroplasty. *J Arthroplasty*. 2012;27:1027–1032.e1–2.
 92. Dorman RB, Miller CJ, Leslie DB, et al. Risk for hospital readmission following bariatric surgery. *PLoS One*. 2012;7:e32506.
 93. Chia P, Seet E, Macachor J, Iyer US, Wu D. The association of pre-operative STOP-BANG questionnaire scores with postoperative critical care admission. *Anaesthesia*. 2013;68:950–952.

94. Grover BT, Priem DM, Mathiason MA, Kallies KJ, Thompson GP, Kothari SN. Intensive care unit stay not required for patients with obstructive sleep apnea after laparoscopic Roux-en-Y gastric bypass. *Surg Obes Relat Dis*. 2010;6:165–170.
95. Sabers C, Plevak DJ, Schroeder DR, Warner DO. The diagnosis of obstructive sleep apnea as a risk factor for unanticipated admissions in outpatient surgery. *Anesth Analg*. 2003;96:1328–1335.
96. Bryson GL, Gomez CP, Jee RM, Blackburn J, Taljaard M, Forster AJ. Unplanned admission after day surgery: a historical cohort study in patients with obstructive sleep apnea. *Can J Anaesth*. 2012;59:842–851.
97. Masoomi H, Kim H, Reavis KM, Mills S, Stamos MJ, Nguyen NT. Analysis of factors predictive of gastrointestinal tract leak in laparoscopic and open gastric bypass. *Arch Surg*. 2011;146:1048–1051.
98. Andrews KL, Dib M, Shives TC, Hoskin TL, Liedl DA, Boon AJ. The effect of obstructive sleep apnea on amputation site healing. *J Vasc Nurs*. 2012;30:61–63.
99. Kurrek MM, Cobourn C, Wojtasik Z, Kiss A, Dain SL. Morbidity in patients with or at high risk for obstructive sleep apnea after ambulatory laparoscopic gastric banding. *Obes Surg*. 2011;21:1494–1498.
100. Coté GA, Hovis CE, Hovis RM, et al. A screening instrument for sleep apnea predicts airway maneuvers in patients undergoing advanced endoscopic procedures. *Clin Gastroenterol Hepatol*. 2010;8:660–665.e1.
101. Goudra BG, Singh PM, Penugonda LC, Speck RM, Sinha AC. Significantly reduced hypoxemic events in morbidly obese patients undergoing gastrointestinal endoscopy: predictors and practice effect. *J Anaesthesiol Clin Pharmacol*. 2014;30:71–77.
102. Kim GH, Lee JJ, Choi SJ, et al. Clinical predictors of apnoea-hypopnoea during propofol sedation in patients undergoing spinal anaesthesia. *Anaesthesia*. 2012;67:755–759.
103. Adler DG, Kawa C, Hilden K, Fang J. Nurse-administered propofol sedation is safe for patients with obstructive sleep apnea undergoing routine endoscopy: a pilot study. *Dig Dis Sci*. 2011;56:2666–2671.
104. Cha JM, Jeun JW, Pack KM, et al. Risk of sedation for diagnostic esophagogastroduodenoscopy in obstructive sleep apnea patients. *World J Gastroenterol*. 2013;19:4745–4751.
105. Gill J, Vidyarthi G, Kulkarni P, Anderson W, Boyd W. Safety of conscious sedation in patients with sleep apnea in a veteran population. *South Med J*. 2011;104:185–188.
106. Khiani VS, Salah W, Maimone S, Cummings L, Chak A. Sedation during endoscopy for patients at risk of obstructive sleep apnea. *Gastrointest Endosc*. 2009;70:1116–1120.
107. Boese ML, Ransom RK, Roadfuss RJ, Todd A, McGuire JM. Utility of the Berlin Questionnaire to screen for obstructive sleep apnea among patients receiving intravenous sedation for colonoscopy. *AANA J*. 2014;82:38–45.
108. Mehta PP, Kochhar G, Kalra S, et al. Can a validated sleep apnea scoring system predict cardiopulmonary events using propofol sedation for routine EGD or colonoscopy? A prospective cohort study. *Gastrointest Endosc*. 2014;79:436–444.
109. Mador MJ, Abo Khamis M, Nag N, Mreyoud A, Jallu S, Mehboob S. Does sleep apnea increase the risk of cardiorespiratory complications during endoscopy procedures? *Sleep Breath*. 2011;15:393–401.
110. Mador MJ, Nadler J, Mreyoud A, et al. Do patients at risk of sleep apnea have an increased risk of cardio-respiratory complications during endoscopy procedures? *Sleep Breath*. 2012;16:609–615.
111. Lockhart EM, Willingham MD, Abdallah AB, et al. Obstructive sleep apnea screening and postoperative mortality in a large surgical cohort. *Sleep Med*. 2013;14:407–415.
112. Abdelsattar ZM, Hendren S, Wong SL, Campbell DA Jr, Ramachandran SK. The impact of untreated obstructive sleep apnea on cardiopulmonary complications in general and vascular surgery: a cohort study. *Sleep*. 2015;38:1205–1210.
113. Chung F, Nagappa M, Singh M, Mokhlesi B. CPAP in the perioperative setting: evidence of support. *Chest*. 2016;149:586–597.
114. Nagappa M, Mokhlesi B, Wong J, Wong DT, Kaw R, Chung F. The effects of continuous positive airway pressure on postoperative outcomes in obstructive sleep apnea patients undergoing surgery: a systematic review and meta-analysis. *Anesth Analg*. 2015;120:1013–1023.
115. Ramachandran SK, Kheterpal S, Haas CF, Saran KA, Tremper KK. Automated notification of suspected obstructive sleep apnea patients to the perioperative respiratory therapist: a pilot study. *Respir Care*. 2010;55:414–418.
116. Chung F, Yegneswaran B, Liao P, et al. STOP questionnaire: a tool to screen patients for obstructive sleep apnea. *Anesthesiology*. 2008;108:812–821.
117. Chung F, Subramanyam R, Liao P, Sasaki E, Shapiro C, Sun Y. High STOP-Bang score indicates a high probability of obstructive sleep apnoea. *Br J Anaesth*. 2012;108:768–775.
118. Ghiasi F, Ahmadpoor A, Amra B. Relationship between obstructive sleep apnea and 30-day mortality among patients with pulmonary embolism. *J Res Med Sci*. 2015;20:662–667.
119. Xará D, Santos A, Abelha F. Adverse respiratory events in a post-anesthesia care unit. *Arch Bronconeumol*. 2015;51:69–75.
120. Acar HV, Yarkan Uysal H, Kaya A, Ceyhan A, Dikmen B. Does the STOP-Bang, an obstructive sleep apnea screening tool, predict difficult intubation? *Eur Rev Med Pharmacol Sci*. 2014;18:1869–1874.
121. Veenstra A, Untalan E. Implications and interventions related to obstructive sleep apnea. *Crit Care Nurs Clin North Am*. 2014;26:499–509.
122. Louis JM, Mogos MF, Salemi JL, Redline S, Salihu HM. Obstructive sleep apnea and severe maternal-infant morbidity/mortality in the United States, 1998–2009. *Sleep*. 2014;37:843–849.
123. Pamidi S, Pinto LM, Marc I, Benedetti A, Schwartzman K, Kimoff RJ. Maternal sleep-disordered breathing and adverse pregnancy outcomes: a systematic review and metaanalysis. *Am J Obstet Gynecol*. 2014;210:52.e1–52.e14.
124. Xu T, Feng Y, Peng H, Guo D, Li T. Obstructive sleep apnea and the risk of perinatal outcomes: a meta-analysis of cohort studies. *Sci Rep*. 2014;4:6982.
125. Ding XX, Wu YL, Xu SJ, et al. A systematic review and quantitative assessment of sleep-disordered breathing during pregnancy and perinatal outcomes. *Sleep Breath*. 2014;18:703–713.
126. Abdullah HR, Nagappa M, Siddiqui N, Chung F. Diagnosis and treatment of obstructive sleep apnea during pregnancy. *Curr Opin Anaesthesiol*. 2016. in print.
127. Antony KM, Agrawal A, Arndt ME, et al. Association of adverse perinatal outcomes with screening measures of obstructive sleep apnea. *J Perinatol*. 2014;34:441–448.
128. Khazaie H, Heidarpour A, Nikray R, et al. Evaluation of sleep problems in preeclamptic, healthy pregnant and non-pregnant women. *Iran J Psychiatry*. 2013;8:168–171.
129. Champagne K, Schwartzman K, Opatrny L, et al. Obstructive sleep apnoea and its association with gestational hypertension. *Eur Respir J*. 2009;33:559–565.
130. Izci Balsearak B, Jackson N, Ratcliffe SA, Pack AI, Pien GW. Sleep-disordered breathing and daytime napping are associated with maternal hyperglycemia. *Sleep Breath*. 2013;17:1093–1102.
131. Williams MA, Gelaye B, Qiu C, Fida N, May Cripe S. Habitual snoring and asthma comorbidity among pregnant women. *J Asthma*. 2011;48:91–97.
132. Facco FL, Ouyang DW, Zee PC, Grobman WA. Development of a pregnancy-specific screening tool for sleep apnea. *J Clin Sleep Med*. 2012;8:389–394.
133. Wilson DL, Walker SP, Fung AM, O'Donoghue F, Barnes M, Howard M. Can we predict sleep-disordered breathing in pregnancy? The clinical utility of symptoms. *J Sleep Res*. 2013;22:670–678.
134. Fernández Alonso AM, Chedraui P, Pérez-López FR. Assessment of obstructive sleep apnea-hypopnea syndrome risk at the end of pregnancy using the Berlin Questionnaire. *Gynecol Endocrinol*. 2015;31:715–719.
135. Tantrakul V, Sirijanchune P, Panburana P, et al. Screening of obstructive sleep apnea during pregnancy: differences in

- predictive values of questionnaires across trimesters. *J Clin Sleep Med*. 2015;11:157–163.
136. Lockhart EM, Ben Abdallah A, Tuuli MG, Leighton BL. Obstructive sleep apnea in pregnancy: assessment of current screening tools. *Obstet Gynecol*. 2015;126:93–102.
 137. Lee W, Nagubadi S, Kryger MH, Mokhlesi B. Epidemiology of obstructive sleep apnea: a population-based perspective. *Expert Rev Respir Med*. 2008;2:349–264.
 138. Sood A, Abdollah F, Sammon JD, et al. Effect of body mass index on perioperative outcomes after major surgery: results from the National Surgical Quality Improvement Program (ACS-NSQIP) 2005–2011. *World J Surg*. 2015; 39:2376–2385.
 139. Schumann R, Shikora SA, Sigl JC, Kelley SD. Association of metabolic syndrome and surgical factors with pulmonary adverse events, and longitudinal mortality in bariatric surgery. *Br J Anaesth*. 2015;114:83–90.
 140. Chung F, Yang Y, Liao P. Predictive performance of the STOP-Bang score for identifying obstructive sleep apnea in obese patients. *Obes Surg*. 2013;23:2050–2057.
 141. Memtsoudis SG, Stundner O, Rasul R, et al. Sleep apnea and total joint arthroplasty under various types of anesthesia: a population-based study of perioperative outcomes. *Reg Anesth Pain Med*. 2013;38:274–281.
 142. Ramachandran SK, Khetarpal S, Consens F, et al. Derivation and validation of a simple perioperative sleep apnea prediction score. *Anesth Analg*. 2010;110:1007–1015.
 143. Ramachandran SK, Josephs LA. A meta-analysis of clinical screening tests for obstructive sleep apnea. *Anesthesiology*. 2009;110:928–939.
 144. Abrishami A, Khajehdehi A, Chung F. A systematic review of screening questionnaires for obstructive sleep apnea. *Can J Anaesth*. 2010;57:423–438.
 145. Chung F, Abdullah HR, Liao P. STOP-Bang questionnaire: a practical approach to screen for obstructive sleep apnea. *Chest*. 2016;149:631–638.
 146. Chung F, Liao P, Farney R. Correlation between the STOP-Bang score and the severity of obstructive sleep apnea. *Anesthesiology*. 2015;122:1436–1437.
 147. Chung F, Yang Y, Brown R, Liao P. Alternative scoring models of STOP-bang questionnaire improve specificity to detect undiagnosed obstructive sleep apnea. *J Clin Sleep Med*. 2014;10:951–958.
 148. Nagappa M, Liao P, Wong J, et al. Validation of the STOP-Bang questionnaire as a screening tool for obstructive sleep apnea among different populations: a systematic review and meta-analysis. *PLoS One*. 2015;10:e0143697.
 149. Nunes FS, Danzi-Soares NJ, Genta PR, Drager LF, Cesar LA, Lorenzi-Filho G. Critical evaluation of screening questionnaires for obstructive sleep apnea in patients undergoing coronary artery bypass grafting and abdominal surgery. *Sleep Breath*. 2015;19:115–122.
 150. Ong TH, Raudha S, Chong S, Lew N, Hsu AA. Simplifying STOP-BANG score of a simple questionnaire to screen for OSA in an Asian population. *Sleep Breath*. 2010;14:371–376.
 151. Farney RJ, Walker BS, Farney RM, Snow GL, Walker JM. The STOP-Bang equivalent model and prediction of severity of obstructive sleep apnea: relation to polysomnographic measurements of the apnea/hypopnea index. *J Clin Sleep Med*. 2011;7:459–465B.
 152. Boynton G, Vahabzadeh A, Hammo, Ruzicka DL, Chervin RD. Validation of the STOP-BANG questionnaire among Patients Referred for Suspected Obstructive Sleep Apnea. *J Sleep Disord Treat Care*. 2013;2.
 153. Pereira EJ, Driver HS, Stewart SC, Fitzpatrick MF. Comparing a combination of validated questionnaires and level III portable monitor with polysomnography to diagnose and exclude sleep apnea. *J Clin Sleep Med*. 2013;9:1259–1266.
 154. Vana KD, Silva GE, Goldberg R. Predictive abilities of the STOP-Bang and Epworth Sleepiness Scale in identifying sleep clinic patients at high risk for obstructive sleep apnea. *Res Nurs Health*. 2013;36:84–94.
 155. Cowan DC, Allardice G, Macfarlane D, et al. Predicting sleep disordered breathing in outpatients with suspected OSA. *BMJ Open*. 2014;4:e004519.
 156. Ha SC, Lee DL, Abdullah VJ, van Hasselt CA. Evaluation and validation of four translated Chinese questionnaires for obstructive sleep apnea patients in Hong Kong. *Sleep Breath*. 2014;18:715–721.
 157. Luo J, Huang R, Zhong X, Xiao Y, Zhou J. STOP-Bang questionnaire is superior to Epworth sleepiness scales, Berlin questionnaire, and STOP questionnaire in screening obstructive sleep apnea hypopnea syndrome patients. *Chin Med J (Engl)*. 2014;127:3065–3070.
 158. Luo J, Huang R, Zhong X, Xiao Y, Zhou J. Value of STOP-Bang questionnaire in screening patients with obstructive sleep apnea hypopnea syndrome in sleep disordered breathing clinic. *Chin Med J (Engl)*. 2014;127:1843–1848.
 159. Reis R, Teixeira F, Martins V, et al. Validation of a Portuguese version of the STOP-Bang questionnaire as a screening tool for obstructive sleep apnea: analysis in a sleep clinic. *Rev Port Pneumol. (2006)* 2015;21:61–68.
 160. Silva GE, Vana KD, Goodwin JL, Sherrill DL, Quan SF. Identification of patients with sleep disordered breathing: comparing the four-variable screening tool, STOP, STOP-Bang, and Epworth Sleepiness Scales. *J Clin Sleep Med*. 2011;7:467–472.
 161. Menown IB, Allen J, Anderson JM, Adgey AA. ST depression only on the initial 12-lead ECG: early diagnosis of acute myocardial infarction. *Eur Heart J*. 2001;22:218–227.
 162. Okin PM, Roman MJ, Devereux RB, Kligfield P. Electrocardiographic identification of left ventricular hypertrophy: test performance in relation to definition of hypertrophy and presence of obesity. *J Am Coll Cardiol*. 1996;27:124–131.
 163. Chung F, Chau E, Yang Y, Liao P, Hall R, Mokhlesi B. Serum bicarbonate level improves specificity of STOP-Bang screening for obstructive sleep apnea. *Chest*. 2013;143:1284–1293.
 164. Chung F, Zhou L, Liao P. Parameters from preoperative overnight oximetry predict postoperative adverse events. *Minerva Anestesiol*. 2014;80:1084–1095.
 165. Guralnick AS, Pant M, Minhaj M, Sweitzer BJ, Mokhlesi B. CPAP adherence in patients with newly diagnosed obstructive sleep apnea prior to elective surgery. *J Clin Sleep Med*. 2012;8:501–506.
 166. Memtsoudis SG, Besculides MC, Mazumdar M. A rude awakening—the perioperative sleep apnea epidemic. *N Engl J Med*. 2013;368:2352–2353.
 167. Rennotte MT, Baele P, Aubert G, Rodenstein DO. Nasal continuous positive airway pressure in the perioperative management of patients with obstructive sleep apnea submitted to surgery. *Chest*. 1995;107:367–374.
 168. Liao P, Luo Q, Elsaid H, Kang W, Shapiro CM, Chung F. Perioperative auto-titrated continuous positive airway pressure treatment in surgical patients with obstructive sleep apnea: a randomized controlled trial. *Anesthesiology*. 2013;119:837–847.
 169. Kohler M, Stoewhas AC, Ayers L, et al. Effects of continuous positive airway pressure therapy withdrawal in patients with obstructive sleep apnea: a randomized controlled trial. *Am J Respir Crit Care Med*. 2011;184:1192–1199.
 170. Yadollahi A, Gabriel JM, White LH, Taranto Montemurro L, Kasai T, Bradley TD. A randomized, double crossover study to investigate the influence of saline infusion on sleep apnea severity in men. *Sleep*. 2014;37:1699–1705.
 171. Ramachandran SK. Can intravenous fluids explain increased postoperative sleep disordered breathing and airway outcomes? *Sleep*. 2014;37:1587–1588.
 172. Lam T, Singh M, Yadollahi A, Chung F. Is perioperative fluid and salt balance a contributing factor in postoperative worsening of obstructive sleep apnea? *Anesth Analg*. 2016;122:1335–1339.
 173. Kaw R, Bhateja P, Paz Y, et al. Postoperative complications in patients with unrecognized obesity hypoventilation syndrome undergoing elective noncardiac surgery. *Chest*. 2016;149:84–91.

174. Ramakrishna G, Sprung J, Ravi BS, Chandrasekaran K, McGoon MD. Impact of pulmonary hypertension on the outcomes of noncardiac surgery: predictors of perioperative morbidity and mortality. *J Am Coll Cardiol*. 2005;45:1691–1699.
175. Minai OA, Ricaurte B, Kaw R, et al. Frequency and impact of pulmonary hypertension in patients with obstructive sleep apnea syndrome. *Am J Cardiol*. 2009;104:1300–1306.
176. Ireland CJ, Chapman TM, Mathew SF, Herbison GP, Zacharias M. Continuous positive airway pressure (CPAP) during the postoperative period for prevention of postoperative morbidity and mortality following major abdominal surgery. *Cochrane Database Syst Rev*. 2014;8:CD008930.
177. Taenzer AH, Pyke JB, McGrath SP, Blike GT. Impact of pulse oximetry surveillance on rescue events and intensive care unit transfers: a before-and-after concurrence study. *Anesthesiology*. 2010;112:282–287.
178. Stoelting RK. Continuous postoperative electronic monitoring and the will to require it. *Anesth Analg*. 2015;121:579–581.
179. Doufas AG, Tian L, Davies MF, Warby SC. Nocturnal intermittent hypoxia is independently associated with pain in subjects suffering from sleep-disordered breathing. *Anesthesiology*. 2013;119:1149–1162.
180. Doufas A. Obstructive sleep apnea, pain, and opioid analgesia in the postoperative Patient. *Curr Anesthesiol Rep*. 2014;4:1–9.
181. Turan A, You J, Egan C, et al. Chronic intermittent hypoxia is independently associated with reduced postoperative opioid consumption in bariatric patients suffering from sleep-disordered breathing. *PLoS One*. 2015;10:e0127809.
182. Lam KK, Kunder S, Wong J, Doufas AG, Chung F. Obstructive sleep apnea, pain, and opioids: is the riddle solved? *Curr Opin Anaesthesiol*. 2016;29:134–140.
183. Parthasarathy S, Haynes PL, Budhiraja R, Habib MP, Quan SF. A national survey of the effect of sleep medicine specialists and American Academy of Sleep Medicine Accreditation on management of obstructive sleep apnea. *J Clin Sleep Med*. 2006;2:133–142.
184. Pamidi S, Knutson KL, Ghods F, Mokhlesi B. The impact of sleep consultation prior to a diagnostic polysomnogram on continuous positive airway pressure adherence. *Chest*. 2012;141:51–57.
185. Parthasarathy S, Subramanian S, Quan SF. A multicenter prospective comparative effectiveness study of the effect of physician certification and center accreditation on patient-centered outcomes in obstructive sleep apnea. *J Clin Sleep Med*. 2014;10:243–249.
186. Russell T. Enhancing adherence to positive airway pressure therapy for sleep disordered breathing. *Semin Respir Crit Care Med*. 2014;35:604–612.
187. Chong CT, Tey J, Leow SL, et al. Management plan to reduce risks in perioperative care of patients with obstructive sleep apnoea averts the need for presurgical polysomnography. *Ann Acad Med Singapore*. 2013;42:110–119.
188. Fonseca MI, Pereira T, Caseiro P. Death and disability in patients with sleep apnea—a meta-analysis. *Arq Bras Cardiol*. 2015;104:58–66.
189. Gami AS, Olson EJ, Shen WK, et al. Obstructive sleep apnea and the risk of sudden cardiac death: a longitudinal study of 10,701 adults. *J Am Coll Cardiol*. 2013;62:610–616.
190. Bradley TD, Floras JS. Obstructive sleep apnoea and its cardiovascular consequences. *Lancet*. 2009;373:82–93.
191. Sharma SK, Agrawal S, Damodaran D, et al. CPAP for the metabolic syndrome in patients with obstructive sleep apnea. *N Engl J Med*. 2011;365:2277–2286.
192. Marin JM, Agustí A, Villar I, et al. Association between treated and untreated obstructive sleep apnea and risk of hypertension. *JAMA*. 2012;307:2169–2176.
193. Mehta V, Subramanyam R, Shapiro CM, Chung F. Health effects of identifying patients with undiagnosed obstructive sleep apnea in the preoperative clinic: a follow-up study. *Can J Anaesth*. 2012;59:544–555.
194. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, Gawande AA. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet*. 2008;372:139–144.
195. An advisory on safety and quality issue. The Joint Commission, Division of Health Care Improvement. At risk: Obstructive sleep apnea patients. *Quick Safety*. 2015;14:1–2.
196. Eckert DJ, White DP, Jordan AS, Malhotra A, Wellman A. Defining phenotypic causes of obstructive sleep apnea. Identification of novel therapeutic targets. *Am J Respir Crit Care Med*. 2013;188:996–1004.
197. Malhotra A, Orr JE, Owens RL. On the cutting edge of obstructive sleep apnoea: where next? *Lancet Respir Med*. 2015;3:397–403.
198. Chung F, Hillman D, Lydic R. Sleep medicine and anesthesia: a new horizon for anesthesiologists. *Anesthesiology*. 2011;114:1261–1262.
199. Potts KJ, Butterfield DT, Sims P, Henderson M, Shames CB. Cost savings associated with an education campaign on the diagnosis and management of sleep-disordered breathing: a retrospective, claims-based US study. *Popul Health Manag*. 2013;16:7–13.
200. Skaer TL, Sclar DA. Economic implications of sleep disorders. *Pharmacoeconomics*. 2010;28:1015–1023.
201. Memtsoudis SG, Sun X, Chiu YL, et al. Perioperative comparative effectiveness of anesthetic technique in orthopedic patients. *Anesthesiology*. 2013;118:1046–1058.
202. Ankichetty S, Wong J, Chung F. A systematic review of the effects of sedatives and anesthetics in patients with obstructive sleep apnea. *J Anaesthesiol Clin Pharmacol*. 2011;27:447–458.
203. McEntire DM, Kirkpatrick DR, Kerfeld MJ, et al. Effect of sedative-hypnotics, anesthetics and analgesics on sleep architecture in obstructive sleep apnea. *Expert Rev Clin Pharmacol*. 2014;7:787–806.
204. Molnar MZ, Mucsi I, Novak M, et al. Association of incident obstructive sleep apnoea with outcomes in a large cohort of US veterans. *Thorax*. 2015;70:888–895.
205. Dwarakanath A, Palissery V, Elliott MW. Prevalence of and treatment outcomes for patients with obstructive sleep apnoea identified by preoperative screening compared with clinician referrals. *Eur Respir J*. 2016. 