

Lawrence Berkeley National Laboratory

Recent Work

Title

COUPLINGS OF VECTOR MESONS TO TENSOR MESONS AND THE POMERON

Permalink

<https://escholarship.org/uc/item/5gj9s09d>

Author

Vasavada, Kashyap V.

Publication Date

1976-09-01

Submitted to Physical Review

LBL-5571
Preprint c.


COUPLINGS OF VECTOR MESONS TO TENSOR
MESONS AND THE POMERON

Kashyap V. Vasavada

September 2, 1976

For Reference

Not to be taken from this room



Prepared for the U. S. Energy Research and
Development Administration under Contract W-7405-ENG-48

01100940110

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

COUPLINGS OF VECTOR MESONS TO TENSOR MESONS

AND THE POMERON[†]

by

Kashyap V. Vasavada

Department of Physics and Lawrence Berkeley Laboratory
University of California
Berkeley, California
94720

Department of Physics
Indiana-Purdue University
Indianapolis, Indiana 46205*

ABSTRACT

By saturating matrix elements of the stress tensor between vector meson states with f , f' and the Pomeron, we find that (i) nonet mass formulas are obtained (ii) vector meson-nucleon total cross sections are determined, with a $1/m_V^2$ dependence on the vector meson masses and (iii) tensor-vector-vector couplings are obtained. All of these are in agreement with the experiments.

1. Introduction

The idea of tensor meson dominance of energy-momentum tensor $\theta_{\mu\nu}$ has been explored for a number of years⁽¹⁾. Recently this concept has become even more attractive because of its possible connection with a gravitation type theory of strongly interacting particles. In previous papers^(2,3) we have attempted to build in Pomeron (P) contribution along with that of f and f' mesons in $\theta_{\mu\nu}$. In particular, in ref. 3 (I)

September 2, 1976

-2-

matrix elements of $\theta_{\mu\nu}$ between the octet baryon states were saturated with f , f' and P to obtain the Gell-Mann-Okubo (GMO) mass formula, D/F ratio for tensor meson-baryon-baryon couplings, values of P and f -nucleon-nucleon couplings, all in good agreement with the experiments. In this note we apply a similar procedure to the matrix elements of $\theta_{\mu\nu}$ between vector meson states and find very interesting consequences for the tensor couplings and total cross-sections of the old (ρ , K^* , ω , ϕ) and the new (ψ , ψ' , D^* etc.) vector mesons.

II. Tensor Meson and Pomeron Dominance

The matrix elements of $\theta_{\mu\nu}$ between two identical vector meson states contain six form factors⁽⁴⁾. In the present work we will be concerned with only one of these:

$$\langle V_2(p_2) | \theta_{\mu\nu} | V_1(p_1) \rangle = \frac{G_1(q^2)}{2} \epsilon_1 \cdot \epsilon_2 P_\mu P_\nu + \dots \quad (1)$$

where $P = p_1 + p_2$, $q = p_1 - p_2$. ϵ_1 and ϵ_2 are the polarization vectors of the vector mesons. The condition

$$\langle V(p=0) | \int \theta_{00}(x) d^3x | V(p=0) \rangle = m_V \langle V | V \rangle \quad (2)$$

gives $G_1(0) = -1$.

Similarly the tensor meson matrix element between the vector meson states is given by four coupling constants⁽⁵⁾. Again we will need only one of these:

$$\langle V(p_2) | T | V(p_1) \rangle = \frac{g_{TVV}}{2m_T} \left(\frac{m_V}{m_T}\right)^{-2} \epsilon_1 \cdot \epsilon_2 P_\mu P_\nu \epsilon^{\mu\nu} + \dots \quad (3)$$

$\epsilon^{\mu\nu}$ and m_T are respectively the polarization tensor and the mass of the tensor meson. m_V is the mass of the vector meson and \bar{m} is some average mass. The factor $\frac{\bar{m}^{-2}}{m^2}$ is explicitly extracted from the coupling constants. g_{TVV} will be assumed to obey SU_3 symmetry relations⁽⁶⁾. The tensor-meson couplings to $\theta_{\mu\nu}$ are defined by

$$\langle T | \theta_{\mu\nu} | 0 \rangle = m_T^3 g_T \epsilon_{\mu\nu} \quad (4)$$

As in I and ref. 2, we saturate the form factor $G_1(0)$ with f , f' (or singlet f_1 and octet f_8) and P . Now the nature of the Pomeron has remained quite mysterious over the years. For our purpose we treat it as a factorizable SU_3 singlet Regge pole with a linearly rising trajectory ($\alpha_P = 1 + \alpha'_P t$) and the contribution to G_1 is evaluated by introducing a spin 2 particle with mass $M_P = 1/\sqrt{\alpha'_P}$. The singlet nature and factorization are probably correct within about 20%. To obtain cross-sections, the spin 2 particle pole will be Reggeized. So the actual existence of such a particle may not be crucial. The small rise in total cross-sections found at NAL and ISR energies could be built in if necessary by taking $\alpha_P(0)$ to be slightly larger than 1. Such a procedure has been followed by several authors recently in their fits and the fundamental problem remains common to many such approaches. At any rate we regard the results obtained here and in I as a posteriori justification for such an effective parametrization of the Pomeron contribution.

Then, assuming D type SU_3 symmetric coupling for vector-vector-tensor octet vertex, saturation of $G_1(0)$ leads to the following relations:

$$\frac{m_\rho^2}{-2} = g_P g_{P\rho\rho} + g_1 g_{1\rho\rho} + g_8 g_{8\rho\rho} \quad (5)$$

$$\frac{m_{K^*}^2}{-2} = g_P g_{P\rho\rho} + g_1 g_{1\rho\rho} - \frac{g_8}{2} g_{8\rho\rho} \quad (6)$$

$$\frac{m_{\omega_8}^2}{-2} = g_P g_{P\rho\rho} + g_1 g_{1\rho\rho} - g_8 g_{8\rho\rho} \quad (7)$$

$$\frac{m_{\omega_1}^2}{-2} = g_P g_{P\omega_1\omega_1} + g_1 g_{1\omega_1\omega_1} \quad (8)$$

$$\frac{m_{\omega_1\omega_8}^2}{-2} = g_8 g_{8\omega_1\omega_8} \quad (9)$$

Here $g_{1\rho\rho}$ (g_1) and $g_{8\rho\rho}$ (g_8) are respectively the singlet and octet tensor meson couplings to the ρ -meson⁽⁷⁾ ($\theta_{\mu\nu}$). ω_1 and ω_8 are the singlet and the octet components of the ω and ϕ mesons. Note that Eq. (5), (6) and (7) automatically satisfy the GMO mass formula

$$4m_{K^*}^2 = 3m_{\omega_8}^2 + m_\rho^2 \quad (10)$$

Similar equations can be written for the diagonal representation with physical masses m_ϕ^2 , m_ω^2 .

$$\frac{m_\phi^2}{-2} = g_P g_{P\phi\phi} + g_1 g_{1\phi\phi} + g_8 g_{8\phi\phi} \quad (11)$$

$$\frac{m_\omega^2}{-2} = g_P g_{P\omega\omega} + g_1 g_{1\omega\omega} + g_8 g_{8\omega\omega} \quad (12)$$

The simplest assumption would be that the same matrix

$\begin{pmatrix} \cos \theta_V & \sin \theta_V \\ -\sin \theta_V & \cos \theta_V \end{pmatrix}$ which diagonalizes m^2 in (7), (8) and (9) also diagonalizes all the couplings of P, I and B separately. A more complicated possibility would be that the mixing angles are different and diagonalization is achieved by conspiracy between different terms. We consider the first alternative. Then it readily follows that

$$g_{P\phi\phi} = g_{P\omega\omega} = g_{P\omega_1\omega_1} = g_{P\omega_8\omega_8} \quad (13)$$

$$g_{I\phi\phi} = g_{I\omega\omega} = g_{I\omega_1\omega_1} = g_{I\omega_8\omega_8} \quad (14)$$

$$g_{B\phi\phi} = -g_{B\omega\omega} \cot^2 \theta_V = g_{B\omega_8\omega_8} \left(\frac{\cos^2 \theta_V}{\cos 2\theta_V} \right) \quad (15)$$

We have used the fact that $g_{P\omega_1\omega_8} = g_{I\omega_1\omega_8} = g_{B\omega_1\omega_8} = 0$. Then

$$m_{\omega_1}^2 + m_{\omega_8}^2 = m_{\phi}^2 + m_{\omega}^2 \quad (16)$$

and

$$\frac{m_{\omega_8}^2 - m_{\omega_1}^2}{m_{\phi}^2 - m_{\omega}^2} = \cos 2\theta_V \quad (17)$$

as it should be.

We need one more input to solve the equations for the couplings.

In accordance with Okubo-Zweig-Iizuka (OZI) -rule⁸ we assume that which are supposed to $g_{f'_{pp}} = 0$. OZI rule forbids couplings between particles contain only strange (or charmed) quarks with the ones without these quarks. It seems to be approximately correct for couplings like $\phi\rho\pi$, $f'\pi\pi$, $f'NM$ etc.

In quark model non vanishing values of these couplings (violation of OZI rule) may just mean some small admixture of the other kind of quarks. Later we will consider deviation from the OZI rule. Presently we obtain

$$g_{f_{pp}} = \frac{g_{1pp}}{\cos \theta_T} = \frac{g_{8pp}}{\sin \theta_T} \quad (18)$$

From Eq. (5), (6) and (18) one finds

$$g_{f_{pp}} = \frac{2}{3g_8} \frac{m_p^2 - m_{K^*}^2}{m^2} \quad (19)$$

Now, as in I, principle of universality of scalar and tensor couplings to the stress tensor⁽⁹⁾ fixes g_f , $g_{f'}$, g_p and hence g_1 and g_8 for a given mixing angle θ_T . We take the values of these constants from I. So these are not new parameters here. $g_{f_{pp}}$ is then determined from (19) and $g_{P_{pp}}$, $g_{P_{\phi\phi}}$, $g_{P_{\omega\omega}}$ from (5), (13), and (18).

Other couplings like $g_{f_{\omega\omega}}$, $g_{f'\phi\phi}$ etc. are now readily given by (14) and (15). Since $g_{f'_{pp}} = 0$ and $g_{8\omega_8\omega_8} = -g_{8pp}$ by SU_3 , we get

$$g_{f_{\omega\omega}} = \left[\cos^2 \theta_T + \frac{\sin^2 \theta_T \sin^2 \theta_V}{\cos 2\theta_V} \right] g_{f_{pp}} \quad (20)$$

$$g_{f'\phi\phi} = -\frac{\sin 2\theta_T}{2} \frac{3 \cos^2 \theta_V - 1}{\cos 2\theta_V} g_{f_{pp}} \quad (21)$$

When $\theta_V = \theta_T = 35.3^\circ$ (ideal), (20) and (21) reduce to the well-known relations: $g_{f_{\omega\omega}} = g_{f_{pp}}$ and $g_{f'\phi\phi} = -\sqrt{2}g_{f_{pp}}$. Now we consider some numerical values of the coupling constants in this approach. Taking $g_f = 0.076$, $g_{f'} = -0.064$ and $g_p = 0.097/\sqrt{a'_p}$ (a'_p in GeV^{-2})

0000460012

from paper I, we find, for ideal $f - f'$ mixing $\theta_T = 35.3^\circ$

$$g_{f\rho\rho} = 38.84, g_{\rho\rho\rho} = g_{\rho\omega\omega} = g_{\rho\phi\phi} = 22.02/\sqrt{\alpha'_P} \quad (22)$$

These are independent of the mixing angle θ_V . As in I, however, the results are sensitive to the value of θ_T . This happens because of the large cancellation between g_f and $g_{f'}$ terms in the equation for g_θ . As we have used the OZI rule, it is perhaps more appropriate (although not necessary) to use the ideal (canonical) value for θ_T . But for the sake of completeness, we quote the values for $\theta_T = 31^\circ$ (which is in the range predicted by the mass formulas of the tensor mesons). We have

$$\theta_T = 31^\circ \rightarrow g_{f\rho\rho} = 23.01, g_{\rho\rho\rho} = 9.63/\sqrt{\alpha'_P} \quad (23)$$

To estimate deviation from the OZI rule will be very model dependent. For the ϕ meson the OZI suppression factor $\frac{g_{\phi\rho\pi}}{g_{\omega\rho\pi}}$ is believed to be typically about 1/10. If in the case of f' , there is a similar factor, $\frac{g_{f'\rho\rho}}{g_{f\rho\rho}} = 1/10$. Solving the equations with this input, we find

$$\theta_T = 35.3^\circ \rightarrow g_{f\rho\rho} = 34.02, g_{\rho\rho\rho} = 16.01/\sqrt{\alpha'_P} \quad (24)$$

In this way all the coupling constants can be determined in terms of masses and α'_P . Next we discuss applications of these results.

III. Decays of Tensor Mesons

From the above values of $g_{fV\bar{V}}$ and $g_{f'V\bar{V}}$, all the other $g_{TV\bar{V}}$ couplings also

can be readily determined by using SU_3 symmetry. These couplings become relevant in the models of decays of tensor mesons. Since this involves a detailed program by itself, we hope to present such calculations separately in future. Here we just make few remarks based on comparison with the already existing works. In ref. 4, Renner studied tensor dominance of vector mesons using f and f' intermediate states and determined $g_{f\rho\rho}$. Then using vector dominance he related the other three couplings to $g_{f\rho\rho}$. In our notation he found $g_{f\rho\rho} = 9.44$ which is smaller by a factor of about 2.5 to 4 when compared with our values (Eq. (22), (23) and (24)). He has calculated radiative decay widths of f mesons, for which, however, unfortunately as yet no experimental data are available. But some time in future the situation could change. In a recent work Levy, Singer and Toaff⁽¹⁰⁾ consider dual amplitudes for pseudoscalar-vector scattering and obtain $g_{TV\bar{V}}$ couplings. In spite of the fact that their model is entirely different from that of Renner, the values of the dominant couplings obtained were very close in the two models. It appears that the different values obtained for $g_{f\rho\rho}$ in the present model stems from the breaking of duality by the introduction of the Pomeron. With their values of $g_{f\rho\rho}$ (comparable to Renner's) authors of ref. 10 calculate the decay width of $f \rightarrow \rho^0 \pi^+ \pi^-$ to be 0.49 MeV. Experimental decay width for $f \rightarrow \pi^+ \pi^- \pi^0$ is given to be 6.2 ± 1.5 MeV. Even granting that the latter width includes more than just $\rho\pi\pi$ events, there is a discrepancy by a factor of about 6 to 10. It is very interesting that the larger value of $g_{f\rho\rho}$ required to remove this discrepancy is indeed provided by the present model. Similarly Γ theory for $A_2 \rightarrow \omega \pi^+ \pi^-$ comes out to be smaller than

Γ_{expt} by a factor of about 4 in ref. 10. Also in another recent work Novikov and Eidelman⁽¹¹⁾ estimate g_{fpp} by assuming that the entire $f \rightarrow 2\pi^+ 2\pi^-$ decay proceeds through $\rho^0 \rho^0$ intermediate states and obtain a value which is about a factor of 2 larger than that of Renner. The motivation in ref. 11 was to estimate the charge asymmetry of π mesons in $e^+e^- \rightarrow \pi^+\pi^-$ reaction for which g_{fpp} is clearly relevant. Thus the coupling constants g_{TVV} obtained here, which are larger than the ones determined by dual or f-f' tensor dominance models, seem to be required by the experimental data. Now we consider the vector meson-nucleon total cross-sections.

IV. Vector Meson-Nucleon Cross-Sections

The couplings constant g_{pVV} is clearly relevant to the calculation of vector meson-nucleon total cross-sections for which data are becoming available. We evaluate the cross-sections in a very simple model in which the spin-2 contribution is Reggeized by assuming structureless residue functions. We consider only the asymptotic Pomeron contribution although the present model also makes definite predictions for the secondary contributions from f and f'.

Let $A(s,t)$ be the spin averaged amplitude for V-N scattering $(V_1(p_1, \epsilon_1) + N_1(q_1) + V_2(p_2, \epsilon_2) + N_2(q_2))$ as $s \rightarrow \infty$. Only the first term in the vertex function (3) contributes as $t \rightarrow 0$ ($p_1 = p_2, \epsilon_1 = \epsilon_2$). The PNM vertex is taken from (1) (Eq. (3))

$$\langle N(q_2) | T | N(q_1) \rangle = \frac{g_{PNN}}{4m_T} \left(\frac{m_B}{m_N}\right) \epsilon^{\mu\nu} \bar{u}(q_2) (\gamma_\mu Q_\nu + \gamma_\nu Q_\mu) u(q_1) \quad (25)$$

where $Q = q_1 + q_2$.

Then

$$A(s,t) = \frac{g_{PVV} g_{PNN}}{m_p^2} \left(\frac{m_B}{m_N}\right) \left(\frac{m}{m_V}\right)^2 \frac{2s^2}{m_p^2 - t} \quad (26)$$

Reggeizing

$$\frac{s^2}{m_p^2 - t} \rightarrow \frac{-\pi \alpha_p' s^2 (1 + e^{-i\pi \alpha_p})}{2 \sin \pi \alpha_p} \left(\frac{s}{s_0}\right) \alpha_p(t)$$

Now the optical theorem

$$\sigma_{VN}^{\text{tot}} = \frac{1}{s} \text{Im} A(s,0) \quad (27)$$

gives

$$\sigma_{VN}^{\text{tot}} = \frac{\pi}{2} s_0 \alpha_p' (\sqrt{\alpha_p'} g_{PVV}) (\sqrt{\alpha_p'} g_{PNN}) \left(\frac{m_B}{m_N}\right) \left(\frac{m}{m_V}\right)^2 \quad (28)$$

where we have used $M_p^2 = 1/\alpha_p'$.

Similarly, nucleon-nucleon total cross-section is given by

$$\sigma_{NN}^{\text{tot}} = \frac{\pi}{2} s_0 \alpha_p' (\sqrt{\alpha_p'} g_{PNN})^2 \left(\frac{m_B}{m_N}\right)^2 \quad (29)$$

g's have been combined with $\sqrt{\alpha_p'}$ since their product is determined by our tensor-dominance relations. As usual $s_0 = 1 \text{ GeV}^2$. All the coupling constants have been already determined. So α_p' remains the only free parameter. This can be fixed from say σ_{NN}^{tot} . Then Eq. (28) gives absolute predictions for σ_{VN}^{tot} .

So far we have considered only the well established SU_3 nonet of vector mesons (ρ, K^*, ω, ϕ). Recently, however some new vector mesons like ψ, D^*, ψ' etc. have been discovered. All of these could be part of some SU_4 multiplets like 1, 15 etc. One can extend the present formalism

to such cases. However, in view of the current uncertainty in SU_4 mass formulas we limit our discussion to some general remarks. By OZI rule presumably f and f' will not contribute to the mass equation for say ψ meson but the corresponding charmed tensor mesons will. Their contributions could be similarly evaluated. Now from (13) we see that the values of $g_{\rho\rho\rho}$, $g_{\rho\omega\omega}$ and $g_{\rho\phi\phi}$ are same in spite of mass breaking and $\phi - \omega$ mixing. So it is plausible that $g_{\rho\psi\psi}$ (or $g_{\rho D^*D^*}$) will also have comparable value. The main difference in the effective coupling will then arise from the factor $1/m_V^2$. An immediate interesting consequence from Eq. (28) is that σ_{VN} will go like $1/m_V^2$. This fact seems to be consistent with the experiments. Vector-meson-nucleon total cross sections have been determined by studying the A -dependence of the diffractive photo-production of vector mesons on different nuclear targets. There are a number of uncertainties. But the typical current values are (12)

$$\sigma_{\rho N} = \sigma_{\omega N} = 25 - 30 \text{ mb}, \quad \sigma_{\phi N} = 13 \text{ mb} \quad (30)$$

$$\sigma_{\psi N} = 3.48 \pm 0.79 \text{ mb}$$

since $\frac{m_\rho^2}{m_\omega^2} = 0.97$, $\frac{m_\rho^2}{m_\phi^2} = 0.57$ and $\frac{m_\rho^2}{m_\psi^2} = 0.062$, the agreement is very good, considering the simplicity of the model. In particular the present model does explain smaller value of $\sigma_{\psi N}$ and a drastically smaller value for $\sigma_{\psi N}$ as compared to $\sigma_{\rho N}$. Similar arguments would lead to

$$\frac{\sigma_{D^*N}}{\sigma_{\rho N}} = 0.15 \text{ and } \frac{\sigma_{\psi'N}}{\sigma_{\rho N}} = 0.044, \text{ assuming that } P \text{ couplings without the mass}$$

factors are same. It will be interesting to test these predictions when data become available.

While this work was being completed, we became aware of a recent work by Carlson and Freund⁽¹³⁾ in which such a $1/m_V^2$ dependence is obtained from the assumptions of tensor meson-dominated Pomeron⁽¹⁴⁾ exchange degeneracy and equality of slopes of the vector meson trajectories ($\alpha'_\rho = \alpha'_\phi = \alpha'_\psi$). In our model such a dependence, however, comes from an entirely different assumption of quadratic mass formulas for vector mesons. Also in the tensor meson-dominated Pomeron model, the following equality holds good.

$$\frac{g_{f\rho\rho}}{g_{\rho\rho\rho}} = \frac{g_{fNN}}{g_{PNN}} \quad (31)$$

Comparing the values of f and P couplings found in this paper with the values from I, we find that the equality is satisfied very well by the couplings for any value of θ_V . In our model however the couplings are determined by the masses. Looking at the equations, it is seen that the result comes out because the experimental masses satisfy the relation:

$$\frac{\bar{m}^2}{m_\rho^2} = \frac{\bar{m}_B}{m_N} \text{ to an extremely good accuracy } (\bar{m}^2 \text{ and } \bar{m}_B \text{ are respectively}$$

the average (mass)² of the vector octet and the average mass of the baryon octet and not just the scale factors introduced in Eq. (3) and (25)). Thus, although the results are similar, the two models obtain them in an entirely different manner. In the present model there is no constraint on the Pomeron couplings except the satisfaction of the mass relations.

Now, relations between cross-sections are given by models such as quark model. But the present approach gives also the absolute magnitude of the cross-sections. To give some numerical examples, let $\theta_T = 35.3^\circ$ ($g_{f'pp} = g_{f'NN} = 0$). Then Eq. (19) of I gives $g_{pNN} = 21.6/\sqrt{\alpha'_p}$. Taking g_{ppp} from Eq. (22) of the present work we have

$$\begin{aligned}\sigma_{\rho N} &= 438.3 \alpha'_p \text{ mb } (\alpha'_p \text{ in GeV}^{-2}) \\ \sigma_{NN} &= 428.5 \alpha'_p \text{ mb}\end{aligned}\quad (32)$$

Taking σ_{NN} to be 39 mb we have $\alpha'_p = 0.09 \text{ GeV}^{-2}$

Hence

$$\begin{aligned}\sigma_{\rho N} &= 39.9 \text{ mb}, \quad \sigma_{\omega N} = 38.7 \text{ mb}, \quad \sigma_{\phi N} = 22.7 \text{ mb} \\ \text{and } \sigma_{\psi N} &= 2.5 \text{ mb}\end{aligned}\quad (33)$$

On the other hand, if we have $\frac{g_{f'pp}}{g_{fpp}} = 1/10$ while still taking $g_{f'NN} = 0$ and $\theta_T = 35.3^\circ$, Eq. (24) gives

$$\sigma_{\rho N} = 29.0 \text{ mb}, \quad \sigma_{\omega N} = 28.1 \text{ mb}, \quad \sigma_{\phi N} = 16.5 \text{ mb} \text{ and } \sigma_{\psi N} = 1.8 \text{ mb} \quad (34)$$

Also because the ratios in (31) remain equal as we change θ_T , $\frac{\sigma_{\rho N}}{\sigma_{NN}}$ has the same value for $\theta_T = 35.3^\circ$ and 31° , only difference being that

$\sigma_{NN} = 39 \text{ mb}$ would need $\alpha'_p = 0.48 \text{ GeV}^{-2}$ in the second case.

In the case of OZI violation, even if $\frac{g_{f'pp}}{g_{fpp}} = \frac{g_{f'NN}}{g_{fNN}}$, $\sigma_{\rho N}$ will be smaller than σ_{NN} if the value of α'_p for the vector case is smaller than that in the nucleon case. Although this is not particularly appealing for the interpretation of a Pomeron trajectory, it should be noted that, in data fitting⁽¹²⁾, α'_p obtained from photo-production of vector mesons is found to be smaller than the one obtained from the

nucleon-nucleon scattering. In addition, some deviations from the universal values of g_f , $g_{f'}$, g_p and F_σ (see I) will also result in changes in absolute magnitudes of the cross-sections.

It is also interesting to note that the linear mass formulas do not work. If we use the factor $\left(\frac{m}{m_V}\right)$ in Eq. (3) instead of $\left(\frac{m}{m_V}\right)^2$, we find that

$$\sigma_{\rho N} : \sigma_{\phi N} : \sigma_{\psi N} : \frac{1}{m_\rho} : \frac{1}{m_\phi} : \frac{1}{m_\psi} = 4:3:1 \quad (35)$$

which is in clear contradiction with the data. In addition, the ratio

$\frac{\sigma_{\rho N}}{\sigma_{NN}}$ comes out to be 0.3 which is also completely off. Thus we can regard the present results as supporting the case for quadratic mass formulas. Because of the ω - ϕ mixing problem, from masses alone one can not distinguish between the linear and the quadratic forms.

In the above we have considered $\sigma_{\psi N}$ and σ_{NN} . However recently, some data on Σ -N and Ξ -N scattering have also become available^(15,16). The model in I, gives for the asymptotic (Pomeron) contributions,

$$\frac{\sigma_{\Sigma N}}{\sigma_{NN}} = \frac{m_N}{m_\Sigma} = 0.78 \quad \text{and} \quad \frac{\sigma_{\Xi N}}{\sigma_{NN}} = \frac{m_N}{m_\Xi} = 0.71 \quad (36)$$

The currently available data^(15,16) indicate that

$$\frac{\sigma_{\Sigma-P}}{\sigma_{pp}} = 0.85 - 0.87 \quad \text{and} \quad \frac{\sigma_{\Xi-P}}{\sigma_{pp}} \approx 0.7 \quad (37)$$

Thus the data are quite consistent with our prediction that

$\sigma_{\Xi N} < \sigma_{\Sigma N} < \sigma_{NN}$ and even the $1/m_B$ dependence also could be approximately true.

V. Concluding Remarks

It might be argued that, because of the relative crudeness with

which the Pomeron contribution has been handled, the striking numerical results may not have particular significance. Even if this is true, some general results should remain valid. In particular, the $1/m_V^2$ behavior of σ_{VN} and the fact that σ_{NN} and σ_{VN} come out to have reasonable magnitude with the usually accepted small values of α'_p ($< 0.5 \text{ GeV}^{-2}$) should be regarded as successes of this approach. Another significant prediction lies in the larger value of g_{TVV} obtained in the present work, as compared to the dual or non-Pomeron tensor dominance models. As we have mentioned in the text the data on tensor meson decays seem to be requiring such larger values. Also, as mentioned in Sec IV, $\sigma_{BN} \propto 1/m_B$ (Baryon mass) may be also approximately true. The results of this paper taken together with paper I, lead us to conclude that the dominant matrix elements of $\theta_{\mu\nu}$ between vector mesons and baryons can be rather well saturated with the known tensor mesons and the Pomeron without requiring any subtraction or some hypothetical mesons. Furthermore this procedure does provide insight into the dynamics of symmetry breaking in masses and leads to a number of experimentally verifiable predictions. As we have already remarked in I, the saturation procedure does not work in the case of pseudoscalar mesons because of the large amount of symmetry breaking in masses. The mass relations probably need subtraction constants or extra tensor mesons. In ref. 2, the masses were absorbed in the definition of the coupling constants and the satisfaction of mass formulas was not required. Then, using non universal values for g_f , g_p etc. consistent solutions were obtained. In ref. 3 (I) and here satisfaction of the mass formulas has been the starting point. It should be noted that similar difficulty for the pseudo scalar case appears in the study of scalar dominance⁽¹⁷⁾.

Finally, as for the role of the Pomeron in the present saturation scheme, one can simply regard it as the contribution which remains after the usual tensor meson pole contributions are taken out from the particle-antiparticle cross-channel amplitude. It would then include continuum also. Of course, factorization will be difficult to understand in such a case. However possibly an effective pole somehow takes care of such contributions. In such a scheme, then, masses of the particles can be consistently expressed as dynamical quantities related to the tensor meson and Pomeron exchange forces.

VI. Acknowledgements

The author wishes to thank Prof. S. Mandelstam and Prof. J. D. Jackson for their kind hospitality at the Physics Dept. and the Lawrence Berkeley Laboratory of University of California at Berkeley where this work was done. He is also thankful to Prof. M. Suzuki for helpful discussions and to the N.S.F. for the summer support.

References and Footnotes

[†]This work has been supported in part by the National Science Foundation.

^{*}Permanent address, on sabbatical leave fall semester 1976.

1. P. G. O. Freund, Phys. Lett. 2, 136 (1962); R. Delbourgo, A. Salam and J. Strathdee, Nuovo Cimento 49A, 593 (1967). H. Pagels Phys. Rev. 144, 1250 (1966) and Univ. of North Carolina technical report (unpublished); B. Renner, Phys. Lett. 33B, 599 (1970); K. Raman, Phys. Rev. D3, 2900 (1971).
2. L. R. Ram Mohan and K. V. Vasavada, Phys. Rev. D9, 2627 (1974).
3. K. V. Vasavada, Phys. Rev. D12, 2918 (1975). This paper is referred to as I in the text.
4. See, for example, B. Renner, Nuclear Physics B30, 634 (1971). A factor of 2 is introduced here for convenience. The full matrix element, for completeness, is given by

$$\begin{aligned}
 2\langle V_2(p_2) | \theta_{\mu\nu} | V_1(p_1) \rangle &= g_1(q^2) \epsilon_1 \cdot \epsilon_2 P_\mu P_\nu \\
 &+ g_2(q^2) (\epsilon_1 \cdot P)(\epsilon_2 \cdot P) P_\mu P_\nu \\
 &+ g_3(q^2) [(\epsilon_1 \cdot P)\epsilon_{2\mu} P_\nu + (\epsilon_1 \cdot P)\epsilon_{2\nu} P_\mu \\
 &\quad + (\epsilon_2 \cdot P)\epsilon_{1\mu} P_\nu + (\epsilon_2 \cdot P)\epsilon_{1\nu} P_\mu] \\
 &+ g_4(q^2) [(\epsilon_1 \cdot q)\epsilon_{2\mu} q_\nu + (\epsilon_1 \cdot q)\epsilon_{2\nu} q_\mu \\
 &\quad + (\epsilon_2 \cdot q)\epsilon_{1\mu} q_\nu + (\epsilon_2 \cdot q)\epsilon_{1\nu} q_\mu]
 \end{aligned}$$

$$- 2(\epsilon_1 \cdot q)(\epsilon_2 \cdot q)g_{\mu\nu} - q^2(\epsilon_{1\mu}\epsilon_{2\nu} + \epsilon_{2\mu}\epsilon_{1\nu})$$

$$+ g_5(q^2)(\epsilon_1 \cdot \epsilon_2)(q_\mu q_\nu - q^2 g_{\mu\nu})$$

$$+ g_6(q^2)(\epsilon_1 \cdot P)(\epsilon_2 \cdot P)(q_\mu q_\nu - q^2 g_{\mu\nu})$$

5. As in ref. 4, the full matrix element in our notation, is given by

$$\begin{aligned}
 \langle V(p_2) | T | V(p_1) \rangle &= \frac{e^{i\nu}}{2} [\] , \text{ where} \\
 [\] &= \frac{g^{(1)}_{TVV}}{m_T} (\epsilon_1 \cdot \epsilon_2) P_\mu P_\nu + \frac{g^{(2)}_{TVV}}{m_T} (\epsilon_1 \cdot P)(\epsilon_2 \cdot P) P_\mu P_\nu \\
 &+ \frac{g^{(3)}_{TVV}}{m_T} [(\epsilon_1 \cdot P)\epsilon_{2\mu} P_\nu + (\epsilon_1 \cdot P)\epsilon_{2\nu} P_\mu \\
 &\quad + (\epsilon_2 \cdot P)\epsilon_{1\mu} P_\nu + (\epsilon_2 \cdot P)\epsilon_{1\nu} P_\mu] \\
 &+ \frac{g^{(4)}_{TVV}}{m_T} [(\epsilon_1 \cdot q)\epsilon_{2\mu} q_\nu + (\epsilon_1 \cdot q)\epsilon_{2\nu} q_\mu + (\epsilon_2 \cdot q)\epsilon_{1\mu} q_\nu \\
 &\quad + (\epsilon_2 \cdot q)\epsilon_{1\nu} q_\mu - 2(\epsilon_1 \cdot q)(\epsilon_2 \cdot q)g_{\mu\nu} \\
 &\quad - q^2(\epsilon_{1\mu}\epsilon_{2\nu} + \epsilon_{1\nu}\epsilon_{2\mu})]
 \end{aligned}$$

In the present paper only $g^{(1)}_{TVV}$ is needed and the superscript is dropped.

6. As it was emphasized in I, the question as to which coupling constants obey SU_3 symmetry is entirely a dynamical one and can not be

9 0 0 0 4 6 0 0 1 3

settled a priori without appealing to the experiments. The definition (3) will give rise to GMO mass formulas for $(\text{mass})^2$ of the vector mesons. Previous authors like Renner in ref. 4 did not define their couplings in this manner and accordingly did not get the mass formulas. If we pull out (\bar{m}/m_V) , linear mass formulas will be obtained. This fact has nothing to do with the kind of normalization one uses for the states. m^{-2} is introduced to make the couplings dimensionless and is taken to be 0.728 GeV^2 (average mass² of the vector octet). However it can be chosen as completely arbitrary since only the product of g's with m^{-2} will appear in any application.

7. We use the convention : $f = f_1 \cos \theta_T + f_8 \sin \theta_T$,
 $f' = -f_1 \sin \theta_T + f_8 \cos \theta_T$
 $\omega = \omega_1 \cos \theta_V + \omega_8 \sin \theta_V$, $\phi = -\omega_1 \sin \theta_V + \omega_8 \cos \theta_V$
8. S. Okubo, Phys. Lett. 5, 165 (1963); G. Zweig, CERN Report No. 8419/TH 412 1964 (unpublished); J. Iizuka, Suppl. Prog. Theor. Phys. 37-38, 21 (1966).
9. P. Nath, R. Arnowitt and M. H. Friedman, Phys. Rev. D6, 1572 (1972); Phys. Lett. 42B, 361 (1972).
10. N. Levy, P. Singer and S. Toaff, Phys. Rev. D13, 2662 (1976).
11. V. N. Novikov and S. I. Eldelman, Sov. J. Nucl. Phys. 21, 529 (1976).
12. A. Silverman, Proceedings of the 1975 International Symposium on Lepton and Photon Interactions at High Energies, Stanford, edited by W. I. Kirk (SLAC, Stanford, 1976); D. M. Ritson, SLAC-PUB-1728, 1976 (unpublished)
13. C. E. Carlson and P. G. O. Freund, Phys. Rev. D11, 2453 (1975);
 C. E. Carlson and P. G. O. Freund, Phys. Lett. 39B, 349 (1972). A

number of other authors have also used similar models to explain suppression of $\sigma_{\phi N}$ and $\sigma_{\psi N}$ relative to $\sigma_{\rho N}$. Some of these are: Chan Hong-Mo, J. Kwiecinski and R. G. Roberts, Phys. Lett. 60B, 367 (1976); C. Rosenzweig and G. F. Chew, Phys. Lett. 58B 93 (1975); T. Inami, Phys. Lett. 56B, 291 (1975); N. Papadopoulos, C. Schmid, C. Sorensen and D. M. Webber, Nucl. Phys. B101, 189 (1975). It is also interesting to note that an entirely different model-the generalized vector dominance model also gives $1/m_V^2$ dependence of $\sigma_{\nu N}^{\text{tot}}$ (transverse part) when scaling behavior is imposed on e^+e^- annihilation cross-section and the transverse virtual photon total cross-section. See, for example, S. Chavin and J. D. Sullivan, Phys. Rev. D13, 2990 (1976) which gives earlier references.

14. R. Carlitz, M. B. Green and A. Zee Phys. Rev. D4, 3439 (1971).
15. J. Badler et al, Phys. Lett. 41B, 387 (1972).
16. R. Majka et al, Phys. Rev. Lett. 37, 413 (1976). Although the data sample for Ξ^- is too small, a rough value for $\sigma_{\Xi^- p}^{\text{tot}}$ can be determined by using the value of $b_{\Xi^- p}$ and the optical model relation

$$\frac{b_{\Xi^- p}}{b_{pp}} = \frac{\sigma_{\Xi^- p}^{\text{tot}}}{\sigma_{pp}^{\text{tot}}}$$
 given in this reference. This relation has been shown to be consistent with the data on $\Xi^- p$ scattering.
17. See for example, M. Gell-Mann, in Proceedings of the Third Hawaii Topical Conference on Particle Physics. (Western Periodicals, North Hollywood, Calif. 1969); P. Carruthers, Phys. Rev. D2, 2265 (1970) and Phys. Rev. D3, 959 (1971).

This report was done with support from the United States Energy Research and Development Administration. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the United States Energy Research and Development Administration.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720