

UCLA

UCLA Previously Published Works

Title

Early diagnosis of Alzheimer's disease: update on combining genetic and brain-imaging measures

Permalink

<https://escholarship.org/uc/item/5gt3278m>

Journal

Dialogues in Clinical Neuroscience, 2(3)

ISSN

1294-8322

Author

Small, Gary W

Publication Date

2000-09-30

DOI

10.31887/dcns.2000.2.3/gsmall

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial-NoDerivatives License, available at

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

Peer reviewed

Clinical research

Early diagnosis of Alzheimer's disease: update on combining genetic and brain-imaging measures

Gary W. Small, MD



Diagnosis of Alzheimer's disease is often missed or delayed in clinical practice; thus, methods to improve early detection would provide opportunities for early intervention, symptomatic treatment, and improved patient function. Emerging data suggest that the disease process begins years before clinical diagnostic confirmation. This paper reviews current research focusing on methods for more specific and sensitive early detection using measures of genetic risk for Alzheimer's disease and functional brain imaging. This approach aims to identify patients in a presymptomatic stage for early treatment to delay progressive cognitive decline and disease onset.

Advances in medical technology have led to an aging population. A major impact of this “age revolution” is a dramatic increase in persons afflicted with Alzheimer's disease (AD), the most common form of late-life mental decline. In the United States, the disease strikes approximately 4 million persons.¹ Patients and

their caregivers suffer emotionally and bear a large proportion of the economic burden, estimated to approach \$90 billion each year.²

Despite the prevalence of AD, the diagnosis is often overlooked or mistaken.³ Barrett and associates⁴ reported that as few as 40% of primary care physicians even knew that AD was the most common cause of dementia in late life. Several types of diagnostic error may occur, including incorrectly applying a dementia diagnosis, positively diagnosing the disease when, in fact, it is not present, or not recognizing dementia when it is present. Such errors may result from a lack of attention to cognitive functioning in routine medical examinations and to misperceptions about the normal aging process.⁵ Early AD detection advances would not only offer presymptomatic disease recognition but likely improve diagnostic accuracy once cognitive impairment progresses to a dementia diagnosis.

Although no cure exists for AD, early disease detection offers a number of potential benefits. Accurate differential diagnosis will identify patients with depression, thyroid disease, or other treatable conditions. Once diagnosed, AD patients can then receive treatment for cognitive losses as well as associated behavioral problems. Drug treatments for cognitive impairment and non-pharmacological and pharmacological treatments for the behavioral problems associated with dementia can also enhance quality of life.¹

Several lines of research suggest that AD actually begins years before its clinical manifestations are obvious. Positron emission tomography (PET) studies of glucose metabolism combined with genetic risk assessment show

Keywords: Alzheimer's disease; brain imaging; genetic risk; positron emission tomography; magnetic resonance imaging

Author affiliations: Department of Psychiatry and Biobehavioral Sciences and the Center on Aging, University of California, Los Angeles (UCLA) School of Medicine; and the Veterans Affairs Medical Center, West Los Angeles, Calif, USA

Address for correspondence: Dr Gary W. Small, UCLA Neuropsychiatric Institute, 760 Westwood Plaza, Los Angeles, CA 90024-1759, USA (e-mail: gsmall@mednet.ucla.edu)

Clinical research

Selected abbreviations and acronyms

AD	<i>Alzheimer's disease</i>
FDG-PET	<i>fluorodeoxyglucose positron emission tomography</i>
fMRI	<i>functional magnetic resonance imaging</i>
MCI	<i>mild cognitive impairment</i>
PET	<i>positron emission tomography</i>
SP	<i>senile plaque</i>
NFT	<i>neurofibrillary tangle</i>
NP	<i>neuritic plaque</i>

regional glucose abnormalities in middle-aged persons with a genetic risk for AD.⁶ Studies of structural images suggest that regional atrophy of hippocampus and other medial temporal regions may be an early predictor of future cognitive decline.⁷ Brain autopsy studies of normal aging and older persons with mild cognitive impairment also indicate very early, preclinical accumulation of neuritic plaques (NPs) and neurofibrillary tangles (NFTs), the neuropathological hallmarks of AD, years before a clinical diagnosis can be confirmed.^{8,9} Finally, findings from a study of 93 nuns also support the notion of subtle preclinical functional abnormalities. In that study,¹⁰ a systemic assessment of these nuns' early autobiographies (mean age 22 years) and their later (age 75-95) cognitive performances found that low idea density and lack of grammatical complexity in early life predicted low cognitive test scores in late life.

Identifying AD in a preclinical state before the condition can be confirmed using consensus diagnostic criteria also has several potential benefits. When early detection assessments are negative, people with mild memory complaints can be reassured that their forgetfulness reflects a normal age-related change that probably will not progress. In addition, many people would like to know about a poor prognosis while still in a mildly impaired state in order to plan their futures while mental faculties remain. The most compelling argument for preclinical detection strategies is to identify candidates for antidementia treatments before extensive neuronal death develops, since new antidementia treatments are more likely to delay disease progression than to reverse neuronal death. Patients with mild memory losses, who are at risk for AD progression are ideal candidates for antidementia interventions. Although current cholinergic treatments result in symptomatic rather than disease-altering or structural effects, it would certainly be of




Figure 1. Hypothetical decline curve for two patients with Alzheimer's disease, one who received treatment (dotted line) another who did not (solid line). A major goal of early detection is to intervene early in the course of the disease, even in presymptomatic stages.

interest to initiate treatments very early, as disease-modifying interventions emerge. Moreover, both the expense and potential risks of treatment make it reasonable to reserve treatment only for those people who are at the greatest risk for developing the disease (*Figure 1*).

Our group has focused on early detection of AD by combining measures of genetic risk and brain imaging. This paper summarizes information on genetic discoveries and recent work using three different brain-imaging approaches: measures of glucose metabolism using fluorodeoxyglucose positron emission tomography (FDG-PET) during mental rest, functional magnetic resonance imaging (fMRI) during memory performance tasks, and in vivo imaging of NPs and NFTs using PET and small molecule probes (*Figure 2*).

Genetic issues

Genetic studies have found an association between the apolipoprotein E-4 (*APOE-4*) allele on chromosome 19 and the common form of AD that begins after age 60.¹¹ *APOE* has three allelic variants (2, 3, and 4) and five common genotypes (2/3, 3/3, 2/4, 3/4, and 4/4). The *APOE-4* allele has a dose-related effect on increasing risk and lowering onset age for late-onset familial and sporadic AD,^{11,12} while *APOE-2* confers protection.¹³ The *APOE-4* allele is associated with decline in delayed recall in older adults¹⁴ and is a strong predictor of future




Figure 2. Technological developments (blue) offer the potential of identifying Alzheimer's disease prior to the time point when clinicians currently arrive at a clinical diagnosis. Combining various approaches is likely to improve early diagnostic accuracy.

cognitive decline in people with mild cognitive impairment (MCI).¹⁵ However, *APOE-4* may have only a modest effect in predicting cognitive decline in many older persons; thus, *APOE* genotype alone is not considered a useful predictor in nondemented persons.¹⁶ The *APOE* gene explains only 50% of the genetic variability in AD,¹⁷ and investigators have searched for other linkage regions¹⁸ and candidate genes.^{19,20}

Combining PET scanning and genetic risk measures

The discovery of the *APOE-4* genetic risk for AD was extremely helpful in our early AD detection efforts with PET imaging. In 1995, our group first reported results combining PET imaging and *APOE* genetic risk in people with a family history of AD.²¹ We studied 12 nondemented relatives with *APOE-4* and 19 relatives without *APOE-4*, and compared them with 7 probable AD patients. "At-risk" subjects had mild memory complaints, normal cognitive performance, and at least two relatives with AD. Subjects with *APOE-4* did not differ from those without *APOE-4* in mean age at exam (56.4 vs 55.5 years) or in neuropsychological performance. Parietal metabolism was significantly lower and left-right parietal asymmetry higher in at-risk subjects with *APOE-4* than in those without *APOE-4*. Patients with

dementia had significantly lower parietal metabolism than did at-risk subjects with *APOE-4*. We concluded that the *APOE-4* allele was associated with reduced cerebral parietal metabolism and increased asymmetry in relatives at risk for AD.

The following year, Reiman and associates²² replicated these results and extended them to other brain regions, including temporal, prefrontal, and posterior cingulate in a study of 11 *APOE-4* homozygotes (4/4 genotype) and 22 *APOE-3* homozygotes (3/3 genotype) of similar ages to those in our own initial study (mid-fifties). They also applied an automated image analysis method, wherein metabolic reductions were standardized using three-dimensional (3D) stereotactic surface projections from FDG-PET scans of AD patients compared with controls. The results from these two studies^{21,22} provided independent confirmation of an association between genetic risk and regional cerebral glucose hypometabolism.

More recently, our group²³ performed FDG-PET scans during mental rest on 65 middle-aged and older persons (mean, SD: 67.3, 9.4 years; range 50-84 years) with mild memory complaints or probable AD.²⁴ Subjects (61 Caucasians, 3 Asians, and 1 Hispanic) were right-handed, in the 50 to 84 year age range (mean, SD: 67.3, 9.4 years), and recruited through newspaper ads and various referrals. Of the 573 persons who volunteered because of minor memory concerns (eg, difficulty remembering names), 508 were excluded for a variety of reasons (eg, medication use, concurrent medical illness). The 27 subjects with age-associated memory impairment (AAMI) and *APOE-4* were matched according to age and educational level to 27 AAMI subjects without *APOE-4*. The 11 AD patients were included as a comparison group without regard to *APOE* status, since cerebral metabolic patterns do not vary according to *APOE* genotype in AD patients.²⁵ Subject groups were similar in mean age at examination, sex ratio, frequency of family history of AD, dementia onset age within families, and educational achievement level. Both verbal (Buschke-Fuld)²⁶ and visual (Benton)²⁷ memory performance scores were significantly lower in the demented group but not significantly different between the two nondemented groups (verbal: $t=0.51$, $df=52$, $P=0.18$; visual: $t=1.05$, $df=52$, $P=0.61$).

Comparisons among the three subject groups (region of interest [ROI] analysis) indicated the lowest metabolic rates in the AD group, intermediate rates in the nondemented group with *APOE-4*, and highest rates in

Clinical research

the nondemented group without *APOE-4*. These differences were bilateral and significant (ANOVAs; $df=2,59$) in inferior parietal (left hemisphere: $F=9.2$, $P=0.0003$; right hemisphere: $F=15.6$, $P<0.0001$), posterior cingulate (left: $F=14.6$, $P<0.0001$; right: $F=17.7$, $P<0.0001$), dorsolateral prefrontal (left: $F=13.7$, $P<0.0001$; right: $F=5.6$, $P=0.006$), and inferior temporal regions (left: $F=4.3$, $P=0.018$; right: $F=3.4$, $P=0.040$). Significant group differences present only in the left hemisphere were found in the medial temporal ($F=4.9$, $P=0.011$) and superior temporal ($F=6.0$, $P=0.004$) regions. Further comparisons between the two nondemented groups indicated significantly lower metabolism in subjects with *APOE-4* in the inferior parietal region for both the right ($t=2.6$, $df=52$, $P=0.011$) and left ($t=2.2$, $df=52$, $P=0.035$) hemispheres compared with those without *APOE-4*. These differences remained significant even if we eliminated from the comparison the two subjects homozygous for *APOE-4*. Statistical parametric mapping (SPM) analysis comparing nondemented groups showed similar results with *APOE-4* subjects having significantly lower metabolism than those without *APOE-4*, particularly in the left inferior parietal, lateral temporal, and posterior cingulate regions.²⁸ The peak voxels were in Brodmann's area 21 at (-68, -38, -16) with a secondary focus at (-70, -48, 0).

Genetic risk and fMRI results

Activation imaging during memory task performance may reveal subtle alterations in brain function, perhaps prior to the emergence of mild memory impairments. This approach has been described as a "cognitive stress test" for the brain. The approach is based on the hypothesis that cognitive activity may uncover subtle brain functional abnormalities not present during resting cognitive states, similar to how the electrocardiogram stress test reveals subtle cardiac abnormalities when the heart is stressed from treadmill exercise. Functional MRI (fMRI) provides measures of relative cerebral blood flow (rCBF) during memory or other cognitive task performance, and has the advantages of high resolution in space and time and lack of radiation exposure.²⁹ Thirty middle-aged and older subjects with mild memory complaints but normal memory performance received *APOE* genotyping. The 14 subjects with the *APOE-4* genetic risk for AD did not differ significantly from the 16 subjects without *APOE-4* group in age, prior educa-

tional achievement, or rates of AD family history. During fMRI scanning on a 3-tesla unit, subjects performed an unrelated paired associate learning task. Brain activation patterns were determined during both learning and retrieval task periods and analyzed using both between-group and within-subject approaches. Compared with subjects without *APOE-4*, those at genetic risk showed significantly greater magnitude and spatial extent of rCBF during memory retrieval in regions affected by AD: left medial temporal and bilateral parietal and prefrontal. Longitudinal data indicated that baseline brain activation correlated with verbal memory decline assessed 2 years later. Relative cerebral blood flow responses to a memory challenge may reflect compensatory cognitive efforts for emerging functional deficits in people at genetic risk for AD. These results suggest that combining brain activation and genetic risk measures may provide information that eventually predicts future cognitive decline.

PET imaging of plaques and tangles in AD

New research is under way to develop additional early detection strategies with greater sensitivity and specificity, including studies aimed at imaging the neuropathological hallmarks of AD. Intraneuronal NFTs and extracellular β -amyloid-rich senile plaques (SPs) have been implicated as central components of the pathogenic cascade in AD, which highlights the importance of non-invasive in vivo assessment of SP and NFT deposition. A hydrophobic radiofluorinated derivative of 1,1-dicyano-2-[6-(dimethylamino)naphthalen-2-yl]propene (FDDNP) was used in conjunction with PET to determine the localization and load of NFTs and SPs in the living brain of AD patients ($n=7$) and controls ($n=3$).³⁰ Fluorescence microscopy also was used to determine SP and NFT binding in AD brain specimens. Greater accumulation and slower FDDNP clearance was observed in SP/NFT-rich brain areas, particularly the hippocampus-amygdala-entorhinal complex, but also temporal and parietal cortex in advanced disease stages. In vitro fluorescence microscopy showed excellent visualization of NFTs, SPs, and diffuse amyloid in AD, matching results with thioflavin T obtained in the same specimens. The availability of this noninvasive technique may allow longitudinal evaluation of SP and NFT deposition, permitting more accurate diagnosis and evaluation of therapies.

Conclusions

Evidence is emerging that suggests the eventual utility of combining genetic risk and brain-imaging measures to identify people with very mild memory loss that may be progressive. With such surrogate markers, clinical trials will be possible to determine efficacy of antidementia treatments in presymptomatic stages. The goal is to delay

progressive memory loss upon dementia onset so that older people can live longer lives with improved functioning and mental capacities. □

Supported by the Montgomery Street Foundation, San Francisco, Calif; the Fran and Ray Stark Foundation Fund for Alzheimer's Disease Research, Los Angeles, Calif; the Department of Energy; NIH grants MH52453, AG10123, AG13308, and the Alzheimer's Association grant IIRG94101. The views expressed are those of the authors and do not necessarily represent those of the Department of Veterans Affairs.

Diagnóstico precoz de la enfermedad de Alzheimer: actualización acerca de la combinación de mediciones genéticas y de imágenes cerebrales

El diagnóstico de la enfermedad de Alzheimer habitualmente es erróneo o se realiza tardíamente en la práctica clínica; por lo tanto, los métodos que ayuden a la detección precoz facilitarán las opciones de una intervención oportuna, de un tratamiento sintomático y de una mejoría en el funcionamiento del paciente. La información reciente sugiere que el proceso patológico comienza años antes de la confirmación del diagnóstico clínico. Este artículo revisa la investigación actual, focalizándose en métodos para una detección precoz más específica y sensible, a través de mediciones del riesgo genético para la enfermedad de Alzheimer y neuroimágenes cerebrales funcionales. Esta aproximación ayuda a identificar pacientes en una etapa presintomática, para un tratamiento precoz que ayude a retardar el deterioro cognitivo progresivo y la aparición de la enfermedad.

Le point sur l'association de l'évaluation génétique et de l'imagerie cérébrale dans le diagnostic précoce de la maladie d'Alzheimer

En pratique clinique, dans la maladie d'Alzheimer, le diagnostic n'est pas toujours posé ou est souvent retardé ; le développement de méthodes diagnostiques précoces pourrait permettre une prise en charge dès le début de la maladie, la prescription d'un traitement symptomatique et l'amélioration de l'état fonctionnel du patient. Les nouvelles données dont nous disposons suggèrent que le processus pathologique débute plusieurs années avant que le diagnostic clinique ne soit confirmé. Cet article concerne les recherches en cours sur des techniques de diagnostic précoce plus spécifiques et plus sensibles associant l'évaluation du risque génétique de développer la maladie et l'imagerie cérébrale. Cette approche a pour objectif d'identifier les patients atteints à un stade présymptomatique afin de les traiter précocement pour retarder l'apparition du déficit cognitif et le début de la maladie.

Clinical research

REFERENCES

1. Small GW, Rabins PV, Barry PP, et al. Diagnosis and treatment of Alzheimer disease and related disorders. Consensus statement of the American Association for Geriatric Psychiatry, the Alzheimer's Association, and the American Geriatrics Society. *JAMA*. 1997;278:1363-1371.
2. Ernst RL, Hay JW. The US economic and social costs of Alzheimer's disease revisited. *Am J Pub Health*. 1994;84:1261-1264.
3. Ross GW, Abbott RD, Petrovitch H, et al. Frequency and characteristics of silent dementia among elderly Japanese-American men: the Honolulu-Asia Aging Study. *JAMA*. 1997;277:800-805.
4. Barrett JJ, Haley WE, Harrell LE, Powers RE. Knowledge about Alzheimer disease among primary care physicians, psychologists, nurses, and social workers. *Alzheimer Dis Assoc Disord*. 1997;11:99-106.
5. Mant A, Eyland EA, Pond DC, Saunders NA, Chancellor AH. Recognition of dementia in general practice: comparison of general practitioners' opinions with assessments using the Mini-Mental State Examination and the Blessed dementia rating scale. *Fam Pract*. 1988;5:184-188.
6. Small GW. Neuroimaging and genetic assessment for early diagnosis of Alzheimer's disease. *J Clin Psychiatry*. 1996;57(suppl 14):9-13.
7. Jack CR, Peterson RC, Xu YC, et al. Medial temporal atrophy on MRI in normal aging and very mild Alzheimer's disease. *Neurology*. 1997;49:786-794.
8. Braak H, Braak E. Neuropathological staging of Alzheimer-related changes. *Acta Neuropathol*. 1991;82:239-259.
9. Price JL, Morris JC. Tangles and plaques in nondemented aging and "pre-clinical" Alzheimer's disease. *Ann Neurol*. 1999;45:358-368.
10. Snowdon DA, Kemper SJ, Mortimer JA, Greiner LH, Wekstein DR, Markesbery WR. Linguistic ability in early life and cognitive function and Alzheimer's disease in late life. Findings from the Nun Study. *JAMA*. 1996;275:528-532.
11. Saunders AM, Strittmatter WJ, Schmechel D, et al. Association of apolipoprotein E allele E4 with late onset familial and sporadic Alzheimer's disease. *Neurology*. 1993;43:1467-1472.
12. Corder EH, Saunders AM, Strittmatter WJ, et al. Gene dose of apolipoprotein E type allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993;261:921-923.
13. Corder EH, Saunders AM, Risch MJ, et al. Apolipoprotein E type 2 allele and the risk of late onset Alzheimer's disease. *Nat Genet*. 1994;7:180-183.
14. O'Hara R, Yesavage JA, Kraemer HC, Mauricio M, Friedman LF, Murphy GMJ. The APOE epsilon4 allele is associated with decline on delayed recall performance in community-dwelling older adults. *J Am Geriatr Soc*. 1998;46:1493-1498.
15. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment: clinical characterization and outcome. *Arch Neurol*. 1999;56:303-308.
16. Relkin NR, Tanzi R, Breitner J, et al. Apolipoprotein E genotyping in Alzheimer's disease: position statement of the National Institute on Aging/Alzheimer's Association Working Group. *Lancet*. 1996;347:1091-1095.
17. Roses AD, Devlin B, Conneally PM, et al. Measuring the genetic contribution of APOE in late-onset Alzheimer disease (AD). *Am J Hum Genet*. 1995;57(suppl):A202.
18. Pericak-Vance MA, Bass MP, Yamaoka LH, et al. Complete genomic screen in late-onset familial Alzheimer disease. Evidence for a new locus on chromosome 12 [see comments]. *JAMA*. 1997;278:1237-1241.
19. Scott WK, Yamaoka LH, Locke PA, et al. No association or linkage between an intronic polymorphism of presenilin-1 and sporadic or late-onset familial Alzheimer disease. *Genet Epidemiol*. 1997;14:307-315.
20. Small GW, Noble EP, Matsuyama SS, et al. D₂ dopamine receptor A1 allele in Alzheimer disease and aging. *Arch Neurol*. 1997;54:281-285.
21. Small GW, Mazziotta JC, Collins MT, et al. Apolipoprotein E type 4 allele and cerebral glucose metabolism in relatives at risk for familial Alzheimer disease. *JAMA*. 1995;273:942-947.
22. Reiman EM, Caselli RJ, Yun LS, et al. Preclinical evidence of Alzheimer's disease in persons homozygous for the epsilon 4 allele for apolipoprotein E [see comments]. *N Engl J Med*. 1996;334:752-758.
23. Small GW, Ercoli LM, Huang SC, et al. PET and genetic risk for Alzheimer disease. *J Nucl Med*. 1999;40(suppl):70p.
24. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984;34:939-944.
25. Corder EH, Jelic V, Basun H, et al. No difference in cerebral glucose metabolism in patients with Alzheimer disease and differing apolipoprotein E genotypes. *Arch Neurol*. 1997;54:273-277.
26. Buschke H, Fuld PA. Evaluating storage, retention and retrieval in disordered memory and learning. *Neurology*. 1974;24:1019-1025.
27. Benton AL. *The Revised Visual Retention Test*. New York: Psychological Corporation; 1974.
28. Silverman DHS, Ercoli LM, Komo S, Huang SC, Phelps ME, Small GW. Influence of single apolipoprotein E type 4 allele on regional brain metabolism of healthy subjects. *J Nucl Med*. 1999;40(suppl):70P.
29. Small GW, Bookheimer SY, Strojwas MH, et al. Functional MRI of memory tasks in older persons with APOE-4. *Biol Psychiatry*. 1999;45:49S.
30. Barrio JR, Huang SC, Cole GM, Satyamurthy N, Petric A, Small GW. PET imaging of tangles and plaques in Alzheimer disease. *J Nucl Med*. 1999;40(suppl):70P-71P.