

Lawrence Berkeley National Laboratory

Recent Work

Title

PREPARATION OF TRITIATED CHOLESTEROL

Permalink

<https://escholarship.org/uc/item/5gx7w5z9>

Authors

Kritchevsky, David
Biggs, Max W.
Freeman, Norman K.

Publication Date

1950-03-29

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

UCRL-644
C.2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

copy 1:

UNIVERSITY OF CALIFORNIA
Radiation Laboratory

Contract No. W-7405-eng-48

UNCLASSIFIED

Preparation of Tritiated Cholesterol

David Kritchevsky, Max W. Biggs, and Norman K. Freeman

March 29, 1950

Berkeley, California

<u>INSTALLATION</u>	<u>No. of Copies</u>
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission, Washington	2
Battelle Memorial Institute	1
Brookhaven National Laboratory	8
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide & Carbon Chemicals Corp. (K-25)	4
Carbide & Carbon Chemicals Corp. (Y-12)	4
Chicago Operations Office	1
Cleveland Area Office	1
Columbia University (Dunning)	2
Columbia University (Failla)	1
Dow Chemical Company	1
General Electric Company, Richland	6
Idaho Operations Office	1
Iowa State College	2
Kansas City	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (Gaudin)	1
Massachusetts Institute of Technology (Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	2
National Bureau of Standards	2
Naval Radiological Defense Laboratory	2
NEPA Project	2
New Brunswick Laboratory	1
New York Operations Office	3
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Advisor, Washington	1
Rand Corporation	1
Sandia Base	1
Sylvania Electric Products, Inc.	1
Technical Information Branch, ORE	15
U. S. Public Health Service	1
UCLA Medical Research Laboratory (Warren)	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University (Friedell)	2
Westinghouse	4
	<hr/>
Total	117
Information Division	
Radiation Laboratory	
Univ. of California	

PREPARATION OF TRITIATED CHOLESTEROL

by

David Kritchevsky, Max W. Biggs(*) and Norman K. Freeman

Radiation Laboratory and Division of Medical Physics,
University of California, Berkeley, California(**)

March 29, 1950

In order to carry out metabolic studies in experimental arteriosclerosis it was desirable to obtain cholesterol labeled with a radioactive atom. Inasmuch as random labeling was sufficient for the purpose of our proposed experiments, preparation of tritiated cholesterol was undertaken.

The method employed by Bloch and Rittenberg (1) for the introduction of deuterium into the cholesterol molecule was used with a few variations. Cholesterol (12.5 gm) was shaken in a sealed tube with acetic acid (40 ml.), reduced platinum oxide (1.25 gm.) and water (13 ml. containing 300 mc. of tritium). The reaction mixture was maintained at 125° C for four days, then the organic material was extracted, saponified with alcoholic KOH and purified by chromatography on an alumina column. The cholesterol was eluted with benzene and benzene-ether 1:1. The amount of cholesterol eluted initially amounted to 7 gm. (56%). The eluted material was purified through the dibromide and several recrystallizations from acetone yielded a product, m.p. 148°, $[\alpha]_D^{25} = 39.3^\circ$ (c, 0.4500 in CHCl₃).

Two grams of the highly purified product were found to contain 0.48 mc. of tritium per gram.

The infrared spectra of the tritiated material and of a sample of purified cholesterol were compared and found to be identical (Fig. 1). The amount of tritium incorporated in the molecule is too small to cause any changes in the infrared spectrum.

-
- (*) Atomic Energy Commission Postdoctoral Research Fellow in the Medical Sciences of the National Research Council.
- (**) The work described in this paper was sponsored by the Atomic Energy Commission.
- (1) Bloch, K. and Rittenberg, D., J. Biol. Chem., 149, 505 (1943).
-

For publication in Science




FIG. 1

Mu 113