

# UC Irvine

## UC Irvine Previously Published Works

### Title

Developing an Internet- and Mobile-Based System to Measure Cigarette Use Among Pacific Islanders: An Ecological Momentary Assessment Study

### Permalink

<https://escholarship.org/uc/item/5j81q5r8>

### Journal

JMIR mHealth and uHealth, 4(1)

### ISSN

2291-5222

### Authors

Pike, James Russell  
Xie, Bin  
Tan, Nasya  
[et al.](#)

### Publication Date

2016

### DOI

10.2196/mhealth.4437

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Original Paper

# Developing an Internet- and Mobile-Based System to Measure Cigarette Use Among Pacific Islanders: An Ecological Momentary Assessment Study

James Russell Pike<sup>1\*</sup>, BA; Bin Xie<sup>1\*</sup>, PhD; Nasya Tan<sup>1</sup>, MPH; Melanie Dee Sabado-Liwag<sup>1</sup>, MPH; Annette Orne<sup>2</sup>; Tupou Toilolo<sup>3</sup>; Steven Cen<sup>4</sup>, PhD; Vanessa May<sup>5</sup>, BA; Cevadne Lee<sup>1</sup>, MPH; Victor Kaiwi Pang<sup>6</sup>, BS; Michelle A Rainer<sup>1</sup>, MSG; Dorothy Etimani S Vaivao<sup>7</sup>; Jonathan Tana Lepule<sup>8</sup>, BA; Sora Park Tanjasiri<sup>8</sup>, MPH, DrPH; Paula Healani Palmer<sup>1</sup>, PhD

<sup>1</sup>School of Community and Global Health, Claremont Graduate University, Claremont, CA, United States

<sup>2</sup>Guam Communications Network, Long Beach, CA, United States

<sup>3</sup>Union of Pan Asian Communities, San Diego, CA, United States

<sup>4</sup>University of Southern California, Los Angeles, CA, United States

<sup>5</sup>Tongan Community Service Center, Hawthorne, CA, United States

<sup>6</sup>Pacific Islander Health Partnership, Santa Ana, CA, United States

<sup>7</sup>Samoan National Nurses Association, Carson, CA, United States

<sup>8</sup>California State University, Fullerton, Fullerton, CA, United States

\*these authors contributed equally

**Corresponding Author:**

James Russell Pike, BA  
School of Community and Global Health  
Claremont Graduate University  
675 West Foothill Boulevard  
Suite 310  
Claremont, CA, 91711-3475  
United States  
Phone: 1 818 406 0286  
Fax: 1 909 607 9655  
Email: [James.Pike@cgu.edu](mailto:James.Pike@cgu.edu)

## Abstract

**Background:** Recent prevalence data indicates that Pacific Islanders living in the United States have disproportionately high smoking rates when compared to the general populace. However, little is known about the factors contributing to tobacco use in this at-risk population. Moreover, few studies have attempted to determine these factors utilizing technology-based assessment techniques.

**Objective:** The objective was to develop a customized Internet-based Ecological Momentary Assessment (EMA) system capable of measuring cigarette use among Pacific Islanders in Southern California. This system integrated the ubiquity of text messaging, the ease of use associated with mobile phone apps, the enhanced functionality offered by Internet-based Cell phone-optimized Assessment Techniques (ICAT), and the high survey completion rates exhibited by EMA studies that used electronic diaries. These features were tested in a feasibility study designed to assess whether Pacific Islanders would respond to this method of measurement and whether the data gathered would lead to novel insights regarding the intrapersonal, social, and ecological factors associated with cigarette use.

**Methods:** 20 young adult smokers in Southern California who self-identified as Pacific Islanders were recruited by 5 community-based organizations to take part in a 7-day EMA study. Participants selected six consecutive two-hour time blocks per day during which they would be willing to receive a text message linking them to an online survey formatted for Web-enabled mobile phones. Both automated reminders and community coaches were used to facilitate survey completion.

**Results:** 720 surveys were completed from 840 survey time blocks, representing a completion rate of 86%. After adjusting for gender, age, and nicotine dependence, feeling happy ( $P<.001$ ) or wanting a cigarette while drinking alcohol ( $P<.001$ ) were

positively associated with cigarette use. Being at home ( $P=.02$ ) or being around people who are not smoking ( $P=.01$ ) were negatively associated with cigarette use.

**Conclusions:** The results of the feasibility study indicate that customized systems can be used to conduct technology-based assessments of tobacco use among Pacific Islanders. Such systems can foster high levels of survey completion and may lead to novel insights for future research and interventions.

(*JMIR mHealth uHealth* 2016;4(1):e2) doi:[10.2196/mhealth.4437](https://doi.org/10.2196/mhealth.4437)

## KEYWORDS

Pacific Islander; tobacco use; cigarette use; mobile phone; text message; ecological momentary assessment

## Introduction

### Assessing Health Disparities among Pacific Islanders

Pacific Islander refers to Chamorros, Marshallese, Native Hawaiians, Samoans, Tongans, and other related groups who share a common origin, culture, and customs. These communities face a wide range of social, economic, and health-related challenges. Educational attainment among Pacific Islanders residing in the United States is low, with only 14.4% obtaining bachelor's degrees as compared to the national average of 27.9% [1]. Per capita income among Pacific Islanders is US \$19,051 whereas the national average is US \$27,334 [1]. Smoking-related conditions, such as cardiovascular and respiratory diseases, are disproportionately high among Pacific Islanders [2]. This fact is often overlooked within epidemiological studies that aggregate Pacific Islander data with those of Asian Americans [3]. Disaggregated Asian-Pacific Islander data from the National Adult Tobacco Survey between 2009 and 2010 revealed past month smoking rates of 20.0% compared with 4.7% for Chinese, 5.5% for Asian Indians, 7.2% for Vietnamese, 13.6% for Filipinos, 15.3% for Koreans, and 18.8% for Japanese [4]. Disaggregated data also indicate that Pacific Islanders have smoking rates of 21.5% for males and 18.4% for females compared with respective averages for the general population of 15.7% and 12.8% [5].

One promising assessment technique that may lead to an improved understanding of the factors that contribute to tobacco use among this population is Ecological Momentary Assessment (EMA). Prior tobacco use research has demonstrated that EMA can generate novel insights for future research and interventions [6-26]. Unfortunately, conducting technology-based research among Pacific Islanders involves numerous challenges. Factors that have hindered past research efforts include Pacific Islanders' broad geographic dispersion and their distrust of academic researchers who often fail to employ culturally-tailored methodologies [27,28]. The objective of the current feasibility study was to overcome these barriers by developing a customized EMA system that facilitated high survey completion rates and fostered new insights into the factors that contribute to cigarette use in this at-risk population.


### Developing an EMA System for Pacific Islanders

EMA is a technique that involves the repeated sampling of participants' behaviors and experiences in real time within their

natural environment [6]. EMA has been used for more than twenty years to measure such behaviors as smoking [6-26,29,30], exercise [31-37], diet [38-42], substance use [43-50], and health information seeking [51]. Historically, these studies have been conducted using electronic diaries [7,9,13-21,24,30,34,39,40,52] that often facilitate survey completion rates greater than 85% [7,13,14,17] but also require extensive in-person training [40,52]. More recently, researchers have reduced respondent burden by utilizing mobile phone apps [53]. However, these programs are often restricted to specific operating systems [11,25,26,31,35,36,50] and are not always programmed to facilitate real-time, remote monitoring of participant responses [31,35,36,50]. Such barriers can be overcome through text messaging [8,44,51,54-56] which is not linked to a specific operating system and typically has higher completion rates. The drawback is that text messaging uses an open-ended format that permits nonstandard responses that can require extensive, time-consuming data cleaning [54]. Another alternative is to employ Internet-based Cell phone-optimized Assessment Techniques (ICAT) to administer online surveys through the Internet browser of Web-enabled mobile phones. This technique is becoming increasingly popular, yet many studies that utilize this approach fail to achieve survey completion rates above 55% [10,29,57]. Moreover, recent ICAT studies of tobacco use [10,12,29] have not capitalized on the unique features offered by this approach, including the ability to use participant responses from earlier in the day to generate tailored survey questions.

In 2011, the Weaving an Islander Network for Cancer Awareness, Research, and Training (WINCART) Center set out to create a customized EMA system that integrated the ubiquity of text messaging, the ease of use associated with mobile phone apps, the enhanced functionality offered by ICAT systems, and the high survey completion rates exhibited by EMA studies that utilized electronic diaries. The WINCART Center is a community-based participatory research [58] consortium that endeavors to reduce cancer health disparities among Pacific Islanders [59] throughout Southern California (see Figure 1). Over a period of several months, members of the consortium collaboratively developed a culturally-tailored system. A feasibility study was then conducted to determine whether young, adult Pacific Islander smokers would be able to effectively utilize the system and whether the resulting data would offer new insights into the intrapersonal, social, and ecological factors associated with cigarette use.

**Figure 1.** Geographic distribution of Pacific Islanders in Southern California.



## Methods

### Participants

Using strategies employed in prior studies [60], 5 community-based organizations recruited 208 young adults who self-identified as Chamorro, Marshallese, Native Hawaiian, Samoan, Tongan, or other Pacific Islander. A 12-item screening survey administered in person or over the phone was used to select 61 current smokers who (a) were between 18 and 29 years old, (b) resided in Southern California, and (c) had smoked at least 100 cigarettes in their lifetime. Trained research staff met with each participant at a mutually agreed-upon location. During this meeting, written consent was obtained using a protocol approved by the Institutional Review Boards at Claremont Graduate University and California State University, Fullerton. The participant completed a computer-based questionnaire that assessed their demographic characteristics, tobacco use behavior, and nicotine dependence [61-63]. Research staff also conducted

a brief, one-to-one semi-structured interview about the intrapersonal, social, and ecological factors that influenced the participant’s cigarette use. Data from both assessments were used to develop a series of EMA measures for the feasibility study.

Due to the potential burden placed on community coaches tasked with ensuring EMA survey completion during the 7-day feasibility study, the members of the WINCART Center voted to restrict enrollment to 20 participants (see Table 1). These participants were selected to parallel the larger sample in terms of gender, age, tobacco use, and geographic distribution. Of these participants, 60% (12/20) were male. Over half (55%, 11/20) of the participants reported smoking more than 10 cigarettes per day. A mid-point recoding strategy (i.e. 0 for the response “did not smoke,” 0.5 cigarettes for the response “less than 1 cigarette,” 3.5 cigarettes for the response “2-5 cigarettes,” etc) was applied to estimate that an average of 11.8 cigarettes (SD 5.75) were smoked per day per participant.

**Table 1.** Descriptive statistics for demographics and tobacco use from 20 participants.

General characteristics	Male N (%)	Female N (%)
<b>Ethnic identification</b>		
Chamorro	2 (16.7)	0 (0)
Native Hawaiian	1 (8.3)	0 (0)
Marshallese	0 (0)	0 (0)
Samoan	5 (41.7)	2 (25)
Tongan	4 (33.3)	5 (62.5)
Other Pacific Islander	0 (0)	1 (12.5)
<b>Age</b>		
18-20	1 (11.2)	0 (0)
21-23	2 (22.2)	1 (16.7)
24-26	2 (22.2)	2 (33.3)
27-30	4 (44.4)	3 (50)
<b>Education</b>		
Less than high school	0 (0)	0 (0)
High school or GED	8 (72.7)	4 (50)
Some college/trade school	1 (9.1)	3 (37.5)
2-year college	1 (9.1)	1 (12.5)
4-year college or above	1 (9.1)	0 (0)
<b>Employment status</b>		
Employed	7 (58.3)	4 (57.1)
Unemployed	5 (41.7)	3 (42.9)
<b>Days smoked in past 30 days</b>		
0 days	0 (0)	0 (0)
1 or 2 days	0 (0)	0 (0)
3 to 5 days	1 (9.1)	1 (12.5)
6 to 9 days	0 (0)	0 (0)
10 to 19 days	0 (0)	0 (0)
20 to 29 days	3 (27.3)	1 (12.5)
All 30 days	7 (63.6)	6 (75)
<b>Cigarettes smoked per day</b>		
Less than 1 cig	0 (0)	0 (0)
1 cig	0 (0)	0 (0)
2-5 cig	3 (25)	2 (25)
6-10 cig	2 (16.7)	2 (25)
11-20 cig	5 (41.7)	4 (50)
More than 20 cig	2 (16.7)	0 (0)
<b>FTND nicotine dependence score</b>		
Low ( $\leq 5$ )	12 (60)	8 (40)
High ( $\geq 6$ )	0 (0)	0 (0)
<b>Lifetime use of alternative tobacco products</b>		
Hookah	8 (72.7)	6 (75)

General characteristics	Male	Female
	N (%)	N (%)
Cigars	8 (72.7)	4 (50)
Pipe	4 (36.4)	1 (12.5)
Smokeless (chew, betel nut, etc.)	4 (36.4)	1 (12.5)
Cloves	3 (27.3)	0 (0)
Bidis	1 (9.1)	1 (12.5)
Kreteks	0 (0)	0 (0)
Other	0 (0)	1 (12.5)
None of the above	2 (18.2)	1 (12.5)

## Procedures

Each of the twenty participants enrolled in the EMA study attended a one-on-one, follow-up appointment at which the customized system was presented. Surveys were initiated by sending a text message to the participant. This text message contained a link to a secure SQL Server hosting a real-time, Web-based survey system formatted to work on any Web-enabled mobile phone (see [Figure 2](#)) as well as any Web-enabled tablet or computer. Participants accessed the system with a self-created username and password. After logging in, the system recognized the participant, recalled the responses entered earlier in the day, and presented a tailored survey.

The Web-based survey was programmed so that if a participant did not complete the first question within 15 minutes of the first text message being sent another text message was delivered. This process was repeated every 15 minutes for up to one hour (see [Figure 3](#)). The system then waited an hour before initiating the next survey. If the participant failed to begin a survey after ten text messages, an automated email was sent to a community coach who contacted the participant by phone and reminded the individual to complete the survey. This reminder process was

explained to the participant through 8 one-minute, animated videos (see [Multimedia Appendices 1-8](#)). These videos used Pacific Islander characters to demonstrate key concepts and were posted online [64] so that the participant could review the materials remotely throughout the study.

After watching the animated videos, the participant selected six consecutive two-hour time blocks during which they would be willing to receive automated text messages each day. These text messages were delivered on even-numbered hours (2:00 PM, 4:00 PM, 6:00 PM, etc.) and asked the participant to report their cigarette use since the last time a survey was completed. Text messages were delivered to a Web-enabled mobile phone either owned by the participant or provided to the participant for the duration of the study. The videos informed the participant that 6 surveys would be administered each day for a period of 9 days. The first 2 days were designed to help the participant acclimate to the process of completing the surveys and to resolve any technical problems encountered. The 7 days after that were the critical test days. At the end of this 9-day period, research staff provided each participant with a US \$75 gift card to compensate them for their time and travel.

**Figure 2.** Web-based survey formatted for Web-enabled mobile phones, tablets, and computers.



## Measures

Items for the EMA survey were developed based on data gathered from the computer-based questionnaire and the semi-structured interview. The EMA survey included measures of cigarette use, craving, location, social environment, mood, and rationale for deciding whether or not to smoke. The resulting items were then refined in consultation with the 5 community-based organizations. This process resulted in items that, while unconventional when compared to traditional EMA studies, were more appropriate for the Pacific Islander community.

Participants initially responded to a single question inquiring if they had smoked since the last time they entered a response that day. If they had smoked, the survey asked the number of cigarettes they had ("How many cigarettes have you had since {time of last entered response}?"). This question was designed so that all responses in a single day could be tabulated and compared to self-report measures of daily cigarette use. The question also facilitated the analytic approach which focused on the average number of cigarettes smoked across time blocks within each day. A follow-up question determined when the participant smoked ("When did you smoke?") utilizing four response categories (" $<30$  mins ago," "30-60 mins ago," "60-90


mins ago," and "90-120 mins ago"). This question did not ascertain the number of cigarettes smoked in each 30-minute sub-block but was instead used to gauge how much time had passed since the last instance in which the participant smoked. The remaining questions asked the participant to reflect on this instance ("Now, please think about the last time you smoked...").

After recalling the last instance in which they smoked, the participant answered one question about the extent to which they craved a cigarette beforehand ("How much were you craving a cigarette?"). Responses were rated on a 4-point Likert scale ranging from "1=Not at all" to "4=A lot" (see [Figure 2](#)). The participant then reported their location ("Where were you?") by selecting among the response options "Home," "Work," "School," "At a restaurant/bar," "In a car," "Around church," and "Other (Please type in)." The next question asked the participant to classify their social environment ("Who were you with when you last smoked?") using the response options "Alone," "People who are smoking," "People who are NOT smoking," and "Other (Please type in)." The survey then used a "Check all that apply" format to assess the participant's mood ("How were you feeling?") using the categories "Happy," "Sad," "Angry," "Stressed out," "Anxious," "Bored," "Relaxed," "Fine," and "Other (Please type in)." The final question asked the participant to reflect on why they had chosen to smoke

(“Why did you smoke?”) and respond using a “Check all that apply” format comprised of the categories “Someone offered me a cig,” “Someone was smoking around me,” “Wanted a cigarette while drinking alcohol,” “Wanted a cigarette while

drinking coffee,” “Wanted a cigarette after eating,” “To eat less,” “To relax or calm down,” “Have a good time/celebrate,” “To concentrate/focus,” and “Other (Please type in).”

**Figure 3.** Example timeline of survey reminders.



## Analysis

Cigarette use was reported in 534 time blocks. The analysis focused on the total and average number of cigarettes smoked during each time block. Repeated assessments of factors associated with the most recent cigarette consumed were aggregated by taking the average of these assessments across time blocks within each day. This produced a two-level hierarchical analysis dataset with data on each day (i.e. level-1 data) nested within individual participants (i.e. level-2 data). Multilevel models with both fixed and random effects were then used to quantify between and within subject variability across repeated measurement points [16,65,66].

Variables were created for the associational analysis. These variables included the total number of cigarettes smoked during time blocks per day, the proportion of time blocks per day in which 4 or more cigarettes were smoked, and the proportion of time blocks per day in which the participant reported an intrapersonal, social, or ecological factor related to cigarette use. Multilevel regression models were conducted using SAS Proc Mixed procedure with the participants' age, gender, and nicotine dependence being adjusted in the analysis as covariates [67]. Analysis results with raw data are presented and consistent

parameter estimates were obtained with multiple imputation analysis from SAS PROC MI and MIANALYZE [68].

## Results

### Survey Completion and Reported Cigarette Use

Surveys in which the participant failed to respond to all questions within two hours of the first text message prompt were classified as missed. In total, 20 participants completed 720 surveys from 840 prompted survey time blocks, representing a prompt-based survey completion rate of 86%. Nineteen (95%) participants completed all surveys on the first five days. Sixteen (80%) participants completed all surveys on the sixth day and fifteen (75%) participants completed all surveys on the seventh day. Participants reported smoking in 535 (74%) of the 720 two-hour time blocks with completed surveys. The average total number of cigarettes smoked per day was 13.96 (SD 9.19) which is 18.3% higher than the recorded estimates reported in the computer-based questionnaire.

### Factors Associated With Cigarette Use

For each time block in which smoking was reported, participants had an average of 3.07 cigarettes (SD 1.44). Home was the most common smoking location and attempting to relax was the most commonly cited reason for choosing to smoke (see Table 2).



**Table 2.** Proportions of responses per day during time blocks in which smoking was reported.

Question	Response option	Mean of proportion (SD)
<b>Where were you?</b>		
	At home	0.61 (0.37)
	In a car	0.20 (0.28)
	At work	0.13 (0.28)
	Around church	0.07 (0.23)
	At restaurant/bar	0.05 (0.12)
	At school	0.01 (0.07)
<b>Who were you with when you last smoked?</b>		
	Alone	0.54 (0.35)
	With people who are not smoking	0.41 (0.36)
	With people who are smoking	0.23 (0.29)
<b>How were you feeling?</b>		
	Fine	0.51 (0.39)
	Relaxed	0.37 (0.37)
	Happy	0.18 (0.30)
	Stressed out	0.16 (0.31)
	Bored	0.10 (0.17)
	Anxious	0.09 (0.20)
	Angry	0.03 (0.13)
	Sad	0.02 (0.11)
<b>Why did you smoke?</b>		
	To relax or calm down	0.55 (0.35)
	Wanted a cigarette after eating	0.26 (0.29)
	Someone was smoking around me	0.15 (0.26)
	Someone offered me a cig	0.11 (0.25)
	To concentrate/focus	0.11 (0.24)
	Have a good time/celebrate	0.09 (0.20)
	Wanted a cigarette while drinking alcohol	0.05 (0.14)
	Wanted a cigarette while drinking coffee	0.03 (0.09)
	To eat less	0.03 (0.10)

After adjusting for gender, age, and nicotine dependence, analyses indicated that a one score increase of craving resulted in 3.8 more cigarettes smoked per day on average ( $P < .001$ ). In addition, feeling happy ( $P < .001$ ) or wanting a cigarette while drinking alcohol ( $P < .001$ ) (see [Table 3](#)) were positively associated with the total number of cigarettes smoked during time blocks. Being at home ( $P = .02$ ) or being around people who are not smoking ( $P = .01$ ) were negatively associated with the total number of cigarettes smoked during time blocks. These

associations persisted when participants reported smoking 4 or more cigarettes (see [Table 4](#)). Several other associations were also identified when the proportion of time blocks per day in which 4 or more cigarettes were smoked was used as the outcome variable in the analysis. Positive associations included feeling angry ( $P = .05$ ) and wanting a cigarette while drinking coffee ( $P = .01$ ). Negative associations included feeling bored ( $P = .02$ ) and wanting to eat less ( $P = .02$ ).

**Table 3.** Associations with total number of cigarettes smoked per day.

Question	Response option	$\beta$	SE	<i>P</i>
<b>Where were you?</b>				
	At home	-0.72	0.31	.02
	In a car	-0.02	0.42	.96
	At work	0.49	0.44	.27
	Around church	0.79	0.85	.36
	At restaurant/bar	0.17	0.71	.82
	At school	1.51	1.16	.20
<b>Who were you with when you last smoked?</b>				
	Alone	0.54	0.29	.06
	With people who are not smoking	-0.76	0.28	.01
	With people who are smoking	0.65	0.35	.07
<b>How were you feeling?</b>				
	Fine	0.31	0.29	.30
	Relaxed	0.19	0.37	.60
	Happy	1.41	0.37	< .001
	Stressed out	0.13	0.43	.76
	Bored	-0.85	0.54	.12
	Anxious	0.9	0.55	.11
	Angry	0.88	0.71	.22
	Sad	0.03	0.84	.97
<b>Why did you smoke?</b>				
	To relax or calm down	0.41	0.32	.22
	Wanted a cigarette after eating	0.26	0.39	.50
	Someone was smoking around me	0.15	0.43	.72
	Someone offered me a cig	0.72	0.47	.13
	To concentrate/focus	-0.52	0.53	.33
	Have a good time/celebrate	0.46	0.55	.41
	Wanted a cigarette while drinking alcohol	2.31	0.63	<.001
	Wanted a cigarette while drinking coffee	1.85	1.08	.09
	To eat less	-1.32	1.25	.29

**Table 4.** Associations with proportion of time blocks per day with four or more cigarettes.

Question	Response option	$\beta$	SE	<i>P</i>
<b>Where were you?</b>				
	At home	-0.19	0.08	.02
	In a car	0.06	0.11	.59
	At work	0.09	0.11	.40
	Around church	0.07	0.20	.74
	At restaurant/bar	0.00	0.19	.99
	At school	0.91	0.29	.003
<b>Who were you with when you last smoked?</b>				
	Alone	0.09	0.07	.23
	With people who are not smoking	-0.17	0.07	.03
	With people who are smoking	0.17	0.09	.06
<b>How were you feeling?</b>				
	Fine	0.01	0.08	.92
	Relaxed	0.08	0.09	.42
	Happy	0.29	0.10	.003
	Stressed out	0.14	0.11	.19
	Bored	-0.32	0.14	.02
	Anxious	0.24	0.14	.10
	Angry	0.37	0.18	.05
	Sad	-0.01	0.22	.95
<b>Why did you smoke?</b>				
	To relax or calm down	0.09	0.08	.28
	Wanted a cigarette after eating	0.12	0.10	.23
	Someone was smoking around me	-0.04	0.11	.75
	Someone offered me a cig	0.09	0.12	.44
	To concentrate/focus	-0.21	0.14	.12
	Have a good time/celebrate	0.19	0.14	.19
	Wanted a cigarette while drinking alcohol	0.38	0.17	.03
	Wanted a cigarette while drinking coffee	0.77	0.27	.01
	To eat less	-0.73	0.31	.02

## Discussion

### Implications for Future Research and Interventions

This is the first study, of which we are aware, in which a customized EMA system assessed factors associated with tobacco use among young, adult Pacific Islanders. Prior EMA studies of tobacco use that utilized electronic diaries reported prompt-based survey completion rates of 65% [15], 68% [30], 75% [9], 88% [17], 89% [14], 90% [7], and 91% [13]. ICAT studies that relied upon the use of Web-based surveys administered through mobile phones reported prompt-based survey completion rates of 50% [29], 52% [57], 55% [10], 69% [42], and 83% [12]. The survey completion rate of the current study was 86% (720/840), suggesting that young, adult Pacific

Islanders were able to effectively use the customized EMA system and that this form of assessment holds promise for future research. The survey completion rate also indicates that remotely accessible standardized training videos (see [Multimedia Appendices 1-8](#)) may serve as a reasonable alternative to extensive in-person training and technical support—especially for populations spread out over a large geographic region.

Prior research has suggested that while EMA measures sometimes mirror the results of recall measures they often gather data with less noise and greater sensitivity [6]. The assessment of daily cigarette use provides additional evidence to support this conclusion. Self-report measures administered at the outset of the study indicated that each participant smoked an average of 11.8 (SD 5.75) cigarettes per day. In contrast, the EMA

system cataloged an average of 13.96 (SD 9.19) cigarettes smoked per day. The health implications of this are notable in that they translate to an estimated 769 additional cigarettes, or approximately 38.5 packs, per year per participant.

An analysis of the EMA data also offers new insight into the intrapersonal, social, and ecological factors associated with cigarette use among young, adult Pacific Islanders. Previous studies have identified the home as a place where smoking frequently occurs [13]. Yet, the fact that Pacific Islanders in the current study smoke fewer cigarettes than average at home suggests that this location could serve as a constructive environment for practicing smoking cessation techniques. Past EMA research has also highlighted how smokers are more likely to have a cigarette when in the presence of other smokers [11-15]. However, the protective influence of nonsmokers on Pacific Islander tobacco use suggests that such individuals may function as a deterrent that can be relied upon in social situations where smoking is prevalent. Future smoking cessation interventions tailored for Pacific Islanders may choose to test these concepts by providing educational materials that promote the use of nonsmoking social networks when attempting to quit using tobacco products.

A technological innovation within the current study, the creation of tailored survey questions based on participant responses entered earlier in the day, hints at new opportunities for additional research. Future studies may choose to capitalize on this feature by administering brief, tailored surveys to multiple, linked individuals within the Pacific Islander community each time one individual reports smoking behavior in a prior assessment. Such investigations may lead to an improved understanding of friend and familial perceptions of Pacific Islander tobacco use, which previous research has suggested is highly relevant in explaining both past and current smoking among Native Hawaiian youth [69].

The correlation between increased tobacco craving and increased cigarette use parallels previous findings that cravings precipitate smoking [9] and that higher levels of craving at one time point within a day tend to produce more smoking at the next time point [8]. Similarly, other studies have identified significant positive associations between alcohol consumption and tobacco use [52] and coffee consumption and tobacco use [13]. These similarities suggest that components of existing evidence-based interventions that address these factors may be highly effective for young, adult Pacific Islanders. The results for mood are more ambiguous. Some studies suggest negative affect is predictive of smoking [8,15] while others find no prospective relationship with either positive or negative affect [13,14]. Unfortunately, the measure utilized in the current study does not delineate whether the mood indicated preceded cigarette use or was a direct outcome of it. The correlation between tobacco use and happiness may therefore be the participant experiencing higher

positive affect before or after smoking [16]. The inconclusive nature of this finding exposes the need for additional research within this population.

### Limitations

There are several limitations to the current study. Perhaps the most relevant is the small sample size and homogeneity of the participants. Specifically, the generalizability of the sample to other Pacific Islander communities may be limited by the fact that all participants were identified via nonprobability sampling. Moreover, key subgroups, including female Chamorros, Native Hawaiians, and Marshallese, were not represented nor were Pacific Islanders over the age of thirty. The small sample size also prohibited analytic techniques that have generated valuable insights in prior EMA research. Examples include explorations of how situational and mood factors differ between light and heavy smokers [12] and analyses that reveal situational covariates that influence smoking patterns [17].

Another limitation within the current study is the focus on regular cigarettes. With the advent of electronic cigarettes, as well as the common usage of alternative forms of tobacco within the Pacific Islander community [4], the range of behavior measured by future tobacco use studies may need to be broadened. Within the current sample, lifetime hookah use was reported at 73% for males (8/11) and 75% (6/8) for females (see Table 1). This is significantly greater than the averages for the general population of 42% for current male smokers and 23% for current female smokers [5] and suggests that future research should explore the different factors that may be associated with the use of alternative tobacco products.

An additional limitation is that all surveys were initiated on even-numbered hours and only assessed factors associated with the most recent cigarette smoked. Future studies should consider initiating surveys at random intervals within time blocks to avoid potential time-based covariates. Such methods will help clarify the causal relationship between cigarette use and the factors associated with it. Nevertheless, future interventions may still consider applying the current findings even if their causal relationship remains unclear [70,71].

### Conclusions

This feasibility study provides new insights on factors that contribute to cigarette use among young, adult Pacific Islanders. It suggests new directions for conducting advanced, technology-based research aimed at understanding tobacco use within the Pacific Islander community and offers new possibilities for crafting culturally-tailored interventions, which past research has demonstrated is often more effective within this population [72]. Given the disproportionately high smoking rates among Pacific Islanders [5], these types of research and intervention efforts are sorely needed to address the health disparities evident within this population.

### Acknowledgments

This research was supported by the Weaving an Islander Network for Cancer Awareness, Research, and Training (WINCART) Center funded by Grant Number U54CA153458 from the National Cancer Institute (NCI), Center to Reduce Cancer Health Disparities (CRCHD). The authors thank the directors, community and academic researchers, and project staff at the Guam

Communications Network, Pacific Islander Health Partnership, Samoan National Nurses Association, Tongan Community Service Center/Special Services for Groups, Union of Pan Asian Communities, Orange County Asian Pacific Islander Community Alliance, Orange County Health Care Agency, California State University, Fullerton, St. Joseph Hospital of Orange, University of Southern California, and Claremont Graduate University for their contributions. The contents are solely the responsibility of the authors and do not necessarily represent the official views of the NCI CRCHD.

---

### Authors' Contributions

JRP and BX conceived and drafted the manuscript. BX performed the analyses with support from NT and MAR. NT, MDS, AO, TT, SC, VM, CL, VKP, MAR, DESV, JTL, SPT, and PHP edited the manuscript for content. All authors reviewed and approved the final version of the manuscript.

---

### Conflicts of Interest

None declared.

---

### Multimedia Appendix 1

Introduction to the ecological momentary assessment study.

[[MOV File, 22MB - mhealth\\_v4i1e2\\_app1.mov](#) ]

---

### Multimedia Appendix 2

Using an iPhone.

[[MOV File, 55MB - mhealth\\_v4i1e2\\_app2.mov](#) ]

---

### Multimedia Appendix 3

Accessing a Web-based survey with an iPhone.

[[MOV File, 30MB - mhealth\\_v4i1e2\\_app3.mov](#) ]

---

### Multimedia Appendix 4

Completing a Web-based survey with an iPhone.

[[MOV File, 46MB - mhealth\\_v4i1e2\\_app4.mov](#) ]

---

### Multimedia Appendix 5

Accessing a Web-based survey after receiving a text message.

[[MOV File, 29MB - mhealth\\_v4i1e2\\_app5.mov](#) ]

---

### Multimedia Appendix 6

Selecting times to receive text message survey reminders.

[[MOV File, 19MB - mhealth\\_v4i1e2\\_app6.mov](#) ]

---

### Multimedia Appendix 7

Receiving text message survey reminders.

[[MOV File, 38MB - mhealth\\_v4i1e2\\_app7.mov](#) ]

---

### Multimedia Appendix 8

Contacting technical support.

[[MOV File, 16MB - mhealth\\_v4i1e2\\_app8.mov](#) ]

---

### References

1. U.S. Census Bureau. United States Census Bureau Data. 2010. 2006-2010 American Community Survey 5-Year Estimates URL: [http://www.census.gov/acs/www/data\\_documentation/2010\\_release/](http://www.census.gov/acs/www/data_documentation/2010_release/) [accessed 2015-01-26] [[WebCite Cache ID 6VsR5NaCg](#)]
2. Centers for Disease Control and Prevention. Health disparities among Native Hawaiians and other Pacific Islanders garner little attention. *Chronic Dis Notes Rep* 2002;15(2):14-27.
3. Srinivasan S, Guillermo T. Toward improved health: disaggregating Asian American and Native Hawaiian/Pacific Islander data. *Am J Public Health* 2000 Nov;90(11):1731-1734. [Medline: [11076241](#)]
4. Mukherjea A, Wackowski OA, Lee YO, Delnevo CD. Asian American, Native Hawaiian and Pacific Islander tobacco use patterns. *Am J Health Behav* 2014 May;38(3):362-369. [doi: [10.5993/AJHB.38.3.5](#)] [Medline: [24636032](#)]
5. Centers for Disease Control and Prevention. CDC - National Adult Tobacco Survey (NATS) - Smoking & Tobacco Use. 2014. National Adult Tobacco Survey URL: [http://www.cdc.gov/tobacco/data\\_statistics/surveys/nats/](http://www.cdc.gov/tobacco/data_statistics/surveys/nats/) [accessed 2015-01-26] [[WebCite Cache ID 6VsQvGpNY](#)]
6. Shiffman S, Stone AA, Hufford MR. Ecological momentary assessment. *Annu Rev Clin Psychol* 2008;4:1-32. [Medline: [18509902](#)]
7. Shiffman S. Dynamic influences on smoking relapse process. *J Pers* 2005 Dec;73(6):1715-1748. [doi: [10.1111/j.0022-3506.2005.00364.x](#)] [Medline: [16274451](#)]
8. Berkman ET, Dickenson J, Falk EB, Lieberman MD. Using SMS text messaging to assess moderators of smoking reduction: Validating a new tool for ecological measurement of health behaviors. *Health Psychol* 2011 Mar;30(2):186-194 [[FREE Full text](#)] [doi: [10.1037/a0022201](#)] [Medline: [21401252](#)]
9. Carter BL, Lam CY, Robinson JD, Paris MM, Waters AJ, Wetter DW, et al. Real-time craving and mood assessments before and after smoking. *Nicotine Tob Res* 2008 Jul;10(7):1165-1169 [[FREE Full text](#)] [doi: [10.1080/14622200802163084](#)] [Medline: [18629726](#)]
10. Brodbeck J, Bachmann MS, Znoj H. Distinct coping strategies differentially predict urge levels and lapses in a smoking cessation attempt. *Addict Behav* 2013 Jun;38(6):2224-2229. [doi: [10.1016/j.addbeh.2013.02.001](#)] [Medline: [23501139](#)]
11. Huh J, Shin H, Leventhal AM, Spruijt-Metz D, Abramova Z, Cerrada C, et al. Momentary negative moods and being with friends precede cigarette use among Korean American emerging adults. *Nicotine Tob Res* 2014 Sep;16(9):1248-1254. [doi: [10.1093/ntr/ntu063](#)] [Medline: [24755398](#)]
12. Thrul J, Bühler A, Ferguson SG. Situational and mood factors associated with smoking in young adult light and heavy smokers. *Drug Alcohol Rev* 2014 Jul;33(4):420-427. [doi: [10.1111/dar.12164](#)] [Medline: [24893955](#)]
13. Shiffman S, Gwaltney CJ, Balabanis MH, Liu KS, Paty JA, Kassel JD, et al. Immediate antecedents of cigarette smoking: an analysis from ecological momentary assessment. *J Abnorm Psychol* 2002 Nov;111(4):531-545. [Medline: [12428767](#)]
14. Shiffman S, Paty JA, Gwaltney CJ, Dang Q. Immediate antecedents of cigarette smoking: an analysis of unrestricted smoking patterns. *J Abnorm Psychol* 2004 Feb;113(1):166-171. [doi: [10.1037/0021-843X.113.1.166](#)] [Medline: [14992670](#)]
15. Shapiro D, Jamner LD, Davydov DM, James P. Situations and moods associated with smoking in everyday life. *Psychol Addict Behav* 2002 Dec;16(4):342-345. [Medline: [12503908](#)]
16. Hedeker D, Mermelstein RJ, Berbaum ML, Campbell RT. Modeling mood variation associated with smoking: an application of a heterogeneous mixed-effects model for analysis of ecological momentary assessment (EMA) data. *Addiction* 2009 Feb;104(2):297-307 [[FREE Full text](#)] [doi: [10.1111/j.1360-0443.2008.02435.x](#)] [Medline: [19149827](#)]
17. Shiffman S, Dunbar MS, Li X, Scholl SM, Tindle HA, Anderson SJ, et al. Smoking patterns and stimulus control in intermittent and daily smokers. *PLoS One* 2014;9(3):e89911 [[FREE Full text](#)] [doi: [10.1371/journal.pone.0089911](#)] [Medline: [24599056](#)]
18. Shiffman S, Dunbar MS, Tindle HA, Ferguson SG. Nondaily smokers' experience of craving on days they do not smoke. *J Abnorm Psychol* 2015 Aug;124(3):648-659. [doi: [10.1037/abn0000063](#)] [Medline: [26052617](#)]
19. Shiffman S, Li X, Dunbar MS, Ferguson SG, Tindle HA, Scholl SM. Social smoking among intermittent smokers. *Drug Alcohol Depend* 2015 Sep 1;154:184-191. [doi: [10.1016/j.drugalcdep.2015.06.027](#)] [Medline: [26205313](#)]
20. McCarthy DE, Minami H, Yeh VM, Bold KW. An experimental investigation of reactivity to ecological momentary assessment frequency among adults trying to quit smoking. *Addiction* 2015 Oct;110(10):1549-1560. [doi: [10.1111/add.12996](#)] [Medline: [26011583](#)]
21. Brondolo E, Monge A, Agosta J, Tobin JN, Cassells A, Stanton C, et al. Perceived ethnic discrimination and cigarette smoking: examining the moderating effects of race/ethnicity and gender in a sample of Black and Latino urban adults. *J Behav Med* 2015 Aug;38(4):689-700. [doi: [10.1007/s10865-015-9645-2](#)] [Medline: [26054448](#)]
22. Bujarski S, Roche DJO, Sheets ES, Krull JL, Guzman I, Ray LA. Modeling naturalistic craving, withdrawal, and affect during early nicotine abstinence: A pilot ecological momentary assessment study. *Exp Clin Psychopharmacol* 2015 Apr;23(2):81-89. [doi: [10.1037/a0038861](#)] [Medline: [25844632](#)]
23. Brodbeck J, Bachmann MS, Brown A, Znoj HJ. Effects of depressive symptoms on antecedents of lapses during a smoking cessation attempt: an ecological momentary assessment study. *Addiction* 2014 Aug;109(8):1363-1370. [doi: [10.1111/add.12563](#)] [Medline: [24690068](#)]

24. Piasecki TM, Trela CJ, Hedeker D, Mermelstein RJ. Smoking antecedents: separating between- and within-person effects of tobacco dependence in a multiwave ecological momentary assessment investigation of adolescent smoking. *Nicotine Tob Res* 2014 May;16 Suppl 2:S119-S126 [FREE Full text] [doi: [10.1093/ntr/ntt132](https://doi.org/10.1093/ntr/ntt132)] [Medline: [23990475](https://pubmed.ncbi.nlm.nih.gov/23990475/)]
25. Soong A, Chen JC, Borzekowski DL. Using ecological momentary assessment to study tobacco behavior in urban India: there's an app for that. *JMIR Res Protoc* 2015;4(2):e76 [FREE Full text] [doi: [10.2196/resprot.4408](https://doi.org/10.2196/resprot.4408)] [Medline: [26109369](https://pubmed.ncbi.nlm.nih.gov/26109369/)]
26. Businelle MS, Ma P, Kendzor DE, Reitzel LR, Chen M, Lam CY, et al. Predicting quit attempts among homeless smokers seeking cessation treatment: an ecological momentary assessment study. *Nicotine Tob Res* 2014 Oct;16(10):1371-1378 [FREE Full text] [doi: [10.1093/ntr/ntu088](https://doi.org/10.1093/ntr/ntu088)] [Medline: [24893602](https://pubmed.ncbi.nlm.nih.gov/24893602/)]
27. Fong M, Braun K, Tsark J. Improving native Hawaiian health through community-based participatory research. *Calif J Health Promot* 2013;1:136-148.
28. George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority research participation among African Americans, Latinos, Asian Americans, and Pacific Islanders. *Am J Public Health* 2014 Feb;104(2):e16-e31. [doi: [10.2105/AJPH.2013.301706](https://doi.org/10.2105/AJPH.2013.301706)] [Medline: [24328648](https://pubmed.ncbi.nlm.nih.gov/24328648/)]
29. Thrul J, Bühler A, Ferguson SG. An Internet-based ecological momentary assessment study relying on participants' own mobile phones: insights from a study with young adult smokers. *Eur Addict Res* 2015;21(1):1-5. [doi: [10.1159/000363231](https://doi.org/10.1159/000363231)] [Medline: [25342514](https://pubmed.ncbi.nlm.nih.gov/25342514/)]
30. Sokolovsky AW, Mermelstein RJ, Hedeker D. Factors predicting compliance to ecological momentary assessment among adolescent smokers. *Nicotine Tob Res* 2014 Mar;16(3):351-358 [FREE Full text] [doi: [10.1093/ntr/ntt154](https://doi.org/10.1093/ntr/ntt154)] [Medline: [24097816](https://pubmed.ncbi.nlm.nih.gov/24097816/)]
31. Dunton GF, Liao Y, Intille SS, Spruijt-Metz D, Pentz M. Investigating children's physical activity and sedentary behavior using ecological momentary assessment with mobile phones. *Obesity* 2011 Jun;19(6):1205-1212. [doi: [10.1038/oby.2010.302](https://doi.org/10.1038/oby.2010.302)] [Medline: [21164502](https://pubmed.ncbi.nlm.nih.gov/21164502/)]
32. Gorely T, Marshall SJ, Biddle SJH, Cameron N. Patterns of sedentary behaviour and physical activity among adolescents in the United Kingdom: Project STIL. *J Behav Med* 2007 Dec;30(6):521-531. [doi: [10.1007/s10865-007-9126-3](https://doi.org/10.1007/s10865-007-9126-3)] [Medline: [17712619](https://pubmed.ncbi.nlm.nih.gov/17712619/)]
33. Hausenblas HA, Gauvin L, Symons DD, Duley AR. Effects of abstinence from habitual involvement in regular exercise on feeling states: an ecological momentary assessment study. *Br J Health Psychol* 2008 May;13(Pt 2):237-255. [doi: [10.1348/135910707X180378](https://doi.org/10.1348/135910707X180378)] [Medline: [17535493](https://pubmed.ncbi.nlm.nih.gov/17535493/)]
34. Dunton GF, Whalen CK, Jamner LD, Henker B, Floro JN. Using ecologic momentary assessment to measure physical activity during adolescence. *Am J Prev Med* 2005 Nov;29(4):281-287. [doi: [10.1016/j.amepre.2005.07.020](https://doi.org/10.1016/j.amepre.2005.07.020)] [Medline: [16242590](https://pubmed.ncbi.nlm.nih.gov/16242590/)]
35. Dunton GF, Liao Y, Kawabata K, Intille S. Momentary assessment of adults' physical activity and sedentary behavior: feasibility and validity. *Front Psychol* 2012;3:260 [FREE Full text] [doi: [10.3389/fpsyg.2012.00260](https://doi.org/10.3389/fpsyg.2012.00260)] [Medline: [22866046](https://pubmed.ncbi.nlm.nih.gov/22866046/)]
36. Dunton GF, Huh J, Leventhal AM, Riggs N, Hedeker D, Spruijt-Metz D, et al. Momentary assessment of affect, physical feeling states, and physical activity in children. *Health Psychol* 2014 Mar;33(3):255-263 [FREE Full text] [doi: [10.1037/a0032640](https://doi.org/10.1037/a0032640)] [Medline: [23668846](https://pubmed.ncbi.nlm.nih.gov/23668846/)]
37. Liao Y, Intille SS, Dunton GF. Using ecological momentary assessment to understand where and with whom adults' physical and sedentary activity occur. *Int J Behav Med* 2015 Feb;22(1):51-61. [doi: [10.1007/s12529-014-9400-z](https://doi.org/10.1007/s12529-014-9400-z)] [Medline: [24639067](https://pubmed.ncbi.nlm.nih.gov/24639067/)]
38. Glanz K, Murphy S. Dietary assessment and monitoring in real time. In: Stone AA, Shiffman S, Atienza AA, Nebeling L, editors. *The science of real-time data capture: self-reports in health research*. Oxford: Oxford University Press; 2007:151-168.
39. Greeno CG, Wing RR, Shiffman S. Binge antecedents in obese women with and without binge eating disorder. *J Consult Clin Psychol* 2000 Feb;68(1):95-102. [Medline: [10710844](https://pubmed.ncbi.nlm.nih.gov/10710844/)]
40. Grenard JL, Stacy AW, Shiffman S, Baraldi AN, MacKinnon DP, Lockhart G, et al. Sweetened drink and snacking cues in adolescents: a study using ecological momentary assessment. *Appetite* 2013 Aug;67:61-73 [FREE Full text] [doi: [10.1016/j.appet.2013.03.016](https://doi.org/10.1016/j.appet.2013.03.016)] [Medline: [23583312](https://pubmed.ncbi.nlm.nih.gov/23583312/)]
41. Borgogna N, Lockhart G, Grenard JL, Barrett T, Shiffman S, Reynolds KD. Ecological momentary assessment of urban adolescents' technology use and cravings for unhealthy snacks and drinks: differences by ethnicity and sex. *J Acad Nutr Diet* 2015 May;115(5):759-766. [doi: [10.1016/j.jand.2014.10.015](https://doi.org/10.1016/j.jand.2014.10.015)] [Medline: [25482855](https://pubmed.ncbi.nlm.nih.gov/25482855/)]
42. Zenk SN, Horoi I, McDonald A, Corte C, Riley B, Odoms-Young AM. Ecological momentary assessment of environmental and personal factors and snack food intake in African American women. *Appetite* 2014 Dec;83:333-341. [doi: [10.1016/j.appet.2014.09.008](https://doi.org/10.1016/j.appet.2014.09.008)] [Medline: [25239402](https://pubmed.ncbi.nlm.nih.gov/25239402/)]
43. Buckner JD, Zvolensky MJ, Crosby RD, Wonderlich SA, Ecker AH, Richter A. Antecedents and consequences of cannabis use among racially diverse cannabis users: an analysis from ecological momentary assessment. *Drug Alcohol Depend* 2015 Feb 1;147:20-25. [doi: [10.1016/j.drugalcdep.2014.12.022](https://doi.org/10.1016/j.drugalcdep.2014.12.022)] [Medline: [25578250](https://pubmed.ncbi.nlm.nih.gov/25578250/)]
44. Phillips MM, Phillips KT, Lalonde TL, Dykema KR. Feasibility of text messaging for ecological momentary assessment of marijuana use in college students. *Psychol Assess* 2014 Sep;26(3):947-957. [doi: [10.1037/a0036612](https://doi.org/10.1037/a0036612)] [Medline: [24749751](https://pubmed.ncbi.nlm.nih.gov/24749751/)]
45. Phillips KT, Phillips MM, Lalonde TL, Tormohlen KN. Marijuana use, craving, and academic motivation and performance among college students: An in-the-moment study. *Addict Behav* 2015 Aug;47:42-47. [doi: [10.1016/j.addbeh.2015.03.020](https://doi.org/10.1016/j.addbeh.2015.03.020)] [Medline: [25864134](https://pubmed.ncbi.nlm.nih.gov/25864134/)]

46. Epler AJ, Tomko RL, Piasecki TM, Wood PK, Sher KJ, Shiffman S, et al. Does hangover influence the time to next drink? An investigation using ecological momentary assessment. *Alcohol Clin Exp Res* 2014 May;38(5):1461-1469 [FREE Full text] [doi: [10.1111/acer.12386](https://doi.org/10.1111/acer.12386)] [Medline: [24588377](https://pubmed.ncbi.nlm.nih.gov/24588377/)]
47. Linas BS, Latkin C, Westergaard RP, Chang LW, Bollinger RC, Genz A, et al. Capturing illicit drug use where and when it happens: an ecological momentary assessment of the social, physical and activity environment of using versus craving illicit drugs. *Addiction* 2015 Feb;110(2):315-325 [FREE Full text] [doi: [10.1111/add.12768](https://doi.org/10.1111/add.12768)] [Medline: [25311241](https://pubmed.ncbi.nlm.nih.gov/25311241/)]
48. Treloar H, Piasecki TM, McCarthy DM, Sher KJ, Heath AC. Ecological evidence that affect and perceptions of drink effects depend on alcohol expectancies. *Addiction* 2015 Sep;110(9):1432-1442. [doi: [10.1111/add.12982](https://doi.org/10.1111/add.12982)] [Medline: [25959045](https://pubmed.ncbi.nlm.nih.gov/25959045/)]
49. Moore TM, Seavey A, Ritter K, McNulty JK, Gordon KC, Stuart GL. Ecological momentary assessment of the effects of craving and affect on risk for relapse during substance abuse treatment. *Psychol Addict Behav* 2014 Jun;28(2):619-624. [doi: [10.1037/a0034127](https://doi.org/10.1037/a0034127)] [Medline: [24128286](https://pubmed.ncbi.nlm.nih.gov/24128286/)]
50. Yang C, Linas B, Kirk G, Bollinger R, Chang L, Chander G, et al. Feasibility and acceptability of smartphone-based ecological momentary assessment of alcohol use among African American men who have sex with men in Baltimore. *JMIR Mhealth Uhealth* 2015;3(2):e67 [FREE Full text] [doi: [10.2196/mhealth.4344](https://doi.org/10.2196/mhealth.4344)] [Medline: [26085078](https://pubmed.ncbi.nlm.nih.gov/26085078/)]
51. Schnall R, Okoniewski A, Tiase V, Low A, Rodriguez M, Kaplan S. Using text messaging to assess adolescents' health information needs: an ecological momentary assessment. *J Med Internet Res* 2013;15(3):e54 [FREE Full text] [doi: [10.2196/jmir.2395](https://doi.org/10.2196/jmir.2395)] [Medline: [23467200](https://pubmed.ncbi.nlm.nih.gov/23467200/)]
52. Piasecki TM, Jahng S, Wood PK, Robertson BM, Epler AJ, Cronk NJ, et al. The subjective effects of alcohol-tobacco co-use: an ecological momentary assessment investigation. *J Abnorm Psychol* 2011 Aug;120(3):557-571 [FREE Full text] [doi: [10.1037/a0023033](https://doi.org/10.1037/a0023033)] [Medline: [21443289](https://pubmed.ncbi.nlm.nih.gov/21443289/)]
53. Ginexi EM, Riley W, Atienza AA, Mabry PL. The promise of intensive longitudinal data capture for behavioral health research. *Nicotine Tob Res* 2014 May;16 Suppl 2:S73-S75 [FREE Full text] [doi: [10.1093/ntr/ntt273](https://doi.org/10.1093/ntr/ntt273)] [Medline: [24711629](https://pubmed.ncbi.nlm.nih.gov/24711629/)]
54. Garcia C, Hardeman RR, Kwon G, Lando-King E, Zhang L, Genis T, et al. Teenagers and texting: use of a youth ecological momentary assessment system in trajectory health research with latina adolescents. *JMIR Mhealth Uhealth* 2014;2(1):e3 [FREE Full text] [doi: [10.2196/mhealth.2576](https://doi.org/10.2196/mhealth.2576)] [Medline: [25098355](https://pubmed.ncbi.nlm.nih.gov/25098355/)]
55. Berkman ET, Giuliani NR, Pruitt AK. Comparison of text messaging and paper-and-pencil for ecological momentary assessment of food craving and intake. *Appetite* 2014 Oct;81:131-137 [FREE Full text] [doi: [10.1016/j.appet.2014.06.010](https://doi.org/10.1016/j.appet.2014.06.010)] [Medline: [24930596](https://pubmed.ncbi.nlm.nih.gov/24930596/)]
56. Garcia C, Zhang L, Holt K, Hardeman R, Peterson B. Latina adolescent sleep and mood: an ecological momentary assessment pilot study. *J Child Adolesc Psychiatr Nurs* 2014 Aug;27(3):132-141. [doi: [10.1111/jcap.12082](https://doi.org/10.1111/jcap.12082)] [Medline: [25103724](https://pubmed.ncbi.nlm.nih.gov/25103724/)]
57. Kuntsche E, Labhart F. ICAT: development of an internet-based data collection method for ecological momentary assessment using personal cell phones. *Eur J Psychol Assess* 2013;29(2):140-148 [FREE Full text] [doi: [10.1027/1015-5759/a000137](https://doi.org/10.1027/1015-5759/a000137)] [Medline: [24285917](https://pubmed.ncbi.nlm.nih.gov/24285917/)]
58. Minkler M, Wallerstein N. *Community-based participatory research for health: from process to outcomes*. San Francisco, CA: Jossey-Bass; 2008.
59. Palmer PH, Lee C, Sablan-Santos L, Lepule JT, Pang VK, Tui'one V, et al. Eliminating tobacco disparities among native Hawaiian Pacific Islanders through policy change: the role of community-based organizations. *Health Promot Pract* 2013 Sep;14(5 Suppl):36S-39S [FREE Full text] [doi: [10.1177/1524839913486150](https://doi.org/10.1177/1524839913486150)] [Medline: [23667058](https://pubmed.ncbi.nlm.nih.gov/23667058/)]
60. Yancey AK, Ortega AN, Kumanyika SK. Effective recruitment and retention of minority research participants. *Annu Rev Public Health* 2006;27:1-28. [doi: [10.1146/annurev.publhealth.27.021405.102113](https://doi.org/10.1146/annurev.publhealth.27.021405.102113)] [Medline: [16533107](https://pubmed.ncbi.nlm.nih.gov/16533107/)]
61. Fagerström KO. Measuring degree of physical dependence to tobacco smoking with reference to individualization of treatment. *Addict Behav* 1978;3(3-4):235-241. [Medline: [735910](https://pubmed.ncbi.nlm.nih.gov/735910/)]
62. Prokhorov AV, Pallonen UE, Fava JL, Ding L, Niaura R. Measuring nicotine dependence among high-risk adolescent smokers. *Addict Behav* 1996;21(1):117-127. [Medline: [8729713](https://pubmed.ncbi.nlm.nih.gov/8729713/)]
63. Prokhorov AV, De MC, Pallonen UE, Hudmon KS, Koehly L, Hu S. Validation of the modified Fagerström tolerance questionnaire with salivary cotinine among adolescents. *Addict Behav* 2000;25(3):429-433. [Medline: [10890296](https://pubmed.ncbi.nlm.nih.gov/10890296/)]
64. Pike J. WINCART ecological momentary assessment research on Vimeo. Video 1 - introduction URL: <http://vimeo.com/channels/309124> [accessed 2015-01-26] [WebCite Cache ID 6VsPgKapR]
65. Moghaddam NG, Ferguson E. Smoking, mood regulation, and personality: an event-sampling exploration of potential models and moderation. *J Pers* 2007 Jun;75(3):451-478. [doi: [10.1111/j.1467-6494.2007.00445.x](https://doi.org/10.1111/j.1467-6494.2007.00445.x)] [Medline: [17489888](https://pubmed.ncbi.nlm.nih.gov/17489888/)]
66. Schwartz JE, Stone AA. Strategies for analyzing ecological momentary assessment data. *Health Psychol* 1998 Jan;17(1):6-16. [Medline: [9459065](https://pubmed.ncbi.nlm.nih.gov/9459065/)]
67. Raudenbush SW, Bryk AS. *Hierarchical linear models: applications and data analysis methods*. Thousand Oaks: Sage Publications; 2002.
68. Allison P. *Missing data*. Thousand Oaks, Calif: Sage Publications; 2002.
69. Glanz K, Mau M, Steffen A, Maskarinec G, Arriola KJ. Tobacco use among Native Hawaiian middle school students: its prevalence, correlates and implications. *Ethn Health* 2007 Jun;12(3):227-244 [FREE Full text] [doi: [10.1080/13557850701234948](https://doi.org/10.1080/13557850701234948)] [Medline: [17454098](https://pubmed.ncbi.nlm.nih.gov/17454098/)]



70. Stacy AW, Wiers RW. Implicit cognition and addiction: a tool for explaining paradoxical behavior. *Annu Rev Clin Psychol* 2010;6:551-575 [FREE Full text] [doi: [10.1146/annurev.clinpsy.121208.131444](https://doi.org/10.1146/annurev.clinpsy.121208.131444)] [Medline: [20192786](https://pubmed.ncbi.nlm.nih.gov/20192786/)]
71. Wood W, Neal DT. A new look at habits and the habit-goal interface. *Psychol Rev* 2007 Oct;114(4):843-863. [doi: [10.1037/0033-295X.114.4.843](https://doi.org/10.1037/0033-295X.114.4.843)] [Medline: [17907866](https://pubmed.ncbi.nlm.nih.gov/17907866/)]
72. Braun KL, Fong M, Kaanoi ME, Kamaka ML, Gotay CC. Testing a culturally appropriate, theory-based intervention to improve colorectal cancer screening among Native Hawaiians. *Prev Med* 2005 Jun;40(6):619-627 [FREE Full text] [doi: [10.1016/j.ypmed.2004.09.005](https://doi.org/10.1016/j.ypmed.2004.09.005)] [Medline: [15850857](https://pubmed.ncbi.nlm.nih.gov/15850857/)]

## Abbreviations

**EMA:** Ecological Momentary Assessment

**ICAT:** Internet-Based Cell Phone-Optimized Assessment Technique

**NCI CRCHD:** National Cancer Institute Center to Reduce Cancer Health Disparities

**SQL:** Structured Query Language

**WINCART:** Weaving an Islander Network for Cancer Awareness, Research, and Training

*Edited by G Eysenbach; submitted 20.03.15; peer-reviewed by A David, R Schnall, J Thurl; comments to author 25.06.15; revised version received 02.08.15; accepted 09.10.15; published 07.01.16*

*Please cite as:*

*Pike JR, Xie B, Tan N, Sabado-Liwag MD, Orne A, Toilolo T, Cen S, May V, Lee C, Pang VK, Rainer MA, Vaivao DES, Lepule JT, Tanjasiri SP, Palmer PH*

*Developing an Internet- and Mobile-Based System to Measure Cigarette Use Among Pacific Islanders: An Ecological Momentary Assessment Study*

*JMIR mHealth uHealth* 2016;4(1):e2

URL: <http://mhealth.jmir.org/2016/1/e2/>

doi: [10.2196/mhealth.4437](https://doi.org/10.2196/mhealth.4437)

PMID: [26743132](https://pubmed.ncbi.nlm.nih.gov/26743132/)

©James Russell Pike, Bin Xie, Nasya Tan, Melanie Dee Sabado-Liwag, Annette Orne, Tupou Toilolo, Steven Cen, Vanessa May, Cevadne Lee, Victor Kaiwi Pang, Michelle A Rainer, Dorothy Etimani S Vaivao, Jonathan Tana Lepule, Sora Park Tanjasiri, Paula Healani Palmer. Originally published in *JMIR Mhealth and Uhealth* (<http://mhealth.jmir.org>), 07.01.2016. This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in *JMIR mhealth and uhealth*, is properly cited. The complete bibliographic information, a link to the original publication on <http://mhealth.jmir.org/>, as well as this copyright and license information must be included.