

UC Irvine

Clinical Practice and Cases in Emergency Medicine

Title

Nausea and Dyspnea on Exertion: Left Ventricular Free-wall Rupture

Permalink

<https://escholarship.org/uc/item/5m74j5jh>

Journal

Clinical Practice and Cases in Emergency Medicine, 6(1)

Authors

Wilson, Juliana
Mendes, Matthew
Betz, Marian
[et al.](#)

Publication Date

2022

DOI

10.5811/cpcem.2021.4.50265

Supplemental Material

<https://escholarship.org/uc/item/5m74j5jh#supplemental>

Copyright Information

Copyright 2022 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Nausea and Dyspnea on Exertion: Left Ventricular Free-wall Rupture

Juliana Wilson, DO, MPH^{††}
 Matthew Mendes, MD^{*}
 Marian E. Betz, MD, MPH^{*}
 Juan Galindez Mingo, BS[‡]

^{*}University of Colorado Denver – Anschutz Medical Campus, Department of Emergency Medicine, Aurora, Colorado

[†]Denver Health Medical Center, Department of Emergency Medicine, Denver, Colorado

[‡]University of Colorado School of Medicine, Aurora, Colorado

Section Editor: Rick A. McPheeters, DO

Submission history: Submitted November 17, 2020; Revision received April 7, 2021; Accepted April 1, 2021

Electronically published December 16, 2021

Full text available through open access at http://escholarship.org/uc/uciem_cpem

DOI: 10.5811/cpem.2021.4.50265

Case Presentation: A 53-year-old female presented to the emergency department with three days of nausea and dyspnea on exertion after using methamphetamine. Initial electrocardiogram revealed an ST-elevation myocardial infarction. While awaiting transfer to the cardiac catheterization lab the patient suffered a witnessed cardiac arrest. During resuscitative efforts an enlarging pericardial effusion on point-of-care ultrasound led to the detection of a left ventricular free-wall rupture (LVFWR). This case illustrates the progression of a left ventricular free-wall rupture using point-of-care ultrasound.

Discussion: Left ventricular free-wall rupture has a low incidence rate in the setting of an acute myocardial infarction. Ultrasonography is the tool of choice for detecting a LVFWR. [Clin Pract Cases Emerg Med. 2022;6(1):90–92.]

Keywords: *Left ventricular free-wall rupture; POCUS; acute myocardial infarction.*

CASE PRESENTATION

A 53-year-old female with a past medical history of diabetes mellitus, hypertension, and substance use disorder presented to the emergency department with three days of nausea, dyspnea on exertion, and orthopnea. The patient's initial vital signs showed a heart rate of 104 beats per minute, blood pressure of 193/109 millimeters of mercury, respiratory rate of 18 breaths per minute, and pulse oximetry of 94% oxygen saturation on room air. Her initial workup and evaluation demonstrated ST-segment elevations and Q waves on electrocardiogram (ECG) (Image 1). The cardiac catheterization laboratory was immediately activated upon review of the ECG. Chest radiograph (Image 2) was obtained indicating pulmonary edema and small pleural effusions.

An initial point-of-care ultrasound (POCUS) (Video 1) showed evidence of a very small pericardial effusion. The patient was initially hesitant to undergo cardiac catheterization but 30 minutes after arrival agreed to intervention. Prior to




Image 1. Electrocardiogram from initial workup demonstrating ST-segment elevations (red arrows) and Q waves (blue arrows).

transfer to the catheterization laboratory, she suffered a cardiac arrest. Cardiopulmonary resuscitation and advanced cardiac life support were initiated. During resuscitative efforts POCUS was obtained, which showed a large pericardial effusion with concern for a mechanical complication of acute myocardial




Image 2. Chest radiograph showing pulmonary edema (red arrows) and small pleural effusions (black arrow).

infarction (MI). Left ventricular free-wall rupture (LVFWR) was the most concerning given the enlarging pericardial effusion (Video 2). Emergent pericardiocentesis via the subxiphoid approach was performed and 80 milliliters of bloody fluid was obtained. Repeat POCUS did not show any residual pericardial effusion following pericardiocentesis. Despite pericardiocentesis and resuscitative efforts, the patient did not survive.

DISCUSSION

Diagnosis of LVFWR with hemopericardium following MI was confirmed through autopsy. A number of risk factors including diabetes mellitus, hypertension, and methamphetamine use could have contributed to a subacute MI (three days prior) resulting in LVFWR. Autopsy revealed the patient suffered from a 90% occluded left anterior descending artery. Left ventricular free-wall rupture will typically occur within five days of an MI due to vulnerable necrotic tissue.¹ Due to coronavirus disease 2019-related factors such as fear of virus contraction, patients may be more likely to present later to the hospital, leading to higher incidence of post-MI complications.² The incidence rate of LVFWR in acute MI is 2.2%.³ The use of methamphetamine can also increase the risk of developing atherosclerotic plaque and acute coronary vasospasm.⁴ Ultrasonography is indicated as the primary diagnostic tool if a patient is suspected of suffering from a LVFWR.⁵ Management of a LVFWR includes pericardiocentesis and positive inotropic agents along with emergent surgical repair.⁵

Video 1. Point-of-care ultrasound (parasternal short axis) prior to cardiac arrest showing a small pericardial effusion (black arrowheads) and a small myocardium rupture (white arrow).

CPC-EM Capsule

What do we already know about this clinical entity?

A Left ventricular free wall rupture (LVFWR) can occur due to weakening of the myocardium following a myocardial infarction. Ultrasonography is the primary tool used to diagnose LVFWR.

What is the major impact of the image(s)?

The echocardiograms presented in this case demonstrate the progression of a left ventricular free wall rupture leading to a large pericardial effusion.

How might this improve emergency medicine practice?

Recognizing patients at risk for LVFWR and identifying a pericardial effusion and myocardial rupture through ultrasonography can lead to prompt diagnosis and improved patient outcomes.

Video 2. Point-of-care ultrasound during resuscitative efforts showing a large pericardial effusion (white arrows).

Consent from patient's family has been obtained and filed for the publication of this case report. Documentation on file.

Address for Correspondence: Juan Galindez Mingo, BS, University of Colorado School of Medicine, 13001 E 17th PI, Aurora, CO 80045. Email: juan.galindez@cuanschutz.edu.

Conflicts of Interest: By the CPC-EM article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2022 Wilson et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Okabe T, Julien HM, Kaliyadan AG, et al. Prompt recognition of left ventricular free-wall rupture aided by the use of contrast echocardiography. *Tex Heart Inst J.* 2015;42(5):474-8.
2. Hammad TA, Parikh M, Tashtish N, et al. Impact of COVID-19

- pandemic on ST-elevation myocardial infarction in a non-COVID-19 epicenter. *Catheter Cardiovasc Interv.* 2020;97(2):208-14.
3. Moreno R, López de Sá E, López-Sendón JL, et al. Frequency of left ventricular free-wall rupture in patients with acute myocardial infarction treated with primary angioplasty. *Am J Cardiol.* 2000;85(6):757-60, A8.
 4. Kevil CG, Goeders NE, Woolard MD, et al. Methamphetamine use and cardiovascular disease. *Arterioscler Thromb Vasc Biol.* 2019;39:1739-46.
 5. Amir O, Smith R, Nishikawa A, et al. Left ventricular free wall rupture in acute myocardial infarction: a case report and literature review. *Tex Heart Inst J.* 2005;32(3):424-6.