

UC Riverside

UC Riverside Previously Published Works

Title

Juvenile hormone and its receptor methoprene-tolerant promote ribosomal biogenesis and vitellogenesis in the *Aedes aegypti* mosquito

Permalink

<https://escholarship.org/uc/item/5mc5q7kz>

Journal

Journal of Biological Chemistry, 292(24)

ISSN

0021-9258

Authors

Wang, Jia-Lin
Saha, Tusar T
Zhang, Yang
[et al.](#)

Publication Date

2017-06-01

DOI

10.1074/jbc.m116.761387

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Juvenile hormone and its receptor methoprene-tolerant promote ribosomal biogenesis and vitellogenesis in the *Aedes aegypti* mosquito

Received for publication, September 30, 2016, and in revised form, April 24, 2017. Published, Papers in Press, April 26, 2017, DOI 10.1074/jbc.M116.761387

Jia-Lin Wang^{‡§}, Tusar T. Saha^{‡¶}, Yang Zhang^{‡||}, Changyu Zhang^{‡**}, and Alexander S. Raikhel^{‡¶1}

From the [‡]Department of Entomology, University of California, Riverside, California 92521, [§]Hubei Key Laboratory of Genetic Regulation and Integrative Biology, School of Life Sciences, Central China Normal University, Wuhan 430079, China, [¶]Institute for Integrative Genomic Biology, University of California, Riverside, California 92521, ^{||}Key Laboratory of Tropical Marine Bio-Resources and Ecology, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou 510301, China, and ^{**}Guizhou Key Laboratory for Plant Pest Management of Mountain Region, College of Agriculture, Guizhou University, Guiyang 550025, China

Edited by Ronald C. Wek

Juvenile hormone (JH) controls many biological activities in insects, including development, metamorphosis, and reproduction. In the *Aedes aegypti* mosquito, a vector of dengue, yellow fever, chikungunya, and Zika viruses, the metabolic tissue (the fat body, which is an analogue of the vertebrate liver) produces yolk proteins for developing oocytes. JH is important for the fat body to acquire competence for yolk protein production. However, the molecular mechanisms of how JH promotes mosquito reproduction are not completely understood. In this study we show that stimulation of the JH receptor methoprene-tolerant (Met) activates expression of genes encoding the regulator of ribosome synthesis 1 (RRS1) and six ribosomal proteins (two ribosomal large subunit proteins, two ribosomal small subunit proteins, and two mitochondrial ribosomal proteins). Moreover, RNAi-mediated depletion of *RRS1* decreased biosynthesis of the ribosomal protein L32 (RpL32). Depletion of *Met*, *RRS1*, or *RpL32* led to retardation of ovarian growth and reduced mosquito fecundity, which may at least in part have resulted from decreased vitellogenin protein production in the fat body. In summary, our results indicate that JH is critical for inducing the expression of ribosomal protein genes and demonstrate that *RRS1* mediates the JH signal to enhance both ribosomal biogenesis and vitellogenesis.

As a vector of dengue, yellow fever, chikungunya, and Zika viruses, the *Aedes aegypti* mosquito causes severe morbidity and increased economic costs (1–4). Due to the lack of effective vaccines, controlling these dangerous vector-borne diseases remains a big challenge. Female hematophagous mosquitoes rely on blood intake to activate vitellogenesis, which involves

the production of vitellogenin (Vg)² the major yolk protein precursor (YPP) in the fat body, and is controlled by ecdysteroid 20-hydroxyecdysone (20E) (5, 6). A sesquiterpenoid juvenile hormone (JH) plays an essential role in stimulating mosquito reproduction during the previtellogenic preparatory period (7). Unraveling the molecular mechanisms regarding hormone control of reproduction will aid in development of novel strategies for prevention of vector-borne diseases.

JH, as an endocrine regulator, plays crucial roles in controlling insect development, metamorphosis, and reproduction. JH hemolymph level increases over the first 2 days posteclosion (PE), reaches its peak at 48–54 h PE, and maintains a high level in female *A. aegypti* before a blood meal (8). JH-dependent PE development is vital for a female fat body to become competent for the production of massive YPPs. The process is always associated with dramatic enlargement of nucleoli, development of Golgi complexes, and ribosome proliferation (9, 10). Removal of the corpora allata (CA; the source of JH) in newly emerged female mosquitoes blocks activation of fat body nucleoli for ribosomal and ribosomal RNA production, whereas re-implantation of CA or topical application of JH rescues the phenotypes (10, 11). This indicates that JH controls ribosome proliferation during PE development. However, the molecular mechanisms underlying JH regulation of ribosome proliferation are not well understood.

JH exerts its genomic function through the receptor methoprene-tolerant (Met), a member of the family of the basic helix-loop-helix (bHLH)-Per-Arnt-Sim (PAS) transcription factors (12, 13). A transcriptional steroid receptor coactivator (FISC/SRC/Taiman (Tai)) has been identified in the *A. aegypti* mosquito, the *Tribolium castaneum* beetle, and the *Bombyx mori* silkworm as a functional partner of Met (14–16). JH induces

This work was supported by National Institutes of Health Grant R01 AI036959 (to A. S. R.). This work was also supported by Scholar Fellowship 201506775010 (to J.-L. W.) from the China Scholarship Council of Chinese Ministry of Education. The authors declare that they have no conflicts of interest regarding the content of this article. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

This article contains supplemental Tables S1 and S2.

¹To whom correspondence should be addressed. E-mail: alexander.raikhel@ucr.edu.

²The abbreviations used are: Vg, vitellogenin; YPP, yolk protein precursor; 20E, 20-hydroxyecdysone; JH, juvenile hormone; PE, posteclosion; CA, corpora allata; Met, methoprene-tolerant; bHLH, basic helix-loop-helix; PAS, Per-Arnt-Sim; Tai, Taiman; iMet, Met RNAi depletion; *RRS1*, regulator of ribosome synthesis 1; *RpL32*, ribosomal protein large subunit 32; qRT-PCR, quantitative real-time PCR; PBM, post blood meal; CHX, cycloheximide; Kr-h1, Krüppel homolog 1; Gce, germ cell-expressed; *DmMet*, *Drosophila Met*; *DmTai*, *Drosophila Tai*; dsRNA, double-stranded RNA; NI, non-injected; FISC, β FTZ-F1 interacting steroid receptor coactivator.

the heterodimerization of Met with its partner FISC/SRC/Tai to bind E-box-like motifs in the regulatory regions of JH-target genes and activate the transcriptions (13–17).

The microarray screen combined with Met RNA interference (RNAi) has shown Met repression of early PE (at 6 h PE) and mid PE (at 24 h PE) genes but activation of late PE (at 66 h PE) genes (18). Met involvement in gene repression mediated by Hairy and a corepressor Groucho has been recently clarified (19). In the study, Saha *et al.* (19) performed a RNAi-based transcriptomic screen to identify genes co-regulated by Met and Hairy in the fat body. As a result, 1613 transcripts were shown to be activated, whereas 538 were suppressed after RNAi depletion of Met (iMet). In the current study we have shown that among 538 iMet-suppressed transcripts, many are those involved in ribosome biogenesis. One of them is a gene termed *RRS1* (regulator of ribosome synthesis 1, AAEL012185) that is the homolog protein of RRS1p identified in *Saccharomyces cerevisiae* that regulates ribosome synthesis. RRS1p, as an essential nuclear protein, is required for maturation of 25S rRNA and functions in assembly of the 60S ribosomal subunits (20, 21). RRS1p, along with Rpf2p, recruits ribosomal proteins RpL5 and RpL11 as well as 5S rRNA into preribosomes (22). However, studies of RRS1 in insects are limited. Here, we have demonstrated that *A. aegypti* *RRS1* and six ribosomal protein genes are activated by Met, and Met directly regulates the transcription of *RRS1*. RNAi-mediated depletion of *RRS1* resulted in a decreased level of ribosomal protein large subunit 32 (RpL32) protein. *Met*, *RRS1*, or *RpL32* depletion caused retardation of ovarian growth, which may at least partly result from a decreased Vg expression in the fat body. Our study provides new insight into JH-dependent ribosome proliferation and vitellogenesis.

Results

Met RNAi depletion suppressed the expression of genes involved in ribosome biogenesis

We previously identified 2151 differentially expressed transcripts controlled by Met in the fat body of female *A. aegypti* mosquitoes at the end of the PE phase, with 1613 transcripts down-regulated and 538 up-regulated (19). Among the latter, a total of 91 transcripts (17%) are involved in ribosome biogenesis; of these, 47 encode ribosomal protein large subunits, 30 encode ribosomal protein small subunits, and 14 encode ribosome biogenesis-related factors (Fig. 1A and supplemental Table S1). From these 14 genes, we selected the gene AAEL012185 encoding regulator of ribosome synthesis 1 (*RRS1*), which has been shown to regulate ribosome biosynthesis in *S. cerevisiae* (20).

To validate the RNA-seq results, *RRS1* along with two genes encoding ribosomal protein large subunits (AAEL003396, AAEL003324) and two genes encoding ribosomal protein small subunits (AAEL010168, AAEL014903) were selected for quantitative real-time PCR (qRT-PCR) analysis. As shown in Fig. 1, B–E and H, the qRT-PCR results confirm the RNA-seq experiments, with all tested genes down-regulated after Met RNAi depletion (iMet). We also measured the expression of two mitochondrial ribosomal protein genes (AET-4427, AET-7465), and




Figure 1. Depletion of Met-suppressed genes involved in ribosomal biogenesis. A, pie chart summarizing that transcripts involved in ribosome biogenesis are suppressed in Met-depleted fat bodies (*iMet*) of female mosquitoes. The transcript statistics were accepted with at least a 2-fold decrease of FPKM values. B–I, qRT-PCR confirming that the expression of six ribosomal protein genes; *RRS1* was suppressed in *iMet*. *NI*, mosquitoes without injection. *iLuc*, double-stranded RNA for the luciferase gene. Both *NI* and *iLuc* served as the controls. All expressions were calculated against housekeeping gene β -actin. Data are the mean \pm S.E. from three biological replicates. *, $p < 0.05$; **, $p < 0.01$ (*t* test).

the results indicate that they are suppressed in *iMet* (Fig. 1, F and G). Efficiency of Met RNAi depletion was also confirmed (Fig. 1I).

JH and Met promote ribosomal biogenesis and vitellogenesis




Figure 2. Developmental profiles of six ribosomal protein genes and *RRS1* were analyzed in the fat bodies of adult female *A. aegypti* using qRT-PCR. The color indicates the -fold change of the gene abundance in the form of a logarithm. The dendrograms are constructed based on an algorithm of hierarchical clustering.

Expression profiles of genes involved in ribosome biogenesis in the fat body correlated with the endogenous JH titers of adult female mosquitoes during the PE phase

JH secretion occurs soon after adult emergence. JH titers increase to their peak at 48–54 h PE and remain high until blood feeding (8). To clarify the relationship between the expression of genes involved in ribosome biogenesis and endogenous JH concentration, we measured the expression profiles of these genes at different developmental stages. Transcripts of all genes involved in ribosome biogenesis are closely correlated with the changes of endogenous JH titers. As shown in Fig. 2, expression levels of all ribosomal protein genes increased along with PE development, reached their highest level at 72 h PE or 12 h post blood meal (PBM), and declined to a low level, then began to increase again. *RRS1* transcript exhibited its highest level at 48 h PE and remained at the same high level at 60 h PE, decreasing by 72 h PE. The active transcription of *RRS1* was concomitant with the peak of endogenous JH, suggesting that *RRS1* responds to JH immediately.

RRS1 is transcriptionally regulated by the JH-receptor complex

To understand whether the action of Met on genes involved in ribosome biogenesis is direct or mediated by an intermediate factor, we employed *in vitro* fat body culture analyses and used the protein biosynthesis inhibitor cycloheximide (CHX). The expression levels of six ribosomal protein genes were considerably elevated by JH, and CHX inhibited JH-mediated induction of ribosomal protein genes (Fig. 3, A–F), suggesting that JH induced the expression of these ribosomal protein genes indirectly. However, CHX could not inhibit JH-mediated induction of *RRS1* (Fig. 3G), suggesting that JH induced the expression of *RRS1* directly. Krüppel homolog 1 (*Kr-h1*, AAEL002390), which has previously been identified as a direct target of Met (23), served as a control; its expression, which was activated by JH, was not inhibited by CHX (Fig. 3H).

To further determine whether Met directly regulates *RRS1* transcription, we conducted luciferase reporter assays. Mosquito bHLH-PAS transcription factors Met and FISC form a heterodimer in the presence of JH or JH analogs to act on JH response element (JHRE) and directly activate transcription of JH target genes (14). Here, the full lengths of Met (amino acid 1–977) and FISC (amino acid 1–1488) were expressed in S2 cells, with a C-terminal Myc tag or FLAG tag, respectively (Fig.

4A). Because E-box or E-box-like motifs are characteristic motifs recognized by bHLH-PAS transcription factors (24), we cloned the upstream regulatory sequences of *RRS1*, which included an E-box-like motif (CACGCG) into the pGL4.17 vector. To diminish the possible influence of endogenous Met, *Gce* (germ cell-expressed), and *Tai* in S2 cells, *Drosophila Met* (*DmMet*) and *Tai* (*DmTai*) double-stranded RNAs (dsRNAs) were added to the transfection mixture. After transfection, the expression of Met or FISC or Met plus FISC was confirmed by Western blot using anti-Myc or anti-FLAG antibodies (Fig. 4B). Treatment with *DmMet* and *DmTai* dsRNAs led to a significant reduction of endogenous *Met*, *Gce* (the paralog of Met, the dsRNA sequences targeting *DmMet* sharing 44% identity with that of *DmGce*), and *Tai* in S2 cells (Fig. 4C). However, the transcripts of transfected *Met-Myc* or *FISC-FLAG* were not influenced (Fig. 4D). In endogenous *Met*-, *Gce*-, and *Tai*-suppressed S2 cells, JH did not induce *RRS1* reporter activity in S2 cells. No protein expression or expression of either Met or FISC alone in S2 cells could not induce *RRS1* reporter activity. Co-expression of Met and FISC could not induce reporter activity either. However, after the addition of JH III or JH analog methoprene, co-expression of Met and FISC significantly induced reporter activity (Fig. 4B), which indicates that the JH-Met-FISC complex is required for *RRS1* transcription.

To determine whether the JH-receptor complex physically interacts with the *RRS1* promoter, we performed electrophoretic mobility shift assay (EMSA) using nuclear extracts from S2 cells overexpressing both Met-Myc and FISC-FLAG fusion proteins followed by JH III treatment. As shown in Fig. 4E, a band shift was detected in samples incubated with the labeled *RRS1* probe; this was abolished after preincubation with unlabeled *RRS1* probe, suggesting that the protein-DNA interaction is specific. The band was diminished when nuclear extracts were preincubated with anti-Myc or anti-FLAG antibody, suggesting that Met-Myc and FISC-FLAG constitute the binding complex. The band shift disappeared after mutation of E-box-like motif (Fig. 4F), suggesting this motif is crucial for the interaction.

RRS1 depletion reduced RpL32 protein but not transcript levels in the fat body of adult female mosquitoes

To elucidate the relationship between *RRS1* and ribosomal proteins, we performed RNAi-mediated depletion analyses. When compared with the control groups, newly emerged female mosquitoes (0–6 h PE) treated with *RRS1* dsRNA (*iRRS1*) displayed an efficient depletion of the *RRS1* transcript in the fat body at 72 h PE (Fig. 5B). We selected Rpl32 to test. The abundance of the *Rpl32* transcript showed no significant difference between *iRRS1* and the control groups (Fig. 5A). However, depletion of *RRS1* resulted in decreased Rpl32 protein in fat bodies at both 72-h PE and 24-h PBM (Fig. 5C), which suggests that Rpl32 is regulated by *RRS1*.

RRS1 or Rpl32 depletion resulted in abnormal ovarian development that mimics the phenotype of ovaries caused by depletion of Met

To assess the function of *RRS1* and Rpl32 in the adult female mosquito, we conducted RNAi-mediated depletion of *RRS1* (*iRRS1*) or Rpl32 (*iRpl32*) in newly emerged female mosqui-




Figure 3. Effects of JH, CHX, and JH plus CHX on the expression of six ribosomal protein genes and *RRS1* in the fat body cultured *in vitro*. Fat bodies dissected from newly emerged female mosquitoes were incubated for 12 h under different treatment conditions. Krüppel homolog 1 (AAEL002390) was used as a control. *Solvent*, medium with acetone; *JH*, medium with 1×10^{-6} M JH; *CHX*, medium with 1×10^{-5} M CHX; *JH+CHX*, medium with 1×10^{-6} M JH and 1×10^{-5} M CHX. All expression values were calculated against the housekeeping gene β -actin. Data are the mean \pm S.E. from three biological replicates. *, $p < 0.05$; **, $p < 0.01$ (t test).

toes and checked the ovaries dissected at both 72 h PE and 24 h PBM. We also performed RNAi depletion of Met (iMet) as a control, as it has been reported that iMet results in retardation of ovarian growth at 72 h PE (18, 19). Analysis of ovaries from non-injected (NI) female mosquitoes at 72 h PE and those injected with luciferase dsRNA (iLuc) demonstrated average primary follicle lengths of 105 μ m and 106 μ m, respectively. However, similar to previously shown iMet retardation of ovar-

ian follicle growth (with an average primary follicle length of 67 μ m), iRRS1 and iRpL32 were also shown to stall the growth, with average primary follicle lengths of 76 μ m and 70 μ m, respectively, which is much smaller than that of the controls (Fig. 6A).

In addition to 72 h PE, we evaluated the ovarian development at 24 h PBM. As shown in Fig. 6, B–D, ovaries from NI or iLuc at 24 h PBM developed normally, with follicle lengths reaching an

JH and Met promote ribosomal biogenesis and vitellogenesis




Figure 4. JH-receptor complex regulates the transcription of *RRS1*. A, schematic diagram of Met-Myc and FISC-FLAG fusion protein expressed in S2 cell. B, luciferase reporter assays after co-transfection of pGL4.17/*RRS1* –3389 to –2597 into S2 cells compared with the pAc5.1 empty vector, pAc5.1/Met-Myc, and/or pAc5.1/FISC-FLAG with or without JH III or methoprene. Western blots confirmed the expression of Met-Myc and/or FISC-FLAG in S2 cells. Antibodies against Myc or FLAG were used for detection. β -Actin was used for loading controls. Points with letters *a* and *b* represent significantly different groups ($p < 0.05$). C, efficiency of *Drosophila Met*, *Gce*, and *Tai* depletion in S2 cells after treatment with *DmMet* and *DmTai* dsRNAs. D, treatment with *DmMet* and *DmTai* dsRNAs resulted in no significant variation of Met-Myc or FISC-FLAG mRNA levels. iLuc served as a control. All expressions are calculated against housekeeping gene *Drosophila β -actin*. Data are the mean \pm S.E. from three biological replicates. *, $p < 0.05$; **, $p < 0.01$ (*t* test). E and F, EMSA confirming the binding of JH-Met-FISC complex to the *RRS1* probe. Mutation of E-box-like motif (CACGCG) to (TCAATA) abolished the binding capacity. Assays were tested with nuclear protein extracts from JH-treated S2 cells with overexpressed Met-Myc and FISC-FLAG fusion proteins. The arrow indicates the specific protein-DNA complex.

average of 214 μ m and 211 μ m, respectively. However, iMet, iRRS1, and iRPL32 significantly inhibited the ovarian development, with average primary follicle lengths of 103 μ m, 132 μ m, and 84 μ m, respectively. In addition, mosquitoes of NI and iLuc laid on average 121 and 113 eggs per female, respectively. However, iMet, iRRS1 and iRPL32 dramatically reduced fecundity, resulting in significantly fewer eggs laid: an average of 14 eggs per female with iMet, 26 with iRRS1, and 4 with iRPL32 (Fig. 6E).

Met, RRS1, or Rpl32 RNAi depletions resulted in the decreased level of Vg protein in the fat body of adult female mosquitoes

During mosquito vitellogenesis, Vg (the major YPP) is synthesized in and secreted from the fat body and is subsequently accumulated in developing oocytes (5). To check whether Met, RRS1, or Rpl32 is involved in Vg synthesis and rule out the possible involvement of other tissues after blood feeding, we performed *in vitro* fat body culture analyses. Newly emerged female mosquitoes were injected with dsRNA, and fat bodies isolated at 72 h post injection (during this stage, the fat body is reproductively competent) were subjected to *in vitro* fat body culture. Depletion of Met or Rpl32 was confirmed by means of Western blots (Fig. 7, E and F), whereas depletion of *RRS1* was

confirmed using qRT-PCR (Fig. 7G). Fat bodies from mosquitoes of iMet, iRRS1, or iRPL32 exhibited a much lower Vg protein level than those from mosquitoes of NI or iLuc (Fig. 7D). Depletion of Met significantly suppressed the abundance of Vg transcripts (Fig. 7A). However, iRRS1 and iRPL32 did not result in significant differences of the Vg transcript level when compared with the controls (Fig. 7, B and C).

Met, RRS1, or Rpl32 RNAi depletions resulted in a decreased level of polysomes in the fat body of adult female mosquitoes

To check whether the decreased level of Vg protein in the fat body of iMet, iRRS1, or iRPL32 is due to the decreased ribosomal biogenesis, we isolated polysomes using sucrose gradient sedimentation and determined the amount as the material absorbing at 254 nm. As shown in Fig. 8A, the amount of polysomes from fat bodies of iMet, iRRS1, or iRPL32 is much lower than that of the control group, suggesting the involvement of Met, RRS1, or Rpl32 in ribosomal biogenesis.

Discussion

Vg synthesis can be induced by JH in many insects (25–27). In mosquitoes, the massive production of Vg only occurs after blood feeding and is primarily governed by 20E (28). However,




Figure 5. Depletion of Met or RRS1 results in decreased expression of RpL32 proteins in the female fat body. *A*, the transcripts of Rpl32 exhibited no significant differences in RRS1-depleted (*iRRS1*) fat bodies of female mosquitoes at 72 h PE. *B*, qRT-PCR confirming the depletion of RRS1. *NI*, mosquitoes without injection; *iLuc*, dsRNA for the luciferase gene. All expression values are calculated against the housekeeping gene β -actin. Data are the mean \pm S.E. from three biological replicates. **, $p < 0.01$ (t test). *C*, RNAi-mediated depletion of Met or RRS1 resulted in decreased expression of Rpl32 in the female fat bodies at both 72 h PE and 24 h PBM. Shown are Western blot analyses using antibodies against Rpl32 and β -actin (as control) in fat bodies from non-injected (*NI*), luciferase dsRNA-treated (*iLuc*), Met dsRNA-treated (*iMet*), or *RRS1* dsRNA-treated (*iRRS1*) mosquitoes. Both *NI* and *iLuc* served as the controls.

in vitro fat body culture demonstrated that Vg expression can only be stimulated by 20E in fat bodies dissected from the late stage of PE or from newly emerged mosquitoes on the condition that their fat bodies are preincubated with JH (7). These results suggest that it is JH that prompts the fat body to become fully competent during PE development. The proliferation of ribosomes and ribosomal RNA has been demonstrated to be one of several critical JH-mediated events during PE (10). Because ribosome biogenesis involves production of ribosomal RNA and, finally, assembly with ribosomal proteins to form the subunits, we speculated that the activation of ribosomal protein genes should also be coordinated by JH. Indeed, our RNA-seq results identified 91 transcripts involved in ribosome biogenesis being activated by Met, accounting for 17% of Met-activation transcripts. qRT-PCR confirmed the expression of six ribosomal protein genes suppressed in *iMet*. The transcription of these ribosomal protein genes gradually intensified along with the increased JH titers during PE development, mostly reaching a maximum at 72 h PE. The active transcription of ribosomal protein genes generally correlates positively with the activation of nucleolus and synthetic rate of ribosomal RNA in the fat body cells during PE, suggesting that JH controls the proliferation of ribosomes to sustain an enhanced translation rate after blood feeding.

JH could induce the expression of ribosomal protein genes, whereas CHX suppressed the JH-mediated inductions, suggesting there is an intermediate factor involved that still needs to be identified. Intriguingly, CHX could not suppress JH-mediated induction of *RRS1*, suggesting that JH induces the transcription of this gene directly. *RRS1* reaches its peak expression at 48 h PE, closely correlating with the elevated JH titers and suggesting that expression of *RRS1* responds to JH quicker than that of ribosomal protein genes. Depletion of Met resulted in a




Figure 6. Depletion of Met, RRS1, or Rpl32 resulted in decreased follicle length and dramatically reduced egg number. *A*, primary ovarian follicle length (in μ m) measured in *iMet*, *iRRS1*, *iRpL32*, *iLuc*, and non-injected (*NI*) mosquitoes at 72 h PE. *B* and *C*, female mosquito ovaries at 24 h PBM. *D*, average follicle length at 24 h PBM. *E*, egg numbers deposited by each female mosquito. *NI*, non-injected; *iLuc*, luciferase dsRNA-treated; *iMet*, Met dsRNA-treated; *iRRS1*, *RRS1* dsRNA-treated; *iRpL32*, *Rpl32* dsRNA-treated mosquitoes. Error bars represent \pm S.D. ****, $p < 0.0001$ (t test).

remarkable decrease of *RRS1* abundance. It has been reported that *A. aegypti* Met and FISC form a complex in the presence of JH to activate transcription of JH target genes directly, such as the early trypsin gene (14) and Kr-h1 gene (23). Here, our luciferase reporter assays demonstrated that the JH-receptor complex composed of Met and FISC in the presence of JH or JH analog methoprene significantly induced the transcription of *RRS1*. EMSA using nuclear extract from S2 cells overexpressing Met-Myc and FISC-FLAG confirmed the specific binding of JH-receptor complex to E-box-like motif in the upstream of *RRS1* gene.

RRS1, characterized as an essential nuclear protein in *S. cerevisiae*, plays a critical role in pre-rRNA processing and assembly of ribosomal subunits (20). It was further speculated that *RRS1* is not required for transcription of ribosomal protein genes but plays a role in maintaining the structure of preribosomal subunit particles (20). Our results indicated that *RRS1* RNAi depletion resulted in no significant differences in the Rpl32 transcript level. However, depletion of *RRS1* reduced Rpl32 proteins, similar to the observation in *Drosophila melanogaster* that depletion of Nopp140 (a ribosomal assembly fac-

JH and Met promote ribosomal biogenesis and vitellogenesis




Figure 7. Depletion of Met, RRS1, or RpL32 resulted in decreased expression of Vg proteins in the female fat body. A, depletion of Met led to a dramatic decline in Vg transcripts. Depletion of RRS1 (B) or RpL32 (C) resulted in no significant differences in Vg transcripts. D, depletion of Met, RRS1, or RpL32 resulted in decreased Vg proteins. Western blot confirming the depletion of Met (E) or RpL32 (F) in fat bodies. G, qRT-PCR confirming the depletion of RRS1. Fat bodies from non-injected (NI), luciferase dsRNA (iLuc)-, Met dsRNA (iMet)-, RRS1 dsRNA (iRRS1)-, or RpL32 dsRNA (iRpL32)-treated female mosquitoes were subjected to *in vitro* culture. Shown are Western blot analyses using antibodies against Vg, Met, RpL32, or β -actin (as control). Error bars represent \pm S.E. *, $p < 0.05$; **, $p < 0.01$ (t test).

tor) led to decreased ribosomal protein L34 (29, 30). Mutation of the *RRS1* gene in yeast leads to a compromised 60S subunit production by affecting ribosome assembly and thus decreased polysomal content (20). A similar decrease in the level of polysomes was observed in RRS1-depleted *A. aegypti* in our study. We speculate that the absence of RRS1 leads to impaired ribosomal assembly in mosquitoes as well. The newly synthesized unassembled RpL32 protein is unstable and is rapidly degraded by the system (31), resulting in the observed decreased protein levels. However, this hypothesis needs to be tested in the future. We would like to stress that we presume this to be the general mechanism by which RRS1 affects the level of the ribosomal proteins. Although providing additional immunoblots for other RpLs is beyond the scope of this manuscript, we furnish the results for RpL32 as a representative.

Our observations indicate that RNAi Met depletion not only causes retardation of ovarian growth at 72 h PE, as reported previously (18, 19), but also severely stalls the growth of ovarian follicles at 24 h PBM and dramatically reduces egg production. Depletion of RRS1 or RpL32 resulted in severe defects linked to ovarian development and egg deposition, which mimics the phenotypes caused by depletion of Met, agreeing well with the above conclusion that the JH-receptor complex targets RRS1 to control the expression of RpL32. Different ribosomal proteins may participate in regulating different physiological functions, such as cell growth and apoptosis (32) and reproductive diapause (33). In *A. aegypti*, Estep *et al.* (34) recently demonstrated that depletion of RpL26 or RpS6 could significantly reduce fecundity, whereas depletion of RpL1 or RpS2 could not. Similar to the phenotype caused by depletion of RpL26 or RpS6, our results show that depletion of RpL32 stalls the growth of ovarian follicles and dramatically reduces fecundity.

Drastically smaller primary ovarian follicles usually represent that YPP synthesis, secretion, and/or uptake are compromised. For example, smaller ovarian follicles have resulted from depletion of miR-8, which is responsible for the proper secretion of lipophorin and Vg by the fat body (35). Considering that RpL32 is a component of the translational apparatus, we speculate that the smaller primary ovarian follicles caused by depletion of Met, RRS1, or RpL32 may have resulted from the decreased YPP synthesis in fat bodies. Because RNAi depletion has a systemic effect on the target gene, we performed RNAi combined with *in vitro* fat body culture experiments to rule out the possible involvement of other tissues. The results demonstrated that depletion of Met, RRS1, or RpL32 dramatically reduced the expression of Vg proteins, which may at least partly account for the decreased primary follicles and reduced fecundity. Furthermore, depletion of Met significantly reduced the abundance of Vg transcripts, similar to the observation in the linden bug, where Met RNAi suppressed Vg gene expression in the fat body and blocked ovarian development (36). This result is reasonable as some factors identified downstream of JH-Met signaling pathway control Vg mRNA expression. For example, β FZ-F1, which could be stimulated by JH in the fat body, is involved in regulating the expression of Vg transcripts as a competence factor (7). However, depletion of RRS1 or RpL32 led to insignificant differences in Vg transcripts, suggesting that RRS1 or RpL32 may only contribute to the translational apparatus and be responsible for Vg translation. The protein translation efficiency is associated with the ribosomes bound to form polysomes. Depletion of Met, RRS1, or RpL32 decreased the level of polysomes, indicating that these genes are involved in ribosomal biogenesis and further explaining why a decreased




Figure 8. A, depletion of Met, RRS1, or Rpl32 resulted in decreased levels of polysomes in the female fat body. Polysomes were extracted from fat bodies of luciferase dsRNA (*iLuc*), *Met* dsRNA (*iMet*), *RRS1* dsRNA (*iRRS1*) or *RpL32* dsRNA (*iRpL32*)-treated mosquitoes at 72 h PE, and the absorbance at 254 nm was determined. Error bars represent \pm S.E. *, $p < 0.05$; **, $p < 0.01$ (t test). B, a proposed model for the function of RRS1 in transducing JH signal to mediate ribosomal biogenesis and vitellogenesis. JH regulates many ribosomal protein genes (including *RpL32*). JH-receptor complex acts on the promoter of *RRS1* to regulate its transcription. Afterward, RRS1 mediates ribosomal biogenesis and vitellogenesis through regulation of the expression of Rpl32 proteins.

expression of Vg protein was observed when depleting these genes.

In conclusion, we demonstrated that JH could regulate many ribosomal protein genes, and JH-receptor complex acts on the promoter of *RRS1* to regulate its transcription. RRS1 transduces JH signal in mediating ribosomal biogenesis and vitellogenesis through regulation of the expression of Rpl32 proteins (Fig. 8B). Future studies should focus on identification of genes that mediate the JH signal to regulate the transcripts of ribosomal protein genes, which will further deepen our understanding of JH-mediated ribosomal biogenesis and vitellogenesis.

Experimental procedures

Mosquito rearing

A. aegypti mosquitoes were maintained at 27 °C and 86% relative humidity, as described previously (19). Adult mosquitoes were fed water and a 10% sucrose solution continuously via wicks. Blood feeding was conducted using white Leghorn chickens following instructions approved by the Animal Care and Use Committee, University of California, Riverside, CA. All dissections were carried out in *Aedes* physiological solution (38).

RNAi

The dsRNAs of *Met*, *RRS1*, and *RpL32* were synthesized using the MEGAscript kit (Ambion, Austin, TX) as described previously (38). The bacterial luciferase gene was used as a template to produce control *iLuc* dsRNA, and synthesized dsRNAs were purified using MEGAclear kit (Ambion). Subsequently, the corresponding dsRNA was quantified to 4 $\mu\text{g}/\mu\text{l}$, and 300-nl samples of dsRNA were microinjected into the thoraxes of CO₂-anesthetized newly emerged female mosquitoes (0–6 h PE) using Picospritzer II (General Valve Corp., Fairfield, NJ). Primers used for dsRNA synthesis are listed in supplemental Table S2.

In vitro fat body culture

Fat body culture analyses were performed according to a method described previously (38). To test the effects of JH (Sigma) or CHX (Sigma) on gene expression, fat bodies dissected from newly emerged female mosquitoes (0–6 h PE) were incubated in culture medium with 1×10^{-6} M JH III or solvent (acetone) or incubated in the medium with 1×10^{-5} M CHX combined with or without JH. After incubation for 12 h, samples were harvested for qRT-PCR analysis.

To analyze the expression of Vg resulting from RNAi depletion of *Met*, *RRS1*, or *RpL32*, fat bodies were dissected from female mosquitoes 72 h after dsRNA injection. Subsequently, fat bodies were incubated in a complete culture medium supplemented with 1×10^{-6} M 20E (Sigma) and amino acids. After incubation for 6 h, fat bodies were collected for qRT-PCR or Western blot analyses.

RNAi in S2 cells and luciferase reporter assay

The dsRNAs of *DmMet* and *DmTai* were synthesized using the MEGAscript kit (Ambion). The promoter region of *RRS1* (nucleotides –3389 to –2597) was synthesized (GenScript, Piscataway, NJ) and subcloned into reporter vector pGL4.17 (Promega, Madison, WI). *A. aegypti* *Met* (nucleotides 1–2931) and *FISC* (nucleotides 1–4464) DNA fragments were amplified and subcloned into a pAc5.1 vector (Invitrogen) fused with Myc and FLAG tag, respectively. The primers used for dsRNA synthesis and PCR amplification are listed in supplemental Table S2. All recombinant plasmids were confirmed by sequencing. Transient transfections were performed in *Drosophila* S2 cells using FuGENE HD transfection reagent (Promega) following the protocol. *DmMet* and *DmTai* dsRNAs were added together with plasmids in the transfection mixture according to a method described previously (39). Each well of S2 cells was co-transfected with 100 ng of desired reporter plasmid, 10 ng of the control *Renilla* luciferase reporter plasmid *pCop*_{II}, and 1.5 μg of each dsRNA. In addition, some wells were also co-transfected with pAc5.1/*Met*-Myc or pAc5.1/*FISC*-FLAG or both. JH III or the JH analog methoprene was added to the wells with a final concentration of 20 μM at 42 h post transfection. Luciferase activities were assayed at 48 h post transfection using the Dual Luciferase Assay kit (Promega) according to protocol.

qRT-PCR

Total RNAs of fat bodies from different developmental stages (PE: 0 h, 12 h, 24 h, 36 h, 48 h, 60 h, and 72 h; PBM: 12 h, 24 h,

JH and Met promote ribosomal biogenesis and vitellogenesis

36 h, 48 h, 60 h, and 72 h), RNAi, or *in vitro* culture treatments were extracted using TRIzol (Invitrogen) following the manufacturer's instructions. Total RNAs from S2 cells were also extracted. Approximately 2 μ g of total RNA was first treated with DNase I and then applied to synthesize cDNAs using SuperScript II reverse transcriptase (Invitrogen). qRT-PCR reactions were processed in 96-well plates with iQ SYBR Green Supermix (Bio-Rad) using the iCycler iQ system (Bio-Rad). Quantitative measurements for each gene were performed in triplicate and normalized against the reference gene β -actin using the $2^{-\Delta CT}$ calculation method. The expression profiles of genes at different developmental stages are shown in the heatmap generated by R3.2.0 software based on the mean value of $2^{-\Delta CT}$ of each time point. Primers designed for qRT-PCR are listed in supplemental Table S2.

Western blot

Total proteins from fat bodies after RNAi treatment, *in vitro* culture treatments, or S2 cells were extracted, quantified, and applied to Western blot analyses as described previously (38). Aliquots of protein samples were resolved on 4–20% mini-protein TGX pre-cast gels (Bio-Rad) and transferred to polyvinylidene difluoride membranes. Rabbit polyclonal antibodies against RpL32 (Abcam, Eugene, OR) were used at a 1:5000 dilution to detect RpL32 proteins followed by the secondary anti-rabbit HRP (Abcam). Mouse monoclonal antibodies against Vg (40) were used at a 1:5000 dilution to detect Vg followed by the secondary anti-mouse HRP (Abcam). Rabbit anti-Myc antibodies (Sigma) were used to detect Met-Myc fusion proteins, whereas rabbit anti-FLAG antibodies (Sigma) were used to recognize FISC-FLAG. Monoclonal antibodies against β -actin (Sigma) were used as a loading control.

EMSA

RRS1 probe (5'-CACTTTTGTGTTTCCACGCGAGAGGCAAGA-3') and RRS1 mutant probe (5'-CACTTTTGTGTTTCTCAATAAGAGGCAAGA-3') were designed for EMSA assays. We performed EMSA according to a method described previously (18). Briefly, the annealed DNA oligonucleotides were end-labeled with [γ - 32 P]ATP by T4 polynucleotide kinase (Promega). Nuclear protein extracts from S2 cells were prepared with NE-PER Nuclear and Cytoplasmic Extraction Reagents (Thermo Scientific, Chino, CA). The DNA-protein complex was resolved on 5% TBE Criterion Precast Gel (Bio-Rad). Subsequently, the gel was dried, incubated with phosphor imaging screen, and visualized using autoradiography. A 50-fold amount of unlabeled RRS1 probe was preincubated with the nuclear extracts before the addition of labeled probe in the competition assays. The anti-Myc antibody (Sigma), anti-FLAG antibody (Sigma), or the rabbit IgG (Sigma) was preincubated with nuclear extracts before the addition of labeled probe in the supershift assays.

Determination of polysomes

We isolated polysomes according to a method described previously (37). Fifty fat bodies dissected from female mosquitoes of each dsRNA treatment were immediately frozen in liquid nitrogen and homogenized in 1 ml of lysis buffer (20 mM Tris-

HCl (pH 8.0), 140 mM KCl, 1.5 mM MgCl₂, 1% Triton X-100, 1% sodium deoxycholate, 0.2 units/ml RNase inhibitor, 1 mg/ml heparin, and 1 mM dithiothreitol). The homogenate was then centrifuged for 10 min at 12,000 \times g. The supernatant was layered on a 20–60% linear sucrose gradient prepared in 20 mM Tris-HCl (pH 8.0), 140 mM KCl, 1.5 mM MgCl₂, and 0.1 mg/ml CHX. The gradients were then centrifuged at 120,000 \times g for 3 h at 4 $^{\circ}$ C. The fractions containing polysomes were pooled, and the amount of polysomes was determined as the material absorbing at 254 nm.

Author contributions—J.-L. W., T. T. S., and A. S. R. designed the research. J.-L. W., T. T. S., Y. Z., and C. Z. performed the research. J.-L. W., A. S. R., T. T. S., and Y. Z. contributed new reagents/analytic tools. J.-L. W., A. S. R., T. T. S., and Y. Z. analyzed the data. J.-L. W. and A. S. R. wrote the paper.

References

1. Barrett, A. D., and Higgs, S. (2007) Yellow fever: a disease that has yet to be conquered. *Annu. Rev. Entomol.* **52**, 209–229
2. Ernst, K. C., Walker, K. R., Reyes-Castro, P., Joy, T. K., Castro-Luque, A. L., Diaz-Caravantes, R. E., Gameros, M., Haenchen, S., Hayden, M. H., Monaghan, A., Jeffrey-Gutierrez, E., Carrière, Y., and Riehle, M. R. (2017) *Aedes aegypti* (Diptera: Culicidae) longevity and differential emergence of dengue fever in two cities in Sonora, Mexico. *J. Med. Entomol.* **54**, 204–211
3. Tsetsarkin, K. A., Chen, R., and Weaver, S. C. (2016) Interspecies transmission and chikungunya virus emergence. *Curr. Opin. Virol.* **16**, 143–150
4. Weaver, S. C., Costa, F., Garcia-Blanco, M. A., Ko, A. I., Ribeiro, G. S., Saade, G., Shi, P. Y., and Vasilakis, N. (2016) Zika virus: History, emergence, biology, and prospects for control. *Antiviral Res.* **130**, 69–80
5. Raikhel, A. S., and Dhadialla, T. S. (1992) Accumulation of yolk proteins in insect oocytes. *Annu. Rev. Entomol.* **37**, 217–251
6. Attardo, G. M., Hansen, I. A., and Raikhel, A. S. (2005) Nutritional regulation of vitellogenesis in mosquitoes: implications for anaotogeny. *Insect Biochem. Mol. Biol.* **35**, 661–675
7. Zhu, J., Chen, L., and Raikhel, A. S. (2003) Posttranscriptional control of the competence factor betaFTZ-F1 by juvenile hormone in the mosquito *Aedes aegypti*. *Proc. Natl. Acad. Sci. U.S.A.* **100**, 13338–13343
8. Zhao, B., Hou, Y., Wang, J., Kokoza, V. A., Saha, T. T., Wang, X. L., Lin, L., Zou, Z., and Raikhel, A. S. (2016) Determination of juvenile hormone titers by means of LC-MS/MS/MS and a juvenile hormone-responsive Gal4/UAS system in *Aedes aegypti* mosquitoes. *Insect Biochem. Mol. Biol.* **77**, 69–77
9. Raikhel, A. S., and Lea, A. O. (1983) Previtellogenic development and vitellogenin synthesis in the fat body of a mosquito: an ultrastructural and immunocytochemical study. *Tissue Cell* **15**, 281–299
10. Raikhel, A. S., and Lea, A. O. (1990) Juvenile hormone controls previtellogenic proliferation of ribosomal RNA in the mosquito fat body. *Gen. Comp. Endocrinol.* **77**, 423–434
11. Raikhel, A. S., and Lea, A. O. (1991) Control of follicular epithelium development and vitelline envelope formation in the mosquito; role of juvenile hormone and 20-hydroxyecdysone. *Tissue Cell* **23**, 577–591
12. Miura, K., Oda, M., Makita, S., and Chinzei, Y. (2005) Characterization of the *Drosophila* methoprene-tolerant gene product: juvenile hormone binding and ligand-dependent gene regulation. *FEBS J.* **272**, 1169–1178
13. Charles, J. P., Iwema, T., Epa, V. C., Takaki, K., Rynes, J., and Jindra, M. (2011) Ligand-binding properties of a juvenile hormone receptor, methoprene-tolerant. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 21128–21133
14. Li, M., Mead, E. A., and Zhu, J. (2011) Heterodimer of two bHLH-PAS proteins mediates juvenile hormone-induced gene expression. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 638–643
15. Zhang, Z., Xu, J., Sheng, Z., Sui, Y., and Palli, S. R. (2011) Steroid receptor co-activator is required for juvenile hormone signal transduction through

- a bHLH-PAS transcription factor, methoprene tolerant. *J. Biol. Chem.* **286**, 8437–8447
16. Kayukawa, T., Minakuchi, C., Namiki, T., Togawa, T., Yoshiyama, M., Kamimura, M., Mita, K., Imanishi, S., Kiuchi, M., Ishikawa, Y., and Shinoda, T. (2012) Transcriptional regulation of juvenile hormone-mediated induction of Krüppel homolog 1, a repressor of insect metamorphosis. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 11729–11734
 17. Guo, W., Wu, Z., Song, J., Jiang, F., Wang, Z., Deng, S., Walker, V. K., and Zhou, S. (2014) Juvenile hormone-receptor complex acts on mcm4 and mcm7 to promote polyploidy and vitellogenesis in the migratory locust. *PLoS Genet.* **10**, e1004702
 18. Zou, Z., Saha, T. T., Roy, S., Shin, S. W., Backman, T. W., Girke, T., White, K. P., and Raikhel, A. S. (2013) Juvenile hormone and its receptor, methoprene-tolerant, control the dynamics of mosquito gene expression. *Proc. Natl. Acad. Sci. U.S.A.* **110**, E2173–E2181
 19. Saha, T. T., Shin, S. W., Dou, W., Roy, S., Zhao, B., Hou, Y., Wang, X. L., Zou, Z., Girke, T., and Raikhel, A. S. (2016) Hairy and Groucho mediate the action of juvenile hormone receptor methoprene-tolerant in gene repression. *Proc. Natl. Acad. Sci. U.S.A.* **113**, E735–E743
 20. Tsuno, A., Miyoshi, K., Tsujii, R., Miyakawa, T., and Mizuta, K. (2000) RRS1, a conserved essential gene, encodes a novel regulatory protein required for ribosome biogenesis in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* **20**, 2066–2074
 21. Miyoshi, K., Shirai, C., Horigome, C., Takenami, K., Kawasaki, J., and Mizuta, K. (2004) Rrs1p, a ribosomal protein L11-binding protein, is required for nuclear export of the 60S pre-ribosomal subunit in *Saccharomyces cerevisiae*. *FEBS Lett.* **565**, 106–110
 22. Zhang, J., Harnpicharnchai, P., Jakovljevic, J., Tang, L., Guo, Y., Oeffinger, M., Rout, M. P., Hiley, S. L., Hughes, T., and Woolford, J. L., Jr. (2007) Assembly factors Rpf2 and Rrs1 recruit 5S rRNA and ribosomal proteins rpL5 and rpL11 into nascent ribosomes. *Genes Dev.* **21**, 2580–2592
 23. Cui, Y., Sui, Y., Xu, J., Zhu, F., and Palli, S. R. (2014) Juvenile hormone regulates *Aedes aegypti* Krüppel homolog 1 through a conserved E box motif. *Insect Biochem. Mol. Biol.* **52**, 23–32
 24. Massari, M. E., and Murre, C. (2000) Helix-loop-helix proteins: regulators of transcription in eucaryotic organisms. *Mol. Cell. Biol.* **20**, 429–440
 25. Wyatt, G. R., and Davey, K. G. (1996) Cellular and molecular actions of juvenile hormone. II. Roles of juvenile hormone in adult insects. *Adv. Insect Physiol.* **26**, 1–155
 26. Belles, X. (2005) Vitellogenesis directed by juvenile hormone, in reproductive biology of invertebrate: progress in vitellogenesis (Raikhel, A. S., ed.) Vol. 12, pp 157–197, Science Publishers Inc.
 27. Parthasarathy, R., Sun, Z., Bai, H., and Palli, S. R. (2010) Juvenile hormone regulation of vitellogenin synthesis in the red flour beetle, *Tribolium castaneum*. *Insect Biochem. Mol. Biol.* **40**, 405–414
 28. Roy, S., Smykal, V., Johnson, L., Saha, T. T., Zou, Z., and Raikhel, A. S. (2016) Regulation of reproductive processes in female mosquitoes. In *Advances in Insect Physiology: Progress in Mosquito Research* (Raikhel, A. S., ed) Academic Press, London; Vol. 51, pp. 115–144
 29. James, A., Cindass, R., Jr, Mayer, D., Terhoeve, S., Mumphrey, C., and DiMario, P. (2013) Nucleolar stress in *Drosophila melanogaster*: RNAi-mediated depletion of Nopp140. *Nucleus* **4**, 123–133
 30. He, F., James, A., Raje, H., Ghaffari, H., and DiMario, P. (2015) Deletion of *Drosophila* Nopp140 induces subcellular ribosomopathies. *Chromosoma* **124**, 191–208
 31. Lam, Y. W., Lamond, A. I., Mann, M., and Andersen, J. S. (2007) Analysis of nucleolar protein dynamics reveals the nuclear degradation of ribosomal proteins. *Curr. Biol.* **17**, 749–760
 32. Naora, H. (1999) Involvement of ribosomal proteins in regulating cell growth and apoptosis: translational modulation or recruitment for extraribosomal activity? *Immunol. Cell Biol.* **77**, 197–205
 33. Kim, M., and Denlinger, D. L. (2010) A potential role for ribosomal protein S2 in the gene network regulating reproductive diapause in the mosquito *Culex pipiens*. *J. Comp. Physiol. B* **180**, 171–178
 34. Estep, A. S., Sanscrainte, N. D., and Becnel, J. J. (2016) DsRNA-mediated targeting of ribosomal transcripts RPS6 and RPL26 induces long-lasting and significant reductions in fecundity of the vector *Aedes aegypti*. *J. Insect Physiol.* **90**, 17–26
 35. Lucas, K. J., Roy, S., Ha, J., Gervaise, A. L., Kokoza, V. A., and Raikhel, A. S. (2015) MicroRNA-8 targets the Wingless signaling pathway in the female mosquito fat body to regulate reproductive processes. *Proc. Natl. Acad. Sci. U.S.A.* **112**, 1440–1445
 36. Smykal, V., Bajgar, A., Provaznik, J., Fexova, S., Buricova, M., Takaki, K., Hodkova, M., Jindra, M., and Dolezel, D. (2014) Juvenile hormone signaling during reproduction and development of the linden bug, *Pyrrhocoris apterus*. *Insect Biochem. Mol. Biol.* **45**, 69–76
 37. Bian, G., Raikhel, A. S., and Zhu, J. (2008) Characterization of a juvenile hormone-regulated chymotrypsin-like serine protease gene in *Aedes aegypti* mosquito. *Insect Biochem. Mol. Biol.* **38**, 190–200
 38. Roy, S. G., Hansen, I. A., and Raikhel, A. S. (2007) Effect of insulin and 20-hydroxyecdysone in the fat body of the yellow fever mosquito, *Aedes aegypti*. *Insect Biochem. Mol. Biol.* **37**, 1317–1326
 39. Jindra, M., Uhlírova, M., Charles, J. P., Smykal, V., and Hill, R. J. (2015) Genetic evidence for function of the bHLH-PAS protein Gce/Met as a juvenile hormone receptor. *PLoS Genet.* **11**, e1005394
 40. Raikhel, A. S., Pratt, L. H., and Lea, A. O. (1986) Monoclonal antibodies as probes for processing of yolk protein in the mosquito; production and characterization. *J. Insect Physiol.* **32**, 879–890