

UCLA

Proceedings of UCLA Health

Title

A Case of Sexually Acquired Campylobacter Infection through Analingus

Permalink

<https://escholarship.org/uc/item/5qg356dk>

Journal

Proceedings of UCLA Health, 26(1)

Authors

CHen, Phillip

Yen, George

Publication Date

2022-05-19

CLINICAL VIGNETTE

A Case of Sexually Acquired *Campylobacter* Infection through Analingus

Phillip Chen, MD and George Yen, MD

Case Presentation

A 34-year-old cis-gender gay male with no significant past medical history presents with 4 days of acute diarrhea and abdominal pain. Patient reported up to 6 non-bloody, non-mucous, non-foul-smelling bowel movements per day. They started as loose stools and became clear watery diarrhea. He denies any nausea, vomiting, subjective fevers or chills. Patient also reports a 3/10 paraumbilical abdominal pain that is sharp in quality without radiation. He denies any unusual food intake. He ate a burger and nachos a day ago with others without similar symptoms. He denies consumption of unpasteurized or undercooked foods as well as any recent antibiotics or hospitalizations. He works as an events manager, works primarily from home and has no recent travel or hiking trips. He did note a trip to Hawaii a few months prior. He denies any contact with animals or animal products, denies food allergies, or family history of gastrointestinal or endocrine conditions. His sexual history, includes three recent sexual partners including oral-anal contact, a few days to weeks prior to symptom onset. He was previously on emtricitabine/tenofovir disoproxil (TDF/FTC) for HIV prevention but has stopped this medication.

On examination, his abdomen was soft, nontender, nondistended, without guarding or rebound. He was hemodynamically stable and afebrile. Comprehensive metabolic panel was all within normal limits. Erythrocyte sedimentation rate was normal and C-reactive protein was elevated to 2.7. HIV RNA PCR test was undetectable. A comprehensive stool evaluation included a parasitic enteric pathogen panel which was negative for giardia or entamoeba. A bacterial enteric pathogen panel was negative for shigella or salmonella but returned positive for campylobacter species. He was treated with azithromycin 500mg daily for three days and symptoms resolved.

Discussion

The CDC estimates 1.3 million people in the United States become ill from *Campylobacter* every year.¹ *Campylobacter* organisms are gram-negative, rod-shaped bacteria with an estimated infection incidence of 14.3/100,000 in North America.² There are 26 known species and specifically, *Campylobacter* Jejuni and *Campylobacter* coli represents majority of the gastroenteritis cases worldwide. Symptoms often include acute onset of watery diarrhea or bloody stools, fever, weight loss, and abdominal cramps, nausea, vomiting that lasts up to 6 days. Risk factors for *Campylobacter* infections include younger age (1-4 and 15-24 years old), travel to developing areas with

limited sanitation resources and consumption of undercooked poultry products, unpasteurized milk, and water.³ Immunodeficiency is also a risk factor. For example, HIV-infected patients presenting with diarrhea are more frequently infected with *Campylobacter* than uninfected individuals with diarrhea. Specifically, *campylobacter jejuni* has been known to be associated with the development of Guillain Barre Syndrome and the Miller Fisher variant. Other campylobacter species infections have also been associated with increased incidence of inflammatory bowel disease, gastroesophageal reflux disease, and through modest-sized studies, colorectal cancer.^{3,4}

Recent literature suggests other risk factors for campylobacter infection. Several studies have highlighted campylobacter infections in men who have sex of with men (MSM). A retrospective study in Seattle reports that of the 235 MSM patients who presented with gastroenteritis, 57.5% had a positive stool test with 88.7% of the positive stool tests detecting three bacterial pathogens *Escherichia coli* 33.1%, *Shigella* 30.5%, and *Campylobacter* 17.2% of positive samples.⁴ Another study reported MSM has higher likelihood of campylobacter infections than controls, consistent with sexual contact being a risk factor.⁵ Current IDSA guidelines recommend empiric antibiotics in immunocompetent adults who show signs of bacillary dysentery primarily to treat presumptive shigellosis with macrolides (azithromycin 500mg daily for three days) and fluoroquinolones (levofloxacin 750mg daily or ciprofloxacin 750mg orally twice daily, each for three days until signs and symptoms of disease have improved.³ Individuals with complications or immunosuppression may warrant a longer course (7 to 14 days). Although both antibiotics are highly effective against susceptible isolates, rates of fluoroquinolones resistance are increasing worldwide and generally exceed those for azithromycin.⁶ Specifically, MSMs infected with campylobacter coli, showed high rates of multidrug resistance strains to fluoroquinolones and macrolides circulating within the MSM community. Optimal management of infections caused by macrolide and fluoroquinolone-resistant strains is not established. Fosfomycin has been suggested as a possible alternative therapeutic agent.⁷

Conclusion

Most cases of acute infectious diarrhea are viral. However, in one community study of severe diarrhea ≥ 4 fluid stools per day for more than three days, bacterial causes were responsible for

87% of the cases.⁸ *Campylobacter* species are amongst the most common enteric bacterial pathogens worldwide and most cases are self-limiting and mild. However, since multidrug resistance is common and campylobacteriosis can have associated extra-intestinal manifestations, close attention to patient follow-up is important.

Receptive anal or oral intercourse increases risk of direct inoculation or oral-fecal transmission of bacterial and parasitic pathogens such as *Shigella*, *Giardia* and *Entamoeba histolytica*. We present a case of sexually transmitted campylobacter enteritis. Since men who are sexually active with men (MSM) are at an increased risk of sexually transmitted enteric infections,^{7,9} healthcare providers should obtain thorough, comprehensive and culturally humble sexual histories with their patients to identify other risk factors when evaluating infectious causes of acute diarrhea.

REFERENCES

1. **Batz MB, Hoffmann S, Morris JG Jr.** Ranking the disease burden of 14 pathogens in food sources in the United States using attribution data from outbreak investigations and expert elicitation. *J Food Prot.* 2012 Jul;75(7):1278-91. doi: 10.4315/0362-028X.JFP-11-418. Erratum in: *J Food Prot.* 2012 Aug;75(8):1366. PMID: 22980012.
2. **Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, Man SM.** Global Epidemiology of *Campylobacter* Infection. *Clin Microbiol Rev.* 2015 Jul;28(3):687-720. doi: 10.1128/CMR.00006-15. PMID: 26062576; PMCID: PMC4462680.
3. **Shane AL, Mody RK, Crump JA, Tarr PI, Steiner TS, Kotloff K, Langley JM, Wanke C, Warren CA, Cheng AC, Cantey J, Pickering LK.** 2017 Infectious Diseases Society of America Clinical Practice Guidelines for the Diagnosis and Management of Infectious Diarrhea. *Clin Infect Dis.* 2017 Nov 29;65(12):e45-e80. doi: 10.1093/cid/cix669. PMID: 29053792; PMCID: PMC5850553.
4. **Newman KL, Newman GS, Cybulski RJ, Fang FC.** Gastroenteritis in Men Who Have Sex With Men in Seattle, Washington, 2017-2018. *Clin Infect Dis.* 2020 Jun 24;71(1):109-115. doi: 10.1093/cid/ciz783. PMID: 31621824; PMCID: PMC7456347.
5. **Kuhn KG, Hvass AK, Christiansen AH, Ethelberg S, Cowan SA.** Sexual Contact as Risk Factor for *Campylobacter* Infection, Denmark. *Emerg Infect Dis.* 2021 Apr;27(4):1133-1140. doi: 10.3201/eid2704.202337. PMID: 33754996; PMCID: PMC8007285.
6. **Kuschner RA, Trofa AF, Thomas RJ, Hoge CW, Pitarangsi C, Amato S, Olafson RP, Echeverria P, Sadoff JC, Taylor DN.** Use of azithromycin for the treatment of *Campylobacter* enteritis in travelers to Thailand, an area where ciprofloxacin resistance is prevalent. *Clin Infect Dis.* 1995 Sep;21(3):536-41. doi: 10.1093/clinids/21.3.536. PMID: 8527539.
7. **Centers for Disease Control and Prevention.** CDC Yellow Book 2020: Health Information for International Travel. New York: *Oxford University Press*; 2017
8. **Greninger AL, Addetia A, Starr K, Cybulski RJ, Stewart MK, Salipante SJ, Bryan AB, Cookson B, Gaudreau C, Bekal S, Fang FC.** International Spread of Multidrug-Resistant *Campylobacter coli* in Men Who Have Sex With Men in Washington State and Québec, 2015-2018. *Clin Infect Dis.* 2020 Nov 5;71(8):1896-1904. doi: 10.1093/cid/ciz1060. PMID: 31665255; PMCID: PMC7643735.
9. **Dryden MS, Gabb RJ, Wright SK.** Empirical treatment of severe acute community-acquired gastroenteritis with ciprofloxacin. *Clin Infect Dis.* 1996 Jun;22(6):1019-25. doi: 10.1093/clinids/22.6.1019. PMID: 8783703.