

UC Irvine

UC Irvine Previously Published Works

Title

Why Are Children in Urban Neighborhoods at Increased Risk for Psychotic Symptoms? Findings From a UK Longitudinal Cohort Study

Permalink

<https://escholarship.org/uc/item/5qv9c2j7>

Journal

Schizophrenia Bulletin, 42(6)

ISSN

0586-7614

Authors

Newbury, Joanne
Arseneault, Louise
Caspi, Avshalom
et al.

Publication Date

2016-11-01

DOI

10.1093/schbul/sbw052

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Why Are Children in Urban Neighborhoods at Increased Risk for Psychotic Symptoms? Findings From a UK Longitudinal Cohort Study

Joanne Newbury¹, Louise Arseneault¹, Avshalom Caspi^{1,2}, Terrie E. Moffitt^{1,2}, Candice L. Odgers^{3,4}, and Helen L. Fisher^{*,1,4}

¹MRC Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK; ²Departments of Psychology and Neuroscience, Psychiatry and Behavioral Sciences, and Centre for Genomic and Computational Biology, Duke University, Durham, NC; ³Center for Child and Family Policy and the Sanford School of Public Policy, Duke University, Durham, NC

⁴These authors are joint senior authors.

*To whom correspondence should be addressed; MRC Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, 16 De Crespigny Park, London SE5 8AF, UK; tel: +44-(0)207-848-5430, fax: +44-(0)207-848-0866, e-mail: helen.2.fisher@kcl.ac.uk

Background: Urban upbringing is associated with a 2-fold adulthood psychosis risk, and this association replicates for childhood psychotic symptoms. No study has investigated whether specific features of urban neighborhoods increase children's risk for psychotic symptoms, despite these early psychotic phenomena elevating risk for schizophrenia and other psychiatric disorders in adulthood. **Methods:** Analyses were conducted on over 2000 children from the Environmental Risk (E-Risk) Longitudinal Twin Study, a nationally-representative cohort of UK-born twins. Neighborhood-level characteristics were assessed for each family via: a geodemographic discriminator indexing neighborhood-level deprivation, postal surveys of over 5000 residents living alongside the children, and in-home interviews with the children's mothers. Children were interviewed about psychotic symptoms at age 12. Analyses were adjusted for important family-level confounders including socioeconomic status (SES), psychiatric history, and maternal psychosis. **Results:** Urban residency at age-5 (OR = 1.80, 95% CI = 1.16–2.77) and age-12 (OR = 1.76, 95% CI = 1.15–2.69) were both significantly associated with childhood psychotic symptoms, but not with age-12 anxiety, depression, or antisocial behavior. The association was not attributable to family SES, family psychiatric history, or maternal psychosis, each implicated in childhood mental health. Low social cohesion, together with crime victimization in the neighborhood explained nearly a quarter of the association between urbanicity and childhood psychotic symptoms after considering family-level confounders. **Conclusions:** Low social cohesion and crime victimization in the neighborhood

partly explain why children in cities have an elevated risk of developing psychotic symptoms. Greater understanding of the mechanisms leading from neighborhood-level exposures to psychotic symptoms could help target interventions for emerging childhood psychotic symptoms.

Key words: childhood psychotic symptoms/neighborhood characteristics/social cohesion/psychosis/urbanicity

Introduction

Urban vs rural upbringing doubles a child's odds of developing schizophrenia in adulthood.¹ The association between urbanicity and psychosis has been frequently replicated,^{2–10} shows a degree of specificity to non-affective psychoses,^{4,7,11,12} and is not explained by a range of potential confounding factors^{2,13–15} including migration of individuals with schizophrenia into cities.¹⁶ These converging lines of evidence suggest that the association between urbanicity and psychosis has genuine aetiological underpinnings,^{16–19} though the mechanisms driving the association are currently unknown. Urbanicity is therefore a key area for psychosis research, considering that over two-thirds of the world's population are predicted to live in cities by 2050.^{20,21}

The vast majority of urbanicity-psychosis research has focused on adult psychosis. Yet urban residency from birth to adolescence, rather than during adulthood, appears to be more strongly associated with adult

psychosis.^{6,10,13} Consistent with the neurodevelopmental model of schizophrenia, this suggests that the processes leading from urban exposure to psychosis begin in adolescence, childhood, or earlier. Notably, positive psychotic symptoms, such as hallucinations and delusions, are surprisingly prevalent among children in the general population.^{22–26} These early psychotic phenomena share familial and environmental risk factors with psychotic disorders,^{27–29} and whilst they are usually transitory,^{22,23,30} children who experience psychotic symptoms have a significantly elevated risk for schizophrenia and other psychoses in adulthood.^{31,32} Additionally, childhood psychotic symptoms have broad psychiatric relevance as they significantly heighten risk for other subsequent mental health difficulties including substance abuse,²⁶ depression,²⁶ PTSD,³² and suicidal behavior.^{32,33} Childhood psychotic symptoms are therefore a useful marker of early-life risk indicators for psychosis and general psychopathology. Childhood psychotic symptoms could also shed light on the urbanicity-psychosis association: a handful of studies have shown that these symptoms occur more frequently²⁹ and are more likely to persist into adulthood among youth living in urban vs nonurban settings.^{34,35} However, no studies have tested whether specific aspects of the urban environment increase risk for psychotic symptoms among children.

Indeed, urbanicity is only a proxy for the currently unknown operative risk factor(s) for psychosis.^{18,36} More recently, attention has turned to potential urban characteristics³⁷ operating at the neighborhood-level. Neighborhood-level deprivation has been frequently implicated in adult psychosis.^{17,38–43} However, modern urban neighborhoods are very mixed in terms of poverty and affluence,⁴⁴ whilst adult psychosis risk increases incrementally through increasing levels of urbanicity.^{2,3,6,13} Furthermore, the association between urbanicity and psychosis appears stronger in more recent generations,^{9,45} despite urban populations becoming generally wealthier. Thus, the association is difficult to explain through neighborhood-level deprivation alone. Cumulative evidence also supports the importance of neighborhood-level social processes such as crime,^{40,46} disorganization^{46,47} and social fragmentation^{37,39,41,48} in adult psychosis (thoroughly reviewed by March et al³⁶), which are purported to increase adult psychosis risk by heightening childhood exposure to social stressors.^{18,19,49,50} Intriguingly, prodromal status among young adults has been shown to follow spatial patterning in accordance with these kinds of neighborhood-level psychosocial characteristics.⁵¹ However, the longitudinal associations between neighborhood-level social processes and childhood psychotic symptoms are currently unknown. Ultimately, such research could help target social and clinical interventions for early psychotic symptoms.

Here we draw from sociological theory and evidence illustrating that neighborhood-level social processes

mediate the effect of neighborhood structural features (eg, urbanicity) on a range of health outcomes.^{52–54} Guided by this theory and adult psychosis findings, the current study focuses on 4 neighborhood-level social processes: (1) social cohesion, describing the cohesiveness and supportiveness of relationships between neighbors⁵²; (2) social control, describing the likelihood that neighbors would intervene in problems in the neighborhood⁵²; (3) neighborhood disorder, describing physical and social evidence of disorder/threat within the neighborhood⁵³; and (4) crime victimization, representing more direct experiences of victimization in the neighborhood (eg, mugging). The current study investigates the pathways leading from urbanicity to childhood psychotic symptoms, whilst differentiating the effects of specific neighborhood-level social processes from family-level effects. We utilized a cohort of 2232 nationally-representative British twin children who have been followed from birth to age 12 and interviewed for psychotic symptoms at age 12. Our longitudinal neighborhood-level measures were obtained from multiple sources, and neighborhood scores were allocated with fine geographic resolution (ie, postcode-level). With these measures, we asked: (1) Are children in urban vs nonurban neighborhoods at increased risk for psychotic symptoms? (2) Is this association specific to childhood psychotic symptoms? (3) Is the association between urbanicity and childhood psychotic symptoms explained by background characteristics of families living in cities? (4) Are urban neighborhoods more likely to lack social cohesion and social control and be characterized by disorder and crime? (5) Finally, does the level of social cohesion, social control, neighborhood disorder, and crime victimization operating within neighborhoods mediate the effect of urban residency on childhood psychotic symptoms? We hypothesized that the effect of urbanicity on childhood psychotic symptoms would be specific to this phenotype, and mediated via exposure to low social cohesion and social control, and high disorder and crime victimization in the neighborhood (proposed pathways shown in [figure 1](#)).

Methods

Study Cohort

Participants were members of the Environmental Risk (E-Risk) Longitudinal Twin Study, which tracks the development of a nationally-representative birth cohort of 2232 British twin children. The sample was drawn from a larger cohort of twins born in England and Wales in 1994–1995.⁵⁵ Full details about the sample are reported elsewhere.⁵⁶ Briefly, the E-Risk sample was constructed in 1999–2000, when 1116 families with same-sex 5-year-old twins (93% of those eligible) participated in home-visit assessments. Families were recruited to represent the UK population of families with newborns in the 1990s, based on residential location throughout England and Wales and mothers' age




Fig. 1. Conceptualized pathways between urbanicity and childhood psychotic symptoms, with the total effects transmitting both directly (solid line), and indirectly (dashed lines) via neighborhood-level social process mediators (low social cohesion, low social control, high neighborhood disorder, and high crime victimization).

(teenaged mothers with twins were over-selected to replace high-risk families who were selectively lost to the register through non-response. Older mothers having twins via assisted reproduction were under-selected to avoid an excess of well-educated older mothers). E-Risk families are representative of UK households across the spectrum of neighborhood-level deprivation: 25.6% of E-Risk families live in “wealthy achiever” neighborhoods compared to 25.3% of households nation-wide; 5.3% vs 11.6% live in “urban prosperity” neighborhoods; 29.6% vs 26.9% live in “comfortably off” neighborhoods; 13.4% vs 13.9% live in “moderate means” neighborhoods; and 26.1% vs 20.7% live in “hard-pressed” neighborhoods.^{57,58} E-Risk families under-represent “urban prosperity” neighborhoods because such households are likely to be childless. Sex was evenly distributed in the resulting sample (49% male). All families were English speaking, and the majority (93.7%) were White. Follow-up home-visits were conducted when children were aged 7, 10, and 12 (participation rates were 98%, 96%, and 96%, respectively). At age 12, the E-Risk sample comprised 2146 twin children, and the majority of these children had complete data on both psychotic symptoms and urbanicity at age 12 (95.7%; $N = 2054$). Over half of children (56.7%, $N = 1180$) never moved house at all between ages 5 and 12, and of those who did nearly two-thirds (65.0%) moved less than 500 meters. The Joint South London and Maudsley and the Institute of Psychiatry Research Ethics Committee approved each phase of the study. Parents gave informed consent and children gave assent.

Measures

Childhood Psychotic Symptoms. E-Risk families were visited by mental health trainees or professionals when

children were aged 12.²⁹ Each child was privately interviewed about 7 psychotic symptoms pertaining to delusions and hallucinations, with items including “have other people ever read your thoughts?,” “have you ever thought you were being followed or spied on?,” and “have you ever heard voices that other people cannot hear?.” This interview has been described in detail previously.²⁹ The item choice was guided by the Dunedin Study’s age-11 interview protocol³¹ and an instrument prepared for the Avon Longitudinal Study of Parents and Children.⁵⁹ Interviewers coded each experience 0, 1, 2 indicating respectively “not a symptom,” “probable symptom,” and “definite symptom.” A conservative approach was taken in designating a child’s report as a symptom. First, the interviewer probed using standard prompts designed to discriminate between experiences that were plausible (eg, “I was followed by a man after school”) and potential symptoms (eg, “I was followed by an angel who guards my spirit”), and wrote down the child’s narrative description of the experience. Second, items and interviewer notes were assessed by a psychiatrist expert in schizophrenia, a psychologist expert in interviewing children, and a child and adolescent psychiatrist to verify the validity of the symptoms. Third, because children were twins, experiences limited to the twin relationship (eg, “My twin and I often know what each other are thinking”) were coded as “not a symptom”. Children were only designated as experiencing psychotic symptoms if they reported at least one definite symptom. At age 12, 5.9% ($N = 125$) of children reported experiencing psychotic symptoms. This is similar to the prevalence of psychotic symptoms in other community samples of children and adolescents.^{22–26} Furthermore, we previously showed that childhood psychotic symptoms in this cohort have good

construct validity, sharing many of the genetic, social, neurodevelopmental, and behavioral risk factors and correlates as adult schizophrenia.²⁹ Additionally, as we focused on psychotic symptoms rather than diagnoses, the present study design avoids confounding by psychiatric service utilization.

Urbanicity. Urban/nonurban classification of E-Risk families' neighborhoods was based on responses from a postal survey sent to residents living alongside E-Risk families when children were aged 12.^{60,61} Questionnaires were sent to every household in the same postcode as the E-Risk families, excluding the E-Risk families themselves (addresses were identified from electoral roll records). The number of surveys sent ranged from 15 to 50 residences per neighborhood (Average = 18.96, SE = 0.21). Excluding undelivered surveys ($N = 600$), the overall response rate was 28.1% (5601/19 926). Survey respondents typically lived on the same street or within the same apartment block as the children in our study. Surveys were returned by an average of 5.18 (SD = 2.73) respondents per neighborhood (range = 0–18 respondents), and there were at least 2 responses from 95% of the neighborhoods ($N = 5601$ respondents).⁶¹ Residents reported whether their neighborhood was in “a city,” “a town,” “a suburb,” “a small village,” or “the countryside.”²⁹ There was high agreement between residents in the same neighborhood, with only 50 neighborhoods returning discordant responses (ie, neighborhoods where residents differed in their urbanicity responses). These 50 ambiguous neighborhoods were clarified by a British researcher (blind to any phenotypic/identifying data) using the children's full postcode, Google Aerial view and the Office of National Statistics' population density map (http://www.neighbourhood.statistics.gov.uk/HTMLDocs/PopulationDensity_2010.html, last accessed April 28, 2016), based on a combination of features including population density, building density, proximity to the countryside or city/town centre, land-use (eg, agriculture, transportation, industry, etc.), and the official definition of the settlement. This same method was used to estimate urbanicity at age-5 for the 35% of children who had moved over 500 metres between ages 5 and 12. For ease of interpretation and to increase analytic power, urbanicity is herein dichotomized as urban (1: city/town) vs nonurban (0: suburb/small village/countryside). At age 12, the sample was split evenly between urban and nonurban neighborhoods, with 51.9% ($N = 1066$) of children living in urban neighborhoods and the remaining 988 children living in nonurban neighborhoods. Similarly, 55.1% ($N = 1117$) lived in urban neighborhoods at age 5.

Neighborhood-Level Deprivation. Neighborhood-level deprivation was constructed using A Classification of Residential Neighbourhoods (ACORN), a geodemographic discriminator developed by CACI Information

Services (<http://www.caci.co.uk/>, last accessed April 28, 2016).⁵⁷ Detailed information about ACORN's classification of neighborhood-level socioeconomic-status (SES) has been provided previously.^{58,60,62} Briefly, CACI utilized over 400 variables from 2001 census data for Great Britain (eg, educational qualifications, unemployment, housing tenure) and CACI's consumer lifestyle database. Following hierarchical-cluster-analysis, 5 distinct and homogeneous ordinal groups were created ranging from “Wealthy Achiever” (coded 1) to “Hard Pressed” (coded 5) neighborhoods. Each family in our sample was matched to the ACORN code for its neighborhood via its postcode (age 5 or age 12 postcode, where relevant).⁵⁸

Neighborhood-Level Social Processes. Social processes included social cohesion, social control, neighborhood disorder and crime victimization, and were measured in both early and late childhood. Social processes were first measured at age 5 via in-home interviews with the children's mothers.⁶³ Social cohesion⁵² (5 items) was assessed by asking mothers whether their neighborhood was close-knit, whether neighbors shared values, and whether neighbors trusted and got along with each other, etc. Higher scores indicate greater social cohesion. Social control⁵² (5 items) was assessed by asking mothers to judge whether people in their neighborhoods would take action against different types of undesirable activities (eg, children skipping school, fights in public places). Higher scores indicate greater social control. For neighborhood disorder,⁵³ mothers were asked whether 13 problems affected their neighborhood, including noisy neighbors, arguments or loud parties, vandalism, graffiti or deliberate damage to property, and cars broken into. Higher scores indicate greater neighborhood disorder. Crime victimization was assessed by asking mothers whether they or their family had been victimized by violent crime (eg, mugging, assault), a burglary, or a theft in the neighborhood. Higher scores indicate greater crime victimization. Items (each coded 0–2) within each social process scale were summed for each mother. Social processes were also measured when children were aged 12 via the resident surveys^{60,61} (survey methodology described in detail under urbanicity heading). Residents were asked the same questions regarding these 4 neighborhood-level social processes. For the resident reports, the social process scales were created in 2 steps. First, items belonging to each social process scale were averaged to create summary scores for each of the 5601 respondents. Second, scores for each E-Risk family were created by averaging the social process scores of respondents within that neighborhood.

Thus, neighborhood-level social processes were estimated both before and contemporaneously to childhood psychotic symptoms, enabling us to triangulate a prospective design with objective neighborhood appraisals. At age 5, mothers' views of the neighborhood were used as mothers are considered more reliable reporters

than children at this age and because their perceptions are likely to influence their children's amount of exposure and experiences in the neighborhood.^{64,65} At age 12, resident reports were used to gain more objective and comprehensive assessments of the neighborhood. As children themselves reported on their own psychotic symptoms at age 12, both our age-5 (mother-reported) and age-12 (resident-reported) assessments of neighborhood-level social processes are obtained from independent sources.

Other Age-12 Outcomes. Anxiety was assessed when children were aged 12, via private interviews using the 10-item version of the Multidimensional Anxiety Scale for Children (MASC).⁶⁶ An extreme anxiety group was formed with children who scored at or above the 95th percentile ($N = 129$, 6.1%). Depression symptoms were assessed at age 12 using the Children's Depression Inventory (CDI).⁶⁷ Children who scored 20 or more⁶⁸ were deemed to have clinically significant depressive symptoms ($N = 74$, 3.5%). Antisocial behavior was assessed using the Achenbach system of empirically-based assessment.⁶⁹ An extreme antisocial behavior group was formed with children who scored at or above the 95th percentile ($N = 110$, 5.1%), based on combined mother and teacher reports at age 12.⁷⁰

Family-Level Confounders. Family SES was measured via a composite of parental income (total household), education (highest mother/father), and occupation (highest mother/father) when children were aged 5, and was categorized into tertiles (ie, low-, medium-, and high-SES). Family psychiatric history and maternal psychosis were both assessed when children were aged 12. In private interviews, mothers reported on family history of DSM disorders,⁷¹ which was converted to a proportion (0–1.0) of family members with a history of psychiatric disorder. For maternal psychosis, mothers were interviewed using the Diagnostic Interview Schedule⁷² for DSM-IV⁷³ which provides a symptom count for characteristic symptoms of schizophrenia (eg, hallucinations, delusions, anhedonia).

Statistical Analysis. Analyses were conducted in STATA 11.2 (Stata-Corp). Firstly, linear regression was used to investigate the association between urbanicity and neighborhood-level social processes (table 1). Secondly, logistic regression was used to investigate the associations between neighborhood-level social processes and childhood psychotic symptoms (table 2). Thirdly, our mediation analyses utilized KHB pathway decomposition (table 3).⁷⁴ This procedure partitions the total effect of one variable (urbanicity) on another variable (childhood psychotic symptoms) into the direct effect (which also includes the effects of unknown/unspecified mediators and measurement error), and indirect effects explained by specified mediators (neighborhood-level social processes). Age-5 urbanicity is used when age-5

social processes are analyzed; age-12 urbanicity is used when age-12 social processes are analyzed. As the scales differed between the age-5 (mother-reported) and age-12 (resident-reported) social process variables, social process variables in steps 2 and 3 were standardized with a mean of 0 and a SD of 1 (subtraction of the mean then division by the SD) to facilitate comparability of the results. Where appropriate, analyses accounted for the non-independence of observations using the "CLUSTER" command because the sample comprised twins. This procedure is derived from the Huber-White variance estimator, and provides robust standard errors adjusted for within-cluster correlated data⁷⁵ (Note: within-pair twin correlations can also be corrected using multi-level approaches. Supplementary table 1 shows that our main logistic regression analyses are highly robust to alternative estimation procedures.).

Results

Are Children in Urban vs Nonurban Neighborhoods at Increased Risk for Psychotic Symptoms?

There was a significant cross-sectional association between age-12 urban residency and childhood psychotic symptoms (OR = 1.76, 95% CI = 1.15–2.69, $P = .009$). Around 7.4% ($N = 79$) of urban-dwelling children compared to 4.4% ($N = 43$) of nonurban-dwelling children experienced at least one definite psychotic symptom at age 12. The association between urbanicity and psychotic symptoms held when analyses were restricted to the 56.6% of children who never moved house between ages 5 and 12 (OR = 2.01, 95% CI = 1.14–3.58, $P = .017$), and when controlling for residential mobility during this period (OR = 1.71, 95% CI = 1.12–2.61, $P = .014$). The association also held for the 93.7% of children who were ethnically White (OR = 1.85, 95% CI = 1.21–2.84, $P = .005$). Although in our sample there was a tendency for urban neighborhoods to be more deprived (OR = 2.57, 95% CI = 1.99–3.32, $P < .001$), half of urban neighborhoods were relatively affluent (ACORN categories 1–3; 50.3%), and over a quarter of nonurban neighborhoods were considered deprived (ACORN categories 4 and 5; 27.8%). Moreover, when urbanicity and neighborhood deprivation were included in a logistic regression model together, they were both significantly associated with childhood psychotic symptoms (OR = 1.62, 95% CI = 1.03–2.56, $P = .039$; OR = 1.62, 95% CI = 1.05–2.50, $P = .029$, respectively), demonstrating that urbanicity is associated with childhood psychotic symptoms largely independently of neighborhood-level deprivation in this sample. Additionally, the association between urbanicity and childhood psychotic symptoms held when earlier urbanicity at age 5 was examined (OR = 1.80, 95% CI = 1.16–2.77, $P = .008$). Therefore, the remaining analyses in this article will focus on tracing the effects of urbanicity (age-5 or age-12, where appropriate) on childhood psychotic symptoms.

Table 1. Bivariate Associations Between Urbanicity and Neighborhood-Level Social Processes

Neighborhood-Level Social Processes	Range	Urban	Nonurban	Standardized Association Between Urbanicity and Social Processes	
		<i>M</i> (SD)	<i>M</i> (SD)	<i>B</i> ^a	<i>P</i> Value
Age-5 (mother reports) ^b					
Social cohesion	0–10	7.11 (2.95)	8.18 (2.32)	–.19	<.001
Social control	0–10	7.04 (2.88)	7.91 (2.41)	–.16	<.001
Neighborhood disorder	0–22	4.40 (4.15)	3.46 (3.24)	.12	<.001
Crime victimization	0–6	1.06 (1.39)	0.75 (1.18)	.12	<.001
Age-12 (resident reports) ^c					
Social cohesion	0–4	2.11 (0.50)	2.36 (0.47)	–.25	<.001
Social control	0–4	2.09 (0.53)	2.33 (0.51)	–.22	<.001
Neighborhood disorder	0–2	0.56 (0.35)	0.40 (0.32)	.23	<.001
Crime victimization	0–2	0.22 (0.24)	0.15 (0.19)	.16	<.001

Note: *B*, standardized beta coefficient; *M*, mean. Social cohesion and social control consistently have negative beta coefficients, demonstrating that urban neighborhoods had lower levels of social cohesion and social control compared to nonurban neighborhoods. In contrast, neighborhood disorder and crime victimization consistently have positive beta coefficients, demonstrating that urban neighborhoods had higher levels of disorder and crime victimization compared to nonurban neighborhoods. All analyses account for the nonindependence of twin observations.

^aThe standardized (*B*) beta coefficients indicate the unit SD change in each social process given 1 unit SD change in urbanicity, and allow comparison between each social process. Standardized betas provide exactly the same point estimates as correlation coefficients and may be interpreted as correlations, with a score of –1.0 indicating a 100% negative correlation and a score of +1.0 indicating a 100% positive correlation.

^bAge-5 urbanicity is used for the bivariate associations between urbanicity and age-5 mother-reported social processes.

^cAge-12 urbanicity is used for the bivariate associations between urbanicity and age-12 resident-reported social processes. Age-12 resident-reported social process scores were imputed for 2 children with missing data.

Table 2. Bivariate Associations Between Neighborhood-Level Social Processes and Childhood Psychotic Symptoms

Neighborhood-Level Social Processes	OR	95% CI	<i>P</i> Value
Age-5 (mother reports)			
Social cohesion	0.68	[0.58, 0.82]	<.001
Social control	0.75	[0.62, 0.91]	.003
Neighborhood disorder	1.26	[1.06, 1.51]	.010
Crime victimization	1.40	[1.19, 1.65]	<.001
Age-12 (resident reports) ^a			
Social cohesion	0.76	[0.65, 0.89]	.001
Social control	0.83	[0.69, 1.00]	.050
Neighborhood disorder	1.27	[1.07, 1.52]	.007
Crime victimization	1.17	[0.96, 1.42]	.123

Note: Social cohesion and social control are consistently associated with odds lower than 1 for childhood psychotic symptoms, demonstrating that children were less likely to experience psychotic symptoms in neighborhoods with higher levels of social cohesion and social control. In contrast, neighborhood disorder and crime victimization are consistently associated with odds greater than 1 for childhood psychotic symptoms, demonstrating that children were more likely to experience psychotic symptoms in neighborhoods with higher levels of neighborhood disorder and crime victimization. All analyses account for the nonindependence of twin observations. All social process variables have been standardized with a mean of 0 and a SD of 1.

^aAge-12 resident-reported social process scores were imputed for 2 children with missing data.

Is Urbanicity Specifically Associated With Childhood Psychotic Symptoms?

Our assumption of specificity to psychotic symptoms was tentatively supported, as associations between age-12 urbanicity and age-12 depression (OR = 1.16, 95% CI = 0.69–1.96, *P* = .571), anxiety (OR = 1.42, 95% CI = 0.95–2.12, *P* = .091) and antisocial behavior (OR = 0.93, 95% CI = 0.59–1.47, *P* = .753) were each nonsignificant, with smaller effect sizes than demonstrated for psychotic symptoms. However, given that the CIs for both depression and anxiety included the point estimate for the association between age-12 urbanicity and childhood psychotic symptoms (OR = 1.76), we cannot be sure that these associations differed significantly. Nevertheless, after simultaneous adjustment for age-12 depression, anxiety and antisocial behavior, urbanicity remained significantly associated with childhood psychotic symptoms (OR = 1.74, 95% CI = 1.15–2.65, *P* = .009), suggesting that urbanicity was independently associated with childhood psychotic symptoms in this sample. Furthermore, the associations between urbanicity and these 3 additional age-12 outcomes remained nonsignificant when they were recategorized at a lower threshold (80th percentile), suggesting that the negative findings were not due to inadequate power (results available upon request).

Table 3. Association Between Urbanicity and Childhood Psychotic Symptoms, Split Into Total Effects, and Direct and Indirect Pathways via Neighborhood-Level Social Process Mediators

Potential Neighborhood-Level Social Process Mediators	Mediation Model 1					Mediation Model 2 ^a				
	Sample Size	Total OR [95% CI]	Direct OR [95% CI]	Indirect OR [95% CI]	% Mediated ^b	Sample Size	Total OR [95% CI]	Direct OR [95% CI]	Indirect OR [95% CI]	% Mediated ^b
Age-5 (mother reports) ^c Social cohesion	2014	1.71* [1.10, 2.64]	1.49† [0.95, 2.35]	1.15** [1.05, 1.25]	25	2005	1.58* [1.00, 2.49]	1.46 [0.92, 2.33]	1.08* [1.02, 1.16]	17
Social control	1998	1.74* [1.13, 2.70]	1.62* [1.04, 2.54]	1.07* [1.00, 1.15]	13	1989	1.62* [1.03, 2.55]	1.57† [0.99, 2.48]	1.04 [0.99, 1.09]	8
Neighborhood disorder	2022	1.72* [1.11, 2.66]	1.65* [1.06, 2.57]	1.04 [0.99, 1.09]	7	2013	1.58* [1.01, 2.49]	1.57† [0.99, 2.47]	1.01 [0.98, 1.04]	2
Crime victimization	2022	1.71* [1.11, 2.65]	1.59* [1.02, 2.48]	1.07* [1.02, 1.13]	13	2013	1.57† [1.00, 2.47]	1.50† [0.95, 2.37]	1.05 [1.00, 1.10]	11
Age-12 (resident reports) ^d Social cohesion	2054	1.76** [1.15, 2.69]	1.58* [1.03, 2.43]	1.11* [1.02, 1.21]	19	2045	1.61* [1.03, 2.50]	1.53† [0.98, 2.40]	1.05 [0.97, 1.13]	10
Social control	2054	1.76** [1.15, 2.69]	1.66* [1.07, 2.57]	1.06 [0.97, 1.16]	11	2045	1.61* [1.03, 2.51]	1.58* [1.00, 2.50]	1.02 [0.94, 1.10]	4
Neighborhood disorder	2054	1.76** [1.15, 2.69]	1.60* [1.01, 2.53]	1.10† [1.00, 1.21]	16	2045	1.61* [1.03, 2.51]	1.55† [0.97, 2.49]	1.04 [0.96, 1.12]	7

Note: The sample sizes and total effect ORs vary slightly for the age-5 mother reports of social processes, due to small numbers of children missing data on neighborhood-level social processes and/or family-level covariates. These sample size differences also account for the difference between the main effect OR (1.80) and the total effect ORs for mother reports in Mediation model 1. Social processes can still be compared for the percentage that they mediate the total effect of urbanicity. We also conducted mediation analyses using Full Information Maximum Likelihood (FIML) in Mplus to include all available cases at age 5 ($N = 2232$ for all Models reported) and found no differences in the size, direction or pattern of effects. Total effect = overall association between urbanicity and childhood psychotic symptoms; direct effect = the part of the overall association that is not explained by the mediator/covariates in the model; and indirect effect = the part of the overall association that is explained by the social process mediator in the model. Bold text denotes significant indirect (mediation) pathways at $P < .05$. All analyses account for the nonindependence of twin observations. All social process variables have been standardized with a mean of 0 and a SD of 1.

^aThe total, direct and indirect ORs in mediation model 2 are adjusted for family-level confounders: family socioeconomic status, family psychiatric history, and maternal psychosis.

^bPercentages rounded to whole numbers.

^cAge-5 urbanicity is used for the mediation analysis of urbanicity and childhood psychotic symptoms via age-5 mother-reported social processes.

^dAge-12 urbanicity is used for the mediation analysis of urbanicity and childhood psychotic symptoms via age-12 resident-reported social processes. Age-12 resident-reported social process scores were imputed for 2 children with missing data.

* $P < .05$, ** $P < .01$, †Nominally significant $P > .05$ and $P < .1$.

Is the Association Between Urbanicity and Childhood Psychotic Symptoms Explained by Background Characteristics of Families Living in Cities?

The association between age-12 urbanicity and childhood psychotic symptoms did not appear to be explained by 3 key potential family-level confounders, namely family SES, family psychiatric history and maternal psychosis. Simultaneous adjustment for these proxy indicators of genetic and environmental risk only slightly attenuated the association between age-12 urbanicity and childhood psychotic symptoms (OR = 1.61, 95% CI = 1.04–2.51, $P = .035$).

Are Urban Neighborhoods More Likely to Lack Social Cohesion and Social Control and Be Characterized by Disorder and Crime?

Associations of urbanicity with neighborhood-level social processes are shown in [table 1](#). At age 5, urban neighborhoods had (ie, mothers reported) significantly lower social cohesion and social control, and significantly higher neighborhood disorder and crime victimization than nonurban neighborhoods (all P s < .001). Similar bivariate associations were found between urbanicity and social processes for age-12 neighborhoods (residents' reports) (all P s < .001; [table 1](#)).

Associations of neighborhood-level social processes with childhood psychotic symptoms are shown in [table 2](#). Children were significantly less likely to experience psychotic symptoms at age 12 if, at age 5, they lived in neighborhoods with higher social cohesion ($P < .001$) and higher social control ($P = .003$). In contrast, children were significantly more likely to experience psychotic symptoms at age 12 if their age-5 neighborhood was characterized by higher neighborhood disorder ($P = .010$) and higher crime victimization ($P < .001$). A comparable cross-sectional pattern was found for the associations between age-12 neighborhood-level social processes and childhood psychotic symptoms, though social control was borderline statistically significant ($P = .050$) and neighborhood-level crime victimization failed to reach conventional levels of statistical significance ($P = .123$; [table 2](#)).

Do Neighborhood-Level Social Processes Mediate the Effect of Urban Residency on Childhood Psychotic Symptoms?

We investigated the extent that neighborhood-level social processes mediated the effect of urban residency on childhood psychotic symptoms ([figure 1](#)). Social processes were only included if they were significantly associated with both urbanicity and childhood psychotic symptoms (ie, age-12 neighborhood-level crime victimization was excluded as it was not associated with childhood psychotic symptoms at $P < .05$). [Table 3](#) shows results as odds ratios with 95% CIs for the total (overall association), direct

(the part of the overall association that is not explained by the mediator/covariates in the model) and indirect (the part of the overall association that is explained by the social process mediator in the model) effects of urbanicity on childhood psychotic symptoms. A model in which the indirect OR is equal to the total OR would indicate that the effect of the predictor on the outcome is entirely (100%) mediated by the specified mediator. Mediation model 1 is unadjusted, and Mediation model 2 is adjusted for family-level confounders (family SES, family psychiatric history, maternal psychosis) (Note: sample size and total effect ORs vary slightly within [table 3](#) due to small numbers of children missing data on age-5 neighborhood-level social processes and/or family-level confounders. Further detail is provided in [Table 3's](#) footnote.). Mediation model 1 shows that neighborhood-level low social cohesion at age 5 significantly mediated the effect of age-5 urbanicity on age-12 psychotic symptoms, explaining 25% of the association. Low social control and high crime victimization in the neighborhood also significantly mediated the effect of age-5 urbanicity on childhood psychotic symptoms, each explaining 13% of the association. These prospective models were somewhat supported by our cross-sectional analysis of age-12 urbanicity and age-12 social processes, in that low social cohesion once again significantly explained the largest proportion of the association between urbanicity and childhood psychotic symptoms (19%). These mediatory effects were slightly attenuated after considering family-level confounders (Mediation model 2, [table 3](#)). Nonetheless, following adjustment, neighborhood-level low social cohesion and high crime victimization at age 5 still significantly mediated the effect of age-5 urbanicity on childhood psychotic symptoms (explaining 17% and 11%, respectively). When age-5 social cohesion and crime victimization were simultaneously modeled, together they explained nearly a quarter of the effect of age-5 urbanicity on age-12 psychotic symptoms (24%: OR = 1.11, 95% CI = 1.03–1.20, $P = .004$).

Discussion

This is the first study to investigate whether specific psychosocial features of the urban environment increase children's risk for psychotic symptoms. Our findings add to existing knowledge in at least 3 ways. First, children living in urban neighborhoods were ~80% more likely to experience psychotic symptoms at age 12 compared to children living in nonurban neighborhoods. This association held in both prospective and cross-sectional models, and was not explained by the socioeconomic or psychiatric composition of urban families. Second, psychotic symptoms were more common among children living in neighborhoods characterized by low social cohesion, low social control, high neighborhood disorder, and where the family had been directly victimized by a crime. Our findings

highlight that these neighborhood-level social processes, which are implicated in adult psychosis,^{17,37,39,40,46–48} may also be relevant to positive psychotic symptoms in childhood. Third, low social cohesion explained the largest proportion of the effect of urbanicity on childhood psychotic symptoms, regardless of reporter (17% for mother reports, 10% for resident reports), and independently of the potential family-level confounders measured in this study. Furthermore, social cohesion together with crime victimization at age-5 explained almost a quarter of the effect of age-5 urbanicity on childhood psychotic symptoms. Though we have investigated childhood psychotic symptoms as the main outcome measure, our findings regarding social cohesion and crime victimization are consistent with previous studies implicating area-level social fragmentation^{39,41,47} (or related constructs) and crime^{40,46} in adult psychosis.

A significant minority of children experience persistent psychotic symptoms and eventual clinical diagnosis.^{31,32} Furthermore, urban upbringing is highly correlated with urban adult residency.⁶ Taken together, ours and previous findings are consistent with the proposal that early-life exposure to neighborhood-level social stressors contributes to the heightened psychosis rates found in cities.^{17–19,49,50} From a child's perspective, growing up in a crowded neighborhood characterized by insecure/nonexistent social support networks, unfriendly/unpredictable interactions between neighbors, and fear of exposure to crime could promote psychotic symptoms in various mutually compatible ways. Prolonged exposure to neighborhood-level social stressors could dysregulate the hypothalamic–pituitary–adrenal axis,⁷⁶ dopaminergic system,⁵⁰ and/or neurodevelopment,²¹ increasing risk for psychotic symptoms particularly among children with genetic predisposition.⁷⁷ A cognitive mechanism, with specific adverse neighborhood-level experiences exacerbating or providing content to emerging delusions and hallucinations⁷⁸ could also explain why urbanicity was associated with positive psychotic symptoms but not significantly with anxiety or depression.⁷⁹ These neighborhood effects could also be transmitted indirectly by heightening children's exposure to family-level stress or even maltreatment. Indeed, low neighborhood-level social cohesion appears to undermine positive parenting practices.⁸⁰ The individual-level factors and potential mechanisms leading from neighborhood-level adverse exposures to childhood psychotic symptoms now require attention. For example, children raised in urban vs non-urban neighborhoods could differ in their neurocognitive reactivity to social stress, as recently demonstrated among healthy adults.^{81,82}

Limitations

Five limitations deserve mention. First, causal inference is limited as families were not randomly selected

into neighborhoods. Whilst we adjusted for a number of important proxy measures of genetic and environmental risk, various non/reverse-causal explanations remain possible. Future research with larger samples, and ideally quasi-experimental designs, are required to more persuasively rule out social selection as an explanation for these findings. The role of gene-environment correlation (eg, individuals with higher genetic risk for psychosis “drifting” into urban neighborhoods) can also now be estimated via emerging methods such as polygenic risk score analysis. Second, childhood psychotic symptoms are relatively rare, with only ~6% of children reporting symptoms in the E-Risk cohort at age 12. Our findings would benefit from replication. Third, this low base-rate also made it necessary to dichotomize urbanicity to increase power. This potentially simplified our findings, particularly given previous evidence for a dose-response urbanicity-psychosis association through the range of urbanicity. Fourth, although childhood psychotic symptoms are thought to lie on a continuum with schizophrenia,⁸³ they are also associated with other psychiatric disorders in adulthood^{26,32,33} and therefore the current findings may extend beyond schizophrenia to risk for serious adult psychopathology in general. Finally, the E-Risk cohort is a twin sample, and whether findings from twin studies generalize to singletons is sometimes contested. However, the children in our study are representative of singletons for the prevalence of psychotic symptoms,^{22–26} and representative of UK families in terms of geographic and socioeconomic distribution.^{57,58}

Importantly, neighborhood-level social processes did not completely explain the effect of urbanicity in our analyses. Future investigations should consider a wider range of potential social and physical neighborhood-level characteristics when testing for environmental contributions to childhood psychotic symptoms. Neighborhood-level physical exposures such as noise, light, and air pollution, as well as exposure to viral infections warrant research in relation to early psychotic symptoms. The modest mediation could also be partly attributable to measurement error entailed in the neighborhood-level social process measures. Additionally, it is possible that up to age 12 the children in our study were relatively sheltered from certain threats in their neighborhoods. Cumulative neighborhood-level exposures, from childhood, through adolescence and into adulthood, may each contribute in different ways or combine to increase risk for psychotic symptoms. It will therefore be important to investigate the contribution of neighborhood-level social processes to the emergence of psychotic symptoms in late adolescence, when many children will have experienced more direct exposure to adversity in their neighborhood.

Conclusion

In this study, the increased risk for childhood psychotic symptoms in urban neighborhoods was explained, in

part, by lower levels of social cohesion and higher levels of crime victimization operating within these neighborhoods. If these novel findings are replicated, they could support the role of exposure to neighborhood-level social stressors in the aetiology of childhood psychotic symptoms.^{17,18,46} Populations are becoming increasingly urban, and child and adolescent psychopathology represents a growing proportion of the global burden of disease.⁸⁴ The present findings therefore underscore the emerging need to identify the social, psychological, and biological pathways leading from neighborhood-level exposures to childhood psychotic symptoms.

Supplementary Material

Supplementary material is available at <http://schizophreniabulletin.oxfordjournals.org>.

Funding

The E-Risk study is funded by the UK Medical Research Council (G1002190, G9806489). This work was supported by the Economic and Social Research Council (RES-177-25-0013, and a Multidisciplinary Studentship to J.N.); National Institute of Child Health and Development (HD061298); MQ: Transforming Mental Health (MQ14F40); Google Streetview; British Academy (SQ140024); William T. Grant Foundation; and Jacobs Foundation. C.L.O. is a Jacobs Foundation Advanced Research Fellow.

Acknowledgments

We are grateful to the study mothers and fathers, the twins, and the twins' teachers for their participation. Our thanks to members of the E-Risk team for their dedication, hard work, and insights, and to CACI Inc. for use of their consumer lifestyle databases. We also thank Emma Hedman, Jill Collins and Paul Langston for their geocoding assistance. The authors have declared that there are no conflicts of interest in relation to the subject of this study.

References

- Vassos E, Pedersen CB, Murray RM, Collier DA, Lewis CM. Meta-analysis of the association of urbanicity with schizophrenia. *Schizophr Bull.* 2012;38:1118–1123.
- Lewis G, David A, Andréasson S, Allebeck P. Schizophrenia and city life. *Lancet.* 1992;340:137–140.
- van Os J, Hanssen M, Bijl RV, Vollebergh W. Prevalence of psychotic disorder and community level of psychotic symptoms: an urban-rural comparison. *Arch Gen Psychiatry.* 2001;58:663–668.
- Sundquist K, Frank G, Sundquist J. Urbanisation and incidence of psychosis and depression: follow-up study of 4.4 million women and men in Sweden. *Br J Psychiatry.* 2004;184:293–298.
- Allardyce J, Boydell J, Van Os J, et al. Comparison of the incidence of schizophrenia in rural Dumfries and Galloway and urban Camberwell. *Br J Psychiatry.* 2001;179:335–339.
- Pedersen CB, Mortensen PB. Evidence of a dose-response relationship between urbanicity during upbringing and schizophrenia risk. *Arch Gen Psychiatry.* 2001;58:1039–1046.
- Kelly BD, O'Callaghan E, Waddington JL, et al. Schizophrenia and the city: a review of literature and prospective study of psychosis and urbanicity in Ireland. *Schizophr Res.* 2010;116:75–89.
- Lundberg P, Cantor-Graae E, Rukundo G, Ashaba S, Ostergren PO. Urbanicity of place of birth and symptoms of psychosis, depression and anxiety in Uganda. *Br J Psychiatry.* 2009;195:156–162.
- Marcelis M, Navarro-Mateu F, Murray R, Selten JP, Van Os J. Urbanization and psychosis: a study of 1942-1978 birth cohorts in The Netherlands. *Psychol Med.* 1998;28:871–879.
- Marcelis M, Takei N, van Os J. Urbanization and risk for schizophrenia: does the effect operate before or around the time of illness onset? *Psychol Med.* 1999;29:1197–1203.
- Pedersen CB, Mortensen PB. Urbanicity during upbringing and bipolar affective disorders in Denmark. *Bipolar Disord.* 2006;8:242–247.
- Vassos E, Agerbo E, Mors O, Pedersen CB. Urban-rural differences in incidence rates of psychiatric disorders in Denmark [published online ahead of print December 17, 2015]. *Br J Psychiatry.* doi:10.1192/bjp.bp.114.161091.
- Mortensen PB, Pedersen CB, Westergaard T, et al. Effects of family history and place and season of birth on the risk of schizophrenia. *N Engl J Med.* 1999;340:603–608.
- van Os J, Hanssen M, Bak M, Bijl RV, Vollebergh W. Do urbanicity and familial liability coparticipate in causing psychosis? *Am J Psychiatry.* 2003;160:477–482.
- Harrison G, Fouskakis D, Rasmussen F, et al. Association between psychotic disorder and urban place of birth is not mediated by obstetric complications or childhood socio-economic position: a cohort study. *Psychol Med.* 2003;33:723–731.
- Pedersen CB. Persons with schizophrenia migrate towards urban areas due to the development of their disorder or its prodromata. *Schizophr Res.* 2015;168:204–208.
- Kirkbride JB, Jones PB, Ullrich S, Coid JW. Social deprivation, inequality, and the neighborhood-level incidence of psychotic syndromes in East London. *Schizophr Bull.* 2014;40:169–180.
- Krabbendam L, van Os J. Schizophrenia and urbanicity: a major environmental influence—conditional on genetic risk. *Schizophr Bull.* 2005;31:795–799.
- Heinz A, Deserno L, Reininghaus U. Urbanicity, social adversity and psychosis. *World Psychiatry.* 2013;12:187–197.
- Dye C. Health and urban living. *Science.* 2008;319:766–769.
- Lederbogen F, Kirsch P, Haddad L, et al. City living and urban upbringing affect neural social stress processing in humans. *Nature.* 2011;474:498–501.
- Kelleher I, Connor D, Clarke MC, et al. Prevalence of psychotic symptoms in childhood and adolescence: a systematic review and meta-analysis of population-based studies. *Psychol Med.* 2012;42:1857–1863.
- Scott J, Chant D, Andrews G, McGrath J. Psychotic-like experiences in the general community: the correlates of CIDI psychosis screen items in an Australian sample. *Psychol Med.* 2006;36:231–238.
- Yoshizumi T, Murase S, Honjo S, Kaneko H, Murakami T. Hallucinatory experiences in a community sample of

- Japanese children. *J Am Acad Child Adolesc Psychiatry*. 2004;43:1030–1036.
25. Horwood J, Salvi G, Thomas K, et al. IQ and non-clinical psychotic symptoms in 12-year-olds: results from the ALSPAC birth cohort. *Br J Psychiatry*. 2008;193:185–191.
 26. Dhossche D, Ferdinand R, Van der Ende J, Hofstra MB, Verhulst F. Diagnostic outcome of self-reported hallucinations in a community sample of adolescents. *Psychol Med*. 2002;32:619–627.
 27. Kelleher I, Cannon M. Psychotic-like experiences in the general population: characterizing a high-risk group for psychosis. *Psychol Med*. 2011;41:1–6.
 28. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med*. 2009;39:179–195.
 29. Polanczyk G, Moffitt TE, Arseneault L, et al. Etiological and clinical features of childhood psychotic symptoms: results from a birth cohort. *Arch Gen Psychiatry*. 2010;67:328–338.
 30. Kelleher I, Keeley H, Corcoran P, et al. Clinicopathological significance of psychotic experiences in non-psychotic young people: evidence from four population-based studies. *Br J Psychiatry*. 2012;201:26–32.
 31. Poulton R, Caspi A, Moffitt TE, et al. Children's self-reported psychotic symptoms and adult schizophreniform disorder: a 15-year longitudinal study. *Arch Gen Psychiatry*. 2000;57:1053–1058.
 32. Fisher HL, Caspi A, Poulton R, et al. Specificity of childhood psychotic symptoms for predicting schizophrenia by 38 years of age: a birth cohort study. *Psychol Med*. 2013;43:2077–2086.
 33. Kelleher I, Lynch F, Harley M, et al. Psychotic symptoms in adolescence index risk for suicidal behavior: findings from 2 population-based case-control clinical interview studies. *Arch Gen Psychiatry*. 2012;69:1277–1283.
 34. Dragt S, Nieman DH, Veltman D, et al. Environmental factors and social adjustment as predictors of a first psychosis in subjects at ultra high risk. *Schizophr Res*. 2011;125:69–76.
 35. Spauwen J, Krabbendam L, Lieb R, Wittchen HU, van Os J. Evidence that the outcome of developmental expression of psychosis is worse for adolescents growing up in an urban environment. *Psychol Med*. 2006;36:407–415.
 36. March D, Hatch SL, Morgan C, et al. Psychosis and place. *Epidemiol Rev*. 2008;30:84–100.
 37. Zammit S, Lewis G, Rasbash J, et al. Individual, schools, and neighbourhood. *Arch Gen Psychiatry*. 2010;67:914–922.
 38. Silver E, Mulvey EP, Swanson JW. Neighborhood structural characteristics and mental disorder: Faris and Dunham revisited. *Soc Sci Med*. 2002;55:1457–1470.
 39. Allardyce J, Gilmour H, Atkinson J, et al. Social fragmentation, deprivation and urbanicity: relation to first-admission rates for psychoses. *Br J Psychiatry*. 2005;187:401–406.
 40. Bhavsar V, Boydell J, Murray R, Power P. Identifying aspects of neighbourhood deprivation associated with increased incidence of schizophrenia. *Schizophr Res*. 2014;156:115–121.
 41. Omer S, Kirkbride JB, Pringle DG, et al. Neighbourhood-level socio-environmental factors and incidence of first episode psychosis by place at onset in rural Ireland: The Cavan-Monaghan First Episode Psychosis Study [CAMFEPS]. *Schizophr Res*. 2014;152:152–157.
 42. Koppel S, McGuffin P. Socio-economic factors that predict psychiatric admissions at a local level. *Psychol Med*. 1999;29:1235–1241.
 43. Goldsmith HF, Holzer CE, Manderscheid RW. Neighborhood characteristics and mental illness. *Eval Program Plann*. 1998;21:211–225.
 44. Massey DS. The age of extremes: concentrated affluence and poverty in the twenty-first century. *Demography*. 1996;33:395–412.
 45. Chan KY, Zhao FF, Meng S, et al. Urbanization and the prevalence of schizophrenia in China between 1990 and 2010. *World Psychiatry*. 2015;14:251–252.
 46. Lögdberg B, Nilsson LL, Levander MT, Levander S. Schizophrenia, neighbourhood, and crime. *Acta Psychiatr Scand*. 2004;110:92–97.
 47. Veling W, Susser E, Selten JP, Hoek HW. Social disorganization of neighborhoods and incidence of psychotic disorders: a 7-year first-contact incidence study. *Psychol Med*. 2015;45:1789–1798.
 48. Kirkbride JB, Boydell J, Ploubidis GB, et al. Testing the association between the incidence of schizophrenia and social capital in an urban area. *Psychol Med*. 2008;38:1083–1094.
 49. Lederbogen F, Haddad L, Meyer-Lindenberg A. Urban social stress—risk factor for mental disorders. The case of schizophrenia. *Environ Pollut*. 2013;183:2–6.
 50. Selten JP, van der Ven E, Rutten BP, Cantor-Graae E. The social defeat hypothesis of schizophrenia: an update. *Schizophr Bull*. 2013;39:1180–1186.
 51. Kirkbride JB, Stochl J, Zimbrón J, et al. Social and spatial heterogeneity in psychosis proneness in a multilevel case-prodrome-control study. *Acta Psychiatr Scand*. 2015;132:283–292.
 52. Sampson RJ, Raudenbush SW, Earls F. Neighborhoods and violent crime: a multilevel study of collective efficacy. *Science*. 1997;277:918–924.
 53. Sampson RJ, Raudenbush SW. Systematic social observation of public spaces: a new look at disorder in urban Neighborhoods 1. *Am J Sociol*. 1999;105:603–651.
 54. Sampson RJ, Morenoff JD, Gannon-Rowley T. Assessing “neighborhood effects”: Social processes and new directions in research. *Annu Rev Sociol*. 2002;28:443–478.
 55. Trouton A, Spinath FM, Plomin R. Twins early development study (TEDS): a multivariate, longitudinal genetic investigation of language, cognition and behavior problems in childhood. *Twin Res*. 2002;5:444–448.
 56. Moffitt TE; E-Risk Study Team. Teen-aged mothers in contemporary Britain. *J Child Psychol Psychiatry*. 2002;43:727–742.
 57. CACI Information Services. *ACORN User Guide*. London, UK: CACI; 2006.
 58. Caspi A, Taylor A, Moffitt TE, Plomin R. Neighborhood deprivation affects children's mental health: environmental risks identified in a genetic design. *Psychol Sci*. 2000;11:338–342.
 59. Schreier A, Wolke D, Thomas K, et al. Prospective study of peer victimization in childhood and psychotic symptoms in a nonclinical population at age 12 years. *Arch Gen Psychiatry*. 2009;66:527–536.
 60. Odgers CL, Moffitt TE, Tach LM, et al. The protective effects of neighborhood collective efficacy on British children growing up in deprivation: a developmental analysis. *Dev Psychol*. 2009;45:942–957.
 61. Odgers CL, Caspi A, Bates CJ, Sampson RJ, Moffitt TE. Systematic social observation of children's neighborhoods using Google Street View: a reliable and cost-effective method. *J Child Psychol Psychiatry*. 2012;53:1009–1017.
 62. Odgers CL, Caspi A, Russell MA, et al. Supportive parenting mediates neighborhood socioeconomic disparities in children's

- antisocial behavior from ages 5 to 12. *Dev Psychopathol.* 2012;24:705–721.
63. Jaffee SR, Caspi A, Moffitt TE, Polo-Tomás M, Taylor A. Individual, family, and neighborhood factors distinguish resilient from non-resilient maltreated children: a cumulative stressors model. *Child Abuse Negl.* 2007;31:231–253.
 64. Veitch J, Bagley S, Ball K, Salmon J. Where do children usually play? A qualitative study of parents' perceptions of influences on children's active free-play. *Health Place.* 2006;12:383–393.
 65. Kohen DE, Leventhal T, Dahinten VS, McIntosh CN. Neighborhood disadvantage: pathways of effects for young children. *Child Dev.* 2008;79:156–169.
 66. March JS, Parker JD, Sullivan K, Stallings P, Conners CK. The Multidimensional Anxiety Scale for Children (MASC): factor structure, reliability, and validity. *J Am Acad Child Adolesc Psychiatry.* 1997;36:554–565.
 67. Kovacs M. *Children's Depression Inventory: Manual.* North Tonawanda, NY: Multi-Health Systems; 1992.
 68. Rivera CL, Bernal G, Rosello J. The Children Depression Inventory (CDI) and the Beck Depression Inventory (BDI): their validity as screening measures for major depression in a group of Puerto Rican adolescents. *Int J Clin Health Psychol.* 2005;5:485.
 69. Achenbach T. *Manual for the Child Behavior Checklist and 1991 Profile.* Burlington, VT: University of Vermont Department of Psychiatry; 1991.
 70. Odgers CL, Donley S, Caspi A, Bates CJ, Moffitt TE. Living alongside more affluent neighbors predicts greater involvement in antisocial behavior among low-income boys. *J Child Psychol Psychiatry.* 2015;56:1055–1064.
 71. Milne BJ, Caspi A, Crump R, et al. The validity of the family history screen for assessing family history of mental disorders. *Am J Med Genet B Neuropsychiatr Genet.* 2009;150B:41–49.
 72. Robins L, Cottler L, Bucholz K, Compton W. *Diagnostic Interview Schedule for DSM-IV (DIS-IV).* St Louis, MO: Washington University School of Medicine; 1995.
 73. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders.* 4th ed. Washington, DC: American Psychiatric Association; 1994.
 74. Breen R, Karlson KB, Holm A. Total, direct, and indirect effects in logit and probit models. *Sociol Methods Res.* 2013;42:164–191.
 75. Rogers W. Regression standard errors in clustered samples. *STB.* 1994;3:19–23.
 76. van Winkel R, Stefanis NC, Myin-Germeys I. Psychosocial stress and psychosis. A review of the neurobiological mechanisms and the evidence for gene-stress interaction. *Schizophr Bull.* 2008;34:1095–1105.
 77. van Os J, Pedersen CB, Mortensen PB. Confirmation of synergy between urbanicity and familial liability in the causation of psychosis. *Am J Psychiatry.* 2004;161:2312–2314.
 78. Garety PA, Kuipers E, Fowler D, Freeman D, Bebbington PE. A cognitive model of the positive symptoms of psychosis. *Psychol Med.* 2001;31:189–195.
 79. Oher FJ, Demjaha A, Jackson D, et al. The effect of the environment on symptom dimensions in the first episode of psychosis: a multilevel study. *Psychol Med.* 2014;44:2419–2430.
 80. Chung HL, Steinberg L. Relations between neighborhood factors, parenting behaviors, peer deviance, and delinquency among serious juvenile offenders. *Dev Psychol.* 2006;42:319–331.
 81. Lederbogen F, Kirsch P, Haddad L, et al. City living and urban upbringing affect neural social stress processing in humans. *Nature.* 2011;474:498–501.
 82. Haddad L, Schäfer A, Streit F, et al. Brain structure correlates of urban upbringing, an environmental risk factor for schizophrenia. *Schizophr Bull.* 2015;41:115–122.
 83. van Os J, Linscott RJ, Myin-Germeys I, Delespaul P, Krabbendam L. A systematic review and meta-analysis of the psychosis continuum: evidence for a psychosis proneness-persistence-impairment model of psychotic disorder. *Psychol Med.* 2009;39:179–195.
 84. Costello EJ, Mustillo S, Erkanli A, Keeler G, Angold A. Prevalence and development of psychiatric disorders in childhood and adolescence. *Arch Gen Psychiatry.* 2003;60:837–844.