

# UC San Diego

## UC San Diego Previously Published Works

**Title**

Mechanical Regulation of Cardiac Aging in Model Systems

**Permalink**

<https://escholarship.org/uc/item/5r31t78d>

**Journal**

Circulation Research, 118(10)

**ISSN**

0009-7330

**Authors**

Sessions, Ayla O  
Engler, Adam J

**Publication Date**

2016-05-13

**DOI**

10.1161/circresaha.116.307472

Peer reviewed

This Review is in a thematic series on **Aging and the Cardiovascular System**, which includes the following articles:

**Mechanical Regulation of Cardiac Aging in Model Systems**

Aging and Autophagy in the Heart

Macromolecular Degradation Systems and Cardiovascular Aging

Mitochondrial Metabolism in Aging Heart

Dietary Interventions, Cardiovascular Aging, and Disease: Animal Models and Human Studies

Pharmacological Strategies to Retard Cardiovascular Aging

*Guest Editors: Guido Kroemer and Lorenzo Galluzzi*

## Mechanical Regulation of Cardiac Aging in Model Systems

Ayla O. Sessions, Adam J. Engler

**Abstract:** Unlike diet and exercise, which individuals can modulate according to their lifestyle, aging is unavoidable.

With normal or healthy aging, the heart undergoes extensive vascular, cellular, and interstitial molecular changes that result in stiffer less compliant hearts that experience a general decline in organ function. Although these molecular changes deemed cardiac remodeling were once thought to be concomitant with advanced cardiovascular disease, they can be found in patients without manifestation of clinical disease. It is now mostly acknowledged that these age-related mechanical changes confer vulnerability of the heart to cardiovascular stresses associated with disease, such as hypertension and atherosclerosis. However, recent studies have aimed at differentiating the initial compensatory changes that occur within the heart with age to maintain contractile function from the maladaptive responses associated with disease. This work has identified new targets to improve cardiac function during aging. Spanning invertebrate to vertebrate models, we use this review to delineate some hallmarks of physiological versus pathological remodeling that occur in the cardiomyocyte and its microenvironment, focusing especially on the mechanical changes that occur within the sarcomere, intercalated disc, costamere, and extracellular matrix.

(*Circ Res.* 2016;118:1553-1562. DOI: 10.1161/CIRCRESAHA.116.307472.)

**Key Words:** cardiomyocyte hypertrophy ■ cardiomyopathy ■ cardiovascular disease ■ *Drosophila* ■ pathophysiology ■ physiology

The majority of people afflicted by cardiovascular disease (CVD) and heart failure are of advanced age. Although aging specifically is not thought to be the cause, it is an independent risk factor for CVD because both the incidence and the prevalence of CVD increases dramatically after the age of 50 years for men and women.<sup>1</sup> The heart undergoes extensive structural changes with aging independent of disease; this is marked by an asymmetrical increase in septum thickness and overall decrease in ventricular tissue compliance.<sup>2</sup> However, when changes are superimposed with cardiac stresses associated with disease, the occurrence and severity of disease increases. Regardless of pathology, however, these global age-related, organ-level structural changes are all rooted in intra- and extracellular remodeling, leading

to hypertrophied cardiomyocytes regardless of sex or disease state. This remodeling is thought at least, in part, to reinforce organ structure under increased systolic pressure and afterload experienced with age.<sup>3,4</sup> Cardiomyocytes must maintain function with increasing loads via compensatory molecular mechanisms that span the sarcomere, intercalated disc (ID), and the sarcolemma. Proteins within these units remodel with increased stress and strain associated with aging to maintain function, but the accumulation of remodeling events may lead to a pathological response resulting to heart failure. Recent studies have focused on differentiating which specific molecular changes that accumulate with age help maintain cardiomyocyte function, and can be deemed hallmarks of physiological aging, versus those that aid in

Original received February 4, 2016; revision received March 29, 2016; accepted April 4, 2016.

From the Biomedical Sciences Program (A.O.S., A.J.E.) and Department of Bioengineering, University of California, San Diego, La Jolla (A.J.E.); and Sanford Consortium for Regenerative Medicine, La Jolla, CA (A.J.E.).

Correspondence to Adam J. Engler, PhD, Sanford Consortium for Regenerative Medicine, 2880 Torrey Pines Scenic Dr, Room 2005, La Jolla, CA 92037. E-mail aengler@ucsd.edu

© 2016 American Heart Association, Inc.

*Circulation Research* is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.116.307472

**Nonstandard Abbreviations and Acronyms**

<b><math>\alpha</math>-CAT</b>	$\alpha$ -catenin
<b><math>\beta</math>-CAT</b>	$\beta$ -catenin
<b>BM</b>	basement membrane
<b>CVD</b>	cardiovascular disease
<b>cTnT</b>	cardiac tropomyosin-binding troponin
<b>DCM</b>	dilated cardiomyopathy
<b>ECM</b>	extracellular matrix
<b>HF</b>	heart failure
<b>ID</b>	intercalated disc
<b>LV</b>	left ventricle
<b>MHC</b>	myosin heavy chain
<b>MLC</b>	myosin light chain
<b>MMP</b>	matrix metalloproteinase
<b>N-CAD</b>	N-cadherin

disease progression during advanced age, and can be deemed pathological aging.

The past few decades have seen the emergence of many likely and unlikely animal models used for their unique ability to study certain aspects of physiological and pathological cardiac aging, disease, or both. Recent aging studies have used large mammalian models such as simians, pigs, and dogs because they may more closely model functional decline and respond similarly to treatments. However, these models are costly and slow to age. Smaller, cheaper, and easier to use models are ideal to elucidate cellular and molecular mechanisms of aging. For that reason, the bulk of cardiac aging and disease research has focused on rat and mouse models, rather than larger mammalian counterparts. Yet, mice and rats have significantly increased heart rates over humans, and still require 2 years to age from young to old. The need for a rapidly aging, genetically tractable, and genetically homologous model has led to the recent popularity of the fruit fly, *Drosophila melanogaster*, as a model for age-related CVD. The fruit fly is uniquely suited to study cardiovascular aging and disease because of its relatively short lifespan  $\approx$ 5 to 7 weeks, conservation of cardiac disease genes,<sup>5</sup> the breadth of transgenic systems,<sup>6–8</sup> and ease of in situ measurement of cardiac parameters.<sup>9–12</sup> With all of these models and tools, the scientific community can use multivariate approaches to explore how the age-related mechanical changes that occur in cardiomyocytes progress from maintenance of physiological function to those that induce pathological disease.

Using the model systems introduced above, this review highlights a range of protein complexes implicated in regulating the mechanical sensing and transmitting of forces necessary for physiological cellular remodeling under increased load and stress,<sup>13–15</sup> which with age may become increasingly dysregulated. We will first focus on the intracellular cytoskeletal and sarcomeric remodeling that leads to age-related cardiac hypertrophy. In this context, we will pay close attention to separate physiological and pathological cardiac aging to highlight differences, which to date have been less appreciated.


## Intracellular Remodeling as a Result of Stress, Age, or Disease

Cardiomyocytes undergo extensive morphological changes during development, stress, and aging due in large part to outside-in transduction of mechanical signals to the sarcomeres, cytoskeleton, and junctions, such as IDs and costameres. Internal remodeling gives cardiomyocytes plasticity to adjust to increased mechanical loads, especially when cells must compensate to preserve function. Here, we detail each structure and outline its pathological remodeling with disease and its potential physiological remodeling with age. For reference, Figure 1 contains many of the proteins referenced in this section and is color-coded based on whether expression changes have been shown to change with physiological aging before disease progression or still only associated with pathological cardiac aging; proteins in black are mentioned in the text without indicating conclusively if they contribute to either, whereas those in gray are shown to indicate overall structure.

## Sarcomeric Remodeling and Maintenance

The sarcomere is a complex assembly of thick (myosin) and thin (actin) filaments responsible for force generation within the cardiomyocyte. Sarcomeres generate the contractile force within myocytes together with regulatory proteins, such as troponins and tropomyosin, and cytoskeletal components, such as titin and myosin-binding protein C. Sarcomere function is spatially and temporally regulated by post-translational modifications and alternative splice isoforms. Therefore, aberrant expression of certain isoforms with age may result in reduced contraction, reduced mechanical response, and disease.

In mammalian hearts, 2 myosin heavy chain (MHC) isoforms,  $\alpha$  and  $\beta$ , exist.  $\alpha$ -MHC is associated with higher ATPase activity, greater contractile velocity,<sup>16</sup> and it is the dominant form of MHC in rodent left ventricle (LV), that is,  $>90\%$ .<sup>17</sup> In adult humans,  $\beta$ -MHC comprises  $\approx 95\%$  of MHC found in ventricles<sup>18</sup> and produces more force per power stroke.<sup>16</sup> Failing human myocardium experiences a further decrease in  $\alpha$ -MHC expression (Figure 1, blue).<sup>19,20</sup> Rat hearts experience a similar decrease in  $\alpha$ -MHC with aortic banding,<sup>21</sup> suggesting that cardiomyocyte regulation of MHC isoforms during pressure overload is not retarded by age and is more closely associated with pathology. Because MHC isoforms have unique effects on contractility, decreasing  $\alpha$ -MHC could negatively affect contractile velocity and impair adaptation to increased cardiac demand. Unfortunately, age-related MHC expression data outside of disease are lacking, which could further clarify if this is a response to pressure changes or downstream activator of pathological remodeling. Similarly, essential myosin light chain (MLC) undergoes isoform shifts during disease, with an increase in the ratio of atrial to ventricular isoforms expressed with hypertrophic cardiomyopathy.<sup>22</sup> In addition to isoform switches, phosphorylation of ventricular MLC-2 regulates cross-bridge cycling; when ventricular MLC-2 is phosphorylated, hearts exhibit greater force production per myosin power stroke by slowing down cycling kinetics and prolonging the duty cycle, allowing more myosin heads to come into contact with actin.<sup>23,24</sup> Conversely, reduced phosphorylation in mice exhibit dilated cardiomyopathy (DCM)-like



**Figure 1. Physiological vs pathological age-related changes of the sarcomere and intercalated disc (ID).** Schematic representation of sarcomere and ID components in the cardiomyocyte implicated in physiological aging (red, upregulated; green, downregulated), pathological aging (yellow, upregulated; blue, downregulated), or yet undetermined (black). Arrows to nucleus indicate translocation of protein into the nucleus with age.  $\alpha$ -CAT indicates  $\alpha$ -catenin; ACTN1,  $\alpha$ -actinin-1;  $\alpha$ -MHC,  $\alpha$ -myosin heavy chain;  $\beta$ -CAT,  $\beta$ -catenin; c-TnT4, cardiac troponin T-4; F-actin, filamentous actin; MLP, muscle-binding limb protein; N-CAD, N-cadherin; P-MLC-2, phosphorylated myosin light chain-2; and Vinc, vinculin.

remodeling.<sup>24,25</sup> Humans also shift to more dephosphorylated MLC-2 in diseased and failing myocardium,<sup>26–28</sup> suggesting that cross-bridge cycling becomes increasingly dysregulated with disease (Figure 1, blue). Yet much of how both MHC and MLC isoforms change with physiological aging have yet to be discovered, owing largely to a focus on how mutations cause familial hypertrophic cardiomyopathy. Because of the strong correlation of heart failure and the decline of  $\alpha$ -MHC and phosphorylated MLC-2 across species, these data suggest that a threshold of  $\alpha$ -MHC expression and MLC-2 phosphorylation is needed for proper cardiomyocyte contraction; thus declining levels of either protein may be a marker for pathology.

In addition to thick filaments, some thin filament proteins change with aging and disease. For example, cardiac troponin-binding protein troponin T (cTnT) undergoes alternative splicing to confer increased myofilament  $\text{Ca}^{2+}$  sensitivity in failing human hearts.<sup>29,30</sup> However, a cTnT3 to cTnT4 isoform switch does not enhance disease progression (Figure 1, red),<sup>31</sup> suggesting that the switch to more  $\text{Ca}^{2+}$ -sensitive isoform might compensate for increased hemodynamic stress. In

*Drosophila*, mutations adjacent to a highly conserved TnT1 tropomyosin-binding domain result in excessive cross-bridge formation at rest even under low  $\text{Ca}^{2+}$ , prolonged muscle activation, and restrictive cardiomyopathy.<sup>32</sup> Because these cTnT changes result in diastolic dysfunction for fly hearts, this model could explain why aging hearts experience increased resting tension and stiffness and ultimately impaired function. However a cTnT isoform expression profile with age is needed to confirm that the isoform switch is part of physiological compensation under increased stress because prolonged stress could have adverse consequences.

Maintaining myofilament structure and function with age and increased load is a combinatory process across the sarcomere. Major mechanical regulators of sarcomere structure, such as titin<sup>14</sup>—a giant sarcomeric protein that spans from Z-disc to M-band,<sup>33</sup> seem to contribute to this balance between physiological compensation and pathological remodeling. Titin acts as a sarcomeric ruler,<sup>33,34</sup> generates resting stiffness, and transduces signals in times of mechanical overload.<sup>35</sup> The role of titin in mechanically regulating the switch from physiological

aging to pathological remodeling, and heart failure has been extensively reviewed (Figure 1, yellow).<sup>36</sup> Recent work studying titin-associated proteins demonstrated that only patients with heart failure with preserved ejection fraction exhibited a significant increase in titin-dependent stiffness,<sup>37</sup> reinforcing the importance of this protein and proteins in complex with it in regulating the switch from normal to pathological function. As another example, the Z-disc protein  $\alpha$ -actinin anchors actin filaments at the Z-disc.<sup>14,38,39</sup> Mutations in  $\alpha$ -actinin are associated with familial hypertrophic cardiomyopathy and arrhythmias, suggesting that it is integral for proper sarcomeric growth and force propagation.<sup>40</sup> Patients who recover enough LV function to warrant explantation of their left ventricular assist device often experience a significant increase in  $\alpha$ -actinin,<sup>41</sup> implying that  $\alpha$ -actinin could scale with sarcomeric function. In addition to concentration,  $\alpha$ -actinin lattice organization has been implicated in disease severity; canine heart failure models show partial rescue of  $\alpha$ -actinin lattice structure post cardiac resynchronization therapy (Figure 1, blue).<sup>42</sup> However, it remains undetermined if these changes precede pathology because aging or stress-induced studies involving  $\alpha$ -actinin are lacking (Figure 1, black). However, nonsarcomeric  $\alpha$ -actinin, which aids costameric and ID adhesion complexes, does experience an increase in aged diseased hearts (Figure 1 and 2, yellow),<sup>43</sup> but how these levels change with physiological aging have not yet been studied. Beyond these most commonly discussed proteins, a variety of other models and proteins have been studied to a lesser extent, such as *Drosophila* and muscle limb protein expression and localization changes (Figure 1, yellow). Common muscle limb protein polymorphisms result in DCM,<sup>44</sup> creating cardiomyocytes unable to sense load.<sup>45</sup> Muscle limb protein is also shuttled to the nucleus on hypertrophic response,<sup>46,47</sup> suggesting that mislocalization results in impaired mechanosensing (Figure 1, arrow).<sup>48</sup> These data clearly imply that muscle limb protein and  $\alpha$ -actinin are tied to mechanical sensing, but future work needs to explore their changes with age outside of disease.

### ID Reinforcement and Gap Junction Remodeling With Age

The ID forms a specialized junction between adjacent cardiomyocytes to propagate longitudinal forces from sarcomeres to adjacent cells. Coupling is achieved through actin cytoskeleton bridges that occur specifically within the fascia adherens and reinforced by the desmosome; electric coupling is achieved through gap junctions.<sup>13</sup> In recent years, the ID has emerged as a key mechanical component, permitting the cell to sense and respond to stress, and load via ultrastructural changes as in volume-overloaded rabbit<sup>49</sup> and mouse hearts<sup>50</sup> as well as human DCM patients.<sup>50</sup> These correlations imply that IDs may have a more regulatory role in cardiomyocyte hypertrophy, which is evident when examining key components of ID and gap junction substructures. Both are regulated with age across animal models,<sup>51–53</sup> suggesting that cardiomyocytes transition from adaptive remodeling to pathological loss of sarcomere regulation.

Within these junctions, N-cadherins (N-CAD), vinculin, and the catenins form  $\text{Ca}^{2+}$ -dependent cell–cell adhesions that

are linked to the cortical actin cytoskeleton.<sup>54</sup> By forming a mechanical link to adjacent cardiomyocytes,<sup>36</sup> they may help to induce pathological remodeling in the heart. For example, N-CAD knockout mice lose ID and gap junction structures but exhibit growth in the LV long axis,<sup>55</sup> suggesting that N-CAD could regulate eccentric or concentric sarcomeric growth. Yet, mouse aging studies show that although other junction components change expression with age, such as connexin 43 and  $\beta$ -catenin ( $\beta$ -CAT), N-CAD expression remains unchanged.<sup>52</sup> These data imply that although N-CAD may mechanically regulate differential cell growth (Figure 1, black), other junctional proteins may regulate disease. The catenins, for example, link N-CAD's cytoplasmic tail to actin via vinculin, and  $\beta$ -CAT in particular changes its localization to the nucleus with age (Figure 1, arrow),<sup>56</sup> mediating fetal gene reactivation through novel transcriptional activity (Figure 1, yellow).<sup>57</sup> Recent work showed that although  $\beta$ -CAT overexpression has no measurable effects, its knockdown induces significant dysfunction.<sup>58</sup> Although  $\beta$ -CAT could have a role in pathological disease, we have recently demonstrated that its counterpart,  $\alpha$ -catenin ( $\alpha$ -CAT), was upregulated in aged, nondiseased simian, and fly hearts using proteomic and genetic analyses. Data suggest that  $\alpha$ -CAT seems upregulated with physiological aging<sup>51</sup> and may precede pathological remodeling (Figure 1, red). Indeed mechanical stimulation of  $\alpha$ -CAT via N-CAD triggers vinculin recruitment to junctions, demonstrating that  $\alpha$ -CAT is necessary for force-mediated remodeling.<sup>59,60</sup> Although consensus is building for force-dependent regulation of the junctions,<sup>61,62</sup> further work is needed to explore how  $\alpha$ -CAT combines with other signals to affect function with age.

Finally vinculin, which is another critical linker protein within the ID, binds to  $\alpha$ -CAT to translate sarcomeric contraction into cell shortening. In aging patients with heart failure, vinculin undergoes remodeling,<sup>53,63</sup> which can be reversed with left ventricular assist devices.<sup>64,65</sup> In a recent comprehensive study across simian, rat, and fly, we demonstrated that upregulation of vinculin with age is conserved from invertebrates to mammals and that its role is compensatory rather than deleterious in aging (Figure 1, red).<sup>51</sup> Vinculin overexpression resulted in increased cortical stiffness most significantly at the ID where it increased longitudinal myofibril anchoring between adjacent cells and increased force transmission. Increased cardiac vinculin correlated with better force production and fractional shortening in fly hearts,<sup>51</sup> further supporting the idea that vinculin upregulation and localization to the ID with age aids the cardiomyocyte in preserving contraction, even under mechanical load. Metavinculin, a muscle-specific alternative splicing, is more associated to actin bundling<sup>66</sup> and involved in DCM versus aging.<sup>66,67</sup> Metavinculin expression levels have not been examined as a function of age, but these molecular interactions could play a large role in the mechanical regulation of vinculin-mediated cytoskeletal reinforcement mentioned above, ultimately adding another potential switch from normal to maladaptive remodeling.

Whether considering cardiomyocyte hypertrophy or subcellular remodeling, it is important to keep in mind that such mechanisms are often interrelated and complex, especially because multiple proteins within a single complex, such as



the ID, can act in opposing directions with age and disease (Figure 1). Thus, we think that although it is difficult to dissect and identify, it is critical to recognize the existence of intracellular mechanisms that counterbalance aging without necessarily contributing to disease.

### Costamere and Extracellular Matrix Remodeling in Response to Age, Stress, and Disease

In addition to internal remodeling, cardiomyocytes experience changes in their surrounding extracellular environment with age. Under physiological conditions, cardiac extracellular matrix (ECM) provides structural support for cardiomyocytes and facilitates force transduction. However, increased ECM deposition and crosslinking along with decreased ECM degradation have all been implicated in aged patients.<sup>68–70</sup> These changes affect cardiomyocyte function<sup>71</sup> not only by changing the mechanical properties of the tissue but also by altering transmembrane receptors.<sup>72–74</sup> Cardiac ECM has been recently reviewed,<sup>75–77</sup> so we will focus on compositional changes that have been implicated in switching from adaptive to maladaptive remodeling (Figure 2, top). On the opposing side of the sarcolemma (Figure 2, bottom), costameres aid in cell–ECM adhesion by adhering Z-discs to ECM.<sup>78,79</sup> Given their changes with age, we will begin our discussion with costamere complexes, specifically the vinculin/talin/integrin structure given its known changes with age,<sup>80</sup> and transition to ECM using Figure 2 to illustrate our conclusions on physiological and pathological changes.

#### Costamere Complex Reinforcement

Costameres are periodic vinculin-containing structures on the lateral surface of cardiomyocytes<sup>81</sup> that link Z-discs to integrins via talin. Vinculin is integral in responding to mechanical stimulus, and patients with DCM and hypertrophic cardiomyopathy often have vinculin mutations.<sup>66</sup> In addition to disease, we have shown that as in humans,<sup>53,63</sup> age-related increases in cardiac vinculin are observed in simians and flies. Vinculin overexpression in *Drosophila* increased cortical stiffness at both the ID and the costameres. Improved assembly maintained myofilament lattice spacing but significantly improved lattice order (Figure 2, red).<sup>51</sup> These findings confirmed data in mice, which showed that cardiac vinculin knockout resulted in significantly more compliant cardiomyocytes and greater myofilament spacing before dysfunction.<sup>82</sup> Improved myofilament spacing may enhance cross-bridge cycling, which increases force production. Future work examining aged, non-diseased human myocardium will be necessary to corroborate vinculin upregulation with age in simian and *Drosophila* models, but these data establish a link between vinculin and compensatory nonmaladaptive response within aged tissue.


Together with vinculin, talin links the actin cytoskeleton to integrins and regulates their binding affinity across species.<sup>83–85</sup> Talin has 2 genes in vertebrates, with talin-2 largely expressed in adult myocardium.<sup>86</sup> Until recently, their role in the heart was unknown, but pharmacological and mechanical stress of mouse hearts revealed an upregulation of talin-1, which may be a compensatory response to stress.<sup>87</sup> However

in pressure-overloaded hearts, talin-1 ablation reduced hypertrophy and improved function,<sup>87</sup> suggesting that pressure-induced upregulation of talin-1 may be deleterious and thus a pathological marker (Figure 2, yellow). Conversely talin-2 is elevated in aged mice (Figure 2, red),<sup>88</sup> specifically between middle aged (10–16 months) and aged (20–24 months) mice.<sup>88</sup> When hearts are unloaded, for example, via left ventricular assist device, talin-2 is downregulated.<sup>41</sup> Although these data suggest that talin is involved in remodeling, aged nondiseased patient samples are needed to better explore if regulation is age-associated or merely a response to external load. Invertebrate studies may prove useful for future work because they only have 1 talin gene and age rapidly.

A final major component in this complex is integrin, which occurs as a heterodimer of  $\alpha$  and  $\beta$  subunits<sup>89,90</sup> and directly transduces signals to and from ECM and costameres. The range of integrin dimer pairs confer ECM ligand specificity,<sup>91</sup> so their modulation with age could affect costamere function.  $\alpha7\beta1$  heterodimer is the most prevalent pair in adult cardiomyocytes,<sup>92</sup> but additional  $\alpha$  and  $\beta$  subunits have been detected after pressure overload, post myocardial infarction, and with age.<sup>93–95</sup> In *Drosophila*,  $\beta1$ -integrin increases with age, but moderate attenuation correlates with improved cardiac function and significantly increased organismal lifespan.<sup>73</sup> Ventricular expression of  $\beta1$ -integrin is also increased in aged monkeys;<sup>51</sup> however, cardiac ablation of  $\beta1$ -integrin results in DCM during pressure overload,<sup>74</sup> suggesting that integrins are an integral regulator of stretch and load sensing. Knockdown of integrin-linked kinase, a  $\beta1$ -integrin binding protein, also correlated with improved cardiac function and lifespan.<sup>73</sup> These results confirmed the correlation of integrin-linked kinase inhibition and extended lifespan in *Caenorhabditis elegans*.<sup>96,97</sup> Although increased expression of  $\beta1$ -integrin and integrin-linked kinase with age correlate with impaired function, a moderate reduction in their expression can be cardioprotective, which suggests that tightly controlling  $\beta1$ -integrin (Figure 2, red and yellow stripes) and integrin-linked kinase (Figure 2, yellow) expression could have beneficial effects on function of aged hearts.

#### ECM Protein Composition and Remodeling Changes With Age

Cardiac ECM consists of a complex arrangement of interacting structural proteins, which include a basement membrane (BM) lattice of laminin and collagen IV linked together with fibronectin-mediated attachment to collagen I and III.<sup>98</sup> Accumulation of collagen I and collagen III with age has been documented across mammals<sup>52,53,99,100</sup> and including aged healthy human myocardium, suggesting it occurs with physiological aging.<sup>101</sup> Although these changes are thought to increase LV tissue stiffness<sup>37</sup> and impair ventricular relaxation, the propensity of data in patients with no known CVD suggests that an increase per se is not indicative of pathology. In addition, accumulation with age can occur in the absence of excessive mRNA<sup>102</sup> unlike in CVD,<sup>103</sup> suggesting that genetic regulation correlates with pathological conditions, but not necessary in physiological aging (Figure 2, yellow). However, an increase of collagen III relative to collagen I has been implicated in disease, but it is not clear how collagen III changes



**Figure 2. Physiological vs pathological age-related changes of costamere and extracellular matrix (ECM).** Schematic representation of Costamere and ECM of the cardiomyocyte implicated in physiological aging (red, upregulated; green, downregulated), pathological aging (yellow, upregulated; blue, downregulated), or yet undetermined (black). Stripes of 2 colors indicate lack of consensus on expression levels across species or involvement in physiological vs pathological aging.  $\alpha 7$  indicates  $\alpha 7$ -integrin;  $\beta 1$ ,  $\beta 1$ -integrin; ACTN1,  $\alpha$ -actinin-1; AGE's, advanced glycosylation endproducts; Col I, collagen type 1; Col IV, collagen type IV; F-actin, filamentous actin; ID, intercalated disc; ILK, integrin-linked kinase; MMP2, matrix metalloproteinase-2; MMP9, matrix metalloproteinase-9; MMP14, matrix metalloproteinase-14; and Vinc, vinculin.

with age.<sup>104,105</sup> If anything, an increase in collagen III has been linked to increased tissue plasticity and not rigidity as with collagen I.<sup>106</sup> Classically though, interstitial ECM accumulation is implicated in less compliant myocardium, and excessive collagen I can induce diastolic dysfunction making its role seem more context specific (Figure 2, red and yellow stripes). In addition to accumulation with age, the degree of crosslinking can have a significant effect on tissue stiffness and function.<sup>107</sup> Collagen crosslinking, measured by changes in hydroxylated lysine and proline levels, increases significantly with age in sedentary rats.<sup>108</sup> Collagen crosslinking via advanced glycosylation endproducts also increases with age in canine and human myocardium (Figure 2, yellow),<sup>109,110</sup> which when inhibited, improved tissue compliance.<sup>110</sup> Such pathological crosslinking is at least partly reversible with exercise as well.<sup>107,108</sup>

Fibronectin, an adhesion molecule that links collagen fibers to BM, changes its expression with collagen, increasing

in mice from 12 to 20 months (Figure 2, yellow).<sup>95</sup> When fibronectin is knocked out, mice experience less cardiac remodeling and hypertrophy with transaortic constriction,<sup>111</sup> which together suggest that fibronectin is associated with pathological remodeling with age. In addition to fibronectin, cardiac BM has an important role in transducing mechanical signals between cells. For example, laminin  $\alpha 2$  decreases as a function of age in both human and rat myocardium, but only in rats it is associated with pathological remodeling because of a decrease in myocyte number (Figure 2, black).<sup>112</sup> Conversely, BM collagen IV increases with age in mice, suggesting an overall age-related BM thickening,<sup>113</sup> although it is not clear if age-related changes contribute to pathological remodeling because their response to stress is not clear (Figure 2, black). However, collagen IV is downregulated after left ventricular assist device unloading, providing evidence that collagen IV expression is responding to pressure overload.<sup>64</sup> Regardless, extensive

studies are necessary to explore how reinforcement of the BM with age affects signal transduction across the sarcolemma.

Finally, all of these age-associated ECM changes—especially collagen remodeling—occur in part because matrix metalloproteinases (MMPs), a large family of endopeptidases, degrade ECM.<sup>114</sup> MMP expression changes with age, for example, upregulation of MMPs, such as MMP-2, MMP-9, and MMP-14 (Figure 2, yellow), modulate collagen persistence, turnover, and fibrotic remodeling, but there is not yet consensus on whether the remodeling is maladaptive. Aged rats have significant interstitial fibrosis and depressed MMP-2 levels, which suggested that decreased MMP-2–mediated proteolysis led to ECM accumulation.<sup>115</sup> However in pressure overload, MMP-2 loss improved diastolic dysfunction and suppressed fibrosis.<sup>116</sup> These studies highlight the different actions by which MMP-2 mediates ECM remodeling with age: MMP-2 decrease eventually leads to ECM accumulation, but acute upregulation and chronic pressure overload initiates pathological activation of matrix degradation (Figure 2, blue and yellow stripes). Indeed, mice with impaired MMP expression experience reduced age-related fibrosis but total loss results in DCM.<sup>100</sup> More specifically, MMP-14 is an important regulator of ECM degradation with age via its sensitivity to mechanical load and its ability to activate MMP-2. Increased myocardial MMP-14 has been noted in patients with LV pressure overload,<sup>117</sup> and in vitro studies have shown MMP-14 upregulation with increased wall tension.<sup>118</sup> Induction of pressure overload in mice also leads to increased MMP-14 activation, which correlates with increased ECM accumulation.<sup>119</sup> Thus, it would seem that MMP-14, and subsequent MMP-2 activation, in part regulate adverse cardiac remodeling under stress conditions that mimic aging (Figure 2, yellow). MMP-9 also increases with age in conjunction with diastolic dysfunction (Figure 2, yellow),<sup>120</sup> whereas its deletion prevents age-related myocardial stiffening.<sup>88</sup> Yet in the larger discussion of whether MMPs contribute to pathological or physiological ECM remodeling, it is critical to note that MMPs degrade matrix, preventing accumulation, which is physiological for collagen I but pathological for collagen III. Thus, the context of all of these changes matter significantly to their effect(s) on aging.

### Conclusions

Cardiac aging is a complex process that involves many molecular changes both inside and outside of the cardiomyocyte. In this review, we have highlighted some of the age-related structural changes that occur within the sarcomere and cytoskeletal complexes that longitudinally transmit forces between adjacent cardiomyocytes. For example, we have indicated that age-related vinculin reinforcement of the ID and costamere is beneficial to cardiomyocyte function, whereas decreasing proportion of  $\alpha$ -MHC and phosphorylated MLC-2 correlates strongly with disease progression. However, because of the overlap in function of multiple complexes and proteins within the sarcomere and fascia adherens junction, conclusive age-related data for other cytoskeletal components has been lacking. Although matrix changes often seem associated with pathological changes, we also noted the underappreciated role it has during normal aging, for example, the increase in interstitial collagens that

can precede heart disease. To that end, we have provided evidence that although collagen accumulation can lead to pathological fibrosis, genetic upregulation of collagen mRNA is clearly implicated in disease progression. Despite the evidence we presented, question remains about what constitutes causal evidence of physiological responses such as: how much fibrillar collagen is too much, or is it the extent of crosslinking that determines pathology? How do BM components affect lateral signal transduction from the sarcomere to the ECM? As aging studies increase, phenotypes that were once thought to be solely maladaptive have become increasingly complex because of some instances in which they are actually mechanisms of physiological adaptation of cardiomyocytes with age. We will need more multi-model approaches to stratify some of these molecular changes that have been previously noted with aging in the context of advance disease into hallmarks of physiological reinforcement versus pathological progression to provide causal evidence for our hypotheses.

### Sources of Funding

We thank the National Institutes of Health for funding support (R01AG045428 to A.J. Engler) and fellowship support from the American Heart Association (15PRE25090334 to A.O. Sessions).

### Disclosures

None.

### References

1. Mozaffarian D, Benjamin EJ, Go AS, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2016 Update: A Report From the American Heart Association. *Circulation*. 2016;133:e38–e360. doi: 10.1161/CIR.0000000000000350.
2. Khouri MG, Maurer MS, El-Khoury Rumbarger L. Assessment of age-related changes in left ventricular structure and function by freehand three-dimensional echocardiography. *Am J Geriatr Cardiol*. 2005;14:118–125.
3. Lakatta EG, Levy D. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part II: the aging heart in health: links to heart disease. *Circulation*. 2003;107:346–354.
4. Lakatta EG. Arterial and cardiac aging: major shareholders in cardiovascular disease enterprises: Part III: cellular and molecular clues to heart and arterial aging. *Circulation*. 2003;107:490–497.
5. Cammarato A, Ahrens CH, Alayari NN, Qeli E, Rucker J, Reedy MC, Zmasek CM, Gucek M, Cole RN, Van Eyk JE, Bodmer R, O'Rourke B, Bernstein SI, Foster DB. A mighty small heart: the cardiac proteome of adult *Drosophila melanogaster*. *PLoS One*. 2011;6:e18497. doi: 10.1371/journal.pone.0018497.
6. Wolf MJ, Rockman HA. *Drosophila*, genetic screens, and cardiac function. *Circ Res*. 2011;109:794–806. doi: 10.1161/CIRCRESAHA.111.244897.
7. Wessells RJ, Bodmer R. Age-related cardiac deterioration: insights from *Drosophila*. *Front Biosci*. 2007;12:39–48.
8. Orfanos Z. Transgenic tools for *Drosophila* muscle research. *J Muscle Res Cell Motil*. 2008;29:185–188. doi: 10.1007/s10974-009-9166-0.
9. Ocorr K, Reeves NL, Wessells RJ, Fink M, Chen HS, Akasaka T, Yasuda S, Metzger JM, Giles W, Posakony JW, Bodmer R. KCNQ potassium channel mutations cause cardiac arrhythmias in *Drosophila* that mimic the effects of aging. *Proc Natl Acad Sci U S A*. 2007;104:3943–3948. doi: 10.1073/pnas.0609278104.
10. Cammarato A, Ocorr S, Ocorr K. Enhanced assessment of contractile dynamics in *Drosophila* hearts. *Biotechniques*. 2015;58:77–80. doi: 10.2144/000114255.
11. Kaushik G, Zamboni AC, Fuhrmann A, Bernstein SI, Bodmer R, Engler AJ, Cammarato A. Measuring passive myocardial stiffness in *Drosophila melanogaster* to investigate diastolic dysfunction. *J Cell Mol Med*. 2012;16:1656–1662. doi: 10.1111/j.1582-4934.2011.01517.x.
12. Kaushik G, Fuhrmann A, Cammarato A, Engler AJ. In situ mechanical analysis of myofibrillar perturbation and aging on soft, bilayered



- Drosophila* myocardium. *Biophys J*. 2011;101:2629–2637. doi: 10.1016/j.bpj.2011.10.042.
13. Sheikh F, Ross RS, Chen J. Cell-cell connection to cardiac disease. *Trends Cardiovasc Med*. 2009;19:182–190. doi: 10.1016/j.tcm.2009.12.001.
  14. Gautel M. The sarcomeric cytoskeleton: who picks up the strain? *Curr Opin Cell Biol*. 2011;23:39–46. doi: 10.1016/jceb.2010.12.001.
  15. Hoshijima M. Mechanical stress-strain sensors embedded in cardiac cytoskeleton: Z disk, titin, and associated structures. *Am J Physiol Heart Circ Physiol*. 2006;290:H1313–H1325. doi: 10.1152/ajpheart.00816.2005.
  16. Holubarsch C, Goulette RP, Litten RZ, Martin BJ, Mulieri LA, Alpert NR. The economy of isometric force development, myosin isoenzyme pattern and myofibrillar ATPase activity in normal and hypothyroid rat myocardium. *Circ Res*. 1985;56:78–86.
  17. Mercadier JJ, Lompré AM, Wisniewsky C, Samuel JL, Bercovici J, Swynghedauw B, Schwartz K. Myosin isoenzyme changes in several models of rat cardiac hypertrophy. *Circ Res*. 1981;49:525–532.
  18. Bouvagnet P, Mairhofer H, Leger JO, Puech P, Leger JJ. Distribution pattern of alpha and beta myosin in normal and diseased human ventricular myocardium. *Basic Res Cardiol*. 1989;84:91–102.
  19. Miyata S, Minobe W, Bristow MR, Leinwand LA. Myosin heavy chain isoform expression in the failing and nonfailing human heart. *Circ Res*. 2000;86:386–390.
  20. Nakao K, Minobe W, Roden R, Bristow MR, Leinwand LA. Myosin heavy chain gene expression in human heart failure. *J Clin Invest*. 1997;100:2362–2370. doi: 10.1172/JCI119776.
  21. Takahashi T, Schunkert H, Isoyama S, Wei JY, Nadal-Ginard B, Grossman W, Izumo S. Age-related differences in the expression of proto-oncogene and contractile protein genes in response to pressure overload in the rat myocardium. *J Clin Invest*. 1992;89:939–946. doi: 10.1172/JCI115675.
  22. Ritter O, Luther HP, Haase H, Baltas LG, Baumann G, Schulte HD, Morano I. Expression of atrial myosin light chains but not alpha-myosin heavy chains is correlated in vivo with increased ventricular function in patients with hypertrophic obstructive cardiomyopathy. *J Mol Med (Berl)*. 1999;77:677–685.
  23. Sheikh F, Lyon RC, Chen J. Functions of myosin light chain-2 (MYL2) in cardiac muscle and disease. *Gene*. 2015;569:14–20. doi: 10.1016/j.gene.2015.06.027.
  24. Sheikh F, Ouyang K, Campbell SG, et al. Mouse and computational models link MLC2v dephosphorylation to altered myosin kinetics in early cardiac disease. *J Clin Invest*. 2012;122:1209–1221. doi: 10.1172/JCI61134.
  25. Warren SA, Briggs LE, Zeng H, Chuang J, Chang EI, Terada R, Li M, Swanson MS, Lecker SH, Willis MS, Spinale FG, Maupin-Furlow J, McMullen JR, Moss RL, Kasahara H. Myosin light chain phosphorylation is critical for adaptation to cardiac stress. *Circulation*. 2012;126:2575–2588. doi: 10.1161/CIRCULATIONAHA.112.116202.
  26. van der Velden J, Papp Z, Boontje NM, Zaremba R, de Jong JW, Janssen PM, Hasenfuss G, Stienen GJ. The effect of myosin light chain 2 dephosphorylation on Ca<sup>2+</sup> -sensitivity of force is enhanced in failing human hearts. *Cardiovasc Res*. 2003;57:505–514.
  27. van der Velden J, Papp Z, Zaremba R, Boontje NM, de Jong JW, Owen VJ, Burton PB, Goldman P, Jaquet K, Stienen GJ. Increased Ca<sup>2+</sup> -sensitivity of the contractile apparatus in end-stage human heart failure results from altered phosphorylation of contractile proteins. *Cardiovasc Res*. 2003;57:37–47.
  28. Scruggs SB, Reisdorph R, Armstrong ML, Warren CM, Reisdorph N, Solaro RJ, Buttrick PM. A novel, in-solution separation of endogenous cardiac sarcomeric proteins and identification of distinct charged variants of regulatory light chain. *Mol Cell Proteomics*. 2010;9:1804–1818. doi: 10.1074/mcp.M110.000075.
  29. Anderson PA, Malouf NN, Oakeley AE, Pagani ED, Allen PD. Troponin T isoform expression in humans. A comparison among normal and failing adult heart, fetal heart, and adult and fetal skeletal muscle. *Circ Res*. 1991;69:1226–1233.
  30. Anderson PA, Greig A, Mark TM, Malouf NN, Oakeley AE, Ungerleider RM, Allen PD, Kay BK. Molecular basis of human cardiac troponin T isoforms expressed in the developing, adult, and failing heart. *Circ Res*. 1995;76:681–686.
  31. Chen Z, Higashiyama A, Yaku H, Bell S, Fabian J, Watkins MW, Schneider DJ, Maughan DW, LeWinter MM. Altered expression of troponin T isoforms in mild left ventricular hypertrophy in the rabbit. *J Mol Cell Cardiol*. 1997;29:2345–2354. doi: 10.1006/jmcc.1997.0468.
  32. Viswanathan MC, Kaushik G, Engler AJ, Lehman W, Cammarato A. A *Drosophila melanogaster* model of diastolic dysfunction and cardiomyopathy based on impaired troponin-T function. *Circ Res*. 2014;114:e6–17. doi: 10.1161/CIRCRESAHA.114.302028.
  33. Tskhovrebova L, Trinick J. Titin: properties and family relationships. *Nat Rev Mol Cell Biol*. 2003;4:679–689. doi: 10.1038/nrm1198.
  34. Tskhovrebova L, Bennett P, Gautel M, Trinick J. Titin ruler hypothesis not refuted. *Proc Natl Acad Sci U S A*. 2015;112:E1172. doi: 10.1073/pnas.1422326112.
  35. Krüger M, Linke WA. Titin-based mechanical signalling in normal and failing myocardium. *J Mol Cell Cardiol*. 2009;46:490–498.
  36. Lyon RC, Zanella F, Omens JH, Sheikh F. Mechanotransduction in cardiac hypertrophy and failure. *Circ Res*. 2015;116:1462–1476. doi: 10.1161/CIRCRESAHA.116.304937.
  37. Zile MR, Baicu CF, Ikonomidis JS, Stroud RE, Nietert PJ, Bradshaw AD, Slater R, Palmer BM, Van Buren P, Meyer M, Redfield MM, Bull DA, Granzier HL, LeWinter MM. Myocardial stiffness in patients with heart failure and a preserved ejection fraction: contributions of collagen and titin. *Circulation*. 2015;131:1247–1259. doi: 10.1161/CIRCULATIONAHA.114.013215.
  38. Goldstein MA, Schroeter JP, Michael LH. Role of the Z band in the mechanical properties of the heart. *FASEB J*. 1991;5:2167–2174.
  39. Goldstein MA, Michael LH, Schroeter JP, Sass RL. Structural states in the Z band of skeletal muscle correlate with states of active and passive tension. *J Gen Physiol*. 1988;92:113–119.
  40. Girolami F, Iascone M, Tomberli B, et al. Novel  $\alpha$ -actinin 2 variant associated with familial hypertrophic cardiomyopathy and juvenile atrial arrhythmias: a massively parallel sequencing study. *Circ Cardiovasc Genet*. 2014;7:741–750. doi: 10.1161/CIRCGENETICS.113.000486.
  41. Latif N, Yacoub MH, George R, Barton PJ, Birks EJ. Changes in sarcomeric and non-sarcomeric cytoskeletal proteins and focal adhesion molecules during clinical myocardial recovery after left ventricular assist device support. *J Heart Lung Transplant*. 2007;26:230–235. doi: 10.1016/j.healun.2006.08.011.
  42. Lichter JG, Carruth E, Mitchell C, Barth AS, Aiba T, Kass DA, Tomaselli GF, Bridge JH, Sachse FB. Remodeling of the sarcomeric cytoskeleton in cardiac ventricular myocytes during heart failure and after cardiac resynchronization therapy. *J Mol Cell Cardiol*. 2014;72:186–195. doi: 10.1016/j.yjmcc.2014.03.012.
  43. Hein S, Block T, Zimmermann R, Kostin S, Scheffold T, Kubin T, Klövekorn WP, Schaper J. Deposition of nonsarcomeric alpha-actinin in cardiomyocytes from patients with dilated cardiomyopathy or chronic pressure overload. *Exp Clin Cardiol*. 2009;14:e68–e75.
  44. Knöll R, Hoshijima M, Hoffman HM, et al. The cardiac mechanical stretch sensor machinery involves a Z disc complex that is defective in a subset of human dilated cardiomyopathy. *Cell*. 2002;111:943–955.
  45. Knöll R, Kostin S, Klede S, et al. A common MLP (muscle LIM protein) variant is associated with cardiomyopathy. *Circ Res*. 2010;106:695–704. doi: 10.1161/CIRCRESAHA.109.206243.
  46. Boateng SY, Senyo SE, Qi L, Goldspink PH, Russell B. Myocyte remodeling in response to hypertrophic stimuli requires nucleocytoplasmic shuttling of muscle LIM protein. *J Mol Cell Cardiol*. 2009;47:426–435. doi: 10.1016/j.yjmcc.2009.04.006.
  47. Gehmlich K, Ehler E, Perrot A, Fürst DO, Geier C. “MLP: a stress sensor goes nuclear” by Sylvia Gunkel, Jörg Heineke, Denise Hilfiker-Kleiner, Ralph Knöll, *J Mol Cell Cardiol*. 2009;47(4):423–5. *J Mol Cell Cardiol*. 2010;48:424–425; author reply 426. doi: 10.1016/j.yjmcc.2009.10.021.
  48. Vafiadaki E, Arvanitis DA, Sanoudou D. Muscle LIM Protein: Master regulator of cardiac and skeletal muscle functions. *Gene*. 2015;566:1–7. doi: 10.1016/j.gene.2015.04.077.
  49. Yoshida M, Sho E, Nanjo H, Takahashi M, Kobayashi M, Kawamura K, Honma M, Komatsu M, Sugita A, Yamauchi M, Hosoi T, Ito Y, Masuda H. Weaving hypothesis of cardiomyocyte sarcomeres: discovery of periodic broadening and narrowing of intercalated disk during volume-load change. *Am J Pathol*. 2010;176:660–678. doi: 10.2353/ajpath.2010.090348.
  50. Wilson AJ, Schoenauer R, Ehler E, Agarkova I, Bennett PM. Cardiomyocyte growth and sarcomerogenesis at the intercalated disc. *Cell Mol Life Sci*. 2014;71:165–181. doi: 10.1007/s00018-013-1374-5.
  51. Kaushik G, Spenlehauer A, Sessions AO, et al. Vinculin network-mediated cytoskeletal remodeling regulates contractile function in the aging heart. *Sci Transl Med*. 2015;7:292ra99. doi: 10.1126/scitranslmed.aaa5843.
  52. Bonda TA, Sznajka B, Sokołowska M, Dziemiłowicz M, Winnicka MM, Chyżewski L, Kamiński KA. Remodeling of the intercalated disc related to aging in the mouse heart [published online ahead of print November 13, 2015]. *J Cardiol*. doi: 10.1016/j.jcc.2015.10.001. [http://www.journal-of-cardiology.com/article/S0914-5087\(15\)00308-1/abstract](http://www.journal-of-cardiology.com/article/S0914-5087(15)00308-1/abstract).

53. Heling A, Zimmermann R, Kostin S, Maeno Y, Hein S, Devaux B, Bauer E, Klöveborn WP, Schlepper M, Schaper W, Schaper J. Increased expression of cytoskeletal, linkage, and extracellular proteins in failing human myocardium. *Circ Res*. 2000;86:846–853.
54. Michaelson JE, Huang H. Cell-cell junctional proteins in cardiovascular mechanotransduction. *Ann Biomed Eng*. 2012;40:568–577. doi: 10.1007/s10439-011-0439-6.
55. Kostetskii I, Li J, Xiong Y, Zhou R, Ferrari VA, Patel VV, Molkentin JD, Radice GL. Induced deletion of the N-cadherin gene in the heart leads to dissolution of the intercalated disc structure. *Circ Res*. 2005;96:346–354. doi: 10.1161/01.RES.0000156274.72390.2c.
56. Marchand A, Atassi F, Gaaya A, LePrince P, Le Feuvre C, Soubrier F, Lompré AM, Nadaud S. The Wnt/beta-catenin pathway is activated during advanced arterial aging in humans. *Aging Cell*. 2011;10:220–232. doi: 10.1111/j.1474-9726.2010.00661.x.
57. Behrens J, von Kries JP, Kühl M, Bruhn L, Wedlich D, Grosschedl R, Birchmeier W. Functional interaction of beta-catenin with the transcription factor Lef-1. *Nature*. 1996;382:638–642. doi: 10.1038/382638a0.
58. Casad ME, Yu L, Daniels JP, Wolf MJ, Rockman HA. Deletion of Siah-interacting protein gene in *Drosophila* causes cardiomyopathy. *Mol Genet Genomics*. 2012;287:351–360. doi: 10.1007/s00438-012-0684-x.
59. Yonemura S, Wada Y, Watanabe T, Nagafuchi A, Shibata M. alpha-Catenin as a tension transducer that induces adherens junction development. *Nat Cell Biol*. 2010;12:533–542. doi: 10.1038/ncb2055.
60. Thomas WA, Boscher C, Chu YS, Cuvelier D, Martinez-Rico C, Seddiki R, Heysch J, Ladoux B, Thiery JP, Mege RM, Dufour S. alpha-Catenin and vinculin cooperate to promote high E-cadherin-based adhesion strength. *J Biol Chem*. 2013;288:4957–4969. doi: 10.1074/jbc.M112.403774.
61. Desai R, Sarpal R, Ishiyama N, Pellikka M, Ikura M, Tepass U. Monomeric alpha-catenin links cadherin to the actin cytoskeleton. *Nat Cell Biol*. 2013;15:261–273. doi: 10.1038/ncb2685.
62. Sheikh F, Chen Y, Chen Y, Liang X, Hirschy A, Stenbit AE, Gu Y, Dalton ND, Yajima T, Lu Y, Knowlton KU, Peterson KL, Perriard JC, Chen J. alpha-E-catenin inactivation disrupts the cardiomyocyte adherens junction, resulting in cardiomyopathy and susceptibility to wall rupture. *Circulation*. 2006;114:1046–1055. doi: 10.1161/CIRCULATIONAHA.106.634469.
63. Hein S, Kostin S, Heling A, Maeno Y, Schaper J. The role of the cytoskeleton in heart failure. *Cardiovasc Res*. 2000;45:273–278.
64. Birks EJ. Molecular changes after left ventricular assist device support for heart failure. *Circ Res*. 2013;113:777–791. doi: 10.1161/CIRCRESAHA.113.301413.
65. Birks EJ, Hall JL, Barton PJ, Grindle S, Latif N, Hardy JP, Rider JE, Banner NR, Khaghani A, Miller LW, Yacoub MH. Gene profiling changes in cytoskeletal proteins during clinical recovery after left ventricular-assist device support. *Circulation*. 2005;112:157–164. doi: 10.1161/CIRCULATIONAHA.104.526137.
66. Olson TM, Illenberger S, Kishimoto NY, Huttelmaier S, Keating MT, Jockusch BM. Metavinculin mutations alter actin interaction in dilated cardiomyopathy. *Circulation*. 2002;105:431–437.
67. Vasile VC, Will ML, Ommen SR, Edwards WD, Olson TM, Ackerman MJ. Identification of a metavinculin missense mutation, R975W, associated with both hypertrophic and dilated cardiomyopathy. *Mol Genet Metab*. 2006;87:169–174. doi: 10.1016/j.ymgme.2005.08.006.
68. Loffredo FS, Nikolova AP, Pancoast JR, Lee RT. Heart failure with preserved ejection fraction: molecular pathways of the aging myocardium. *Circ Res*. 2014;115:97–107. doi: 10.1161/CIRCRESAHA.115.302929.
69. Lieber SC, Aubry N, Pain J, Diaz G, Kim SJ, Vatner SF. Aging increases stiffness of cardiac myocytes measured by atomic force microscopy nanoindentation. *Am J Physiol Heart Circ Physiol*. 2004;287:H645–H651. doi: 10.1152/ajpheart.00564.2003.
70. Hein S, Gaasch WH, Schaper J. Giant molecule titin and myocardial stiffness. *Circulation*. 2002;106:1302–1304.
71. Engler AJ, Carag-Krieger C, Johnson CP, Raab M, Tang HY, Speicher DW, Sanger JW, Sanger JM, Discher DE. Embryonic cardiomyocytes beat best on a matrix with heart-like elasticity: scar-like rigidity inhibits beating. *J Cell Sci*. 2008;121:3794–3802. doi: 10.1242/jcs.029678.
72. Simpson DG, Terracio L, Terracio M, Price RL, Turner DC, Borg TK. Modulation of cardiac myocyte phenotype in vitro by the composition and orientation of the extracellular matrix. *J Cell Physiol*. 1994;161:89–105. doi: 10.1002/jcp.1041610112.
73. Nishimura M, Kumsta C, Kaushik G, Diop SB, Ding Y, Bisharat-Kernizan J, Catan H, Cammarato A, Ross RS, Engler AJ, Bodmer R, Hansen M, Ocorr K. A dual role for integrin-linked kinase and beta1-integrin in modulating cardiac aging. *Aging Cell*. 2014;13:431–440. doi: 10.1111/acel.12193.
74. Shai SY, Harpf AE, Babbitt CJ, Jordan MC, Fishbein MC, Chen J, Omura M, Leil TA, Becker KD, Jiang M, Smith DJ, Cherry SR, Loftus JC, Ross RS. Cardiac myocyte-specific excision of the beta1 integrin gene results in myocardial fibrosis and cardiac failure. *Circ Res*. 2002;90:458–464.
75. Horn MA. Cardiac Physiology of aging: extracellular considerations. *Compr Physiol*. 2015;5:1069–1121. doi: 10.1002/cphy.c140063.
76. Horn MA, Trafford AW. Aging and the cardiac collagen matrix: novel mediators of fibrotic remodelling [published online ahead of print November 11, 2015]. *J Mol Cell Cardiol*. doi: 10.1016/j.yjmcc.2015.11.005. http://www.jmcc-online.com/article/S0022-2828(15)30110-3/abstract.
77. Leask A. Getting to the heart of the matter: new insights into cardiac fibrosis. *Circ Res*. 2015;116:1269–1276. doi: 10.1161/CIRCRESAHA.116.305381.
78. Danowski BA, Imanaka-Yoshida K, Sanger JM, Sanger JW. Costameres are sites of force transmission to the substratum in adult rat cardiomyocytes. *J Cell Biol*. 1992;118:1411–1420.
79. Samarel AM. Costameres, focal adhesions, and cardiomyocyte mechanotransduction. *Am J Physiol Heart Circ Physiol*. 2005;289:H2291–H2301. doi: 10.1152/ajpheart.00749.2005.
80. Anastasi G, Cutroneo G, Gaeta R, Di Mauro D, Arco A, Consolo A, Santoro G, Trimarchi F, Favalaro A. Dystrophin-glycoprotein complex and vinculin-talin-integrin system in human adult cardiac muscle. *Int J Mol Med*. 2009;23:149–159.
81. Pardo JV, Siliciano JD, Craig SW. Vinculin is a component of an extensive network of myofibril-sarcolemma attachment regions in cardiac muscle fibers. *J Cell Biol*. 1983;97:1081–1088.
82. Tangney JR, Chuang JS, Janssen MS, Krishnamurthy A, Liao P, Hoshijima M, Wu X, Meininger GA, Muthuchamy M, Zemljic-Harpe A, Ross RS, Frank LR, McCulloch AD, Omens JH. Novel role for vinculin in ventricular myocyte mechanics and dysfunction. *Biophys J*. 2013;104:1623–1633. doi: 10.1016/j.bpj.2013.02.021.
83. Israeli-Rosenberg S, Manso AM, Okada H, Ross RS. Integrins and integrin-associated proteins in the cardiac myocyte. *Circ Res*. 2014;114:572–586. doi: 10.1161/CIRCRESAHA.114.301275.
84. Brown NH, Gregory SL, Rickoll WL, Fessler LI, Prout M, White RA, Fristrom JW. Talin is essential for integrin function in *Drosophila*. *Dev Cell*. 2002;3:569–579.
85. Critchley DR. Biochemical and structural properties of the integrin-associated cytoskeletal protein talin. *Annu Rev Biophys*. 2009;38:235–254. doi: 10.1146/annurev.biophys.050708.133744.
86. Monkley SJ, Pritchard CA, Critchley DR. Analysis of the mammalian talin2 gene TLN2. *Biochem Biophys Res Commun*. 2001;286:880–885. doi: 10.1006/bbrc.2001.5497.
87. Manso AM, Li R, Monkley SJ, Cruz NM, Ong S, Lao DH, Koshman YE, Gu Y, Peterson KL, Chen J, Abel ED, Samarel AM, Critchley DR, Ross RS. Talin1 has unique expression versus talin 2 in the heart and modifies the hypertrophic response to pressure overload. *J Biol Chem*. 2013;288:4252–4264. doi: 10.1074/jbc.M112.427484.
88. Padmanabhan Iyer R, Chiao YA, Flynn ER, Hakala K, Cates CA, Weintraub ST, de Castro Brás LE. Matrix metalloproteinase-9-dependent mechanisms of reduced contractility and increased stiffness in the aging heart. *Proteomics Clin Appl*. 2016;10:92–107. doi: 10.1002/prca.201500038.88.
89. Imanaka-Yoshida K, Enomoto-Iwamoto M, Yoshida T, Sakakura T. Vinculin, Talin, Integrin alpha6beta1 and laminin can serve as components of attachment complex mediating contraction force transmission from cardiomyocytes to extracellular matrix. *Cell Motil Cytoskeleton*. 1999;42:1–11. doi: 10.1002/(SICI)1097-0169(1999)42:1<1::AID-CM1>3.0.CO;2-0.
90. Schwartz MA. Transmembrane signalling by integrins. *Trends Cell Biol*. 1992;2:304–308.
91. Carter WG, Wayner EA, Bouchard TS, Kaur P. The role of integrins alpha 2 beta 1 and alpha 3 beta 1 in cell-cell and cell-substrate adhesion of human epidermal cells. *J Cell Biol*. 1990;110:1387–1404.
92. Brancaccio M, Cabodi S, Belkin AM, Collo G, Koteliensky VE, Tomatis D, Altruda F, Silengo L, Tarone G. Differential onset of expression of alpha 7 and beta 1D integrins during mouse heart and skeletal muscle development. *Cell Adhes Commun*. 1998;5:193–205.
93. Johnston RK, Balasubramanian S, Kasiganesan H, Baicu CF, Zile MR, Kuppaswamy D. Beta3 integrin-mediated ubiquitination activates survival signaling during myocardial hypertrophy. *FASEB J*. 2009;23:2759–2771. doi: 10.1096/fj.08-127480.
94. Nawata J, Ohno I, Isoyama S, Suzuki J, Miura S, Ikeda J, Shirato K. Differential expression of alpha 1, alpha 3 and alpha 5 integrin subunits in acute and chronic stages of myocardial infarction in rats. *Cardiovasc Res*. 1999;43:371–381.

95. Burgess ML, McCreia JC, Hedrick HL. Age-associated changes in cardiac matrix and integrins. *Mech Ageing Dev.* 2001;122:1739–1756.
96. Hansen M, Hsu AL, Dillin A, Kenyon C. New genes tied to endocrine, metabolic, and dietary regulation of lifespan from a *Caenorhabditis elegans* genomic RNAi screen. *PLoS Genet.* 2005;1:119–128. doi: 10.1371/journal.pgen.0010017.
97. Kumsta C, Ching TT, Nishimura M, Davis AE, Gelino S, Catan HH, Yu X, Chu CC, Ong B, Panowski SH, Baird N, Bodmer R, Hsu AL, Hansen M. Integrin-linked kinase modulates longevity and thermotolerance in *C. elegans* through neuronal control of HSF-1. *Ageing Cell.* 2014;13:419–430. doi: 10.1111/accel.12189.
98. Yurchenco PD, Amenta PS, Patton BL. Basement membrane assembly, stability and activities observed through a developmental lens. *Matrix Biol.* 2004;22:521–538. doi: 10.1016/j.matbio.2003.10.006.
99. Polyakova V, Loeffler I, Hein S, Miyagawa S, Piotrowska I, Dammer S, Risteli J, Schaper J, Kostin S. Fibrosis in endstage human heart failure: severe changes in collagen metabolism and MMP/TIMP profiles. *Int J Cardiol.* 2011;151:18–33. doi: 10.1016/j.ijcard.2010.04.053.
100. Huet E, Gabison E, Vallee B, Mougenot N, Linguet G, Riou B, Jarosz C, Menashi S, Besse S. Deletion of extracellular matrix metalloproteinase inducer/CD147 induces altered cardiac extracellular matrix remodeling in aging mice. *J Physiol Pharmacol.* 2015;66:355–366.
101. Dworatzek E, Baczek I, Kararigas G. Effects of aging on cardiac extracellular matrix in men and women. *Proteomics Clin Appl.* 2016;10:84–91. doi: 10.1002/prca.201500031.
102. Horn MA, Graham HK, Richards MA, Clarke JD, Greensmith DJ, Briston SJ, Hall MC, Dibb KM, Trafford AW. Age-related divergent remodeling of the cardiac extracellular matrix in heart failure: collagen accumulation in the young and loss in the aged. *J Mol Cell Cardiol.* 2012;53:82–90. doi: 10.1016/j.yjmcc.2012.03.011.
103. Querejeta R, López B, González A, Sánchez E, Larman M, Martínez Ubago JL, Díez J. Increased collagen type I synthesis in patients with heart failure of hypertensive origin: relation to myocardial fibrosis. *Circulation.* 2004;110:1263–1268. doi: 10.1161/01.CIR.0000140973.60992.9A.
104. Pauschinger M, Knopf D, Petschauer S, Doerner A, Poller W, Schwimmbeck PL, Kühl U, Schultheiss HP. Dilated cardiomyopathy is associated with significant changes in collagen type I/III ratio. *Circulation.* 1999;99:2750–2756.
105. Medugorac I. Collagen type distribution in the mammalian left ventricle during growth and aging. *Res Exp Med (Berl).* 1982;180:255–262.
106. Jugdutt BI. Remodeling of the myocardium and potential targets in the collagen degradation and synthesis pathways. *Curr Drug Targets Cardiovasc Haematol Disord.* 2003;3:1–30.
107. Thomas DP, McCormick RJ, Zimmerman SD, Vadlamudi RK, Gosselin LE. Aging- and training-induced alterations in collagen characteristics of rat left ventricle and papillary muscle. *Am J Physiol.* 1992;263:H778–H783.
108. Thomas DP, Cotter TA, Li X, McCormick RJ, Gosselin LE. Exercise training attenuates aging-associated increases in collagen and collagen crosslinking of the left but not the right ventricle in the rat. *Eur J Appl Physiol.* 2001;85:164–169.
109. Campbell DJ, Somaratne JB, Jenkins AJ, Prior DL, Yui M, Kenny JF, Newcomb AE, Schalkwijk CG, Black MJ, Kelly DJ. Diastolic dysfunction of aging is independent of myocardial structure but associated with plasma advanced glycation end-product levels. *PLoS One.* 2012;7:e49813. doi: 10.1371/journal.pone.0049813.
110. Asif M, Egan J, Vasani S, Jyothirmayi GN, Masurekar MR, Lopez S, Williams C, Torres RL, Wagle D, Ulrich P, Cerami A, Brines M, Regan TJ. An advanced glycation endproduct cross-link breaker can reverse age-related increases in myocardial stiffness. *Proc Natl Acad Sci U S A.* 2000;97:2809–2813. doi: 10.1073/pnas.040558497.
111. Konstantin MH, Völkers M, Collins B, Quijada P, Quintana M, De La Torre A, Ormachea L, Din S, Gude N, Toko H, Sussman MA. Fibronectin contributes to pathological cardiac hypertrophy but not physiological growth. *Basic Res Cardiol.* 2013;108:375. doi: 10.1007/s00395-013-0375-8.
112. Oliviero P, Chassagne C, Salichon N, Corbier A, Hamon G, Marotte F, Charlemagne D, Rappaport L, Samuel JL. Expression of laminin alpha2 chain during normal and pathological growth of myocardium in rat and human. *Cardiovasc Res.* 2000;46:346–355.
113. de Castro Brás LE, Toba H, Baicu CF, Zile MR, Weintraub ST, Lindsey ML, Bradshaw AD. Age and SPARC change the extracellular matrix composition of the left ventricle. *Biomed Res Int.* 2014;2014:810562. doi: 10.1155/2014/810562.
114. Bonnema DD, Webb CS, Pennington WR, Stroud RE, Leonardi AE, Clark LL, McClure CD, Finklea L, Spinale FG, Zile MR. Effects of age on plasma matrix metalloproteinases (MMPs) and tissue inhibitor of metalloproteinases (TIMPs). *J Card Fail.* 2007;13:530–540. doi: 10.1016/j.cardfail.2007.04.010.
115. Robert V, Besse S, Sabri A, Silvestre JS, Assayag P, Nguyen VT, Swynghedauw B, Delcayre C. Differential regulation of matrix metalloproteinases associated with aging and hypertension in the rat heart. *Lab Invest.* 1997;76:729–738.
116. Matsusaka H, Ide T, Matsushima S, Ikeuchi M, Kubota T, Sunagawa K, Kinugawa S, Tsutsui H. Targeted deletion of matrix metalloproteinase 2 ameliorates myocardial remodeling in mice with chronic pressure overload. *Hypertension.* 2006;47:711–717. doi: 10.1161/01.HYP.0000208840.30778.00.
117. Polyakova V, Hein S, Kostin S, Ziegelhoeffer T, Schaper J. Matrix metalloproteinases and their tissue inhibitors in pressure-overloaded human myocardium during heart failure progression. *J Am Coll Cardiol.* 2004;44:1609–1618. doi: 10.1016/j.jacc.2004.07.023.
118. Ruddy JM, Jones JA, Stroud RE, Mukherjee R, Spinale FG, Ikonomidis JS. Differential effects of mechanical and biological stimuli on matrix metalloproteinase promoter activation in the thoracic aorta. *Circulation.* 2009;120:S262–S268. doi: 10.1161/CIRCULATIONAHA.108.843581.
119. Zile MR, Baicu CF, Stroud RE, Van Laer A, Arroyo J, Mukherjee R, Jones JA, Spinale FG. Pressure overload-dependent membrane type 1-matrix metalloproteinase induction: relationship to LV remodeling and fibrosis. *Am J Physiol Heart Circ Physiol.* 2012;302:H1429–H1437. doi: 10.1152/ajpheart.00580.2011.
120. Chiao YA, Ramirez TA, Zamilpa R, Okoronkwo SM, Dai Q, Zhang J, Jin YF, Lindsey ML. Matrix metalloproteinase-9 deletion attenuates myocardial fibrosis and diastolic dysfunction in ageing mice. *Cardiovasc Res.* 2012;96:444–455. doi: 10.1093/cvr/cvs275.

# Circulation Research

JOURNAL OF THE AMERICAN HEART ASSOCIATION



## Mechanical Regulation of Cardiac Aging in Model Systems

Ayla O. Sessions and Adam J. Engler

*Circ Res.* 2016;118:1553-1562

doi: 10.1161/CIRCRESAHA.116.307472

*Circulation Research* is published by the American Heart Association, 7272 Greenville Avenue, Dallas, TX 75231

Copyright © 2016 American Heart Association, Inc. All rights reserved.

Print ISSN: 0009-7330. Online ISSN: 1524-4571

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://circres.ahajournals.org/content/118/10/1553>

**Permissions:** Requests for permissions to reproduce figures, tables, or portions of articles originally published in *Circulation Research* can be obtained via RightsLink, a service of the Copyright Clearance Center, not the Editorial Office. Once the online version of the published article for which permission is being requested is located, click Request Permissions in the middle column of the Web page under Services. Further information about this process is available in the [Permissions and Rights Question and Answer](#) document.

**Reprints:** Information about reprints can be found online at:  
<http://www.lww.com/reprints>

**Subscriptions:** Information about subscribing to *Circulation Research* is online at:  
<http://circres.ahajournals.org/subscriptions/>