

UC Irvine

UC Irvine Previously Published Works

Title

Monitoring, classification, and characterization of interior Alaska forests using AIRSAR and ERS-1 SAR

Permalink

<https://escholarship.org/uc/item/5r66046g>

Journal

Polar Record, 31(177)

ISSN

0032-2474

Authors

Williams, CL
McDonald, K
Rignot, E
[et al.](#)

Publication Date

1995-04-01

DOI

10.1017/s0032247400013723

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Monitoring, classification, and characterization of interior Alaska forests using AIRSAR and ERS- 1 SAR

C.L. Williams

Institute of Northern Forestry, Pacific Northwest Research Station, USDA Forest Service, Fairbanks, AK 99775, USA

K. McDonald and E. Rignot

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA

L.A. Viereck

Institute of Northern Forestry, Pacific Northwest Research Station, USDA Forest Service, Fairbanks, AK 99775, USA

J.B. Way and R. Zimmermann

Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA

Received July 1994

ABSTRACT. At the Bonanza Creek Experimental Forest (BCEF), past ecological research has been directed at forest successional processes on the floodplain of the Tanana River and adjacent uplands. Research at the Bonanza Creek site continues on the mosaic of forests, shrublands, and wetlands in a wide variety of successional stages on the Tanana floodplain. This paper reviews research since 1988 into the capabilities of Synthetic Aperture Radar (SAR) for monitoring, classification, and characterization of these forests using radar remote sensing and modelling techniques. Classifications of successional stages, obtained by use of different classifiers on multi-frequency and multi-polarimetric AIRSAR data, are contrasted; these classifications have been used to predict classification accuracies obtained with ERS-1 data, and to estimate the utility of an ERS-1 and RADARSAT combination for classification. Forest classifications, used in combination with ground-truth data for more than 50 forest stands, are used to summarize the distribution of biomass on the landscape. This will allow projections of future biomass. Monitoring of forest phenology, seasonality of flooding, and freeze-thaw transitions is ongoing. Also, direct monitoring of dominant tree species is demonstrating diurnal variation and interrelationships among environmental, physiological, and backscatter measurements.

contents

Introduction	22-1
The study Site	22-)
SAR coverage at Bonanza Creek	228
Use of SAR for landscape interpretation	229
Forest classification	230
Biomass estimation and prediction	231
Forest structure and function	231
Summary and conclusions	232
Acknowledgements	234
References	234

Introduction

Landscapes in interior Alaska are representative of the circumboreal taiga; their forests, subject to widespread periodic disturbance, consist of complex mosaics of different stages in vegetational succession (Van Cleve and Viereck 1981). The use of Synthetic Aperture Radar (SAR) for remote sensing of these forests shows particular potential for ecological research. In the sub-Arctic, variations in annual cycles of freezing and thawing have dramatic consequences for ecosystems; these cycles influence both the biotic and the abiotic processes shaping landscapes. Active microwave sensors, such as SAR, are sensitive to the physical states of water and are also able to penetrate cloud cover and darkness.

Work in progress since 1988 has shown Alaska forests to be important silts foresting the ecological capabilities of SAR, because of the availability of both upland sites and relatively level floodplain sites, the existing detailed knowledge of both upland and floodplain successional

processes, the records of stand history, composition, and geometry for a wide variety of stands, and the existing net work of meteorologic monitoring stations. Interdisciplinary ecological research continues into the future at the Bonanza Creek Long Term Ecological Research (LTER) site,

This paper summarizes ongoing research utilizing SAR in forests in interior Alaska. It includes a description of the study site and the available SAR imagery, examples of insights into landscape interpretation and phenology obtained with SAR, AIRSAR-based classifications of forest types, and predictions for the accuracy of classifications based on satellite-borne SAR. Two methods, one direct and one indirect, for estimating forest biomass are described; one of these allows predictions of future biomass. Finally, it briefly describes backscatter modelling and physiological monitoring that are establishing relationships between radar backscatter, forest structure, environmental change, and tree-canopy physiology.

The study site

The LTER site is located within and adjacent to the Bonanza Creek Experimental Forest (BCEF), 20 km southwest of Fairbanks, Alaska. The LTER site includes both the Tanana River floodplain and the adjacent uplands. The Tanana River is one of three major rivers draining interior Alaska; it is primarily glacial-fed, with high suspended sediment loads, and with consequent braiding and meandering along its length. The meandering of the river has created a floodplain landscape of exposed silt bars, islands, terraces, and meander scrolls, with an associated mosaic of




Fig. 1. Location of upland and floodplain forest stands and of weather stations at the Bonanza Creek LTER site. Stand types for floodplain LTER stands are FP1: willow; f P2: alder, FP3: balsam poplar; FP4: white spruce; and FP5: black spruce. The polygons represent stands where the geometry and composition are being measured specifically for use in remote sensing studies and represent areas from where radar backscatter has been extracted for classification analyses. Stand types for these stands are AL: alder; BP: balsam poplar; WS: white spruce; ES: black spruce; and CC: clearcut.

successional vegetation (Van Cleve and others 1993). Floodplain vegetation represents a primary successional sequence starting on newly formed silt bars near the river's edge. Upland vegetation is characterized by a secondary successional sequence initiated by wildfire. The vegetation and landforms of the Bonanza Creek LTER site reflect the major disturbance processes—wild fires, insect infestations, and river geomorphic processes governing the development of boreal forests. Ecological research at BCFE during the last 30 years has resulted in development of detailed successional models and availability of long-term data for vegetation, soils, climate, and river morphology (Van Cleve and others 1991; Viereck and others 1993b; 1993c).

The LTER site includes 24 intensively studied upland and floodplain successional forest stands, with 10 weather stations distributed among the stands. These 24 stands represent eight distinct successional stages of the predominant upland and floodplain successional sequences. In 23 additional successional stands, detailed measurements of forest composition and forest geometry have been made for use in remote sensing studies (Fig. 1). Vegetational

succession is generally similar to that found along all of the major rivers in interior Alaska and adjacent Canada. The general successional pattern begins on exposed silt bars, first colonized by willow species (*Salix* spp.), with subsequent generations dominated by alders (*Alnus tenuifolia*), balsam poplars (*Populus balsamifera*), and white spruce (*Picea glauca*). White spruce stands may persist for several generations before development of permafrost and replacement by black spruce (*Picea mariana*). Twelve stages of succession are recognized. Replacement of white spruce by black spruce and succession between black spruce and open bog are less well understood than earlier stages in succession (Viereck and others 1993b).

SAR coverage at Bonanza Creek

Image data from two currently available SAR sensors are being utilized in studies of interior Alaskan forests. Images from aircraft-borne SAR (AIRSAR) have been collected for three seasons, winter (March), spring (May), and summer (July), with March images taken at temperatures both well above and below 0°C and with variable amounts of river flooding in May. AIRSAR data include backscatter




Fig. 2. Winter and Spring L-band AIRSAR images of Bonanza Creek Experimental Forest. Color bar shows total backscatter, expressed in decibels. The two March images show the dramatic decrease in backscatter as landscapes freeze. The two May images show the influence of ice jams (A) on flooding of the Tanana River floodplain during breakup. Flooding is seen both as increases in the surface area of open water wetlands (B) with low backscatter and as flooding of wooded areas (C), which exhibit high backscatter due to double-bounce backscattering mechanisms. Backscatter variation useful in forest classification and biomass mapping can be seen by comparisons of these images with stand locations in Figure 1.

for three bands (C, L, and P), with four polarizations (HH, HV, VH, VV) for each band (Way and Smith 1991). The multi-frequency, multi-polarization nature of AIRSAR data allows differentiation of radar-scattering mechanisms under different canopy geometries (Van Zyl 1989), assists with backscatter modelling (McDonald 1991), and allows identification of optimal bands and polarizations for different ecological or engineering purposes (Williams and others 1992, 1993; Rignot and others 1994 b). Because of its multi-channel capabilities, AIRSAR may be used to simulate the specific band and polarization combinations of future spaceborne radar configurations (Williams and others 1992, 1993).

The ERS-1 SAR, with its C-band, VV polarization configuration, provides only one observation channel with each sensor pass but provides frequent repeat coverage for examination of seasonal change (Way and Smith 1991).

Use of SAR for landscape interpretation

AIRSAR images from March 1988 (Fig. 2) at Bonanza Creek and ERS-1 images from August-September 1991

further down the Tanana River (Rignot and Way 1994; Rignot and others 1994a) show the dramatic drop in backscatter that occurs as landscapes freeze.

Both ERS-1 and AIRSAR images of the Tanana Valley display ecologically significant features of floodplain geomorphology, including oxbow lakes, meander scrolls, and abandoned channels. In addition, use of SAR to delineate patterns of flooding in May 1991 demonstrates differences between northern and southern portions of the floodplain (Fig. 2). Flooding on 4 May 1991 was caused during river breakup by the formation of ice jams across the main river channels; ice jams had dispersed, and flooding subsided by 7 May. Ice jams are clearly evident as areas of increased backscatter in channels of open water with very low backscatter; this demonstrates the use of SAR for monitoring river freeze and breakup. Flooding is seen both as increases in surface area of lakes and wetlands containing open water (low backscatter) and as dramatic increases in backscatter caused by the presence of standing water in forests. Flooding upstream (to the left) of the ice




Fig. 3. Separation of six vegetation types of the Tanana River floodplain, Bonanza Creek LTER site. Discriminant analysis was applied to C-band AIRSAR backscatter, full polarization,

jam is most dramatic on the south side of the floodplain (top of figure); the north side of the floodplain is narrower and may show a steeper gradient as the river is being pushed north against adjacent uplands by the continued rise of the Alaska Range. Although in a year without ice jam formation, seasonal flooding typically affects treeless seasonal and permanent wetlands, most of the flooding caused here by ice jam formation is that of forested areas.

ERS-1 transects across Alaska provide examples of larger scale environmental monitoring with SAR (Rignot and Way 1994). Such transects show latitudinal differences across Alaska in freeze-thaw cycles, variation among years, and variation in abruptness of transitions between frozen and thawed states. Spring snowmelt is identifiable, as distinctions among landscape components become blurred and images show reduced contrast. Likewise, wet snowfalls in early winter led to image contrast.

Forest classification

Land cover of the world's boreal forests is composed of complex vegetation mosaics subject to periodic disturbance by wildfire, insect infestation, and riverine processes. Long-term monitoring of landscape change requires remote sensing of this mosaic of vegetation types and successional stages.

Discriminant analysis (Fig. 3) demonstrates the separability of successional stages in floodplain forests in interior Alaska using C-band full-polarimetric AIRSAR data. Comparisons of single-channel classifications are possible with subsets of the AIRSAR data; C-band, HV-polarization consistently provides the best classifications (Table 1). Based on these classifications, none of the

single-channel satellite-borne SARs was expected to classify forest types as accurately as sensors with cross-polarization. This prediction was tested for ERS-1 data. From AIRSAR data, classifications based on C-band, VV-polarization average 36% error; classifications based on ERS-1 data showed comparable accuracy: 46% on 25 April 1992 and 39% on 30 May 1992. Combining data from these two dates did not improve classification accuracy significantly.

A striking improvement in classification accuracy was obtained by combining AIRSAR C-band VV and HH polarization data with L-band HH polarization data. For this reason forest classifications based on images from ERS-1, JERS-1, and RADARSAT were expected to be 82% accurate, well within accuracy levels obtained with optical sensors (Williams and others 1993).

Similar classification accuracies have also been obtained on a pixel to pixel basis with a maximum *a posteriori* Bayesian classifier (Rignot and others 1994 b). With this classifier a combination of C-band HV and L-band HV yields the highest overall classification accuracy, with

Table 1. Error rates for single-channel classifications of vegetation types of the Tanana River floodplain. Discriminant analysis is applied to AIRSAR data both for classification and for accuracy assessment. Error rates are expressed as percentages of total numbers of stands.

Band/polarization	May 4	May 6	May 7	Mean
C/HV (consistently best)	18	11	13	14
C/VV (as in ERS-1)	26	50	33	36
C/HH (as in RADARSAT)	29	28	28	28
L/HH (as in JERS-1)	38	25	38	34




Fig. 4. DECORANA of tree and shrub vegetation for 120 Tanana River floodplain stands. Stands are represented by identification numbers. On each figure, distances between stands represent vegetational dissimilarity. Overlays for the top graph are *Populus balsamifera*; overlays for the bottom graph are *Picea glauca*. For each overlay, circle sizes represent percent cover in each stand for the chosen species.

10% error rates in the test stands (Table 2).

In addition, AIRSAR-based classifications showed potential as a means of identifying outlier or unusual forest stands. In Figure 4, detrended correspondence analysis (Hill 1979) of tree and shrub vegetation for 120 stands demonstrated that one stand (847), classified by discriminant analysis on different dates as either a balsam poplar or a white spruce stand, was in fact an intermediate stand with significant cover of both white spruce and balsam poplar.

Biomass estimation and prediction

Monitoring of biomass in boreal forests requires estimation

of biomass levels exceeding 200 tonne ha⁻¹; biomass distributions are highly correlated with the distribution of successional stages on the landscape (Viereck and others 1993.1). AIRSAR imagery has been used to estimate and predict biomass in two ways. Direct estimation of biomass by inversion of regression curves relating backscatter to biomass for different available channels has proven especially useful at low biomass levels (Fig. 5; Rignot and others 1994c). Indirect estimation of biomass is underway using a model (Fig. 6) designed to incorporate existing

long-term data and ecological insight for the floodplain. This model combines classification of land cover with

Table 2. Confusion matrix resulting from vegetation classification with a maximum a posteriori Bayesian classifier (Rignot and others 1994 b). Percentages on the diagonal represent stands correctly classified into each vegetation type. Column labelled 'From' designates identifications from ground-truth data. Overall error rate is 10%.

From\to:	CC	AL	BP	WS	BS	Water
Clearcut	78	0	0	0	2	0
Alder	0	68	3	1	1	0
Balsam poplar	0	31	89	6	0	0
White spruce	0	1	7	91	4	0
Black spruce	2?	0	0	2	93	0
Water	0	0	0	0	0	100

existing successional models, long-term photographic and field records, and known stand variations in biomass and productivity. Changes in the vegetation mosaic and in landscape-scale biomass can be projected into the past or the future (Fig. 7; Williams and others 1994).

Forest structure and function

Although landscape interpretation,





Fig. 5. Predicted biomass levels of forest stands from the Bonanza Creek Experimental Forest. Predictions were obtained from the following regressions relating logarithms of stand biomass to backscatter in decibels, where shv is backscatter of HV polarization, shh is backscatter of HH polarization, and svv is backscatter of VV polarization:
 L-band HV: $\ln \text{biomass} = 9.204 + 0.200 \text{ shv} - 0.011 \text{ shv}^2$
 P-band HV: $\ln \text{biomass} = -3.372 - 1.408 \text{ shv} - 0.057 \text{ shv}^2$
 L band: $\ln \text{biomass} = 9.839 - 0.601 \text{ shv} - 0.037 \text{ shv}^2 - 0.914 \text{ shh} - 0.070 \text{ shh}^2 + 2.396 \text{ svv} + 0.150 \text{ svv}^2$
 P band: $\ln \text{biomass} = 6.598 - 0.129 \text{ shv} - 0.007 \text{ shv}^2 + 0.370 \text{ shh} + 0.011 \text{ shh}^2 - 0.880 \text{ svv} - 0.050 \text{ svv}^2$

forest classification, and biomass estimation may be effective without understanding the exact relationships between backscatter and structure or function of biological systems. Such understanding will ultimately become important for interpretation or predictions at larger spatial and temporal scales and under less controlled conditions. At Bonanza Creek, these relationships are being investigated with backscatter modelling and with physiological monitoring of dominant tree species. Backscatter modelling with the MIMICS model provides information about the kinds of structural and temporally changing biophysical properties that can be detected with spaceborne and airborne SAR (McDonald 1991). Physiological monitoring of three dominant tree species (*Picea glauca*, *Picea mariana*, *P. opulus balsamifera*) has established linkages between water relations of forest canopies, dielectric properties of vegetation, and radar backscatter (Zimmermann and others 1994). The derived relationships between dielectric constant, xylem flux density, and water potential have potential utility for monitoring of forest health.



Summary and conclusions

This research has demonstrated applications for the use of

Fig. 6. Model for mapping of biomass and productivity in boreal forests of interior Alaska (Williams and others 1994),



YEAR 1



YEAR 100



Fig. 7. Present and projected 100-year biomass distribution for the Tanana River floodplain, Bonanza Creek LTER site. This version of the model described in Figure 6 lacks algorithms for recruitment of early successional stages; photographic records will be used for this purpose. The land classification (Rignot and others 1994b) is from May 1991 AIRSAR imagery obtained during spring breakup; it shows an unusually high proportion of landscape as open water. After flooding subsides, most of the areas not in immediate proximity to the river are low biomass bog or black spruce forest. Biomass categories represent means and standard deviations of existing biomass measurements for the different successional stages.

SAR imagery in monitoring, classifying, and characterizing boreal forests of interim Alaska. Use of SAR imagery has obvious advantages in such regions, where there are long dark periods or frequent cloud cover, and where patterns of freezing and thawing of water have such dramatic ecological consequences. Both AIRSAR and ERS-1 images display geomorphological features and annual phenological cycles; flooding of forested and non-forested areas may also be distinguished, and patterns of snowmelt may reflect topography and vegetation pattern. Application of the MIMICS canopy-scattering model to forests at Bonanza Creek aids in interpretation of observed

temporal and spatial changes; physiological monitoring is in place for the establishment of relationships between radar backscatter and canopy physiology.

Classification of at least five of the successional stages of floodplain forests is effective from AIRSAR imagery with both discriminant analysis and Bayesian classifiers; accuracy rates range between 80 and 90%. The availability of cross-polarization is important to these classifications: C-band HV and L- and C-band HV-polarization were the most useful combinations of sensors. As predicted from subsets of AIRSAR hands and polarizations, C-band VV-polarization data from ERS-1 was not su-

ciently powerful for vegetation classifications, but the future combination of data from ERS-1, JERS-1, and RADARSAT may be.

The successional stages distinguished by classification represent the full range of biomass and productivity in floodplain ecosystems. Landscape-scale mapping of biomass is underway, both directly by use of inversion equations and indirectly by using the strong correlations between successional stage and biomass. The relative precision and accuracy of these contrasting approaches are currently under investigation. Use of successional stage classifications to map biomass is making possible projection of landscape-scale biomass patterns into the past or the future.

While continuing work has demonstrated the usefulness of SAR for a wide variety of ecological uses, SAR continues to be under-utilized in practical applications of these uses. An important next step is the development of funding mechanisms and data acquisition protocols making possible increased use of SAR imagery for ecological monitoring and for land management purposes.

Acknowledgements

This work was carried out at the Institute of Northern Forestry and at the Jet Propulsion Laboratory, California Institute of Technology, in part under contract with the National Aeronautics and Space Administration. The authors thank Andrew W. Balser for technical assistance.

References

- Hill, M.O. 1979. *DECORANA: a FORTRAN program for detrended correspondence analysis and reciprocal averaging*. Ithaca, NY: Cornell University.
- McDonald, K. 1991. Modeling microwave backscatter from tree canopies. Unpublished PhD thesis. University of Michigan.
- Rignot, E., and J.B. Way. 1994. Monitoring freeze-thaw cycles along north-south Alaskan transects using ERS-1 SAR. *Remote Sensing of Environment* 49:131-137.
- Rignot, E., and 8 others. 1994a. Monitoring of environmental conditions in taiga forests using ERS-1 SAR. *Remote Sensing of Environment* 49:145-154.
- Rignot, E., C. Williams, J.B. Way, and L. Viereck. 1994b. Mapping of forest types in Alaskan boreal forests using SAR imagery. *IEEE Transactions on Geoscience and Remote Sensing* 32:1051-1059.
- Rignot, E., C. Williams, J.B. Way, J. Yarie, K. McDonald, and L. Viereck. 1994c. Radar mapping of aboveground biomass in boreal forests of interior Alaska. *IEEE Transactions on Geoscience and Remote Sensing* 32: 1117-1124.
- Van Cleve, K., F. Chapin III, C. Dyrness, and L. Viereck. 1991. Element cycling in taiga forests: state factor control. *Bioscience* 41: 78-88.
- Van Cleve, K., and L. Viereck. 1981. Forest succession in relation to nutrient cycling in the boreal forest of Alaska. In: West, C., H. Shugart, and D. Botkin (editors). *Forest succession: concepts and application*. New York: Springer-Verlag: 185-211.
- Van Cleve, K., L. Viereck, and G. Marion. 1993. Introduction and overview of a study dealing with the role of salt-affected soils in primary succession on the Tanana River floodplain, interior Alaska. *Canadian Journal of Forest Research* 23:879-888.
- Van Zyl, J. 1989. Unsupervised classification of scattering behavior using radar polarimetry data. *IEEE Transactions on Geoscience and Remote Sensing* 27:36-45.
- Viereck, L., C. Dyrness, A. Batten, and K. Wenzlick. 1993a. *The Alaska vegetation classification*. Portland, OR: USDA Forest Service, Pacific Northwest Research Station (General 1 technical Report PNW-GTR-286).
- Viereck, L., C. Dyrness, and M. Foote. 1993b. An overview of the vegetation and soils of the floodplain ecosystems of the Tanana River, interior Alaska. *Canadian Journal of Forest Research* 23:889-898.
- Viereck, L., K. Van Cleve, P. Adams, and R. Schlentner. 1993c. Climate of the Tanana River floodplain near Fairbanks, Alaska. *Canadian Journal of Forest Research* 23: 899-913.
- Way, J., and E. Smith. 1991. The evolution of synthetic aperture radar systems and their progression to the EOS SAR. *IEEE Transactions on Geoscience and Remote Sensing* 29: 962-985.
- Williams, C., K. McDonald, E. Rignot, L. Viereck, and J. Way. 1994. An ecological approach to radar mapping of biomass in interior Alaska boreal forests. In: Stein, T.I. *Surface and atmospheric remote sensing: technologies, data analysis and interpretation: proceedings of IGARSS '94 symposium, 8-12 August, California Institute of Technology, Pasadena, California*. Piscataway, NJ: Institute of Electrical and Electronics Engineers: IV, 1856-1859.
- Williams, C., E. Rignot, J. Way, and K. McDonald. 1993. Use of AIRSAR and ERS-1 SAR for classification of successional stage on the Tanana River floodplain, interior Alaska. *Bulletin of the Ecological Society of America* 74:488.
- Williams, C., L. Viereck, K. McDonald, E. Rignot, and J. Way. 1992. Use of AIRSAR in classification of successional stage along the Tanana River, interior Alaska. In: Dean, K. (editor). *Environments/ change: natural and man-made: proceedings AAAS 43rd Arctic Science Conference*. Fairbanks: Geophysical Institute, University of Alaska,
- Zimmerman, R., K. McDonald, J. Way, and R. Oren. 1994. Microclimate, water potential, transpiration, and bole dielectric constant of conifers and deciduous tree species in the continental boreal ecotone in central Alaska. In: Stein, T. I. *Surface and atmospheric remote sensing: technologies, data analysis and interpretation: proceedings of IGARSS '94 symposium, 8-12 August, California Institute of Technology, Pasadena, California*. Piscataway, NJ: Institute of Electrical and Electronics Engineers: 1, 226-228.

The accuracy of references in the text and in this list is the responsibility of the authors, to whom queries should be addressed.