

Lawrence Berkeley National Laboratory

LBL Publications

Title

The NN ® $\pi\pi$ Amplitude

Permalink

<https://escholarship.org/uc/item/5rt1t5v5>

Author

Wong, David Y

Publication Date

1960-08-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

UNIVERSITY OF
CALIFORNIA

Ernest O. Lawrence
**Radiation
Laboratory**

TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Invited paper for Rochester Conference on
High Energy Physics.

UCRL-9374

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California
Contract No. W-7405-eng-48

THE $NN \rightarrow \pi\pi$ AMPLITUDE

David Y. Wong

August 18, 1960


THE $\bar{N}N \rightarrow \pi\pi$ AMPLITUDE*

David Y. Wong

Lawrence Radiation Laboratory
 University of California
 Berkeley, California


August 18, 1960

In the same spirit as the dispersion approach to the pion-pion and the pion-nucleon problems, I shall now describe the determination of the $\bar{N}N$ to $\pi\pi$ amplitude in terms of known singularities of the function. I should mention that this work is done in collaboration with Dr. Ball of Berkeley. We are mainly concerned with the region where the energy of the nucleon-antinucleon system is not too far from twice the pion mass. This region is of some immediate interest since it is expected to give a substantial contribution to the absorptive parts of the nucleon form factor, the pion-nucleon amplitude and also, the nucleon-nucleon amplitude. Let us denote the square of the nucleon-antinucleon center-of-mass energy by t . For any partial wave of a given angular momentum and spin, the singularities in the t -plane are: The branch cut due to the exchange of a single nucleon starting at $t = 4\mu^2(1 - \mu^2/4m^2)$ the exchange of a nucleon plus a pion starting at $t = 0$, etc. The threshold for the production of two pions is at $t = 4\mu^2$. From this threshold to the production of more than two pions, the $\langle \bar{N}N | \pi\pi \rangle$ amplitude has the same phase as the $\langle \pi\pi | \pi\pi \rangle$ amplitude since $\text{Im}(\bar{N}N | \pi\pi) \simeq (\bar{N}N | \pi\pi)^*(\pi\pi | \pi\pi)$ in this region.



* This work was performed under the auspices of the U.S. Atomic Energy Commission.

Frazer and Fulco¹ were the first to write down partial wave dispersion relations for the $N\bar{N}$ to $\pi\pi$ amplitude. They considered the two p-wave amplitudes in some detail. For convenience, they took the combination



$$\Gamma_1 = \alpha_1(+|+) + \beta_1(+|-)$$

$$\Gamma_2 = \alpha_2(+|+) + \beta_2(+|-)$$

where the symbol $(\pm|\pm)$ denotes the nucleon-antinucleon annihilation amplitude in a given helicity state and α, β are known functions of t . They have shown that Γ_1 contributes to the nucleon charge form factor and Γ_2 contributes to the magnetic moment form factor. Frazer and Fulco took a "one-pole" approximation for the $\pi\pi$ p-wave and determine the phase of the $\langle N\bar{N} | \pi\pi \rangle$ amplitude on such basis. They considered the nucleon pole term exactly and estimated the "rescattering" cut (the exchange of a nucleon plus a pion) by a δ-function approximation for the pion-nucleon (π, π) amplitude. They found it necessary to introduce a cut-off on the left hand cut in order to obtain convergent integrals. It turned out that the magnetic amplitude (Γ_2) is insensitive to the cut off. They then chose the p-wave $\pi\pi$ parameters to fit the magnetic form factor. However, the charge amplitude was quite dependent on the cut off. What I would like to report first today are two modifications of the Frazer-Fulco solution. (1) A "two-pole" approximation is taken for the $\pi\pi$ p-wave with a repulsive outer region plus an attractive inner region as discussed by Chew. (2) The values of Γ_1 and Γ_2 in the neighborhood of $t = 0$ are determined

by the pion-nucleon fixed-momentum transfer dispersion relation in the neighborhood of forward scattering. The left hand cut of Γ is calculated in the same manner as Frazer and Pulco except that we terminate the rescattering cut at $t = -26\mu^2$ (where the all partial wave expansion diverges) and replace all remaining cuts by a pole which is adjusted to give the correct value of Γ in the neighborhood of $t = 0$. Of course, this phenomenological pole can also compensate for part of the inaccuracy in the rescattering cut.

I will now give the value and derivative of Γ_1 and Γ_2 at $t = 0$ as calculated from a one-subtraction pion-nucleon dispersion relation. The subtraction constants are related to pion-nucleon scattering lengths and the dispersion integrals involve pion-nucleon partial cross sections which are expressed in terms of combinations of total cross sections and the $(3, 3)$ amplitude in such a way that the $J = 3/2$ states ($P_{3/2}$ and $D_{3/2}$) are taken into account exactly. We believe that such a combination is more accurate than the $(3, 3)$ amplitude alone since the second resonance is probably in the $J = 3/2$ D-state. The following table is a summary of the value and derivative of the Γ 's and the contributions from various terms:

Barn Term	Scattering Lengths			Dispersion Integral	Total
	S	P	D		
$\Gamma_1(0)$	-0.1055	-0.0339		-0.0012	-0.141
$\Gamma_2(0)$	0.0238	-0.0002	-0.0519	0.0041	-0.0042
$\Gamma_1'(0)$	0.00935	0.00155	0.05800	-0.00186	0.0430
$\Gamma_2'(0)$	-0.00562	-0.00010	-0.00258	? -0.00036	?

The scattering lengths are taken from the analysis of Barnes et al.² and Hamilton and Woolcock.³ It is clear that $\Gamma_1(0)$ is most accurately determined. The uncertainty in $\Gamma_2(0)$ and $\Gamma_1'(0)$ mainly comes from the inaccuracy of the small p-wave scattering lengths. If the small p-waves are ignored, we find $\Gamma_2(0) = 0.0033$ and $\Gamma_1'(0) = 0.0348$. The uncertainty from the dispersion integral is considerably smaller. We have checked this point by a comparison with the corresponding integrals where only the (3, 3) p-wave is kept. Since the d-wave scattering lengths are yet unknown, we can only estimate the order of magnitude of $\Gamma_2'(0)$ from the no-subtraction formula which gives $\Gamma_2'(0) \approx -0.005$. Fortunately, it turns out that the phenomenological pole in the Γ_2 amplitude is very weak and the amplitude on the right is quite insensitive to the position of this pole. Hence we can adjust the position and residue of the pole in Γ_1 to give the normalized value and derivative at $t = 0$, and adjust only the residue of the Γ_2 pole to fit the normalized value leaving the position arbitrary as long as it is beyond ~ -15 .

Now that we have the formalism set up, we can compute the Γ 's and the two-pion contribution to the vector part of the nucleon form factors for any given set of $\pi\pi$ parameters in much the same way as the Frazer-Fulco calculation. A typical set that gives the observed magnetic moment form factor is: $\gamma_1 = 60$, $\gamma_2 = 4$, $A_1 = 0.3$, $A_2 = 0.51$ (in pion units) where we have taken the $\pi\pi$ p-wave amplitude to be

$$\sqrt{\frac{\gamma + \mu^2}{\gamma}} e^{i\theta} \sin \theta = N/D$$

with

$$\nu = \frac{e}{4} - \mu^2 ,$$

$$B = \frac{A_1 \nu_1}{\nu + \nu_1} - \frac{A_2 \nu_2}{\nu + \nu_2} ,$$

$$D = 1 - \nu [A_1 \nu_1 K(\nu_1, -\nu) - A_2 \nu_2 K(\nu_2, -\nu)]$$

and $K(a, b)$ is the kernel defined by Chew and Mandelstan. This set of parameters gives a resonance at $t \sim 14$. The two pion contribution to the charge turns out to be $\sim 20\%$ of the total charge. The smallness of this charge is due to the cancellation of the phenomenological pole and all other terms in the normalized Γ_1 amplitude. The pole is found to be situated at a very high energy region but gives far greater (negative) contribution to the electric charge than the contribution (positive) from the rescattering cut. This leads to our belief that although the Γ_1 amplitude is still quite sensitive to the uncertainty in the normalizations, we have at least obtained a $\Gamma_1(t)$ qualitatively more reliable than the function given by Frazer and Fulco.

I shall now turn to the question of the compatibility of our $\pi\pi$ parameters with the dispersion theory of Chew and Mandelstan.⁴ For any given set of p-wave parameters, we can determine one or more sets of s-wave parameters by using the so called "almost exact" crossing conditions in the neighborhood of the $\pi\pi$ symmetry point. Following Chew and Mandelstan, we have chosen an s-wave solution of the form

$$\sqrt{\frac{v + \mu^2}{v}} e^{ia_I} \sin a_I = \frac{a_I + b_I(v - v_0)(\frac{v_s + v_0}{v_s - v})}{1 - (v - v_0)[a_I K(-v_0, -v) + b_I(v_s + v_0)K(v_s, -v)]}$$

where $I = 0, 2$ and $v_0 = -2/3$. The s-wave pole v_s is related to the p-wave resonance energy by $v_s \approx 2(v_r + 1)$. With the p-wave parameters given above, we have found a set of s-wave parameters which gives a repulsive pole for $I = 0$ and an attractive pole for $I = 2$. The parameters are $a_0 = -4.5$, $b_0 = 4.6$, $a_2 = -1.7$, $b_2 = -2.5$. The consistency of this solution and the "approximate" crossing relations of Chew and Mandelstam remains to be seen. Should we be able to obtain an s-wave $\pi\pi$ solution in this way or in any other way, we can immediately construct the s-wave $\langle \bar{N}N | \pi\pi \rangle$ amplitude in much the same manner as the p-wave problem. The normalization at $t = 0$ plays an even more important role in the s-wave problem. In fact, it serves to suppress the nucleon pole term which is known to give a superfluously large fourth order potential in the nucleon-nucleon scattering problem. In closing, I should mention that it is quite probable that a $\pi\pi$ p-wave resonance will give a substantial contribution to the "medium range" attractive force between two nucleons but it is unlikely that the resonance will ever produce a repulsive core. However, there is still a possibility that a one-subtraction formula including the one- and two-pion exchange terms in the nucleon-nucleon problem may assimilate the effect of a hard core in the physical region.

REFERENCES

1. W. R. Frazer and J. R. Fulco, Phys. Rev. 117, 1603 (1960) and Phys. Rev. 117, 1609 (1960).
2. S. W. Barnes, B. Rose, G. Giacomelli, J. Ring, K. Miyabe and K. Kinsey, Phys. Rev. 117, 226 (1960); S. W. Barnes, W. H. Winet, K. Miyabe and K. Kinsey, Phys. Rev. 117, 233 (1960).
3. J. Hamilton and W. S. Woolcock, Phys. Rev. 118, 291 (1960).
4. G. F. Chew and S. Mandelstam, "Theory of the Low-Energy Pion-Pion Interaction," UCRL-9126, March 1960.