

UC Irvine

UC Irvine Previously Published Works

Title

Age, Sex and Weight Effects on Lactate and Leukocyte Response to Exercise in Children and Adolescents.

Permalink

<https://escholarship.org/uc/item/5sf650tt>

Journal

European journal of sports & exercise science, 6(1)

ISSN

2278-005X

Authors

Chen, Jen Jen
Gendy, Elizabeth
Leu, Sharon
[et al.](#)

Publication Date

2018

Peer reviewed



Published in final edited form as:

Eur J Sports Exerc Sci. 2018 ; 6(1): 1–10.

Age, Sex and Weight Effects on Lactate and Leukocyte Response to Exercise in Children and Adolescents

Jen Jen Chen¹, Elizabeth Gendy², Sharon Leu¹, Frank Zaldivar¹, Pietro Gallasseti¹, Elieze Nussbaum², Dan Cooper¹, and Shlomit Random-Aizik¹

¹Pediatric Exercise and Genomics Research Center, University of California, USA

²Miller Children's Hospital, Long Beach, USA

Abstract

Post-exercise leucocytosis has been found in children and adults in response to exercise. When normalized to the work performed, children demonstrate a lower lactate response to exercise than in adults. This study examines the association between leukocyte and lactate response in children following exercise.

148 healthy children participated in this study. Each subject performed a ramp exercise on a cycle ergometer followed by ten 2-minute bouts of constant work rate separated by 1-minute rest intervals. Lactate, leukocyte and leukocyte subtype levels were taken pre- and post-exercise.

Older children showed a significantly higher pre-to post-exercise leukocyte compared to younger children (3599 ± 165 cells/mL, 2544 ± 163 , $p < 0.0001$). Compared to older children, younger children demonstrated a smaller median fold change in lactate (2.7 (1.4-8.1), 4.1 (1.7-29.6)) and leukocyte levels (1.4 (1.0-2.1), 1.6 (1.2-2.8)) after exercise, with a larger leukocyte to lactate fold change ratio.

Conclusion: Younger children have a greater leukocyte to lactate fold change ratio compared to older children. This finding may be due to the lower anaerobic dependence that is found in younger children.

Keywords

Exercise; Paediatrics; Inflammation; Leukocyte; Lactate

INTRODUCTION

It is becoming increasingly recognized that even brief exercise can substantially influence the distribution of white blood cells within the circulation and marginal pools [1,2]. The response of immune cells to physical activity is of growing interest because some key health effects of exercise, both beneficial and detrimental, may well be mediated through the activity of circulating leukocytes [3].

Growth in humans is a dynamic process regulated through the interplay of anabolic and catabolic hormones and cytokines. What is particularly intriguing is that many of these mediators in children, such as growth mediators and their antagonist (often inflammatory mediators), also appear to play substantial roles in the adaptive responses to physical activity

[4,5]. A potential link between inflammatory mediators and exercise may be related to the availability of oxygen to the tissues. When normalized to the amount of work performed, muscle oxygen uptake is greater and blood lactate concentration is lower in children compared to adults. Lactate level in response to exercise reflects the degree of tissue stress, predominantly related to insufficient oxygen transport [6].

The adaptive effects of exercise-induced inflammation have been well studied in adults but less so in children. There has been scant data on the inflammatory response to exercise in children in the literature. These previous studies are limited by small sample sizes and evaluate either sex or age differences alone. Our larger study evaluates relationships between leucocytosis and lactate concentrations following exercise characterized by BMI, age and sex-related differences. We hypothesized that following exercise, the lower lactate levels in younger children would be associated with similarly lower leukocyte levels as compared to older children, and that both responses would be modulated by sex and BMI.

MATERIALS AND METHODS

Participants

148 healthy children ages eight to eighteen years old participated in this study. Table 1 illustrates the anthropometric and physiologic characteristics of the participants. Individuals engaged in competitive sports or with history of chronic medical conditions or use of any medications were excluded from participation. The study was approved by the Institutional Review Board at the University of California, Irvine and written informed assent and consent was obtained from all participants and their parents upon enrolment.

Anthropometric measurements

Standard calibrated scales and stadiometers were used to determine height, body mass and body mass index ($BMI=wt/ht^2$). We also calculated BMI percentile for each child using the published standards from the Centers for Disease Control, National Center for Health Statistics [7].

Pubertal status

Pubertal status was evaluated using a widely used validated self-administered questionnaire [8].

Measurement of fitness

Each participant performed a ramp-type exercise protocol on a cycle ergometer (Ergoline 800S, Yorba Linda, CA) using the Sensor Medics metabolic system. Participants were vigorously encouraged during the high-intensity phases of the exercise protocol. Gas exchange was measured breath-by-breath and the anaerobic (lactate) threshold and peak $\dot{V}O_2$ were calculated using standard methods [9].

At least 48 h, but not exceeding 7 days following the completion of the ramp test; each participant performed an exercise challenge consisting of ten, 2 min bouts, of constant work rate cycle ergometry separated by 1 min rest intervals. The constant work rate was

individualized for each participant by using the work rate at the halfway point between the anaerobic (lactate) threshold and the peak oxygen uptake, which had been determined from the ramp test. This constant work rate was equivalent to approximately 65-75% peak work rate.

Blood sampling and analysis

Prior to performing the interval exercise protocol, an indwelling catheter was inserted into the antecubital vein. A baseline sample was taken after the placement of the catheter and before the onset of exercise. The sample was taken 30 min after catheter placement to ensure that measurable physiological parameters of stress (e.g. heart rate and blood pressure) were at baseline levels. Participants then completed the ten 2 min bouts of constant work rate and additional blood samples were obtained immediately after exercise. Complete blood counts (CBC) and differential for white blood cell analysis were obtained by standard methods from the clinical haematology laboratory at UC Irvine. Plasma lactate was measured before and after exercise using YSI 2300 STAT Plus Glucose and Lactate Analyser.

Statistical analysis

In order to evaluate baseline difference and exercise effect between different sex, age and weight group in the same model, we utilized a mixed model which incorporated repeated measurements from the same participant to construct an appropriate covariance matrix to appropriately examine their fixed effects on the outcome variables. For each of the outcome variables-lactate, leukocyte and leukocyte subtypes-the mixed model evaluated their interactions with sex, age and weight pre- exercise and post-exercise. Repeated measurements at pre- exercise and post-exercise from the same participant were used to construct additional covariance matrices. Descriptive statistics presented in tables are mean and standard deviation (SD), while results from mixed models are mean and standard error (SE). Fold change was calculated as peak-exercise over pre-exercise.

RESULTS

Anthropometric and physiological characteristics

The anthropometric and physiological characteristics of the 148 participants appear in Table 1. 79 (53.4%) were females, 75 (50.7%) aged 8-12 and 66 (44.6%) were overweight or obese (BMI percentile \geq 85%). 21 (14.2%) of participants were Asian, 9 (6.1%) African-American, 86 (58.1%) Caucasian and 32 (21.6%) were Hispanic.

During the ramp test, participants achieved greater than 80% maximum heart rate and a respiratory exchange ratio (RER) equal or greater than 1.1, consistent with current standards that define acceptable efforts for peak $\dot{V}O_2$. No differences in maximal heart rate or RER as percent predicted were found among age groups. The relative exercise intensity determined as the average percentage for HR, WR and $\dot{V}O_2$ was not statistically significant across the groups. Overweight and obese children (BMI \geq 85%) were grouped together for this analysis. A post-hoc analysis of leukocyte and lactate responses between overweight and obese groups ultimately revealed no difference.

Lactate and leukocyte response to exercise

Table 2 presents the observed pre-exercise and post-exercise lactate levels and cell counts of total leukocyte and subtypes among gender, age and weight groups. Figure 1 displays baseline lactate, leukocyte and leukocyte subtype levels within each group and Figure 2 illustrates the effect of exercise on those groups. We found that baseline lactate level was significantly higher in overweight and obese compared to normal weight children (2.11 ± 0.08 vs. 1.68 ± 0.07 mmol/L, respectively, $p=0.0001$) and the pre-exercise to post-exercise lactate change was significantly greater in older children compared to younger children (5.71 ± 0.24 vs. 3.26 ± 0.24 , respectively, $p<0.0001$).

Similarly, the baseline leukocyte level was significantly higher ($p=0.0009$) in overweight (6362 ± 186 cells/ml) than in normal weight children (5511 ± 168) and the pre-exercise to post-exercise WBC change was significantly greater ($p<0.0001$) in older children (3599 ± 165) than in younger children (2544 ± 163).

For leukocyte subtypes, the baseline lymphocyte was significantly lower ($p=0.0044$) in boys (1948 ± 62 cells/ml) than in girls (2193 ± 58) and significantly higher ($p<0.0001$) in overweight children (2241 ± 63) than in normal-weight children (1900 ± 57); the baseline neutrophil was also significantly higher ($p=0.041$) in overweight children (3438 ± 160) than in normal-weight children (2994 ± 143).

In addition, the pre-exercise to post-exercise change in leukocyte levels was significantly higher ($p=0.021$) in girls (3343 ± 160) than in boys (2799 ± 169). After the interval exercise test, the lymphocyte increased significantly more ($p=0.037$) in girls (1547 ± 101) than in boys (1236 ± 107), significantly more ($p=0.004$) in older children (1606 ± 104) than in younger children (1177 ± 104) and also significantly more ($p=0.015$) in overweight children (1573 ± 110) than in normal-weight children (1211 ± 98). The neutrophil and the monocyte increased, after exercise, significantly more ($p=0.001$ & $p=0.0007$) in older children than (1604 ± 95 & 293 ± 23) in younger children (1155 ± 94 & 181 ± 23). Figure 3 shows that the median fold change pre- to post-exercise in lactate levels was 2.7 (range 1.4 to 8.1) in younger children and 4.1 fold (1.7 to 9.6) in older children, while the median fold change in leukocyte levels was 1.4 (1.0 to 2.1) in younger children and 1.6 fold (1.2 to 2.8) in older children, with a significantly higher ($p<0.001$) leukocyte to lactate ratio in younger children than older children.

Figure 4 demonstrates the leukocyte subsets response compared to the lactate response following exercise, with a higher neutrophil and lymphocyte subset response in the younger children when compared to lactate response.

DISCUSSION AND CONCLUSION

Traditionally, exercise response is characterized by either its immunological or metabolic indices, but not both. While lactate and leukocyte levels reflect metabolic and immune behaviours, they also are related to stress. Because we could not completely separate the lactate and leukocyte responses, we also evaluated both markers simultaneously by comparing their responses as a ratio. Our results show that the lactate and leukocyte

responses to the same relative submaximal exercise intensity were significantly greater in older children compared to younger children. Leukocyte response was also higher post-exercise in females compared to males. Younger children demonstrated a larger ratio of leucocytosis to lactate response following exercise. It is often challenging to compare exercise effects in children given varying body mass, maturational differences and fitness levels [10]. We addressed this by scaling the work rate to the physiologic capabilities of each individual. Consequently, our findings add information to the few paediatric studies that have demonstrated growth-related changes in immune cell trafficking in response to physical activity [2].

The mechanisms responsible for the larger exercise-associated elevations in total leukocyte and leukocyte subpopulation counts in older compared to younger children are unclear. We speculate that this observation may be related to maturational differences in stress and inflammation related particularly to oxygen availability and tissue uptake during exercise. It is well established that conditions of reduced oxygen delivery/availability lead to an inflammatory state characterized by higher leukocyte and lactate level [11]. Work from this and other laboratories indicates that younger children utilize more oxygen and produce less lactate during high-intensity exercise than do older children and adults [12]. Consequently, the attenuated leukocyte response to exercise in younger children may parallel the smaller lactate response and indicate relatively lower inflammatory activation during heavy exercise. Along these lines, earlier studies from our exercise lab have shown that brief exercise could lead to alterations in inflammatory gene expression, which may influence the leukocyte response [13]. A study by Dotan et al. [14] speculated that the rate of lactate elimination was similar despite age differences and that higher post-exercise lactate concentrations in older children were related to an increased lactate production in those older children rather than increased lactate clearance in younger children supporting the notion that heavy exercise in younger children is less inflammatory.

We had hypothesized that the association between the lower lactate and leukocyte response in younger children would be proportionally similar to the higher lactate and leukocyte response in older children. We anticipated, but did not find, that the exercise-induced leukocyte/ lactate would be the same in the two groups since both lactate and leukocytes are indicators of metabolic stress and part of the pro-inflammatory state, which, as noted, develops under stressful conditions such as exercise and hypoxia [15]. Whether the unexpected differences in the leukocyte/ lactate ratios resulted from a maturation-related altered pattern of leukocyte trafficking in the younger children is not known. We also examined two other possible factors, BMI and sex, that could influence lactate and leukocyte dynamics in response to exercise. In this study, body weight-related differences were seen in the lactate and leukocyte levels at rest and in response to exercise. Corroborating findings from this and other exercise laboratories, higher BMIs have been associated with higher lactate concentration and higher leukocyte levels at rest [16]. Interestingly, out of the leukocyte subtypes measured in this study, only the increase in lymphocytes seemed to be significant in the overweight group in response to exercise.

There was a small but significant sex effect across all ages with females having a higher leucocytosis in response to exercise compared to males. Interestingly, this sex-related

difference was not accompanied by a corresponding sex-related difference in lactate response to exercise. Although the mechanism related to the higher leucocytosis response in females remains unclear, we speculate that perhaps it may be related to different dynamics in blood flow and oxygen transport in exercise that is suggested by Winsley et al. [17].

These results highlight the need to control for age and sex when interpreting immunologic responses to exercise in a paediatric population. The mechanisms for how these different perturbations in leukocyte and lactate mobilization following exercise contribute to healthy growth and development still need further investigation.

ACKNOWLEDGEMENT

This work was supported in part by National Institutes of Health grants P01HD-048721, the UCI Institute for Clinical and Translational Science (CTSA grant) UL1RR031985 and the UC Irvine Multidisciplinary Exercise Sciences Training Program (5 T32 AR047752-09).

REFERENCES

- [1]. Baeza I, Alvarado C, Ariznavarreta C, Castillo C, Tresguerres JAF and Fuente M, *NeuroimmunoModulation*, 2008 15(6): p. 279–284. [PubMed: 19047805]
- [2]. Timmons BW, Tarnopolsky MA, Snider DP and Bar-Or O, *Med Sci Sports Exerc*, 2006 38(2): p. 293–304. [PubMed: 16531898]
- [3]. Cooper DM, Radom-Aizik S, Schwindt C and Zaldivar F, *J Appl Physiol*, 2007 103(2): p. 700–709. [PubMed: 17495117]
- [4]. Perez CJ, Nemet D, Mills PJ, Scheet TP, Ziegler MG and Cooper DM, *Eur J Appl Physiol*, 2001 86(1): p. 34–39. [PubMed: 11820320]
- [5]. Radom-Aizik S and Cooper DM, *Pediatr Exerc Sci*, 2016 28(2): p. 194–201. [PubMed: 27137166]
- [6]. Beneke R, Hutler M, Jung M and Leithauser RM, *J Appl Physiol*, 1985 99(2): p. 499–504.
- [7]. Kuczumski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R and Johnson CL, *Adv Data*, 2000 314: p. 1–27.
- [8]. Petersen AC, Crockett L, Richards M and Boxer A, *J Youth Adolesc*, 1988 17(2): p. 117–133. [PubMed: 24277579]
- [9]. Cooper D, Weiler-Ravell D, Whipp B and Wasserman K, *J Appl Physiol Respir Environ Exerc Physiol*, 1984 56(3): p. 628–634. [PubMed: 6706770]
- [10]. Cooper DM, Nemet D and Galassetti P, *Curr Opin Pediatr*, 2004 16(3): p. 286–292. [PubMed: 15167015]
- [11]. Chao J, Wood JG and Gonzalez NC *Resp Res*, 2009 10(1): p. 54–63.
- [12]. Zanonato S, Cooper DM and Armon Y, *J Appl Physiol*, 1991 71(3): p. 993–998. [PubMed: 1757338]
- [13]. Radom-Aizik S, Zaldivar F and Leu SY., Cooper DM. *J Appl Physiol*, 2009 107(1): p. 168–175. [PubMed: 19407257]
- [14]. Dotan R, Ohana S, Bediz C and Falk B, *J Pediatr Endocrinol Metab*, 2003 16(3): 419–429. [PubMed: 12705368]
- [15]. Rohm I, Ratka J, Pistulli R, Goebel B, Gecks T, Figulla HR and Jung C., *Clin Lab*, 2015 61(8): p. 1053–1059. [PubMed: 26427151]
- [16]. McNarry MA, Farr C, Middlebrooke A, Welford D, Breese B, Armstrong N and Barker AR, *Med Sci Sports Exerc*. 2015 47(9): p. 1877–1884. [PubMed: 25551403]
- [17]. Winsley RJ, Fulford J, Roberts AC, Welsman JR and Armstrong N, *J Sci Med Sport*, 2009 12(6): p. 647–651. [PubMed: 18768359]




Figure 1: Estimated mean lactate level and leukocyte counts at baseline by gender, age, or weight groups. Estimated mean and corresponding 95% CI are obtained from mixed model




Figure 2: Estimated mean lactate level and leukocyte counts change from pre-exercise to post-exercise by gender, age or weight groups. Estimated mean and corresponding 95% CI are obtained from mixed model




Figure 3: Pre- exercise and post-exercise differences in lactate, leukocyte, and leukocyte/lactate ratio in younger versus older children. When compared to lactate, the leukocyte count is significantly higher in younger children




Figure 4: Comparison of exercise-induced leukocyte subsets/lactate fold change ratio in younger versus older children

Table 1:

Anthropometric and physiological characteristics

		Girls				Boys			
Age		8-12		13-18		8-12		13-18	
BMI %tile		<85	85	<85	85	<85	85	<85	85
N		26	17	21	15	15	17	20	17
Height (cm)		141.5 ± 10.9	147.5 ± 10.8	163.2 ± 5.4	162.7 ± 7.0	142.6 ± 10.1	149.8 ± 10.4	176.1 ± 7.4	176.1 ± 7.6
Weight (kg)		34.5 ± 9.9	57.3 ± 17.7	57.0 ± 6.1	91.9 ± 29.0	34.9 ± 6.6	59.5 ± 16.6	65.0 ± 9.6	96.4 ± 23.1
Tanner stage		1.71 ± 0.95	2.47 ± 0.94	4.29 ± 0.46	4.27 ± 0.59	1.27 ± 0.59	1.76 ± 0.83	4.05 ± 0.52	3.76 ± 0.66
Ramp test	Peak $\dot{V}O_2$ (L/min)	1.21 ± 0.30	1.61 ± 0.38	1.92 ± 0.45	1.98 ± 0.29	1.40 ± 0.41	1.60 ± 0.51	2.68 ± 0.60	2.85 ± 0.67
	Peak $\dot{V}O_2$ (ml/min/kg)	35.7 ± 5.3	29.0 ± 5.7	33.7 ± 6.1	22.6 ± 6.3	39.9 ± 7.6	27.9 ± 8.0	41.4 ± 7.2	30.4 ± 8.0
	Max Heart Rate (bpm)	192.3 ± 12.8	193.9 ± 9.5	189.6 ± 11.9	183.1 ± 11.6	187.4 ± 9.9	186.7 ± 21.0	193.3 ± 11.7	190.5 ± 16.4
	Max RQ	1.13 ± 0.08	1.12 ± 0.06	1.15 ± 0.04	1.11 ± 0.08	1.09 ± 0.10	1.07 ± 0.06	1.17 ± 0.06	1.14 ± 0.05
	Average Heart Rate (bpm)	166.3 ± 10.1	159.2 ± 10.0	158.3 ± 15.0	153.4 ± 15.2	159.5 ± 13.1	161.4 ± 16.5	158.1 ± 9.8	159.4 ± 13.4
Constant Work Rate Protocol	Percent Work Rate *	65.3 ± 9.6	68.1 ± 5.1	64.7 ± 7.9	66.4 ± 5.3	64.8 ± 6.1	73.3 ± 14.2	69.2 ± 4.4	67.2 ± 3.6
	Percent Heart Rate *	86.8 ± 5.2	82.3 ± 5.7	83.6 ± 8.7	83.7 ± 6.1	85.2 ± 6.8	87.0 ± 9.6	82.6 ± 5.9	83.9 ± 6.3
	Percent $\dot{V}O_2$ *	66.0 ± 11.0	65.7 ± 7.4	63.0 ± 7.8	67.8 ± 6.2	63.8 ± 13.8	68.9 ± 16.1	71.9 ± 10.6	68.5 ± 9.3

* Calculated as the proportion of average at the 10 × 2 test over the maximum at the ramp exercise

Table 2:

Lactate level and leukocyte count and absolute change pre- and post- 10 × 2 exercise.

	Girls				Boys			
	08-Dec		13-18		08-Dec		13-18	
BMI %tile	<85	85	<85	85	<85	85	<85	85
N	25	16	21	15	15	17	20	17
Lactate (mmol/L)								
Pre-exercise	1.75 ± 0.54	2.1 ± 0.85	1.65 ± 0.52	2.15 ± 0.74	1.7 ± 0.44	2.23 ± 0.85	1.62 ± 0.58	1.98 ± 0.78
Post-exercise	5.23 ± 2.06	5.56 ± 2.00	6.88 ± 2.52	7.65 ± 2.03	4.35 ± 1.59	5.46 ± 1.71	7.87 ± 2.85	7.62 ± 2.07
Absolute change	3.48 ± 2.08	3.46 ± 1.83	5.23 ± 2.71	5.7 ± 1.65	2.65 ± 1.51	3.23 ± 1.24	6.25 ± 2.62	5.65 ± 1.97
Leukocyte (cells/ml)								
Pre-exercise	5188 ± 1331	6359 ± 1420	6119 ± 1580	6807 ± 1423	5613 ± 1917	6388 ± 1660	5305 ± 1292	5918 ± 1419
Post-exercise	7938 ± 2459	9212 ± 2614	9881 ± 2601	10733 ± 2982	7327 ± 2123	9147 ± 1969	8555 ± 2037	9329 ± 1872
Absolute change	2750 ± 1461	2853 ± 1578	3376 ± 1677	3927 ± 1911	1713 ± 897	2759 ± 925	3250 ± 1076	3412 ± 1176
Neutrophil (cells/ml)								
Pre-exercise	2630 ± 964	3139 ± 1063	3581 ± 1379	3953 ± 1147	3047 ± 2042	3444 ± 1442	2865 ± 1064	3259 ± 1078
Post-exercise	3915 ± 1911	4202 ± 1442	5353 ± 1663	5730 ± 1959	4040 ± 2151	4719 ± 1731	4390 ± 1563	4600 ± 1353
Absolute change	1285 ± 1017	1063 ± 492	1563 ± 1041	1778 ± 955	993 ± 592	1275 ± 467	1525 ± 606	1341 ± 635
Lymphocyte (cells/ml)								
Pre-exercise	1996 ± 538	2597 ± 657	1955 ± 491	2254 ± 411	1773 ± 467	2063 ± 444	1885 ± 384	2035 ± 582
Post-exercise	3194 ± 1054	4161 ± 1479	3530 ± 1255	4065 ± 1406	2393 ± 723	3344 ± 765	3265 ± 824	3659 ± 1374
Absolute change	1198 ± 758	1564 ± 1055	1442 ± 1068	1811 ± 1207	620 ± 409	1281 ± 622	1380 ± 675	1624 ± 1114
Monocyte (cells/ml)								
Pre-exercise	369 ± 142	411 ± 111	413 ± 108	417 ± 176	440 ± 216	456 ± 159	405 ± 182	476 ± 182
Post-exercise	593 ± 201	598 ± 204	708 ± 253	697 ± 255	547 ± 300	644 ± 193	720 ± 161	753 ± 283
Absolute change	224 ± 147	187 ± 178	267 ± 249	280 ± 204	107 ± 191	188 ± 126	315 ± 153	276 ± 278