

# UCLA

## UCLA Previously Published Works

### Title

Predicting Risk of End-Stage Liver Disease in Antiretroviral-Treated Human Immunodeficiency Virus/Hepatitis C Virus-Coinfected Patients

### Permalink

<https://escholarship.org/uc/item/5tz81386>

### Journal

Open Forum Infectious Diseases, 2(3)

### ISSN

2328-8957

### Authors

Re, Vincent Lo  
Kallan, Michael J  
Tate, Janet P  
et al.

### Publication Date

2015-09-01

### DOI

10.1093/ofid/ofv109

Peer reviewed

# Predicting Risk of End-Stage Liver Disease in Antiretroviral-Treated Human Immunodeficiency Virus/Hepatitis C Virus-Coinfected Patients

Vincent Lo Re III,<sup>1,2,3</sup> Michael J. Kallan,<sup>2</sup> Janet P. Tate,<sup>4,5</sup> Joseph K. Lim,<sup>4,5</sup> Matthew Bidwell Goetz,<sup>6</sup> Marina B. Klein,<sup>7</sup> David Rimland,<sup>8</sup> Maria C. Rodriguez-Barradas,<sup>9</sup> Adeel A. Butt,<sup>10,11,12</sup> Cynthia L. Gibert,<sup>13</sup> Sheldon T. Brown,<sup>14</sup> Lesley S. Park,<sup>5,15</sup> Robert Dubrov,<sup>5,15</sup> K. Rajender Reddy,<sup>1</sup> Jay R. Kostman,<sup>1</sup> Amy C. Justice,<sup>4,5</sup> and A. Russell Localio<sup>2</sup>

Departments of <sup>1</sup>Medicine, and <sup>2</sup>Biostatistics and Epidemiology and Center for Clinical Epidemiology and Biostatistics, Perelman School of Medicine, University of Pennsylvania, Philadelphia; <sup>3</sup>Medical Service, Philadelphia VA Medical Center, Pennsylvania; <sup>4</sup>VA Connecticut Healthcare System, West Haven; <sup>5</sup>Yale University School of Medicine, New Haven, Connecticut; <sup>6</sup>VA Greater Los Angeles Healthcare System and David Geffen School of Medicine at UCLA, California; <sup>7</sup>Chronic Viral Illness Service, McGill University Health Centre, Montreal, Canada; <sup>8</sup>Atlanta VA Medical Center and Emory University School of Medicine, Georgia; <sup>9</sup>Infectious Diseases Section, Michael E. DeBakey VA Medical Center and Department of Medicine, Baylor College of Medicine, Houston, Texas; <sup>10</sup>VA Pittsburgh Healthcare System, Pennsylvania; <sup>11</sup>Hamad Healthcare Quality Institute, Doha, Qatar; <sup>12</sup>Hamad Medical Corporation, Doha, Qatar; <sup>13</sup>Washington DC VA Medical Center, George Washington University Medical Center, Washington, District of Columbia; <sup>14</sup>James J. Peters VA Medical Center and Mt. Sinai School of Medicine, New York, New York; and <sup>15</sup>Yale School of Public Health, New Haven, Connecticut

**Background.** End-stage liver disease (ESLD) is an important cause of morbidity among human immunodeficiency virus (HIV)/hepatitis C virus (HCV)-coinfected patients. Quantifying the risk of this outcome over time could help determine which coinfecting patients should be targeted for risk factor modification and HCV treatment. We evaluated demographic, clinical, and laboratory variables to predict risk of ESLD in HIV/HCV-coinfecting patients receiving antiretroviral therapy (ART).

**Methods.** We conducted a retrospective cohort study among 6016 HIV/HCV-coinfecting patients who received ART within the Veterans Health Administration between 1997 and 2010. The main outcome was incident ESLD, defined by hepatic decompensation, hepatocellular carcinoma, or liver-related death. Cox regression was used to develop prognostic models based on baseline demographic, clinical, and laboratory variables, including FIB-4 and aspartate aminotransferase-to-platelet ratio index, previously validated markers of hepatic fibrosis. Model performance was assessed by discrimination and decision curve analysis.

**Results.** Among 6016 HIV/HCV patients, 532 (8.8%) developed ESLD over a median of 6.6 years. A model comprising FIB-4 and race had modest discrimination for ESLD (*c*-statistic, 0.73) and higher net benefit than alternative strategies of treating no or all coinfecting patients at relevant risk thresholds. For FIB-4 >3.25, ESLD risk ranged from 7.9% at 1 year to 26.0% at 5 years among non-blacks and from 2.4% at 1 year to 14.0% at 5 years among blacks.

**Conclusions.** Race and FIB-4 provided important predictive information on ESLD risk among HIV/HCV patients. Estimating risk of ESLD using these variables could help direct HCV treatment decisions among HIV/HCV-coinfecting patients.

**Keywords.** end-stage liver disease; hepatic decompensation; HIV; hepatitis C; HIV/HCV coinfection.

Chronic hepatitis C virus (HCV) coinfection is common among patients with human immunodeficiency

virus (HIV) infection [1]. Liver fibrosis progression is accelerated in HIV/HCV-coinfecting patients [2], and

Received 3 June 2015; accepted 5 July 2015.

Presented in part: Conference on Retroviruses and Opportunistic Infections (CROI) 2014, Boston, Massachusetts, 3–6 March 2014.

Correspondence: Vincent Lo Re III, MD, MSCE, Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania School of Medicine, 836 Blockley Hall, 423 Guardian Drive, Philadelphia, PA (vincentl@mail.med.upenn.edu).

**Open Forum Infectious Diseases**

© The Author 2015. Published by Oxford University Press on behalf of the Infectious

Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com).

DOI: 10.1093/ofid/ofv109

rates of hepatic decompensation are higher among coinfect- ed than HCV-monoinfected patients, even if HIV RNA is fully suppressed on antiretroviral therapy (ART) [3]. Consequently, end-stage liver disease (ESLD) has emerged as an important cause of morbidity in this population [3–5].

Despite the clinical impact of HCV-related ESLD in coinfect- ed patients, no models have been developed using readily avail- able demographic, clinical, and laboratory characteristics to predict the risk of ESLD within a specified time period. Quan- tifying ESLD risk over time could prompt clinicians and pa- tients to aggressively address modifiable risk factors for this condition (e.g., hazardous alcohol use; uncontrolled diabetes; severe anemia; detectable HIV viremia due to ART nonadher- ence; and/or resistance [3]) and identify subgroups of patients who should be followed closely for liver complications. Because highly efficacious and well tolerated, but costly, all-oral direct- acting antivirals for HCV treatment are now available [6, 7], a prognostic model for ESLD in coinfect- ed patients could help prioritize the need for HCV therapy among HIV/HCV patients, who have an urgent need for HCV treatment [3].

We previously identified important risk factors for hepatic de- compensation among HIV/HCV patients on ART [3]. In this study, we evaluated and compared alternative models comprising demographic, clinical, and laboratory variables to determine which combination of predictors can best classify HIV/HCV pa- tients according to their risk of progression to ESLD over time. Because liver fibrosis stage is an important factor used to deter- mine the urgency and timing of HCV therapy [8], we first eval- uated the prognostic ability of 2 noninvasive markers of hepatic fibrosis (FIB-4 and aspartate aminotransferase-to-platelet ratio index [APRI]) that can be calculated from routinely measured laboratory data (i.e., liver aminotransferases, platelet count). We then evaluated whether the addition of clinical and demographic variables improved the predictive ability of these indices.

## METHODS

### Study Design and Data Source

We conducted a retrospective cohort study among ART-treated HIV/HCV-coinfect- ed patients in the Veterans Aging Cohort Study (VACS) between January 1, 1997 and September 30, 2010 [9]. The Veterans Aging Cohort Study consists of electronic medical record data from HIV-infected patients receiving care at Veterans Affairs (VA) medical facilities across the United States. Data include demographic characteristics, hospital and outpatient diagnoses (recorded using International Classifica- tion of Diseases, Ninth Revision [ICD-9] codes), procedures, laboratory results, and dispensed medications. Clinically con- firmed cancer diagnoses are identified using the VA Central Cancer Registry. Date of death was ascertained from the VA Vital Status file [10]. The underlying cause of death was deter- mined by linkage with the National Death Index (NDI). [11]

This study was approved by the Institutional Review Boards of the University of Pennsylvania and Philadelphia VA Medical Center.

### Study Patients

Coinfect- ed patients were included if they had: (1) HIV (identi- fied using a validated algorithm [9]), (2) detectable HCV RNA, and (3) received ART, defined as use of at least 3 antiretrovirals from 2 different classes [12], for at least 12 months. We included only coinfect- ed patients on ART because current guidelines rec- ommend ART in all HIV/HCV patients regardless of CD4 count [12]. The majority of Veterans receive their medications from VA pharmacies [13, 14], and prior studies have shown the validity of VA dispensing data as a surrogate of actual adherence [15–18]. Patients were excluded if prior to start of follow-up (de- fined below), they had ESLD or received interferon-based therapy (because such treatment can affect noninvasive markers of hepatic fibrosis and reduces risk of ESLD [19]).

Follow-up began after 12 months on ART. The 12 months before the start of follow-up represented the baseline period, during which predictors were ascertained. Follow-up continued until study endpoint, death, initiation of HCV therapy, or last visit before September 30, 2010.

### Main Study Outcome

The primary endpoint was ESLD, defined as hepatic decompen- sation, hepatocellular carcinoma, or liver-related death. Hepatic decompensation was determined by 1 hospital discharge ICD-9 diagnosis or 2 or more outpatient ICD-9 diagnoses of ascites, spontaneous bacterial peritonitis, or esophageal variceal hemor- rhage (Supplementary Appendix 1). The requirement of 2 out- patient diagnoses aimed to exclude events that were suspected, but not subsequently confirmed, at follow-up visits. We validat- ed this definition in a prior survey, with 91% of events con- firmed by medical records [20]. Based on the results of the prior validation study [20], we did not include ICD-9 diagnoses for hepatic encephalopathy or jaundice, which could indicate decompensation, because these diagnoses were often linked to unrelated conditions (e.g., narcotic overuse, stroke recorded as encephalopathy; biliary obstruction, atazanavir-associated hyperbilirubinemia recorded as jaundice). The date of decompensation was defined as the hospital discharge date (if identi- fied by hospital diagnosis) or initial outpatient diagnosis date (if identified by outpatient diagnoses). Hepatocellular carcinoma was determined using the VA Central Cancer Registry, which confirms diagnoses by histology, cytology, or consistent radiog- raphy. Deaths were considered liver-related if the underlying cause of death in the NDI was hepatic decompensation, liver cancer, alcoholic liver disease, viral hepatitis, or nonalcoholic liver disease [3]. The date of ESLD was defined as the initial date of hepatic decompensation, hepatocellular carcinoma, or death from liver disease, whichever occurred first.

## Data Collection

Baseline variables evaluated as predictors of ESLD included the following: FIB-4 (<1.45; 1.45–3.25; >3.25), APRI (<1.0; 1.0–2.0; >2.0), age, body mass index ([BMI] <18.5; 18.5–24.9; 25.0–30.0; >30.0 kg/m<sup>2</sup>), severe anemia (hemoglobin <10 g/dL), CD4 cell count <200 cells/mm<sup>3</sup>, HIV RNA >400 copies/mL, alcohol dependence/abuse, diabetes mellitus, hepatitis B coinfection, and nonblack race, which have been shown to be risk factors for advanced liver disease in coinfecting patients [3, 19, 21]. FIB-4 is a measure of liver fibrosis stage that was developed and validated compared with liver biopsy in HIV/HCV-coinfecting patients [22]. It is calculated from age, alanine aminotransferase (ALT), aspartate aminotransferase (AST), and platelet count:  $FIB-4 = (\text{age [years]} \times \text{AST [U/L]}) / ((\text{platelet count [10}^9\text{/L]}) \times (\text{ALT [U/L]})^{1/2})$ . FIB-4 <1.45 identifies no/minimal FIB, whereas results >3.25 indicate advanced hepatic FIB/cirrhosis [22]. APRI, another noninvasive measure of hepatic fibrosis, uses AST and platelet count:  $APRI = ((\text{AST [U/L]}) / \text{upper limit of normal [considered as 40 U/L]}) / \text{platelet count (10}^9\text{/L)} \times 100$  [23]. An APRI >2.0 identifies cirrhosis in chronic HCV, and APRI <0.5 identifies no or minimal FIB [23]. We collected ALT, AST, platelet count, body weight, hemoglobin, CD4 count, and HIV RNA results measured closest to, but prior to, the start of follow-up. Alcohol dependence/abuse was defined by previously validated ICD-9 diagnoses (Supplementary Appendix 1) [24]. Diabetes was defined by random glucose


≥200 mg/dL, ICD-9 diagnosis, and/or antidiabetic medication use [25]. Hepatitis B coinfection was defined by a positive hepatitis B surface antigen. Additional variables collected at baseline for descriptive purposes included sex, HCV genotype, and HCV RNA.

## Statistical Analysis

We used Cox regression to estimate the relative hazards of ESLD (with 95% confidence intervals) for the predictors of interest [26]. We also evaluated FIB-4, APRI, hemoglobin, CD4 count, and HIV RNA level as continuous variables.

Next, because the stage of hepatic fibrosis is a strong determinant of ESLD [3, 19] and an important factor in the decision to initiate HCV therapy [8], we initially created separate models comprising FIB-4 and APRI alone. We then developed 2 separate lines of prognostic models with FIB-4 and APRI as the base models, respectively. Various combinations of the other predictors were added to the base models, including an interaction term for race. Age is a component of FIB-4 and so was not included in FIB-4 models.

Each model's performance was assessed by statistical significance and discrimination, using the concordance statistic (c-statistic) as an index of the model's ability to separate persons who will develop ESLD from those who will not [27]. We favored simpler models with fewer predictors that had appreciably different c-statistics from those of FIB-4 or APRI-based



**Figure 1.** Selection of Veterans Aging Cohort Study (VACS) human immunodeficiency virus (HIV)/hepatitis C virus (HCV)-coinfecting patients for inclusion in the study. Abbreviation: ART, antiretroviral therapy.

**Table 1. Baseline Characteristics of the HIV/Hepatitis C Virus (HCV)-Coinfected Patients Included in the Study Sample, Overall and By End-Stage Liver Disease Status<sup>a</sup>**

Characteristic	Overall (n = 6016)	End-Stage Liver Disease (n = 532)	No End-Stage Liver Disease (n = 5484)	P Value
Median age (years, IQR)	48 (44–53)	49 (45–53)	48 (44–52)	.061
Male sex	5929 (98.6)	532 (100.0)	5397 (98.4)	.003
Race/ethnicity				<.001
Black	3794 (63.1)	264 (49.6)	3530 (64.4)	
Caucasian	1474 (24.5)	164 (30.8)	1310 (23.9)	
Hispanic	576 (9.6)	89 (16.7)	487 (8.9)	
Other/Unknown	172 (2.9)	15 (2.8)	157 (2.9)	
Body mass index				.64
<18.5 kg/m <sup>2</sup>	179 (3.0)	13 (2.4)	166 (3.0)	
18.5–24.9 kg/m <sup>2</sup>	3045 (50.6)	259 (48.7)	2786 (50.8)	
25–29.9 kg/m <sup>2</sup>	1969 (32.7)	188 (35.3)	1781 (32.5)	
≥30 kg/m <sup>2</sup>	661 (11.0)	56 (10.5)	605 (11.0)	
Missing weight and/or height	162 (2.7)	16 (3.0)	146 (2.7)	
Diabetes mellitus	483 (8.0)	47 (8.8)	436 (8.0)	.47
History of alcohol dependence/abuse	1506 (25.0)	132 (24.8)	1374 (25.1)	.90
HCV genotype				.18
1 or 4	3553 (59.1)	306 (57.5)	3247 (59.2)	
2 or 3	456 (7.6)	31 (5.8)	425 (7.7)	
Other	14 (0.2)	2 (0.4)	12 (0.2)	
No result available	1993 (33.1)	193 (36.3)	1800 (32.8)	
HCV RNA <sup>b</sup>				.005
≥400 000 IU/mL and/or ≥1 000 000 copies/mL	4514 (75.0)	371 (69.7)	4143 (75.5)	
<400 000 IU/mL and/or <1 000 000 copies/mL	647 (10.8)	62 (11.7)	585 (10.7)	
Qualitative HCV RNA result only	855 (14.2)	99 (18.6)	756 (13.8)	
Hepatitis B surface antigen				.031
Positive	263 (4.4)	33 (6.2)	230 (4.2)	
Negative	5268 (87.6)	466 (87.6)	4802 (87.6)	
No result available at baseline	485 (8.1)	33 (6.2)	452 (8.2)	
Median HIV RNA (copies/mL, IQR)	400 (75–4870)	400 (179–3726)	400 (75–4905)	.080
HIV RNA				.66
>400 copies/mL	2534 (42.1)	233 (43.8)	2301 (42.0)	
≤400 copies/mL	3245 (54.0)	277 (52.1)	2968 (54.1)	
No result available at baseline	237 (3.9)	22 (4.1)	215 (3.9)	
Median CD4 cell count (cells/mm <sup>3</sup> , IQR)	320 (180–491)	294 (159–484)	322 (182–492)	.043
CD4 cell count				.23
≥500 cells/mm <sup>3</sup>	1377 (22.9)	113 (21.2)	1264 (23.0)	
350–499 cells/mm <sup>3</sup>	1156 (19.2)	94 (17.7)	1062 (19.4)	
200–349 cells/mm <sup>3</sup>	1544 (25.7)	135 (25.4)	1409 (25.7)	
<200 cells/mm <sup>3</sup>	1629 (27.1)	166 (31.2)	1463 (26.7)	
No result available at baseline	310 (5.2)	24 (4.5)	286 (5.2)	
Hemoglobin				.29
<10 g/dL	161 (2.7)	18 (3.4)	143 (2.6)	
≥10 g/dL	5820 (96.7)	509 (95.7)	5311 (96.8)	
No result available at baseline	35 (0.6)	5 (0.9)	30 (0.5)	
APRI				<.001
<1.0	4095 (68.1)	209 (39.3)	3886 (70.9)	
1.0–2.0	1003 (16.7)	153 (28.8)	850 (15.5)	
>2.0	591 (9.8)	142 (26.7)	449 (8.2)	
No result available at baseline	327 (5.4)	28 (5.3)	299 (5.5)	

Table 1 continued.

Characteristic	Overall (n = 6016)	End-Stage Liver Disease (n = 532)	No End-Stage Liver Disease (n = 5484)	P Value
FIB-4				<.001
<1.45	2160 (35.9)	92 (17.3)	2068 (37.7)	
1.45–3.25	2561 (42.6)	205 (38.5)	2356 (43.0)	
>3.25	968 (16.1)	207 (38.9)	761 (13.9)	
No result available at baseline	327 (5.4)	28 (5.3)	299 (5.5)	

Abbreviations: APRI, aspartate aminotransferase-to-platelet ratio index; HCV, hepatitis C virus; HIV, human immunodeficiency virus; IQR, interquartile range.

<sup>a</sup> Data are presented as number (%) unless otherwise specified.

<sup>b</sup> Based on highest baseline HCV RNA result.

models alone. Performance of promising models was further compared by decision curves [28, 29], a graphical approach for evaluating and comparing prognostic models that considers the inherent trade-offs of false-positive versus false-negative classifications arising out of varying risk thresholds. Details appear in [Supplementary Appendix 2](#). From the final model selected, we plotted Kaplan-Meier failure curves. Differences in ESLD by FIB-4 categories were estimated using log-rank tests.

To ensure that missing data did not affect the validity of our results, in a sensitivity analysis, we generated 10 imputed datasets, each with completed data on all predictors [30]. Data were analyzed using SAS 9.4 (SAS Institute Inc., Cary, NC) and Stata 13.1 (Stata Corporation, College Station, TX).

## RESULTS

### Patient Characteristics

Between 1997 and 2010, a total of 6016 ART-treated HIV/HCV-coinfected patients met our inclusion criteria (Figure 1). Absence of HCV RNA assessment was a common reason for exclusion (18%), but there were no clinically relevant differences in the characteristics between included patients and those excluded due to missing HCV RNA (data not shown).

The cohort (Table 1) was predominantly male and black, and one quarter of the patients had a history of alcohol dependence/abuse. Most patients were infected with HCV genotype 1 and had a high baseline HCV RNA level. The patients were followed for a median of 6.6 (interquartile range, 3.4–9.8) years. During follow-up, 482 (8.0%) patients initiated HCV therapy during and were censored. End-stage liver disease occurred in 532 (8.8%) patients during follow-up. At the time of the first ESLD event, a diagnosis indicative of hepatic decompensation was recorded in 397 (6.6%) patients, hepatocellular carcinoma was recorded in 63 (1.0%) patients, and liver-related death was recorded in 72 (1.2%) patients. Among the 397 patients with an incident hepatic decompensation event, 335 (84.4%) had ascites, 71 (17.9%) had spontaneous bacterial peritonitis, and 110 (27.7%) had variceal hemorrhage. Overall, 1936 (32.2%)

patients died during follow-up, of which 164 (8.5%) were liver-related deaths ([Supplementary Appendix 3](#)).

### Performance and Comparison of Prognostic Models for End-Stage Liver Disease

Baseline FIB-4, baseline APRI, and nonblack race were the strongest factors associated with ESLD among the variables evaluated in Cox regression analyses ([Supplementary Appendix 4](#)). Interactions

**Table 2. Candidate Prediction Models for End-Stage Liver Disease Among 6016 Antiretroviral-Treated HIV/Hepatitis C Virus-Co-infected Patients<sup>a</sup>**


Prediction Model	C-Statistic	
	FIB-4-Based Model	APRI-Based Model
No additional predictors included (FIB-4 or APRI alone)	0.70	0.71
Alcohol included	0.71	0.72
Age included	– <sup>b</sup>	0.73
Race included	0.73	0.72
Age + race included	– <sup>b</sup>	0.74
Race + diabetes + hepatitis B + anemia included	0.74	0.74
Age + race + diabetes + hepatitis B + anemia included	– <sup>b</sup>	0.75
All predictors included <sup>c</sup>	0.74 <sup>b</sup>	0.75

Abbreviations: APRI, aspartate aminotransferase-to-platelet ratio index; HCV, hepatitis C virus; HIV, human immunodeficiency virus.

<sup>a</sup> C-statistics for the base prognostic models consisting of FIB-4 (categorized as follows: <1.45; 1.45–3.25; >3.25) and APRI (categorized as follows: <1.0; 1.0–2.0; >2.0) alone are presented in the top row. C-statistics for alternative prognostic models comprising FIB-4 or APRI plus select combinations of the other predictors are presented in subsequent rows below. Models with FIB-4 did not include age because age is a component of FIB-4.

<sup>b</sup> C-statistic was not evaluated within the FIB-4-based models because age is a component of FIB-4.

<sup>c</sup> Covariates included age (for APRI-based model only), body mass index, anemia (hemoglobin <10 g/dL), CD4 cell count <200 cells/mm<sup>3</sup>, HIV RNA >400 copies/mL, history of alcohol dependence/abuse, diabetes mellitus, hepatitis B coinfection, and non-black race. FIB-4, APRI, hemoglobin, CD4 cell count, and HIV RNA level were also evaluated as continuous variables.



**Figure 2.** Risk of end-stage liver disease among antiretroviral-treated non-black (A) and black (B) HIV/hepatitis C virus-coinfected patients, by fibrosis (FIB)-4 score. Abbreviations: ALT, alanine aminotransferase; AST, aspartate aminotransferase.

with race were not observed for FIB-4 or APRI. Discrimination was similar for models with FIB-4 (c-statistic, 0.70) and APRI (c-statistic, 0.71) alone (Table 2). Alternative models comprising FIB-4 and race (c-statistic, 0.73) and APRI, age, and race (c-statistic, 0.74) had higher discrimination, but further addition of other combinations of predictors did not result in appreciably increased model discrimination (Table 2). Results of the decision curve analysis are shown in Supplementary Appendix 5. Because the model with FIB-4 + race had comparable discriminative ability and was simpler than the one with APRI + age + race, the model with FIB-4 + race was selected as the final model.

The risk of ESLD among non-blacks and blacks by category of FIB-4 is shown in Figure 2A and B, respectively (log-rank test among non-blacks,  $P < 0.001$ ; log-rank test among blacks,  $P < 0.001$ ). The risk at 5 years ranged from 2.6% to 26% among non-blacks (Figure 2A) and 2.0% to 14.0% among blacks (Figure 2B) across FIB-4 levels. Analyses using imputed data produced similar results.

## DISCUSSION

This study evaluated prognostic models for determining the risk of developing ESLD in ART-treated HIV/HCV-coinfected patients. A prognostic model for risk of developing ESLD comprising FIB-4 and race had better discriminative accuracy than FIB-4 or APRI alone, and it was comparable to the APRI, age, and race model, but simpler. Using established cut-points of FIB-4, we determined the risk of ESLD over time in non-blacks and blacks. Despite including a number of additional

variables associated with progression to ESLD (Supplementary Appendix 3), a model with all candidate predictors did not appreciably increase prognostic ability.

More importantly, our analyses in Figure 2 demonstrated that for HIV/HCV-coinfected individuals with moderate fibrosis by FIB-4 (i.e., results ranging from 1.45–3.25), there remains a significant risk of ESLD for non-blacks (>16% at 10 years) and blacks (>10% at 10 years). These results provide additional evidence that coinfecting patients with at least moderate (F2) or higher liver fibrosis should be strongly considered for HCV treatment and should not be deferred until the development of more advanced liver disease, where the risk of ESLD will be even higher.

With the availability of all-oral direct-acting anti-HCV regimens that have high virologic cure rates and tolerability, but which are very costly, determining the rational use of these agents is imperative [31]. Quantifying the absolute risk of ESLD over time using a simple prognostic model comprising FIB-4 and race could have important clinical implications for the management of coinfecting patients. Identifying coinfecting patients at higher risk of ESLD could help prioritize HCV treatment in those who would benefit from it most and inform the need for close monitoring for liver-related complications. Alternatively, in those with a low risk of ESLD, emphasis could be placed on the modification of other risk factors for ESLD, including the reduction of alcohol consumption, control of diabetes, and HIV suppression, as well as clinical monitoring for liver disease progression. Analogous to other quantitative risk assessment tools, such as the Framingham Coronary Heart Disease Risk Score [32] and World Health Organization Fracture Risk Assessment Tool [33], which estimate the

10-year risk of cardiovascular disease and fracture, respectively, use of this ESLD risk model could aid the management of chronic HCV in patients with HIV, more than simply knowing which factors are predictive of developing ESLD.

If validated, this model could contribute to an individualized approach to HCV treatment among HIV/HCV patients. We sought to develop a simple model that reports ESLD risk over a 10-year period in both graphical and tabular form to allow easy interpretation by coinfecting patients and their providers. Because the exact risk of ESLD that should prompt initiation of HCV treatment remains unclear, providers could use these results to discuss a patient's likelihood of developing ESLD over time and consider what risk of ESLD, in conjunction with other clinical and demographic factors, would prompt their initiation of HCV therapy.

FIB-4 is clinically credible as a predictor, easy to compute, and could be used in clinical practice in developed and developing world settings. Prior studies in coinfecting patients have demonstrated that the baseline stage of hepatic fibrosis is strongly associated with ESLD [3, 19]. A separate analysis among coinfecting women showed that FIB-4 was associated with all-cause mortality [34]. Among HIV-infected patients, baseline FIB-4 is associated with hepatocellular carcinoma risk [35]. In addition, FIB-4 uses a limited set of variables that are routinely and inexpensively measured in the outpatient setting, making this a useful option for resource-limited settings and practices where other noninvasive modalities of liver fibrosis assessment (e.g., transient elastography) might not be available.

Race is also credible as a predictor of ESLD. Recent studies in coinfecting patients reveal that persons of non-black race have a higher risk of hepatic decompensation [3] and liver-related death [36]. A recent cohort study among 149 407 chronic HCV patients showed that blacks had lower rates of cirrhosis and hepatocellular carcinoma than non-Hispanic whites, even after adjustment for HCV genotype, HCV treatment, diabetes, and BMI [37]. Non-black HCV-infected patients have been reported to have stronger HCV-specific immunity than blacks [38], which could result in increased immune-mediated hepatic inflammation and accelerated liver fibrosis progression. Recent analyses in HCV-monoinfected and HIV/HCV-coinfecting patients suggest that interleukin-28B CC genotype, which is common in non-blacks, is associated with increased hepatic inflammation and worse hepatic outcomes than those with CT or TT genotypes [39, 40]. Differences in the distribution of other genetic and non-genetic factors may also explain the disparity in ESLD risk between blacks and nonblacks. In future analyses, such factors may prove to be better indicators of ESLD risk than race.

This study has several potential limitations. First, we might have missed hepatic decompensation events and underestimated ESLD risk, because our decompensation definition did not include hepatic encephalopathy and because some patients

did not have a confirmed cause of death. However, the incidence of ESLD among HIV/HCV patients in this study (8.8%) was similar to that reported in other cohort studies of coinfecting patients [5, 19]. Furthermore, we previously evaluated the incidence rate of decompensation within VACS, US Medicare, and US Medicaid data using the validated ICD-9 definition (to capture outcomes occurring at non-VA hospitals that did not result in transfer to a VA facility). The incidence rate of decompensated cirrhosis was almost identical to that within the VACS alone, suggesting that few ESLD events were missed [3]. Second, our sample included mostly male US Veterans, and results may not be generalizable to women or to non-Veteran populations. Future analyses should evaluate the ability of FIB-4 and race to predict ESLD among coinfecting women and non-Veterans. Third, differences by race in receiving HCV treatment (and being censored) should not affect our results because we stratified our analysis by race. Fourth, external validation of the prognostic model to predict ESLD in other coinfecting populations is essential to ensure reproducibility of our findings across varied settings. Finally, although we considered many potential predictors during model development, data on liver stiffness measurement (using transient elastography) and genetic markers, such as interleukin-28B genotype and interferon lambda-4 deltaG genotype [40, 41], were not assessed. Future models evaluating ESLD risk in coinfecting patients could determine whether these factors improve accuracy and utility. However, FIB-4 and race have the advantage of being simple predictors. Furthermore, because our final model included 2 categorical predictors, a simple Kaplan-Meier plot allows users to estimate risk easily for any given time frame.

## CONCLUSIONS

In summary, FIB-4 and race can be used to quantify the risk of ESLD among HIV/HCV-coinfecting patients on ART. Such information could prompt coinfecting patients to reduce their risk factors for liver disease progression and aid HCV treatment decisions for both clinicians and patients. Future studies should evaluate the classification performance of FIB-4 and race in other coinfecting populations, compare the performance of FIB-4 using the established cutoff points versus alternative thresholds, and determine whether its use improves outcomes.

## Supplementary Material

Supplementary material is available online at *Open Forum Infectious Diseases* (<http://OpenForumInfectiousDiseases.oxfordjournals.org/>).

## Acknowledgments

We are grateful to the VA Central Cancer Registry for permitting us to link the Veterans Aging Cohort Study data with the VA Central Cancer Registry data and for providing us with a dataset containing the linked records.



**Disclaimers.** The contents of this article do not represent the views of the Department of Veterans Affairs, the US Food and Drug Administration, or the United States Government.

**Financial Support.** This study was supported by research grant funding from the National Institute of Allergy and Infectious Diseases (K01 AI070001; to V. L. R.), National Institute on Alcohol Abuse and Alcoholism (U01 AA13566; to A. C. J.), and National Cancer Institute (F31 CA180775; to L. S. P.).

**Potential conflicts of interest.** J. K. L. has received research grant support (to Yale University) from Abbott, Boehringer-Ingelheim, Bristol-Myers Squibb, Genentech, Gilead, Glaxo-Smith Kline, Globeimmune, Janssen, and Vertex and has served as a consultant to Boehringer-Ingelheim, Bristol-Myers Squibb, Gilead, Merck, and Vertex. M. B. K. has received research grant support from the Fonds de la Recherche en Santé du Québec, served as a consultant to Glaxo-SmithKline and Viv, received honoraria for lectures from Bristol-Myers Squibb, and received honoraria for the development of educational materials from Gilead. V. L. R. has received investigator-initiated research grant support (to the University of Pennsylvania) from AstraZeneca and Gilead Sciences.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

## References

1. Sherman KE, Rouster SD, Chung RT, Rajicic N. Hepatitis C virus prevalence among patients infected with human immunodeficiency virus: a cross-sectional analysis of the US adult AIDS Clinical Trials Group. *Clin Infect Dis* **2002**; 34:831–7.
2. Macias J, Berenguer J, Japon MA, et al. Fast fibrosis progression between repeated liver biopsies in patients coinfecting with human immunodeficiency virus/hepatitis C virus. *Hepatology* **2009**; 50:1056–63.
3. Lo Re V III, Kallan MJ, Tate JP, et al. Hepatic decompensation in antiretroviral-treated patients co-infected with HIV and hepatitis C virus compared with hepatitis C virus-monoinfected patients: a cohort study. *Ann Intern Med* **2014**; 160:369–79.
4. Weber R, Sabin CA, Friis-Moller N, et al. Liver-related deaths in persons infected with the human immunodeficiency virus: the D:A:D study. *Arch Intern Med* **2006**; 166:1632–41.
5. Pineda JA, Garcia-Garcia JA, Aguilar-Guisado M, et al. Clinical progression of hepatitis C virus-related chronic liver disease in human immunodeficiency virus-infected patients undergoing highly active antiretroviral therapy. *Hepatology* **2007**; 46:622–30.
6. Dieterich D, Rockstroh JK, Orkin C, et al. Simeprevir (TMC435) with pegylated interferon/ribavirin in patients coinfecting with HCV genotype 1 and HIV-1: a phase 3 study. *Clin Infect Dis* **2014**; 59:1579–87.
7. Sulkowski MS, Naggie S, Lalezari J, et al. Sofosbuvir and ribavirin for hepatitis C in patients with HIV coinfection. *JAMA* **2014**; 312:353–61.
8. Shiffman ML, Benhamou Y. Patients with HCV and F1 and F2 fibrosis stage: treat now or wait? *Liver Int* **2013**; 33(Suppl 1):105–10.
9. Fultz SL, Skanderson M, Mole LA, et al. Development and verification of a “virtual” cohort using the National VA Health Information System. *Med Care* **2006**; 44(8 Suppl 2):S25–30.
10. Fisher SG, Weber L, Goldberg J, Davis F. Mortality ascertainment in the veteran population: alternatives to the National Death Index. *Am J Epidemiol* **1995**; 141:242–50.
11. Sathiakumar N, Delzell E, Abdalla O. Using the National Death Index to obtain underlying cause of death codes. *J Occup Environ Med* **1998**; 40:808–13.
12. Gunthard HF, Aberg JA, Eron JJ, et al. Antiretroviral treatment of adult HIV infection: 2014 recommendations of the International Antiviral Society-USA Panel. *JAMA* **2014**; 312:410–25.
13. Elixhauser A, Eisen SA, Romeis JC, Homan SM. The effects of monitoring and feedback on compliance. *Med Care* **1990**; 28:882–93.
14. Steiner JF, Koepsell TD, Fihn SD, Intui TS. A general method of compliance assessment using centralized pharmacy records. Description and validation. *Med Care* **1988**; 26:814–23.
15. Braithwaite RS, Kozal MJ, Chang CC, et al. Adherence, virological and immunological outcomes for HIV-infected veterans starting combination antiretroviral therapies. *AIDS* **2007**; 21:1579–89.
16. Grossberg R, Zhang Y, Gross R. A time-to-prescription-refill measure of antiretroviral adherence predicted changes in viral load in HIV. *J Clin Epidemiol* **2004**; 57:1107–10.
17. Heisler M, Hogan MM, Hofer TP, et al. When more is not better: treatment intensification among hypertensive patients with poor medication adherence. *Circulation* **2008**; 117:2884–92.
18. Steiner JF, Prochazka AV. The assessment of refill compliance using pharmacy records: methods, validity, and applications. *J Clin Epidemiol* **1997**; 50:105–16.
19. Limketkai BN, Mehta SH, Sutcliffe CG, et al. Relationship of liver disease stage and antiviral therapy with liver-related events and death in adults coinfecting with HIV/HCV. *JAMA* **2012**; 308:370–8.
20. Lo Re V III, Lim JK, Goetz MB, et al. Validity of diagnostic codes and liver-related laboratory abnormalities to identify hepatic decompensation events in the Veterans Aging Cohort Study. *Pharmacoepidemiol Drug Saf* **2011**; 20:689–99.
21. Kirk GD, Mehta SH, Astemborski J, et al. HIV, age, and the severity of hepatitis C virus-related liver disease: a cohort study. *Ann Intern Med* **2013**; 158:658–66.
22. Sterling RK, Lissen E, Clumeck N, et al. Development of a simple non-invasive index to predict significant fibrosis in patients with HIV/HCV coinfection. *Hepatology* **2006**; 43:1317–25.
23. Wai CT, Greenson JK, Fontana RJ, et al. A simple noninvasive index can predict both significant fibrosis and cirrhosis in patients with chronic hepatitis C. *Hepatology* **2003**; 38:518–26.
24. Justice AC, McGinnis KA, Atkinson JH, et al. Psychiatric and neurocognitive disorders among HIV-positive and negative veterans in care: Veterans Aging Cohort Five-Site Study. *AIDS* **2004**; 18(Suppl 1):S49–59.
25. Butt AA, Fultz SL, Kwok CK, et al. Risk of diabetes in HIV infected veterans pre- and post-HAART and the role of HCV coinfection. *Hepatology* **2004**; 40:115–9.
26. Collett D. *Modeling Survival Data in Medical Research*. 2nd ed. New York: Chapman and Hall, **2003**.
27. Harrell FE Jr, Lee KL, Mark DB. Multivariable prognostic models: issues in developing models, evaluating assumptions and adequacy, and measuring and reducing errors. *Stat Med* **1996**; 15:361–87.
28. Vickers AJ, Elkin EB. Decision curve analysis: a novel method for evaluating prediction models. *Med Decis Making* **2006**; 26:565–74.
29. Moons KG, Altman DG, Reitsma JB, et al. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): explanation and elaboration. *Ann Intern Med* **2015**; 162:W1–73.
30. Royston P. Multiple imputation of missing values: update. *The Stata Journal* **2005**; 5:188–201.
31. Steinbrook R, Redberg RF. The high price of the new hepatitis C virus drugs. *JAMA Intern Med* **2014**; 174:1172.
32. Wilson PW, D’Agostino RB, Levy D, et al. Prediction of coronary heart disease using risk factor categories. *Circulation* **1998**; 97:1837–47.
33. Kanis JA, Johnell O, Oden A, et al. FRAX and the assessment of fracture probability in men and women from the UK. *Osteoporos Int* **2008**; 19:385–97.
34. Bambha K, Pierce C, Cox C, et al. Assessing mortality in women with hepatitis C virus and HIV using indirect markers of fibrosis. *AIDS* **2012**; 26:599–607.
35. Park LS, Tate JP, Justice AC, et al. FIB-4 index is associated with hepatocellular carcinoma risk in HIV-infected patients. *Cancer Epidemiol Biomarkers Prev* **2011**; 20:2512–7.
36. Sarkar M, Bacchetti P, French AL, et al. Lower liver-related death in African-American women with human immunodeficiency virus/hepatitis C virus coinfection, compared to Caucasian and Hispanic women. *Hepatology* **2012**; 56:1699–705.

37. El-Serag HB, Kramer J, Duan Z, Kanwal F. Racial differences in the progression to cirrhosis and hepatocellular carcinoma in HCV-infected veterans. *Am J Gastroenterol* **2014**; 109:1427–35.
38. Rosen HR, Weston SJ, Im K, et al. Selective decrease in hepatitis C virus-specific immunity among African Americans and outcome of antiviral therapy. *Hepatology* **2007**; 46:350–8.
39. Noureddin M, Wright EC, Alter HJ, et al. Association of IL28B genotype with fibrosis progression and clinical outcomes in patients with chronic hepatitis C: a longitudinal analysis. *Hepatology* **2013**; 58:1548–57.
40. Barreiro P, Pineda JA, Rallon N, et al. Influence of interleukin-28B single-nucleotide polymorphisms on progression to liver cirrhosis in human immunodeficiency virus-hepatitis C virus-coinfected patients receiving antiretroviral therapy. *J Infect Dis* **2011**; 203:1629–36.
41. Meissner EG, Bon D, Prokunina-Olsson L, et al. IFNL4-deltaG genotype is associated with slower viral clearance in hepatitis C, genotype-1 patients treated with sofosbuvir and ribavirin. *J Infect Dis* **2014**; 209:1700–4.