

UCSF

UC San Francisco Previously Published Works

Title

Sleep duration in Mexican American children: Do mothers' and fathers' parenting and family practices play a role?

Permalink

<https://escholarship.org/uc/item/5xn0w1n6>

Journal

Journal of Sleep Research, 28(4)

ISSN

0962-1105

Authors

Martinez, Suzanna M

Tschann, Jeanne M

Butte, Nancy F

et al.

Publication Date

2019-08-01

DOI

10.1111/jsr.12784

Peer reviewed



Published in final edited form as:

J Sleep Res. 2019 August ; 28(4): e12784. doi:10.1111/jsr.12784.

Mothers' and fathers' parent feeding practices, family dinner and TV practices related to weeknight sleep duration in Mexican American children

SM Martinez¹, JM Tschann², NF Butte³, S Gregorich⁴, C Penilla⁵, E Flores⁶, LA Pasch⁷, LC Greenspan⁸, and J Deardorff⁹

¹University of California, Nutrition Policy Institute, 1111 Franklin St, 10th Floor, Oakland, CA, USA, 94607;

²Department of Psychiatry, University of California at San Francisco, San Francisco, CA, USA 94143-0848,

³Baylor College of Medicine, USDA/ARS Children's Nutrition Research Center, Department of Pediatrics, 1100 Bates Street Houston, TX, USA 77030-2600,

⁴Department of Medicine, University of California at San Francisco, San Francisco, CA, USA 94143-0856,

⁵School of Public Health, University of California at Berkeley, 50 University Hall, University of California at Berkeley, Berkeley, CA, USA 94720-7360,

⁶Counseling Psychology Department, School of Education, University of San Francisco, 2130 Fulton Street, San Francisco, CA, USA 94118,

⁷Department of Psychiatry, University of California at San Francisco, San Francisco, CA, USA 94143-0848,

⁸Kaiser Permanente, 2200 O'Farrell Street, San Francisco, CA, USA 94115,

⁹Division of Community Health and Human Development, School of Public Health, 50 University Hall, University of California at Berkeley, Berkeley, CA, USA 94720-7360,

Summary

This study examined parenting styles, parenting practices, and family practices that may be associated with weeknight sleep duration among 8–10-year-old Mexican American (MA) children. This cross-sectional study of MA children used baseline data from a two-year cohort study of mother-child pairs (n=308) with additional data on fathers (n=166). Children's weeknight sleep duration was accelerometer-estimated and averaged for 2 weeknights. Parents reported on their parenting styles and practices regarding food, and family food-related practices. Multivariable linear regression analysis was used to examine sleep duration with parenting styles and practices,

Corresponding author: suzanna.martinez@ucop.edu.

Conflict of Interest Statement:

The authors have no conflicts of interest to disclose.

Financial Disclosure: The authors have no financial relationships to disclose.

and family practices, and adjusting for child gender and BMI. Model 1 included mothers' parenting styles and practices; Model 2 included both mothers' and fathers' parenting styles and practices. Children's average sleep duration was 9.5 (SD= 0.8) hours. Mothers who used pressure to eat and those who used food to control behavior had children with longer sleep duration ($\beta=0.21$, $P<0.01$; $\beta=0.15$, $P=0.03$, respectively). Mothers who reported their child ate dinner with the TV on and those who valued eating dinner as a family had children with shorter sleep duration ($\beta=-0.16$, $P=0.01$, $\beta=-0.18$, $P=0.01$, respectively). Fathers who restricted their child's food amount had children with shorter sleep duration ($\beta=-0.27$, $P=0.01$). Mothers' and fathers' feeding practices, the child's eating dinner with the TV on, and valuing family dinners played a role in children's weeknight sleep duration among Mexican American families. Parental feeding practices and family mealtime contexts may matter regarding children's weeknight sleep duration.

INTRODUCTION

The optimal recommended duration of sleep for children aged 6–12 years, is 9 hours (American Academy of Sleep Medicine, 2016). However, 36%–41% of elementary school-age children are not obtaining the recommended sleep amount (Hawkins and Takeuchi, 2016).

There has been a decline in sleep duration (1980–2008), which parallels the rise in obesity (1985–2012) in the U.S. (Finucane et al., 2011, Ford et al., 2015). The elevated risks for short sleep and obesity have been observed in Latino children (Peña et al., 2016, Skinner and Skelton, 2014), with concomitant calls for a more comprehensive approach to obesity prevention, especially among Latinos (US Department of Health and Human Services, 2011). Targeting sufficient sleep duration may be a potential intervention to reduce childhood obesity (Haines et al., 2013). However, little is known regarding modifiable factors for improving sleep duration, particularly among elementary school-age children, and few interventions have focused on improving sleep duration to prevent childhood obesity (Hamre et al., 2006, Yoong et al., 2016).

We examined potential parenting and family factors related to children's sleep duration. Parenting styles in the domains of parental warmth (e.g., supportiveness, responsiveness, acceptance) and parental control (e.g., demandingness, monitoring) have often been examined in the context of child functioning, such as academic performance (Aunola and Nurmi, 2004, Darling and Steinberg, 1993), but few studies have examined parenting styles in relation to children's sleep. In one study, a parenting style that encouraged social maturity was related to longer sleep duration, less night-to-night variability in sleep (for girls), and a bedtime before 11:00 PM (Spilsbury et al., 2005). Another study of low-income adolescents found that parental control was related to fewer difficulties initiating sleep, and that Latino adolescents had fewer difficulties maintaining sleep, which resulted in longer sleep duration than their non-Latino counterparts (Roblyer and Grzywacz, 2015). Authors attributed longer sleep in Latino adolescents to family stability, specifically a two-parent household, which is similar to findings in Black and White adolescents (Troxel et al., 2014).

Parenting practices, such as instilling family meals and routines, as compared with parenting styles, have also been examined in relation to a range of lifestyle-related health behaviors,

including children's eating and diet (Shloim et al., 2015). Studies have found that family practices around eating overlap with children's sleep practices. Family meals have been shown to be related to more healthful dietary intake and longer sleep in children (Ray and Roos, 2012). In addition, instilling routines (e.g., schedules, family meals) and limits around screen time has shown to be protective of children's sleep duration (Koulouglioti et al., Ray et al., 2013). One relationship that has not been examined is that of parental feeding practices in relation to children's sleep duration. Parental feeding practices refer to the goal-directed behaviors used by parents to influence child eating (Shloim et al., 2015). Certain parental feeding practices, specifically restricting certain foods and using food to control behavior, have been shown to increase risk for child obesity (Tschann et al., 2015). Moreover, there is little known about the role of fathers' parenting practices in addition to mothers' in the relationship between parental feeding practices and children's sleep duration, especially among Latino families.

We posited that parental styles, parent feeding practices, and family practices related to meals and television associate with children's sleep, as they do for eating behaviors. Because parenting styles and practices in different domains might be interconnected (Jones et al., 2014), it is also possible that these factors are related to longer sleep duration in children. We explored the association of mothers' parenting styles, parenting practices, and family practices with weeknight sleep duration among 8-to 10-year-old Mexican American children (N=308). Additionally, we explored these relationships in families in which both mothers and fathers participated in the research (n=166). This research has the potential to shed light on parenting and family contextual factors to improve sleep duration, and possibly reduce the burden of obesity among Mexican American children.

METHODS

Participants and Data Collection

A total of 322 Mexican American children and their mothers who were members of Kaiser Permanente Northern California, an integrated health care delivery organization, were recruited to participate in a cohort study to understand parental influences on obesity in Mexican American children. Letters were sent to parents, who were then invited to participate in a 2-year study if: 1) the mother was of Mexican decent (born in the U.S./ Mexico), and 2) their child was 8–10 years of age and had no major illnesses. Bilingual interviewers obtained parental informed consent and child assent to participate in the research. Parents and children were interviewed in their homes in the participants' preferred language. Of the 322 mother-child pairs, 174 fathers opted to participate and were also interviewed. For the current cross-sectional study, we used baseline data and included families with complete information on children's sleep duration, resulting in 308 mother-child pairs. Of these mother-child pairs, a total of 166 fathers had complete information on parenting. The study was approved by the University of California San Francisco and Kaiser Permanente Northern California Research Foundation Institutional Review Boards.

Dependent variable

Child sleep duration.—Sleep duration was assessed for three consecutive nights (2 weeknights and 1 weekend night), and as previously reported, was estimated using the hip-worn Actical accelerometer (Philips Respironics, Bend, OR) (Martinez et al., 2014b). The monitor was attached to an elastic belt and positioned on the child above the iliac crest of the right hip. Research assistants provided both verbal and written instructions for care and placement of the monitor at the time of the home visit. Children were instructed to wear the monitor at all times for three consecutive days, with the exception of bathing. Data were collected at 1-minute intervals at a specified start time. (Butte et al., 2014) Data were downloaded to a computer, and data output included the time stamp and total accelerometer counts. Data completeness was verified against the participant's log, and times and reasons for monitor removal were coded in the file. A single trained research assistant conducted visual inspections and cross-checked bedtimes and wake times with the child-reported diaries, using a plot of activity counts per minute for each 24-hour period. A plot of activity counts per minute for each 24-hour period was used to identify the time of sleep onset and end. During the nighttime sleep period, activity counts were usually zero. Sleep periods were cross-checked with the participants' wear log for "off" times. Any minutes scored as awake were removed from the sleep duration variable. For the current study, we used weeknight sleep duration because parents are more likely to engage in weekday routines on weekdays compared to weekends, which may be less structured. We observed a higher mean in sleep duration on the weekend than the weekday, with weekend sleep averaging 23 minutes longer (SD=78 minutes).

Independent variables

Parenting styles and practices.—Parenting styles of warmth (10 items; *e.g.*, tells child that what he/she tries to do or achieve is appreciated) and control (6 items; *e.g.*, child is punished if he/she misbehaves) were assessed using the Blocks' Child Rearing Practices Report as adapted by Aunola & Nurmi (Aunola and Nurmi, 2004). Response options ranged from never/strongly disagree [1] to always [5]/strongly agree [4]. For each subscale, mean scores were calculated ($\alpha_{\text{mothers}} = 0.71$, $\alpha_{\text{fathers}} = 0.78$), and centered with scores ranging from -2 to 2; higher scores represented more use of that parenting style.

Mothers and fathers reported on their own child-feeding behaviors using the 55-item Parental Feeding Practices (PFP) Questionnaire (Musher-Eizenman and Holub, 2007). In our prior research with Mexican American children 8–10 years old, we validated the PFP questionnaire, and identified the following four parental feeding practices: use of food to control behavior, pressure to eat, restriction of food amount, and positive involvement in child eating (Musher-Eizenman and Holub, 2007). Using this questionnaire, we have found that restrictive feeding practices and pressure to eat predicted future increases in child weight (Tschann et al., 2015). The PFP subscale items include: 9 items on use of food to control behavior (*e.g.*, gives child food in exchange for good behavior or to lift mood), 24 items on positive involvement in child eating (*e.g.*, encourages child to healthy foods before less healthy foods; sets limits on sweets, junk food, and sodas), 10 items on pressure to eat (*e.g.*, child has to eat everything on his/her plate), and 12 items on restriction of food amount (*e.g.*, tells child when he/she has eaten enough) ($\alpha_{\text{mothers}} = .77-.88$; $\alpha_{\text{Fathers}} = 0.70-0.91$). All

questions were worded in terms of frequency of behavior, and response options range from never [1] to always [5] (Tschann et al., 2013). For each subscale, mean scores ranged from 1 to 5; higher scores represented more use of that feeding practice.

Lastly, parental screen time monitoring was assessed using the scale developed by Robinson et al. (Robinson, 1999) and included 8 items (*e.g.*, keeps track of the amount of TV/ videos child watches; limits the amount of time child watches TV/videos during the week). Response options ranged from never [1] to always [5]. A mean score was computed ($\alpha_{\text{mothers}} = 0.84$, $\alpha_{\text{fathers}} = 0.86$), ranging from 1 to 5; a higher score indicated more TV monitoring.

Family practices regarding TV.—Mothers and fathers were asked to report on the number of TVs in the home, whether there was a TV where the child sleeps, and whether there was a computer where the child sleeps (no/yes).

Mothers and fathers were asked to report the number of days in a week the child ate breakfast or dinner in a room where a TV was on. Response categories were never, 1–2, 3–4, 5–6, and 7 days per week. Numeric values were assigned to the categories 0, 1.5, 3.5, 5.5, and 7, respectively to construct approximately continuous measures in days per week.

Family practices regarding dinner.—Two items measured family practices regarding dinner. First, mothers and fathers reported on the *value of family dinners*, which included ‘*In my family, we expect everyone to be home for dinner.*’ Response options ranged from strongly disagree [1] to strongly agree [4], and were collapsed into categories of disagree [1] and agree [2], given the very low frequency of responses in the strongly disagree category. Mothers and fathers also reported on the *frequency of shared family dinners* in the last 7 days. Response categories were never, 1–2, 3–4, 5–6, and 7 days per week. Numeric values were assigned to the categories 0, 1.5, 3.5, 5.5, and 7, respectively to construct approximately continuous measures in days per week.

Covariates and descriptive variables.—Mothers and fathers reported on parent age, education, household income, marital status, single-parent household, foreign-born status, years in the U.S., and number of hours worked per week. Parent participation was tested as a covariate and was coded as follows: married mothers with fathers’ participation (reference), married mothers without fathers’ participation, single mothers with fathers’ participation, and single mothers without fathers’ participation. Research assistants collected child gender from mothers, and measured child height and weight. Height and weight were obtained using standard procedures in duplicate while the participants were wearing light indoor clothing and no shoes (Lohman et al., 1989, Stallings and Fung, 1999). BMI was calculated [$\text{BMI} = \text{weight (kg)} / \text{height (m)}^2$] converted to age- and gender-specific percentiles, and converted to z-scores using Centers for Disease Control (CDC) growth charts. Cutoffs at the 85th percentiles for overweight status, and 95th percentiles for obese status were used for descriptive purposes (Centers for Disease Control and Prevention).

We tested neighborhood environment as a potential covariate of children’s sleep duration. Studies have found that neighborhood crime and disorder are elements within the

environmental domain that potentially disrupt children's sleep.(Martinez and Thompson-Lastad, 2015) Mothers and fathers reported on 27 items regarding neighborhood perceptions of disorder (13 items; *e.g.*, conditions such as drugs, litter, and presence of gangs.) and victimization (14 items; *e.g.*, fear of going out in the neighborhood during the day or at night, fear of being robbed and/or raped).(Coulton et al., 1996) Response options ranged from never [1] to always [5]. A mean score was computed for neighborhood disorder ($\alpha_{\text{mothers}}= 0.91$; $\alpha_{\text{fathers}}= 0.91$) and neighborhood victimization ($\alpha_{\text{mothers}}= 0.92$; $\alpha_{\text{fathers}}= 0.92$). Mean scores ranged from 1 to 5; a higher score represented more disorder and victimization.

Analysis

All statistics were computed using IBM SPSS 23 Statistics for Windows (Armonk, NY: IBM Corp). Descriptive statistics were computed using means (SD) and frequencies for all study variables. Bivariate correlations were examined between sleep duration and all independent variables, including parental demographic factors, parenting styles and practices, and family/home practices regarding food and TV.

A multivariable linear regression model was first examined using the full sample of mother-child pairs with mothers' report of parenting styles and practices, and family/home practices regarding TV and dinner as independent variables, in relation to sleep duration (hours) as the dependent variable (Model 1). Second, we fit data on a subsample of families in which fathers also participated in the research; this model used the same variables for fathers that had been included in the model for the full sample of mother-child pairs (Model 2). Associations were significant at $P < 0.05$.

RESULTS

Participant characteristics

Table 1 presents the descriptive information for each study variable for mother-child pairs and father-child pairs. On average, children were 8.9 (SD= 0.8) years old, 53% female, and had a sleep duration of 9.5 (SD= 0.8) hours. T-tests comparing mothers and fathers on all variables showed that fathers were older (df= 165, $t=-6.23$, $P < 0.001$), had lived in the U.S. longer (df= 113, $t=-3.65$, $P < 0.001$), and engaged more in the parental feeding practices of positive involvement in child eating (df= 165, $t= 4.56$, $P < 0.001$) and use of food to control their child's behavior (df= 165, $t=-2.52$, $P = 0.01$).

Bivariate correlations between child sleep duration and all variables were examined. Mothers' positive involvement in child eating, use of pressure to eat and use of food to control behavior were correlated with longer sleep duration ($\rho= 0.12$, $P=0.04$; $\rho=0.27$, $P < 0.001$; $\rho= 0.23$, $P < 0.001$ respectively). Mothers' reports of child more often eating dinner with the TV on was correlated with shorter sleep duration ($\rho=-0.13$; $P= 0.05$). For fathers, pressure to eat and use of food to control behavior were correlated with longer sleep duration ($\rho= 0.23$, $P < 0.01$; $\rho= 0.16$, $P= 0.05$), but restriction of food amount was nearly-significantly related to shorter sleep duration ($\rho=-0.14$, $P= 0.07$). Fathers' reports that the

family had dinner together more often was correlated with shorter sleep duration ($\rho = -0.17$, $P = 0.03$).

Parent characteristics, including age, education, household income, marital status, single-parent household, foreign-born status, years in the U.S., hours worked per week, neighborhood disorder and victimization, and parent participation were not significant at the bivariate level. Family/home characteristics including number of TVs in home, TV in room where child sleeps, child eats breakfast with TV on were also not significant at the bivariate level. Variables not significant at the bivariate level and with a P value > 0.20 were not included in final analyses for parsimony. Child gender ($\rho = -0.21$, $P = 0.01$) and BMI z-score ($\rho = -0.17$, $P = 0.03$) given were significant at included in the final models.

Multivariable linear regression analyses

Mothers.—Using the full sample of mother-child pairs in Model 1 (Table 2), mothers' parenting styles, parenting practices, and family practices were examined with sleep duration. Pressure to eat and use of food to control behavior duration were associated with longer sleep duration ($\beta = 0.19$, $P = 0.01$; $\beta = 0.15$, $P = 0.03$, respectively). The child eating dinner with the TV on and valuing family dinners were associated with shorter sleep duration ($\beta = -0.15$, $P = 0.03$, $\beta = -0.18$, $P = 0.01$, respectively). Female gender ($\beta = -0.18$, $P = 0.04$) was significantly related to sleep duration while BMI z-score was not.

Fathers.—Using the subsample of mother-child pairs with fathers' participation in Model 2 (Table 2), mothers' and fathers' parenting styles and practices, and family practices were examined with sleep duration. Fathers' use of restriction of child's food amount was associated with shorter sleep duration ($\beta = -0.25$, $P = 0.03$). Mothers' report of valuing family dinners was associated with shorter sleep duration ($\beta = -0.24$, $P = 0.02$). Child gender and BMI z-score were not significantly related to sleep duration.

DISCUSSION

Research is needed to understand how to increase sufficient sleep duration among elementary school-age children. To our knowledge, this is the first study to examine mothers' and fathers' parenting styles and practices, and family/home practices in relation to children's weeknight sleep duration in Mexican American children. In analyses using the full sample of mother-child pairs, we found that mothers who pressured their child to eat and who used food to control behavior had children who had longer sleep duration. Eating dinner with the TV on and valuing family dinners were both related to shorter sleep duration. In the subsample that included both mothers and fathers, we found that fathers who restricted their child's food amount had children with shorter sleep duration. Findings suggest that use of pressure to eat and controlling behavior with food (parent directives), but not restriction and certain family practices, may positively impact children's sleep, and result in longer sleep duration.

We found that mothers' feeding practices previously found to be associated with children's weight status were also related to children's longer sleep duration (Tschann et al., 2015). We used parenting practices specific to child feeding as proxies for parenting in the context of

sleep, given that more direct measures of parenting practices regarding children's sleep were not available in the dataset used in this study. Nevertheless, it is likely that parenting in one domain is closely related to parenting in other domains, such as eating, screen time, and sleep (Jones et al., 2014). In the current study, mothers used parent-centered feeding directives that may be parallel to parent-centered directives regarding bedtime. Pressuring a child to eat is indicative of feeding a child even when the child is not hungry, and it is possible that mothers may also pressure children to sleep even when they are not willing or sleepy. Controlling behavior with food may also translate to controlling behavior with children's sleep, in that mothers may enforce child bedtime schedules and/or reward their child for going to bed. Future studies should assess parent-centered directives related to sleep and their role in improving children's sleep duration.

Given that fathers are not well represented in prior pediatric research, we examined fathers' parenting practices in addition to mothers' parenting practices in a subset of families. In families with both mothers' and fathers' participation, we did not find an association between mothers' parenting styles and practices and children's sleep duration. Rather we found that fathers who used a restrictive feeding practice had children with shorter sleep duration. Interestingly, this feeding practice has been shown to predict greater child weight over time in the same study sample (Tschann et al., 2015). It is possible that restrictive parenting practices, especially coming from fathers, in general lead to negative health outcomes. Interestingly, fathers' influence was significant over and beyond that of mothers' parenting practices. Similarly, Tavassolie et al. found that when fathers were more strict and harsh than mothers, children experienced more behavior problems (Tavassolie et al., 2016). More research that includes fathers is needed to understand how differences between mothers' and fathers' parenting may influence children's sleep.

Several studies focusing specifically on parenting practices and children's sleep have found bedtime reading to children at bedtime (Brown et al., 2016) and instilling bedtime rules to be protective of longer sleep in preschool age children (Jones et al., 2014). In Finnish 10–11 year olds, parents who practiced more rules around eating (e.g., eating at the table), bedtime, and screen time limits had children who slept longer (Ray et al., 2013). Similarly, we found several findings suggesting that family dinnertime practices are related to children's sleep duration. First, we found that mothers who valued eating dinner as a family had children with shorter sleep duration, and this was true for all mother-child pairs as well as in the subsample of families with both mother and father participation. This finding was unexpected, given previous research reporting that family meals as part of children's daily routines are related to longer sleep duration, less screen time, and higher fruit and vegetable intake (Ray and Roos, 2012). In focus groups of Latino preschoolers, we found that children wait up for their parent to get home when they work late schedules (Martinez and Thompson-Lastad, 2015). In the full study sample, only 5% of mothers reported having a late night/swing work shift. Therefore, it is possible that mothers/fathers arrive home past the typical dinnertime for children due to their work schedules and commute, which would therefore result in late night dinners that delay bedtime. It is also possible that some children stay up to spend time (e.g., watching TV and/or eating) with the entire family when everyone is home.

Our second finding was that children who ate dinner in a room with a TV on had shorter sleep duration. Combining family meals with screen time potentially results from hectic family schedules, which has resulted in a family-oriented ritual that may extend late into the evening. Fiese and colleagues (2012) found that what matters more during mealtime is the time spent gathered together engaged in positive forms of communication among families with healthy weight children compared to overweight or obese children (Fiese et al., 2012). The concept of TV co-viewing as “family time” is a practice that has been described among Latino families (Thompson et al., 2015), and is likely during dinner. Research has also found that more TV time is related to shorter sleep duration in six to seven year olds (Peña et al., 2016). An additional consideration is that late night exposure to blue-light emitting devices (LED) light from TV viewing may delay sleep onset and impact sleep duration and quality (Cajochen et al., 2011). In the current study, we were unable to assess whether TV light exposure while eating dinner was bright enough or close enough to bedtime that it could impact children’s sleep duration. Future research could help to provide light on the effect of LED light exposure during dinner or evening hours on sleep quantity and quality. In the current study, we were unable to assess whether TV light exposure while eating dinner was bright enough or close enough to bedtime that it could impact children’s sleep duration. Future research could help to provide light on the effect of LED light exposure during dinner or evening hours on sleep quantity and quality. Overall, research is needed to better understand whether parental schedules and family routines impact bedtime routines and sleep hygiene in children.

We did not detect significant associations between parenting styles and children’s sleep duration. It remains unclear whether general parenting styles, such as warmth and control, are related to sleep duration among Latino children. A strong emphasis on the cultural values of *familismo* (emphasizes loyalty and solidarity among family members) and *respeto* (emphasizes obedience to authority, deference, and decorum) has been associated with high levels of parental monitoring, the expectation of obedience, and the perceived need to use harsh parenting to teach children these cultural values (Romero and Ruiz, 2007). We did not capture these cultural values in relation to parenting styles. Future studies should consider including such constructs to increase understanding of parenting styles and their role in the development of children’s sleep-related practices.

At the bivariate level, we did not find number of TVs in home, TV in room where child sleeps, or child eats breakfast with TV on to be associated with children’s sleep duration. Interestingly, 70% of children had TVs in the bedroom, however, the average number of TVs in the home was 3 meaning that children might not be watching TV in bed. Lastly, neighborhood environmental factors that were not associated with children’s sleep duration included neighborhood disorder and victimization based on parents’ perceptions. Perhaps neighborhood context should be assessed from a child’s perspective.

A strength of this study is that we examined parenting practices among both mothers and fathers in relation to children’s sleep duration. To date, fathers have been underrepresented in research on health behaviors related to child obesity. Another strength of this research is that we used accelerometer-estimated weeknight sleep duration, which has been used in prior research on children’s sleep duration and is more reliable than parent report (Martinez

et al., 2014a). A limitation of this study is that we did not assess parental rules regarding bedtime or any other child behaviors besides sleep duration. Additionally, this study was conducted in a Mexican American sample of 8-to 10-year-old children, which limits the generalizability of findings to other ages and ethnic/cultural groups. Lastly, this study was cross-sectional and causality cannot be addressed.

Conclusions

The current study suggests that when considering only mothers' behaviors, mothers' pressure to eat, use of food to control behavior, child eating dinner with the TV on, and valuing family dinners may play a role in children's weeknight sleep duration among Mexican American families. Among children with both mothers' and fathers' study participation, fathers' restriction of food was associated with shorter sleep duration. These findings suggest that family context as a whole matters concerning children's sleep duration, which is also in accord with findings from other studies focusing on obesity in Latino children (Berge and Everts, 2011). Future research should examine the role of fathers and parental schedules in relation to children's sleep. Furthermore, interventions should consider tapping into domains of parenting related to family practices and routines to improve children's sleep quantity and quality as a strategy to prevent pediatric obesity.

ACKNOWLEDGEMENTS

This research was supported by grants from the National Heart, Lung and Blood Institute: R01 HL084404 awarded to J.M. Tschann and K01 HL129087 awarded to S.M. Martinez. We thank Jennifer Cho, Irene Takahashi, and the Kaiser Foundation Research Institute, which provided access to members of Kaiser. We would like to extend thanks to the UCSF Clinical & Translational Science Institute K Scholars Program and the BSM PRIDE Program (R25HL105444-08) for providing additional support.

Funding source:

This study was funded by R01HL084404 (PI: Tschann) and K01HL129087 (PI: Martinez).

REFERENCES

- American Academy of Sleep Medicine. Recharge with sleep: Pediatric sleep recommendations promoting optimal health. Available at <http://www.aasmnet.org/articles.aspx?id=6326> (accessed 13 6 2016).
- Aunola K and Nurmi JE Maternal affection moderates the impact of psychological control on a child's mathematical performance. *Dev. Psychol*, 2004, 40: 965–78. [PubMed: 15535751]
- Berge JM and Everts JC Family-based interventions targeting childhood obesity: A meta-analysis. *Childhood Obes.*, 2011, 7: 110–21.
- Brown SJ, Rhee KE and Gahagan S Reading at bedtime associated with longer nighttime sleep in Latino preschoolers. *Clin. Pediatr*, 2016, 55: 525–31.
- Butte NF, Gregorich SE, Tschann JM et al. Longitudinal effects of parental, child and neighborhood factors on moderate-vigorous physical activity and sedentary time in Latino children. *Int. J. Behav. Nutr. Phys. Act*, 2014, 11: 108. [PubMed: 25186810]
- Cajochen C, Frey S, Anders D et al. Evening exposure to a light-emitting diodes (LED)-backlit computer screen affects circadian physiology and cognitive performance. *J. Appl. Physiol*, 2011, 110: 1432–38. [PubMed: 21415172]
- Centers for Disease Control and Prevention Clinical Growth Charts. Available at http://www.cdc.gov/growthcharts/cdc_charts.htm (accessed 15 1 2013).

- Coulton CJ, Korbin JE and Su M Measuring neighborhood context for young children in an urban area. *Am. J. Community Psychol*, 1996, 24: 5.
- Darling N and Steinberg L Parenting style as context: An integrative model. *Psychol Bull.*, 1993, 113: 487–96.
- Fiese BH, Hammons A and Grigsby-Toussaint D Family mealtimes: A contextual approach to understanding childhood obesity. *Econ. Hum. Biol.*, 2012, 10: 365–74. [PubMed: 22652025]
- Finucane MM, Stevens GA, Cowan M et al. National, regional, and global trends in body mass index since 1980: Systematic analysis of health examination surveys and epidemiological studies with 960 country-years and 9.1 million participants. *Lancet*, 2011, 377: 557–67. [PubMed: 21295846]
- Ford ES, Cunningham TJ and Croft JB Trends in self-reported sleep duration among US adults from 1985 to 2012. *Sleep*, 2015, 38: 829–32. [PubMed: 25669182]
- Haines J, McDonald J, O'Brien A et al. Healthy Habits, Happy Homes Randomized Trial to improve household routines for obesity prevention among preschool-aged children. *JAMA Pediatr*, 2013, 167: 1072–79. [PubMed: 24019074]
- Hamre R, Kuester S, Renaud J et al. Improving nutrition, physical activity and obesity prevention: Performance report of the Nutrition and Physical Activity Program to prevent obesity and other chronic diseases: July 1 through December 31, 2005. Government Printing Office, Washington, DC, 2006.
- Hawkins SS and Takeuchi DT Social determinants of inadequate sleep in US children and adolescents. *Public Health*, 2016, 138: 119–26. [PubMed: 27207726]
- Jones CH, Pollard TM, Summerbell CD and Ball H Could parental rules play a role in the association between short sleep and obesity in young children?. *Biosoc. Sci.*, 2014, 46: 405–18.
- Koulouglioti C, Cole R, Moskow M, Mcquillan B, Carno M-A and Grape A The longitudinal association of young children's everyday routines to sleep duration. *J. Pediatr. Health Care*, 28: 80–87. [PubMed: 23360666]
- Lohman TG, Roche AF and Martorell R Anthropometric standardization reference manual. Human Kinetics Books, Champaign, IL, 1989.
- Martinez S, Greenspan L, Tschann J et al. Mother-reported sleep, accelerometer-estimated sleep, and weight status in Mexican American children: Is parent-reported sleep reliable? *J. Sleep Res*, 2014a, 23: 326–34. [PubMed: 24329818]
- Martinez S, Tschann J, Greenspan L et al. Is it time for bed? Short sleep duration increases risk for obesity in Mexican American children. *Sleep Medicine*, 2014b, 15: 1484–89. [PubMed: 25454984]
- Martinez SM and Thompson-Lastad A Latino parents' insight on optimal sleep for their preschool-age child: Does context matter? *Acad Pediatr*, 2015, 15: 636–43. [PubMed: 26547544]
- Musher-Eizenman D and Holub S Comprehensive feeding practices questionnaire: validation of a new measure of parental feeding practices. *J. Pediatr. Psychol*, 2007, 32
- Peña MM, Rifas-Shiman SL, Gillman MW, Redline S and Taveras EM Racial/ethnic and socio-contextual correlates of chronic sleep curtailment in childhood. *Sleep*, 2016, 39: 1653–61. [PubMed: 27306269]
- Ray C, Kalland M, Lehto R and Roos E Does parental warmth and responsiveness moderate the associations between parenting practices and children's health-related behaviors? *J. Nutr. Educ. Behav*, 2013, 45: 602–10. [PubMed: 23850012]
- Ray C and Roos E Family characteristics predicting favourable changes in 10 and 11-year-old children's lifestyle-related health behaviours during an 18-month follow-up. *Appetite*, 2012, 58: 326–32. [PubMed: 22056489]
- Robinson TN Reducing children's television viewing to prevent obesity: A randomized controlled trial. *JAMA*, 1999, 282: 1561–67. [PubMed: 10546696]
- Roblyer MIZ and Grzywacz JG Demographic and Parenting Correlates of Adolescent Sleep Functioning. *J. Child Fam. Stud*, 2015, 24: 3331–40. [PubMed: 26543343]
- Romero AJ and Ruiz M Does familism lead to increased parental monitoring?: Protective factors for coping with risky behaviors. *J. Child Fam. Studies*, 2007, 16: 143–54.

- Shloim N, Edelson LR, Martin N and Hetherington MM Parenting styles, feeding styles, feeding practices, and weight status in 4–12 year-old children: A systematic review of the literature. *Front. Psychol*, 2015, 6: 1849. [PubMed: 26696920]
- Skinner A and Skelton JA Prevalence and trends in obesity and severe obesity among children in the united states, 1999–2012. *JAMA Pediatrics*, 2014
- Spilsbury JC, Storfer-Isser A, Drotar D, Rosen CL, Kirchner HL and Redline S Effects of the home environment on school-aged children’s sleep. *Sleep*, 2005, 28: 1419–27. [PubMed: 16335483]
- Stallings VA and Fung EB Clinical nutritional assessment of infants and children In: Shils ME, Olson JA, Shike M and Ross AC (Eds), *Modern nutrition in health and disease*. Lippincott, Williams & Wilkins, Philadelphia, 1999: 885–93.
- Tavassolie T, Dudding S, Madigan AL, Thorvardarson E and Winsler A Differences in perceived parenting style between mothers and fathers: Implications for child outcomes and marital conflict. *J. Child Fam. Studies*, 2016, 25: 2055–68.
- Thompson DA, Polk S, Cheah CSL et al. Maternal Beliefs and Parenting Practices Regarding Their Preschool Child’s Television Viewing: An Exploration in a Sample of Low-Income Mexican-Origin Mothers. *Clin. Pediatr*, 2015, 54: 862–70.
- Troxel WM, Lee L, Hall M and Matthews KA Single-parent family structure and sleep problems in black and white adolescents. *Sleep Med*, 2014, 15: 255–61. [PubMed: 24424100]
- Tschann JM, Gregorich SE, Penilla C et al. Parental feeding practices in Mexican American families: initial test of an expanded measure. *Int. J. Behav. Nutr. Phys. Act*, 2013, 10: 1–11. [PubMed: 23281722]
- Tschann JM, Martinez SM, Penilla C et al. Parental feeding practices and child weight status in Mexican American families: a longitudinal analysis. *Int. J. Behav. Nutr. Phys. Act*, 2015, 12: 66. [PubMed: 25986057]
- Us Department of Health and Human Services Strategic Plan for NIH Obesity Research. A Report of the NIH Obesity Research Task Force. National Institutes of Health, Rockville, MD, 2006.
- Yoong SL, Chai LK, Williams CM, Wiggers J, Finch M and Wolfenden L Systematic review and meta-analysis of interventions targeting sleep and their impact on child body mass index, diet, and physical activity. *Obesity*, 2016, 24: 1140–47 [PubMed: 27112069]

Table 1.

Mother (n=308), father (n=166) and child characteristics of Mexican American family participants in the San Francisco Bay Area (CA) .

	Mothers	Fathers
Child characteristics	Mean (SD) or %	
Female	53%	50%
Age	8.9 (0.8)	9.6 (0.8)
BMI z-score	0.97 (1.01)	0.92 (1.0)
BMI 85 th %ile	21%	19%
BMI 95 th %ile	30%	27%
Weeknight sleep duration	9.5 (0.8)	9.5 (0.8)
Parent characteristics		
Age *	37.3 (6.3)	39.8 (6.8)
Education (years)	11.4 (3.7)	11.0 (3.7)
Married/living with a partner (n=247) ^A	80%	
Mother-only participation (n=94)	31%	
Mother and father participation (n=153)	50%	
Single parent household (n=57) ^A	19%	
Mother-only participation (n=46)	15%	
Mother and father participation (n=11)	4%	
Annual household income		
\$40,000	41%	26%
\$40,001–70,000	34%	39%
\$70,001	26%	36%
Employed	91%	100%
Hours work per week	34.3 (12.2)	42.4 (9.3)
Years in the U.S. *	17.2 (7.3)	20.4 (7.6)
Born in Mexico	78%	74%
Parenting style and practices		
Warmth ^B	1.1 (0.5)	1.0 (1.1)
Control ^B	0.7 (0.8)	0.6 (0.6)
Positive involvement in eating ^C	3.4 (0.6)	3.1 (0.7)
Pressure to eat ^C	2.3 (0.9)	2.4 (0.9)
Use of food to control behavior ^C	1.5 (0.5)	1.6 (0.5)
Restriction of food amount ^C	2.3 (0.4)	2.3 (0.5)
Monitors screen time ^C	3.2 (0.9)	3.1 (1.0)
Family practices regarding food and TV		
Number of TVs in home	2.9 (1.2)	2.9 (1.2)
TV in room where child sleeps	70%	66%

	Mothers	Fathers
Child characteristics	Mean (SD) or %	
Child eats breakfast with TV on (d/wk)	1.8 (1.2)	1.7 (1.2)
Child eats dinner with TV on (d/wk)	1.8 (1.2)	1.7 (1.1)
Frequency of shared family dinners ^D	5.2 (1.9)	5.1 (2.0)
Values family dinners	76%	80%
Neighborhood environment		
Disorder ^E	1.5 (0.6)	1.6 (0.6)
Victimization ^E	2.4 (1.1)	2.3 (1.0)

na (not applicable): Fathers in this sample were the spouse of mother-child pair participants.

* T-test of differences between mothers and fathers significant at $P < 0.05$.

^A Denominator for percentages is 308; 4 families did not reply, therefore, percentages do not total 100%.

^B Higher score means more use of that parenting style, ranging from -2 to 2.

^C Higher score means more use of that feeding practice, ranging from 1 to 5.

^D Score ranging from 1 to 7 days.

^E Higher score means more perceived disorder or victimization, ranging from 1 to 5.

Table 2.

Multiple variable linear regression models examining parenting styles and practices, and family practices with sleep duration in Mexican American children: Model 1 includes the full sample of mother-child pairs and Model 2 includes a subsample of children with mothers' and fathers' participation (values are standardized regression coefficients).

	Child sleep duration (hours)	
	Model 1: Mothers only (n=308)	Model 2: Mothers + Fathers (n=166)
Mothers' styles and practices		
Warmth	-0.04	-0.11
Control	0.01	<0.01
Positive involvement	-0.07	-0.11
Pressure to eat	0.19*	0.21
Control behavior with food	0.15*	0.18
Restriction of food amount	0.08	0.10
Monitors screen time	0.03	0.06
Fathers' styles and practices		
Warmth		-0.15
Control		0.15
Positive involvement		0.03
Pressure to eat		0.07
Control behavior with food		0.08
Restriction of food amount		-0.25*
Monitors screen time		-0.07
Family practices regarding TV and dinner		
Child eats dinner with the TV on (mother-reported)	-0.15*	-0.03
Frequency of shared family dinners (mother reported)	<0.01	-0.14
Values family dinners (mother reported)	-0.18*	-0.24*
Child eats dinner with the TV on (father-reported)		-0.05
Frequency of shared family dinners (father reported)		-0.05
Values family dinners (father-reported)		0.09
R²	0.10	0.13

Note. Models are adjusted for child gender and child BMI z-score.

* Significant at $P < 0.05$; ** $P < 0.01$