

# UC Davis

## UC Davis Previously Published Works

### Title

The Glucosinolate Biosynthetic Gene AOP2 Mediates Feed-back Regulation of Jasmonic Acid Signaling in Arabidopsis

### Permalink

<https://escholarship.org/uc/item/6055q4b5>

### Journal

Molecular Plant, 8(8)

### ISSN

1674-2052

### Authors

Burow, Meike  
Atwell, Susanna  
Francisco, Marta  
et al.

### Publication Date

2015-08-01

### DOI

10.1016/j.molp.2015.03.001

Peer reviewed

# The Glucosinolate Biosynthetic Gene *AOP2* Mediates Feed-back Regulation of Jasmonic Acid Signaling in *Arabidopsis*

Meike Burow<sup>1,2,\*</sup>, Susanna Atwell<sup>3</sup>, Marta Francisco<sup>3,4</sup>, Rachel E. Kerwin<sup>3</sup>, Barbara A. Halkier<sup>1,2</sup> and Daniel J. Kliebenstein<sup>1,3</sup>

<sup>1</sup>DynaMo Center of Excellence, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

<sup>2</sup>Copenhagen Plant Science Centre, Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Thorvaldsensvej 40, 1871 Frederiksberg C, Denmark

<sup>3</sup>Department of Plant Sciences, University of California at Davis, Davis, CA 95616, USA

<sup>4</sup>Misión Biológica de Galicia, (MBG-CSIC), PO Box 28, 36080 Pontevedra, Spain

\*Correspondence: Meike Burow ([mbu@plen.ku.dk](mailto:mbu@plen.ku.dk))

<http://dx.doi.org/10.1016/j.molp.2015.03.001>

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## ABSTRACT

Survival in changing and challenging environments requires an organism to efficiently obtain and use its resources. Due to their sessile nature, it is particularly critical for plants to dynamically optimize their metabolism. In plant primary metabolism, metabolic fine-tuning involves feed-back mechanisms whereby the output of a pathway controls its input to generate a precise and robust response to environmental changes. By contrast, few studies have addressed the potential for feed-back regulation of secondary metabolism. In *Arabidopsis*, accumulation of the defense compounds glucosinolates has previously been linked to genetic variation in the glucosinolate biosynthetic gene *AOP2*. *AOP2* expression can increase the transcript levels of two known regulators (*MYB28* and *MYB29*) of the pathway, suggesting that *AOP2* plays a role in positive feed-back regulation controlling glucosinolate biosynthesis. We generated mutants affecting *AOP2*, *MYB28/29*, or both. Transcriptome analysis of these mutants identified a so far unrecognized link between *AOP2* and jasmonic acid (JA) signaling independent of *MYB28* and *MYB29*. Thus, *AOP2* is part of a regulatory feed-back loop linking glucosinolate biosynthesis and JA signaling and thereby allows the glucosinolate pathway to influence JA sensitivity. The discovery of this regulatory feed-back loop provides insight into how plants optimize the use of resources for defensive metabolites.

**Keywords:** feed-back regulation, jasmonates, JA signaling, glucosinolates, *Arabidopsis*

Burow M., Atwell S., Francisco M., Kerwin R.E., Halkier B.A., and Kliebenstein D.J. (2015). The Glucosinolate Biosynthetic Gene *AOP2* Mediates Feed-back Regulation of Jasmonic Acid Signaling in *Arabidopsis*. *Mol. Plant*. **8**, 1201–1212.

## INTRODUCTION

All organisms must coordinate a phenomenal array of highly interconnected metabolic processes to efficiently obtain and use resources. This coordination of metabolism is particularly critical to maximize a plant's survival in dynamically changing environments. A key component in the regulation of primary plant metabolism is the fine-tuning provided by feed-back mechanisms (Grant, 2006; de Kraker et al., 2007; Smith and Stitt, 2007; Kauss et al., 2012). These feed-back regulatory processes whereby the output controls a system's input are essential to generate a precise and robust response within

any interconnected process (Freeman, 2000; Yi et al., 2000; Orrell et al., 2006; Lankau, 2007). In contrast, secondary metabolism is predominantly viewed to be under forward transcriptional regulation with minimal analysis of potential feed-back regulation. As secondary metabolites are biosynthesized from primary metabolites, feed-back regulation from secondary to primary metabolisms is likely to play an important role in the optimization of plant metabolism. A recent

## Molecular Plant

analysis within polyphenol metabolism in *Arabidopsis* has suggested the potential for plants to use unknown feed-back regulatory processes to coordinate growth and lignin deposition (Bonawitz et al., 2014).

Key defensive secondary metabolites within *Arabidopsis* are the amino acid-derived glucosinolates, produced predominantly from tryptophan (indole glucosinolates) and chain-elongated methionine (short- or long-chained aliphatic glucosinolates) (Sønderby et al., 2010b). These compounds help to defend *Arabidopsis* against both insect and microbial attackers (Chew, 1988; Brader et al., 2001; Stotz et al., 2001; Tierens et al., 2001; Hopkins et al., 2009). Although all classes of glucosinolates are constitutively present in all *Arabidopsis* tissues (Petersen et al., 2002; Brown et al., 2003), higher levels can be induced by herbivores (Mewis et al., 2005, 2006; Kim and Jander, 2007; Bidart-Bouzat and Kliebenstein, 2008) or upon treatment with jasmonates (Brader et al., 2001; Kliebenstein et al., 2002; Mikkelsen et al., 2003; Jost et al., 2005). Recent studies have linked this constitutive and inductive transcriptional regulatory control to a clade of six R2R3 domain MYB transcription factors and the three basic helix-loop-helix (bHLH) transcription factors MYC2 (bHLH006), MYC3 (bHLH005), and MYC4 (bHLH004) (Gigolashvili et al., 2007a, 2007b; Hirai et al., 2007; Sønderby et al., 2007; Gigolashvili et al., 2008b; Sønderby et al., 2010a; Schweizer et al., 2013; Frerigmann and Gigolashvili, 2014).

In contrast to the beneficial contribution to biotic defense, the accumulation of the amino acid-derived defense compounds glucosinolates has been shown to be metabolically costly using both modelling and empirical studies (Zust et al., 2011; Bekaert et al., 2012). Thus, higher levels of glucosinolates can have both beneficial and detrimental effects on plant fitness (Giamoustaris and Mithen, 1995; Mauricio, 1998; Lankau, 2007; Bidart-Bouzat and Kliebenstein, 2008). These contradictory effects require that the accumulation of these chemical defenses be tightly controlled. Within the modelling and empirical studies, the modelling studies predicted a much larger growth cost of glucosinolate accumulation than was empirically measured in defined mutants lacking glucosinolates (Giamoustaris and Mithen, 1995; Mauricio, 1998; Lankau, 2007; Bidart-Bouzat and Kliebenstein, 2008). One explanation for this contradiction is that *in vivo* glucosinolate accumulation may be regulated differently than implied in the modelling studies. Because the flux model was built to mimic glucosinolate levels, it inherently mimics the transcriptional regulation of the glucosinolate pathway under steady state conditions. The difference from the empirical studies thus indicates that there may be other regulatory processes such as feed-back regulation that can alleviate the costs of glucosinolate production under basal conditions.

Another line of evidence suggesting a potential for feed-back regulation in the glucosinolate pathway came from studies of naturally occurring *Arabidopsis* accessions that differ in their aliphatic glucosinolate structures. This glucosinolate profile variation can be partially explained by genetic variation in the *GS-AOP* locus encoding the two 2-oxo acid-dependent dioxygenases AOP2 and AOP3 (Kliebenstein et al., 2001b). AOP2 and AOP3 convert short-chained aliphatic methylsulfinylalkylglucosinolates to alkenyl- and hydroxyalkylglucosinolates, respectively. Genetic variation at AOP2 is also linked to increased glucosino-

## Feed-back Regulation of Glucosinolate Biosynthesis

late accumulation (Kliebenstein et al., 2001a; Wentzell et al., 2007). Furthermore, expression of AOP2 from *Brassica oleracea* in the *Arabidopsis* accession Col-0, which lacks a functional AOP2, results not only in the accumulation of alkenyl glucosinolates, but also in a two-fold increase in total aliphatic glucosinolate content (Li and Quiros, 2003; Wentzell et al., 2007). Analysis of transcriptome changes in these AOP2 expressing lines indicated that AOP2 positively regulates glucosinolate accumulation by increasing the transcript levels of most biosynthetic genes in the pathway and two known transcription factors, i.e. the R2R3 domain MYB transcription factors MYB28 and MYB29 (Wentzell et al., 2007). Based on these findings, AOP2 has been suggested to play a role in a positive feed-back loop controlling aliphatic glucosinolate biosynthesis via an unknown mechanism (Wentzell et al., 2007; Burou et al., 2010).


With the aim of investigating the regulatory role of AOP2 in aliphatic glucosinolate biosynthesis and deciphering its interplay with MYB28 and MYB29 in the glucosinolate regulatory network, we introduced the AOP2 from *B. oleracea* into different *myb* knockout mutants in the Col-0 background. We show that the AOP2 feed-back mechanism for aliphatic glucosinolate accumulation depends on the presence of MYB28 and MYB29. Transcriptome analysis of mutants affecting AOP2, MYB28/29, or both identified an unrecognized link between AOP2 and jasmonic acid (JA) signaling that was independent of MYB28 and MYB29. Using AOP2 expression lines, we could show that the introduction of this gene into *Arabidopsis* leads to an altered sensitivity to exogenous JA. Our findings describe AOP2 as a biosynthetic gene from a JA-inducible defense pathway that feed-back regulates JA signaling and thereby its own pathway and potentially links to other pathways controlled by JA.

## RESULTS

### AOP2 Depends on MYB28 to Increase Levels of Aliphatic Glucosinolates

Expression of the enzymatically functional AOP2 from *B. oleracea* in *Arabidopsis* Col-0, a natural AOP2 knockout, increased transcript levels of the transcription factors MYB28 and MYB29 and glucosinolate accumulation (Wentzell et al., 2007). Based on this observation, we hypothesized that AOP2 increases glucosinolate levels via the MYB28 and MYB29 transcription factors. To test this, we crossed 35S:AOP2 plants to *myb28-1* and *myb29-2* single knockouts and analyzed leaf glucosinolate accumulation in homozygous progeny obtained from the same maternal lineage (Figure 1). Introduction of AOP2 to Col-0 led to a higher accumulation of short- and long-chained aliphatic glucosinolates. By contrast, no significant changes in aliphatic glucosinolate levels were detected upon AOP2 over-expression in the *myb28-1* knockout background indicating that AOP2 requires MYB28 to up-regulate the biosynthesis of both short- and long-chained aliphatic glucosinolates (Figure 1A and 1B).

While MYB28 is considered a major transcriptional activator of aliphatic glucosinolate biosynthesis, MYB29 has been described as playing a lesser role in regulating basal glucosinolate levels (Gigolashvili et al., 2008a; Sønderby et al., 2010a). Accordingly, AOP2 expression in the *myb29-2* knockout resulted in higher accumulation of short-chained aliphatic



**Figure 1. Effect of Constitutive AOP2 Expression in *myb28-1*, *myb29-2* Single Knockout Mutants, and in the *myb28-1 myb29-1* dko.**

**(A and B)** *myb28-1* single knockout mutants.

**(C and D)** *myb29-2* single knockout mutants.

**(E and F)** *myb28-1 myb29-1* dko.

Glucosinolates were extracted from rosette leaves of 21- to 24-day old plants and quantified as desulfo-glucosinolates by HPLC. Gray

glucosinolates compared with the *myb29-2* knockout and the wild-type (Figure 1C). Short-chained aliphatic glucosinolate over-accumulation upon AOP2 expression in the *myb29-2* knockout was significantly lower than in the wild-type background. MYB29 therefore contributes to the regulatory role of AOP2 but with a lower magnitude than MYB28. Wild-type and *myb29-2* knockout controls segregated out of this cross were found to accumulate relatively high levels of long-chained aliphatic glucosinolates, which were not found to be further increased by AOP2 expression (Figure 1D).


### The Effects of AOP2 and MYB28 on Short-Chained Aliphatic Glucosinolates Are Additive

The elevated levels of aliphatic glucosinolates in plants over-expressing either AOP2 or MYB28 have been shown to be accompanied by increased transcript levels of the majority of the biosynthetic genes in the pathway (Sønderby et al., 2007; Wentzell et al., 2007). This suggests that the positive regulatory effect of AOP2 on aliphatic glucosinolate biosynthesis could be explained by up-regulation of MYB28, which then transcriptionally up-regulates the biosynthetic genes. To further investigate the interplay between AOP2 and MYB28 in the regulation of glucosinolate biosynthesis, we generated lines over-expressing both genes. Analysis of leaf glucosinolates revealed that over-expression of both genes leads to higher short-chained aliphatic glucosinolate accumulation than each transgene alone (Figure 2). In this set of experiments, there were no observed effects of AOP2 and MYB28 on long-chained aliphatic glucosinolates suggesting that this phenotype is more environmentally sensitive (Figure 2). Together with the knockout analysis, the data show that AOP2 acts upstream of MYB28 to regulate short-chain aliphatic glucosinolate accumulation. Furthermore, wild-type levels of MYB28 are a limiting factor in the ability of AOP2 to increase the accumulation of aliphatic glucosinolates.

### AOP2 Cannot Trigger Aliphatic Glucosinolate Accumulation in the Absence of Both MYB28 and MYB29

While *myb28 myb29* double knockout plants have been shown to be devoid of aliphatic glucosinolates (Sønderby et al., 2007; Beekwilder et al., 2008), production of aliphatic glucosinolate is not strictly dependent on MYB28 and MYB29. Aliphatic glucosinolates can be induced in the *myb28 myb29* double knockout background either upon over-expression of MYB76 or upon feeding damage caused by *Mamestra brassicae* caterpillars (Beekwilder et al., 2008; Sønderby et al., 2010a). Thus, we introduced the full-length *B. oleracea* AOP2 to the *myb28-1 myb29-1* double knockout to test if AOP2 can act on regulatory factors that induce aliphatic glucosinolate biosynthesis independently of MYB28 and MYB29.

bars represent control plants not expressing AOP2, black bars 35S:AOP2 lines. Data are shown as means (+SE) obtained in at least two independent experiments ( $n = 12-47$ ). Different letters indicate significant differences between genotypes (ANOVA,  $P < 0.05$ , see Supplemental Tables 1, 2, and 4). Levels of individual glucosinolates can be found in Supplemental Tables 1, 2, and 4, short-chained aliphatic glucosinolates; LC, long-chained aliphatic glucosinolates; FW, fresh weight.



**Figure 2. Effect of Constitutive AOP2 Expression in 35S:MYB28 Plants.**

**(A)** Short-chained aliphatic glucosinolates (SC).

**(B)** Long-chained aliphatic glucosinolates (LC).

Glucosinolates were extracted from 23- to 25-day old plants and quantified as desulfo-glucosinolates by HPLC. Gray bars represent control plants not expressing AOP2, black bars 35S:AOP2 lines. Data are shown as means (+SE) obtained in two independent experiments ( $n = 13-36$ ). Different letters indicate significant differences between genotypes (ANOVA,  $P < 0.05$ , see Supplemental Table 3). Levels of individual glucosinolates can be found in Supplemental Table 3. FW, fresh weight.

As previously found, AOP2 over-expression in the wild-type background led to increased accumulation of short- and long-chained aliphatic glucosinolates as well as indole glucosinolates (Figure 1E and 1F). The effect on indole glucosinolates is less consistent suggesting that it is more environmentally dependent (Supplemental Table 4). In the absence of MYB28 and MYB29, AOP2 was unable to induce a change in glucosinolate accumulation. This agrees with the observation that AOP2 does not increase MYB76 transcript levels in the Col-0 background (Wentzell et al., 2007). Together, this shows that the AOP2-mediated increase in glucosinolate levels is completely dependent on the presence of a functional MYB28 or MYB29.

### AOP2 Causes Distinct Changes in Glucosinolate-Related Transcripts Independently of MYB28 and MYB29

To test for regulatory effects of AOP2 expression in the absence of both MYB28 and MYB29, we analyzed transcript profiles of wild-type, 35S:AOP2, *myb28-1 myb29-1* dko, and 35S:AOP2 in the *myb* dko background. Consistent with the glucosinolate profile of the *myb28-1 myb29-1* dko (Figure 3), transcript levels of nearly all genes involved in aliphatic glucosinolate biosynthesis were reduced compared with wild-type (Figure 3). In contrast to a previous study, which found a slight but consistent increase in most aliphatic glucosinolate biosynthesis genes (Wentzell et al., 2007), AOP2 expression led to altered transcript levels of a distinct subset of glucosinolate biosynthetic genes in the *myb* dko background in this experiment. This discrepancy between the two studies could be due to differences in the growth conditions. As glucosinolate biosynthesis undergoes diurnal

### Feed-back Regulation of Glucosinolate Biosynthesis

rhythm, the direct comparison of both data sets is moreover precluded because the experiments were harvested at different times of the day (Kerwin et al., 2011; Huseby et al., 2013). Both studies, however, agree that introduction of AOP2 into *Arabidopsis* alters the transcript accumulation of some aliphatic glucosinolate biosynthetic genes.

Remarkably, expression of AOP2 changed the expression levels of some genes in the pathway both in the presence (wild-type background) and the absence of MYB28 and MYB29 (*myb28-1 myb29-1* dko background). This is despite the absence of detectable aliphatic glucosinolates in any lines lacking both MYB28 and MYB29. For example, transcript levels of GGP1, CYP83B1, and SOT16 were up-regulated by AOP2 independently of MYB28 and MYB29 (Bak et al., 2001; Hansen et al., 2001; Naur et al., 2003; Piotrowski et al., 2004; Geu-Flores et al., 2011) (Figure 3). Individual genes involved in secondary modification of glucosinolate side chains, i.e. CYP81F4 and GS-OX2 (Li et al., 2008; Kai et al., 2011; Pfalz et al., 2011), showed reduced expression levels in 35S:AOP2 lines independent of MYB28 or MYB29. These five transcripts regulated by AOP2 in the absence of MYB28 and MYB29 were not changed in the *myb28-1 myb29-1* dko indicating that AOP2 can bypass MYB28 and MYB29 to regulate specific glucosinolate biosynthetic genes. As this occurs in the absence of detectable levels of aliphatic glucosinolates, this MYB28/29-independent regulatory function may not require the AOP2-catalyzed formation of alkenyl glucosinolate metabolites.


The finding that AOP2 feeds into regulatory networks that bypass MYB28 and MYB29 is further supported by the negative regulatory effect of AOP2 on genes involved in the enzymatic activation of glucosinolates upon tissue damage (Figure 3), i.e. myrosinase-binding protein 1 (MBP1) and nitrile-specifier proteins (NSP1, NSP3, NSP4) (Capella et al., 2001; Burow et al., 2009). In contrast, these four AOP2-regulated genes involved in glucosinolate activation are not regulated by MYB28 and MYB29. Thus, AOP2 must be affecting an additional regulatory network (Hirai et al., 2007; Sønderby et al., 2007, 2010a).

### AOP2 Feed-Back Regulates JA Biosynthesis and Signaling on the Transcriptional Level

One candidate for the additional regulatory network targeted by AOP2 came from an observed increase in transcript levels of MYC2 (*bHLH006*), known to regulate indole glucosinolate biosynthesis and other JA-responsive genes (Lorenzo et al., 2004; Dombrecht et al., 2007). This prompted us to further investigate the regulatory effect of AOP2 on JA biosynthesis and signaling. Twelve of 25 JA biosynthetic and metabolic genes showed AOP2-induced changes in expression level with LOX3, LOX4, AOC1, AOC3, OPR3, CYP94B3, and CYP94C1 (Sanders et al., 2000; Stintzi and Browse, 2000; Stenzel et al., 2003; Kandel et al., 2007; Koo et al., 2011; Chauvin et al., 2013) being most strongly up-regulated by AOP2 (Figure 4 and Supplemental Table 6). Likewise, AOP2 over-expression was sufficient to generate significantly higher transcript accumulation for eight members of the jasmonate ZIM-domain (JAZ) protein family that serve as key regulators of JA signaling (Chini et al., 2007; Thines et al., 2007). The MYB28/MYB29 genotype had no significant effect on the transcript accumulation of any JA-related gene



GLS BIOSYNTHESIS




**Figure 3. Transcript Levels of Genes Involved in Glucosinolate Biosynthesis, Activation, Regulation and Transport.**


Average transcript levels are shown for Col-0, *myb28-1 myb29-1* dko, and lines constitutively expressing *AOP2* in these two backgrounds. Different colors represent different transcript levels in log CPM from -12 (dark blue) to ≥9 (dark yellow). Underlined gene names depict transcripts altered in lines devoid of MYB28 and MYB29, while bold letters indicate changes in transcript levels in lines expressing *AOP2* (ANOVA, FDR-adjusted  $P < 0.05$ ). Red circles depict transcript levels significantly altered by *AOP2* independent of MYB28 and MYB29 ( $P > 0.05$  for *AOP2*-MYB interaction). Gene names and transcript levels are as listed in Supplemental Table 5. CPM, counts per million. GLS, glucosinolate.

## Molecular Plant



### JA BIOSYNTHESIS



### JA METABOLISM



### JA PERCEPTION




**Figure 4. Transcript Levels of Genes Involved in JA Biosynthesis, Metabolism, and Perception.**

Average transcript levels are shown for Col-0, *myb28-1 myb29-1* dko, and lines constitutively expressing *AOP2* in these two backgrounds (Col-0+AOP2 and *myb dko*+AOP2). Different colors represent different transcript levels in log CPM from  $-12$  (dark blue) to  $\geq 9$  (dark yellow). Underlined gene names depict transcripts altered in lines devoid of MYB28 and MYB29, while bold letters indicate changes in transcript levels in lines expressing *AOP2* (ANOVA, FDR-adjusted  $P < 0.05$ ). Red circles depict transcript levels significantly altered by *AOP2* independent of MYB28 and MYB29 ( $P > 0.05$  for AOP2–MYB interaction). Gene names and transcript levels are as listed in Supplemental Table 6. CPM, counts per million, JA, jasmonic acid.

showing that these effects are specific to *AOP2* and independent of *MYB28* and *MYB29*. This *AOP2*-to-JA link was found even in the absence of glucosinolate substrates for the *AOP2* enzyme (35S:*AOP2* vs. *myb28-1 myb29-1* dko + 35S:*AOP2*; Figure 4 and Supplemental Table 6). This suggests that the enzymatic activity within the glucosinolate pathway is not critical for *AOP2*'s ability to feed-back regulate JA biosynthesis and signaling components on the transcriptional level.

The effect of *AOP2* expression on JA pathway transcript accumulation led us to hypothesize that the genotypes over-expressing *AOP2* may have an altered sensitivity to JA. To test this hypothesis, we germinated seeds from wild-type, 35S:*AOP2*, *myb28-1 myb29-1* dko, and 35S:*AOP2*/dko on media with and without exogenous JA and measured the root length daily after 3–8 days. On control plates without JA, the roots of seedlings

## Feed-back Regulation of Glucosinolate Biosynthesis



**Figure 5. Root Growth of Wild-Type and 35S:*AOP2* Seedlings on JA.**

(A) Data are shown as means (+SE) obtained in three independent experiments. Solid lines, seedlings grown on MS plates containing JA (blue, wild-type, total  $n = 28$ ; red, 35S:*AOP2*, total  $n = 23$ ); dashed lines, seedlings grown on control plates without JA (ctr; blue, wild-type, total  $n = 24$ ; red, 35S:*AOP2*, total  $n = 22$ ).

(B) Root growth of 35S:*AOP2* seedlings shown relative to the wild-type on MS medium supplemented with JA (solid line) and MS control plates (dashed line). For relative root growth of *myb28-1 myb29-1* dko and the *myb28-1 myb29-1* dko+AOP2 seedlings, see Supplemental Figure 1 and Supplemental Table 7.

expressing *AOP2* in the Col-0 wild-type background were shorter (Figure 5A, dashed lines). Relative to wild-type seedlings on control plates, the *AOP2*-expressing seedlings were about 20%–25% shorter. Interestingly, when grown on JA, the root length of *AOP2*-expressing seedlings relative to the wild-type was increased (Figure 5B). These seedlings showed significantly decreased sensitivity to exogenous JA that was strongest on days 4 and 5 but significant at least to day 8, the end of the experiment. As predicted from the transcriptomics, the *MYB28*/*MYB29* genotype had no effect on sensitivity to JA in the growth media and no effect on the JA–*AOP2* interaction (Table 1). Despite the altered JA sensitivity in *AOP2*-expressing seedlings, JA levels were not significantly changed (Figure 6). Thus, *AOP2* leads to altered JA sensitivity in Col-0 seedlings by an unknown mechanism that is independent of *MYB28*/*MYB29* and thereby of glucosinolate precursor availability.

## DISCUSSION

Glucosinolate profiles change dynamically in response to internal and external signals (Petersen et al., 2002; Brown et al.,

Day	Plate	Treatment	AOP2	MYBs	AOP2×MYBs	Treatment×AOP2	Treatment×MYBs	Residuals	
Df	4	1	1	1	1	1	1	78	
3	SS	0.106	0.732	0.063	0.090	0.000	0.011	0.022	0.282
	F	7.328	202.340	17.287	24.790	0.066	2.975	6.215	
	P	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.798	0.089	<b>0.015</b>	
4	SS	0.421	4.302	0.153	0.184	0.000	0.105	0.041	0.886
	F	9.259	378.688	13.455	16.160	0.013	9.242	3.637	
	P	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.909	<b>0.003</b>	0.060	
5	SS	0.595	11.832	0.361	0.102	0.020	0.204	0.018	1.567
	F	7.400	589.072	17.979	5.095	0.997	10.163	0.880	
	P	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>0.027</b>	0.321	<b>0.002</b>	0.351	
6	SS	0.823	20.794	0.531	0.142	0.100	0.164	0.060	3.362
	F	4.772	482.461	12.308	3.298	2.315	3.796	1.393	
	P	<b>0.002</b>	<b>&lt;0.001</b>	<b>0.001</b>	0.073	0.132	0.055	0.242	
7	SS	1.159	34.267	0.798	0.156	0.270	0.287	0.137	3.949
	F	5.722	676.926	15.767	3.078	5.333	5.676	2.701	
	P	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>&lt;0.001</b>	0.083	<b>0.024</b>	<b>0.020</b>	0.104	
8	SS	1.931	56.100	0.915	0.328	0.280	0.319	0.279	6.071
	F	6.202	720.767	11.757	4.211	3.598	4.095	3.587	
	P	<b>&lt;0.001</b>	<b>&lt;0.001</b>	<b>0.001</b>	<b>0.044</b>	0.062	<b>0.046</b>	0.062	

**Table 1. Results of ANOVA Testing of Plate, Treatment ( $\pm$ JA), AOP2 (Presence and Absence of the AOP2 Transgene), MYBs (Presence and Absence of MYB28/29), AOP2×MYBs Interaction, Treatment×AOP2 Interaction, and Treatment×MYBs Interaction for Significant Impact on Root Growth.**

For means and SEs, see Supplemental Table 7. F, F value; P, the statistical significance of each term across the different days (numbers in bold indicate P values <0.05); SS, type III sums-of-squares.

2003); (Hirai et al., 2004; Falk et al., 2007; Hopkins et al., 2009; Huseby et al., 2013). The regulatory network controlling glucosinolate biosynthesis must therefore allow for integration of various signals, both internal and external. This fine-tuning of the pathway likely requires feed-back regulation. Previous findings had suggested that AOP2 mediates feed-back regulation of glucosinolate biosynthesis via MYB28 and MYB29 (Wentzell et al., 2007). In agreement with this hypothesis, expression of AOP2 from *B. oleracea* in the *myb28-1 myb29-1* dko did not result in accumulation of detectable levels of aliphatic glucosinolates (Figure 1). Thus, AOP2 functions to regulate glucosinolate accumulation in a feed-back loop that involves the MYB28/MYB29 transcription factors.

Transcriptomic analysis of plants expressing AOP2 in the *myb28-1 myb29-1* dko background revealed an additional regulatory role for AOP2 bypassing MYB28 and MYB29. Transcript levels of numerous genes involved in JA biosynthesis and signaling were increased by AOP2 irrespective of the MYB28/MYB29 genotype (Figure 4). Likewise, AOP2 alone altered the transcript levels of a distinct subset of genes involved in indole glucosinolate biosynthesis, side chain modification, and activation of glucosinolates (Figure 3). Most of these genes show strongly reduced transcript levels in the *myc234* triple knockout indicating that they are JA-responsive (Schweizer et al., 2013). Thus, AOP2 has a regulatory effect on JA-mediated glucosinolate biosynthesis different from the transcriptional regulators MYB28/29/76 and MYB51/34/122, which ascertain co-


ordinated expression of all genes required for aliphatic and indole glucosinolate biosynthesis, respectively (Sønderby et al., 2010a; Frerigmann and Gigolashvili, 2014). Instead, AOP2 can fine-tune glucosinolate biosynthesis and activation through feed-back regulation of JA signaling independently of MYB28 and MYB29.

One of the JA-related transcripts changed by AOP2 was the bHLH transcription factor MYC2, a master switch in the regulatory network coordinating JA responses and furthermore critical for the crosstalk with the signaling pathways of other phytohormones (Lorenzo et al., 2004; Dombrecht et al., 2007; Kazan and Manners, 2013). In concert with the two related transcription factors, MYC3 (*bHLH005*) and MYC4 (*bHLH004*), MYC2 has recently been shown to be indispensable for the formation of both indole and aliphatic glucosinolates (Schweizer et al., 2013; Frerigmann et al., 2014). The triple knockout mutant *myc234* is nearly devoid of glucosinolates and shows strongly reduced expression of glucosinolate biosynthetic genes including AOP2 (Schweizer et al., 2013; Frerigmann et al., 2014).

By contrast, AOP2 expression levels are unchanged in the *myb28-1 myb29-1* dko that accumulates only trace amounts of aliphatic glucosinolates and shows highly reduced transcript levels of the genes in that pathway (Sønderby et al., 2010a). Interestingly, both MYB28 and MYB29 were shown to bind the AOP2 promoter in a yeast-1-hybrid study, while MYB76, MYC2 and MYC4 did not (Li et al., 2014). Thus,



## Molecular Plant



**Figure 6. JA Levels in 4-day Old Seedlings Grown on MS Medium.**

Gray bars represent control plants not expressing *AOP2*, black bars 35S:*AOP2* lines. Data are shown as means (+SE) obtained in three independent experiments (total  $n = 9$ ). There were no significant differences between genotypes (ANOVA,  $P > 0.05$ ). FW, fresh weight.

*AOP2* expression might require the formation MYC-MYB protein complexes *in vivo*. In turn, *AOP2* feeds into the regulatory network controlling *MYC2*, *MYB28*, and *MYB29* and thereby establishes a positive feed-back loop in the JA-mediated regulation of glucosinolate biosynthesis mediated by *AOP2* (Figure 7). As *AOP2* encodes a cytoplasmic enzyme, it is unlikely that the *AOP2* protein directly or indirectly interferes with transcriptional gene activation in the nucleus. The direct target of *AOP2* as a regulator and the molecular mechanism underlying the regulatory function of this biosynthetic gene remain to be identified.

The observed changes in the transcript levels of genes involved in JA biosynthesis and signaling within the *AOP2* genotypes were accompanied by a decreased JA sensitivity in the *AOP2* plants (Figure 5). This effect of *AOP2* on root growth was independent of *MYB28* and *MYB29*. However, JA levels in the seedlings were not significantly altered (Figure 6) underlining the complexity of JA-mediated regulation *in vivo*. Nevertheless, feed-back regulation of JA signaling mediated by *AOP2* allows a gene within the glucosinolate biosynthetic pathway to influence JA sensitivity by an unknown mechanism. This shows that a downstream defense output, an enzyme in defense metabolism, can feed-back regulate the upstream signaling network that controls the defense output. It remains to be tested if this

## Feed-back Regulation of Glucosinolate Biosynthesis

feed-back regulation is a broader property of plant defense networks.


The observation that the *AOP2* gene can alter JA sensitivity and glucosinolate gene expression in the *myb28 myb29* dko suggests that this regulatory activity is independent of the known enzyme activity. One possibility is that *AOP2* has other, so far unknown enzymatic activities that mediate this regulatory function. There are, however, several arguments against this hypothesis. First, *AOP2* is a newly evolved enzyme that arose via a tandem duplication and neo-functionalization within the *Arabidopsis* clade that has only ever been linked to glucosinolate metabolism and is thus not a conserved gene likely to have any ancestral function remaining (Wentzell et al., 2007; Kliebenstein et al., 2001a). Second, in metabolomics studies of lines that vary for the presence of the *AOP2* gene, there were no metabolites that showed similar variation that were not glucosinolate or precursors (Rowe et al., 2008; Chan et al., 2011; Joseph et al., 2013). This leaves two other possible explanations. First, it is possible that the residual trace level of aliphatic glucosinolates in the *myb28 myb29* dko may be sufficient for the regulatory process. Alternatively, there may be as yet unrecognized regulatory functions of the *AOP2* RNA or protein that remain to be identified. Further work is necessary to address the specific regulatory mechanisms of the *AOP2* regulatory feed-back mechanisms.

As *AOP2* expression significantly varies between naturally occurring accessions of *Arabidopsis* (Kliebenstein et al., 2001a; Wentzell et al., 2007), this regulatory feed-back loop linking *AOP2* expression and JA signaling suggests natural variation in JA-mediated regulation of glucosinolates as well as in glucosinolate-mediated tuning of JA sensitivity. This variation in fine-tuning of glucosinolate biosynthesis allows for the higher accumulation of glucosinolates in *AOP2*-expressing *Arabidopsis* accessions (Kliebenstein et al., 2001a; Burow et al., 2010). Natural variation in *AOP2* may moreover reflect differences in the JA regulatory network among *Arabidopsis* accessions both with regard to the input to and the output from JA signaling. Future efforts will aim to elucidate the upstream mechanistic basis allowing *AOP2*-mediated feed-back regulation of JA signaling. Identification of the specific mechanistic link between *AOP2* and JA signaling will provide new insight into how biological systems integrate internal and external signals to generate a specific phenotypic output in dynamically changing environments.

## METHODS

### Plant Material and Cultivation

The double knockout mutant *myb28-1 myb29-1* (*myb28-1* = At5g61420, line SALK\_136312; *myb29-1* = At5g07690, line GABI\_868E02) and the *MYB28* over-expression lines (35S:*MYB28*) have been described previously (Sonderby et al., 2007, 2010a). To introduce expression of the enzymatically functional *AOP2* allele of the *B. oleracea* BoGSL-ALK under the control of the 35S CaMV promoter (Li and Quiros, 2003), two independent homozygous 35S:*AOP2* lines were crossed to the homozygous *myb28-1 myb29-1* dko and to two independent homozygous 35S:*MYB28* lines. The F1 plants were self-fertilized and progeny in the F2 generation was genotyped by PCR on genomic DNA. For all experiments, F3 seeds were sown in a randomized design and



**Figure 7. Proposed Regulatory Link between JA and Glucosinolate Biosynthesis.**

Herbivory results in increased JA biosynthesis, which triggers increased degradation of JAZ proteins by the 26S proteasome and increased formation of protein complexes comprising MYC (MYC2, MYC3, MYC4) and MYB (MYB28, MYB29, MYB76, MYB34, MYB51, MYB122) transcription factors. Increased expression of glucosinolate pathway genes leads to higher glucosinolate levels. The glucosinolate biosynthetic gene *AOP2* mediates feedback regulation by an unknown mechanism. GLS, glucosinolate; JA, jasmonic acid.

cold stratified at 4°C for at least 2 days before being moved to growth chambers. Plants were grown at 80–120  $\mu\text{E}/(\text{m}^2 \text{ s})$ , 16 h light, 21°C, 70% relative humidity.

### Glucosinolate Analysis

Glucosinolate were analyzed as desulfo-glucosinolates as previously described (Kliebenstein et al., 2001b). 96-well filter plates were charged with 45 mg of DEAE Sephadex A25 and 300  $\mu\text{l}$  of water per well and equilibrated at room temperature for at least 2 h. The water was removed using a vacuum manifold (Millipore). Plant material was harvested in 300  $\mu\text{l}$  of 85% MeOH (v/v) containing 5 nmol *p*-OH-benzyl glucosinolate extracted from seeds of *Sinapis alba* (SeedCom A/S, Vissenbjerg, Denmark) as previously described (Thies, 1979; Zrybko et al., 1997) as an internal standard. The tissue was homogenized with one stainless steel ball by shaking for 2 min at a frequency of 30/s on a Mixer Mill 303 (Retsch, Haan, Germany), centrifuged, and the supernatant was applied to the filter plates and absorbed on the ion exchanger by vacuum filtration for 2–4 s. Sephadex material was washed with 2  $\times$  100 ml of 70% methanol (v/v) and 2  $\times$  100  $\mu\text{l}$  of water and briefly centrifuged before addition of 20  $\mu\text{l}$  of sulfatase solution (1.25 mg/ml, sulfatase type H1, Sigma-Aldrich) on each filter. After incubation at room temperature overnight, desulfo-glucosinolates were eluted with 100  $\mu\text{l}$  of water for 96-well filter plates and 250  $\mu\text{l}$  for media samples. Media sample elute was lyophilized and resuspended in 50  $\mu\text{l}$  of water. All samples were analyzed by HPLC on an Agilent HP1200 Series instrument equipped with a C-18 reversed phase column (Zorbax SB-Aq, 25 cm  $\times$  4.6 mm, 5  $\mu\text{m}$  particle size, Agilent) or Lichrospher RP18-5 25 cm  $\times$  4.6 mm, 5  $\mu\text{m}$  particle size, Supelco) by using a water (solvent A)–acetonitrile (solvent B) gradient at a flow rate of 1 ml/min at 25°C (injection volume 45  $\mu\text{l}$ ). The gradient applied was as follows: 1.5%–7% B

(5 min), 7%–25% (6 min), 25%–80% (4 min), 80% B (3 min), 80%–35% B (2 min), 35%–1.5% B (2 min), and 1.5% B (3 min). The eluent was monitored by diode array detection between 200 and 400 nm (2-nm interval). Desulfo-glucosinolates were identified based on comparison of retention times and UV absorption spectra with those of known standards (Reichelt et al., 2002). Results are given as nmol/(mg fresh weight) calculated relative to response factors (Fiebig and Arens, 1992; Brown et al., 2003). ANOVA was used to test for significant differences.

### RNA Sequencing

To test for differential expression across the genotypes, they were planted in a randomized complete block design with three completely separate outgrowths. Two plants per genotype were used to make independent libraries per experiment providing six samples per genotype. RNA was isolated from individual plants and sequencing libraries created as previously described (Kumar et al., 2012).

The libraries were then sequenced at the Beijing Genome Institute. All reads were mapped against the *Arabidopsis thaliana* Ensembl (TAIR10) genome using TopHat v2.0.8 (Trapnell et al., 2009) and default settings. Mapped reads were counted using HTSeq (Anders et al., 2014) using the setting `-m intersection-nonempty`. Differential expression was analyzed with edgeR (Robinson et al., 2010) using a model that accounted for the genotype at *AOP2*, *MYB28/29*, and directly tested for an interaction of the genotypes. The model also included experiment to allow for this to be included in the statistical assessment. All *P* values were adjusted to a false discovery ratio (FDR) of 0.05 within edgeR using the factorial model and are presented along with the mean corrected counts per million per transcript per genotype (Supplemental Table 8).

## Molecular Plant

### Root Growth Assays

Seeds were sterilized and plated on 1 × MS medium (Murashige & Skoog medium, M0221, Duchefa Biochemie BV) containing 2% (w/v) sucrose. For the JA treatment, (±)-jasmonic acid (J2600, Sigma-Aldrich) was added to a concentration of 50 μM. Thirty seeds were placed in a line 2 cm below the upper plate margin on each plate and cold stratified on the plate for 2 days. The experiment was carried out in two rounds of sowing with two plates per experimental round and treatment. All four genotypes were individually randomized on each plate. The seedlings were grown vertically in a Percival growth chamber at 80–120 μE/(m<sup>2</sup> s), 12 h light, 21°C, and 70% relative humidity. Plates were scanned on days 3–8 after placing them in the chamber. Root length was measured using the SmartRoot image analysis software (Lobet et al., 2011).

### Jasmonic Acid Analysis

JA was extracted and analyzed as described previously (Stingl et al., 2013). The seedlings were sown three times in a random block design. Per experimental round, three pools of seedlings were harvested per combination of genotype and treatment (total *n* = 9). 10–20 mg of 5-day-old seedlings (pools of 15–20 seedlings) were snap frozen in liquid nitrogen and homogenized with two stainless steel balls by shaking for 2 × 30 s minimum at a frequency of 20/s on a Mixer Mill 303 (Retsch, Haan, Germany). 1 ml of cold ethyl acetate/formic acid (99:1 v/v) was added. The samples were mixed and then centrifuged at 16 000 *g* at room temperature for 10 min. The supernatants were transferred to new tubes and the solvents were removed by using a centrifugal vacuum evaporator at 30°C (30–40 min). The residues were dissolved in 80 μl of 5% acetonitrile in water (v/v), the samples were filtered, and stored at –20°C until analysis. The samples were analyzed by UHPLC/TQ-MS on an Advance™-UHPLC/EVOQ™ Elite-TQ-MS instrument (Bruker) equipped with a C-18 reversed phase column (Kinetex 1.7 μ XB-C18, 10 cm × 2.1 mm, 1.7 μm particle size, Phenomenex) by using a 0.05% formic acid in water (v/v) (solvent A)–0.05% formic acid in acetonitrile (v/v) (solvent B) gradient at a flow rate of 0.4 ml/min at 40°C. The gradient applied was as follows: 2% B (0.5 min), 2%–30% (0.7 min), 30%–100% (0.8 min), 100% B (0.5 min), 100%–2% B (0.1 min), and 2% B (1.4 min). Compounds were ionized by electrospray ionization with a spray voltage of –3900 V, heated probe temperature 210°C, cone temperature 250°C. JA was monitored based on the MRM (–)209 > 59 [11V]. Results are given as nmol/(g fresh weight) calculated based on external standard curves.

## SUPPLEMENTAL INFORMATION

Supplemental Information is available at *Molecular Plant Online*.

## FUNDING

Financial support for the work was provided by the Danish National Research Foundation DNRF grant 99 (B.A.H., M.B., D.J.K.) and IEE Marie Curie stipend PIEF-GA-2008-221236 (M.B.).

## ACKNOWLEDGMENTS

Sophie Lambertz and Emma Sofie Timm Aller are thanked for help with the JA analyses. No conflict of interest declared.

Received: September 8, 2014

Revised: March 1, 2015

Accepted: March 2, 2015

Published: March 7, 2015

## REFERENCES

- Anders, S., Pyl, P.T., and Huber, W. (2014). HTSeq – a Python framework to work with high-throughput sequencing data. *Bioinformatics* **31**:166–169.
- Bak, S., Tax, F.E., Feldmann, K.A., Galbraith, D.W., and Feyereisen, R. (2001). CYP83B1, a cytochrome P450 at the metabolic branch point in auxin and indole glucosinolate biosynthesis in *Arabidopsis*. *Plant Cell* **13**:101–111.
- Beekwilder, J., van Leeuwen, W., van Dam, N.M., Bertossi, M., Grandi, V., Mizzi, L., Soloviev, M., Szabados, L., Molthoff, J.W., Schipper, B., et al. (2008). The impact of the absence of aliphatic glucosinolates on insect herbivory in *Arabidopsis*. *PLoS One* **3**:e2068.
- Bekaert, M., Edger, P.P., Hudson, C.M., Pires, J.C., and Conant, G.C. (2012). Metabolic and evolutionary costs of herbivory defense: systems biology of glucosinolate synthesis. *New Phytol.* **196**:596–605.
- Bidart-Bouzat, M.G., and Kliebenstein, D. (2008). Differential levels of insect herbivory in the field associated with genotypic variation in glucosinolates in *Arabidopsis thaliana*. *J. Chem. Ecol.* **34**:1026–1037.
- Bonawitz, N.D., Kim, J.I., Tobimatsu, Y., Ciesielski, P.N., Anderson, N.A., Ximenes, E., Maeda, J., Ralph, J., Donohoe, B.S., Ladisch, M., et al. (2014). Disruption of Mediator rescues the stunted growth of a lignin-deficient *Arabidopsis* mutant. *Nature* **509**:376–380.
- Brader, G., Tas, É., and Palva, E.T. (2001). Jasmonate-dependent induction of indole glucosinolates in *Arabidopsis* by culture filtrates of the nonspecific pathogen *Erwinia carotovora*. *Plant Physiol.* **126**:849–860.
- Brown, P.D., Tokuhisa, J.G., Reichelt, M., and Gershenzon, J. (2003). Variation of glucosinolate accumulation among different organs and developmental stages of *Arabidopsis thaliana*. *Phytochemistry* **62**:471–481.
- Burow, M., Losansky, A., Muller, R., Plock, A., Kliebenstein, D.J., and Wittstock, U. (2009). The genetic basis of constitutive and herbivore-induced ESP-independent nitrile formation in *Arabidopsis*. *Plant Physiol.* **149**:561–574.
- Burow, M., Halkier, B.A., and Kliebenstein, D.J. (2010). Regulatory networks of glucosinolates shape *Arabidopsis thaliana* fitness. *Curr. Opin. Plant Biol.* **13**:348–353.
- Capella, A., Menossi, M., Arruda, P., and Benedetti, C. (2001). COI1 affects myrosinase activity and controls the expression of two flower-specific myrosinase-binding protein homologues in *Arabidopsis*. *Planta* **213**:691–699.
- Chan, E.K., Rowe, H.C., Corwin, J.A., Joseph, B., and Kliebenstein, D.J. (2011). Combining genome-wide association mapping and transcriptional networks to identify novel genes controlling glucosinolates in *Arabidopsis thaliana*. *PLoS Biol.* **9**:e1001125.
- Chauvin, A., Caldelari, D., Wolfender, J.L., and Farmer, E.E. (2013). Four 13-lipoxygenases contribute to rapid jasmonate synthesis in wounded *Arabidopsis thaliana* leaves: a role for lipoxygenase 6 in responses to long-distance wound signals. *New Phytol.* **197**:566–575.
- Chew, F.S. (1988). Biological effects of glucosinolates. In *Biologically Active Natural Products*, H.G. Cutler, ed. (Washington DC: American Chemical Society), pp. 155–181.
- Chini, A., Fonseca, S., Fernández, G., Adie, B., Chico, J.M., Lorenzo, O., García-Casado, G., López-Vidriero, I., Lozano, F.M., Ponce, M.R., et al. (2007). The JAZ family of repressors is the missing link in jasmonate signalling. *Nature* **448**:666–671.
- de Kraker, J.W., Luck, K., Textor, S., Tokuhisa, J.G., and Gershenzon, J. (2007). Two *Arabidopsis* genes (IPMS1 and IPMS2) encode isopropylmalate synthase, the branchpoint step in the biosynthesis of leucine. *Plant Physiol.* **143**:970–986.
- Dombrecht, B., Xue, G.P., Sprague, S.J., Kirkegaard, J.A., Ross, J.J., Reid, J.B., Fitt, G.P., Sewelam, N., Schenk, P.M., Manners, J.M., et al. (2007). MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *Plant Cell* **19**:2225–2245.
- Falk, K.L., Tokuhisa, J.G., and Gershenzon, J. (2007). The effect of sulfur nutrition on plant glucosinolate content: physiology and molecular mechanisms. *Plant Biol.* **9**:573–581.
- Fiebig, H.J., and Arens, M. (1992). Glucosinolate (HPLC-Methode) – Gemeinschaftsarbeiten der DGF, 128. Mitteilung: Deutsche Einheitsmethoden zur Untersuchung von Fetten, Fettprodukten,

## Feed-back Regulation of Glucosinolate Biosynthesis

- Tensiden und verwandten Stoffen, 98. Mitt.: Analyse von Fettrohstoffen XII. *Lipid/Fett* **94**:199–203.
- Freeman, M.** (2000). Feedback control of intercellular signalling in development. *Nature* **408**:313–319.
- Frerigmann, H., and Gigolashvili, T.** (2014). MYB34, MYB51, and MYB122 distinctly regulate indolic glucosinolate biosynthesis in *Arabidopsis thaliana*. *Mol. Plant* **7**:814–828.
- Frerigmann, H., Berger, B., and Gigolashvili, T.** (2014). bHLH05 is an interaction partner of MYB51 and a novel regulator of glucosinolate biosynthesis in *Arabidopsis*. *Plant Physiol.* **166**:349–369.
- Geu-Flores, F., Moldrup, M.E., Bottcher, C., Olsen, C.E., Scheel, D., and Halkier, B.A.** (2011). Cytosolic gamma-glutamyl peptidases process glutathione conjugates in the biosynthesis of glucosinolates and camalexin in *Arabidopsis*. *Plant Cell* **23**:2456–2469.
- Giamoustaris, A., and Mithen, R.** (1995). The effect of modifying the glucosinolate content of leaves of oilseed rape (*Brassica napus* ssp. *oleifera*) on its interaction with specialist and generalist pests. *Ann. Appl. Biol.* **126**:347–363.
- Gigolashvili, T., Berger, B., Mock, H.P., Muller, C., Weisshaar, B., and Flugge, U.I.** (2007a). The transcription factor HIG1/MYB51 regulates indolic glucosinolate biosynthesis in *Arabidopsis thaliana*. *Plant J.* **50**:886–901.
- Gigolashvili, T., Yatusovich, R., Berger, B., Muller, C., and Flugge, U.I.** (2007b). The R2R3-MYB transcription factor HAG1/MYB28 is a regulator of methionine-derived glucosinolate biosynthesis in *Arabidopsis thaliana*. *Plant J.* **51**:247–261.
- Gigolashvili, T., Berger, B., and Flügge, U.-I.** (2008a). Specific and coordinated control of indolic and aliphatic glucosinolate biosynthesis by R2R3-MYB transcription factors in *Arabidopsis thaliana*. *Phytochem. Rev.* **8**:3–13.
- Gigolashvili, T., Engqvist, M., Yatusovich, R., Muller, C., and Flugge, U.I.** (2008b). HAG2/MYB76 and HAG3/MYB29 exert a specific and coordinated control on the regulation of aliphatic glucosinolate biosynthesis in *Arabidopsis thaliana*. *New Phytol.* **177**:627–642.
- Grant, G.A.** (2006). The ACT domain: a small molecule binding domain and its role as a common regulatory element. *J. Biol. Chem.* **281**:33825–33829.
- Hansen, C.H., Du, L., Naur, P., Olsen, C.E., Axelsen, K.B., Hick, A.J., Pickett, J.A., and Halkier, B.A.** (2001). CYP83B1 is the oxime-metabolizing enzyme in the glucosinolate pathway in *Arabidopsis*. *J. Biol. Chem.* **276**:24790–24796.
- Hirai, M.Y., Yano, M., Goodenowe, D.B., Kanaya, S., Kimura, T., Awazuhara, M., Arita, M., Fujiwara, T., and Saito, K.** (2004). Integration of transcriptomics and metabolomics for understanding of global responses to nutritional stresses in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **101**:10205–10210.
- Hirai, M.Y., Sugiyama, K., Sawada, Y., Tohge, T., Obayashi, T., Suzuki, A., Araki, R., Sakurai, N., Suzuki, H., Aoki, K., et al.** (2007). Omics-based identification of *Arabidopsis* Myb transcription factors regulating aliphatic glucosinolate biosynthesis. *Proc. Natl. Acad. Sci. USA* **104**:6478–6483.
- Hopkins, R.J., van Dam, N.M., and van Loon, J.J.** (2009). Role of glucosinolates in insect-plant relationships and multitrophic interactions. *Annu. Rev. Entomol.* **54**:57–83.
- Huseby, S., Koprivova, A., Lee, B.R., Saha, S., Mithen, R., Wold, A.B., Bengtsson, G.B., and Kopriva, S.** (2013). Diurnal and light regulation of sulphur assimilation and glucosinolate biosynthesis in *Arabidopsis*. *J. Exp. Bot.* **64**:1039–1048.
- Joseph, B., Corwin, J.A., Li, B., Atwell, S., and Kliebenstein, D.J.** (2013). Cytoplasmic genetic variation and extensive cytonuclear interactions influence natural variation in the metabolome. *Elife* **2**:e00776.
- Jost, R., Altschmied, L., Bloem, E., Bogs, J., Gershenzon, J., Hähnel, U., Hänsch, R., Hartmann, T., Kopriva, S., Kruse, C., et al.** (2005). Expression profiling of metabolic genes in response to methyl jasmonate reveals regulation of genes of primary and secondary sulfur-related pathways in *Arabidopsis thaliana*. *Photosyn. Res.* **86**:491–508.
- Kai, K., Takahashi, H., Saga, H., Ogawa, T., Kanaya, S., and Ohta, D.** (2011). Metabolomic characterization of the possible involvement of a Cytochrome P450, CYP81F4, in the biosynthesis of indolic glucosinolate in *Arabidopsis*. *Plant Biotechnol.* **28**:379–385.
- Kandel, S., Sauveplane, V., Compagnon, V., Franke, R., Millet, Y., Schreiber, L., Werck-Reichhart, D., and Pinot, F.** (2007). Characterization of a methyl jasmonate and wounding-responsive cytochrome P450 of *Arabidopsis thaliana* catalyzing dicarboxylic fatty acid formation in vitro. *FEBS J.* **274**:5116–5127.
- Kauss, D., Bischof, S., Steiner, S., Apel, K., and Meskauskiene, R.** (2012). FLU, a negative feedback regulator of tetrapyrrole biosynthesis, is physically linked to the final steps of the Mg(++)-branch of this pathway. *FEBS Lett.* **586**:211–216.
- Kazan, K., and Manners, J.M.** (2013). MYC2: the master in action. *Mol. Plant* **6**:686–703.
- Kerwin, R.E., Jimenez-Gomez, J.M., Fulop, D., Harmer, S.L., Maloof, J.N., and Kliebenstein, D.J.** (2011). Network quantitative trait loci mapping of circadian clock outputs identifies metabolic pathway-to-clock linkages in *Arabidopsis*. *Plant Cell* **23**:471–485.
- Kim, J.H., and Jander, G.** (2007). *Myzus persicae* (green peach aphid) feeding on *Arabidopsis* induces the formation of a deterrent indole glucosinolate. *Plant J.* **49**:1008–1019.
- Kliebenstein, D.J., Kroymann, J., Brown, P., Figuth, A., Pedersen, D., Gershenzon, J., and Mitchell-Olds, T.** (2001a). Genetic control of natural variation in *Arabidopsis* glucosinolate accumulation. *Plant Physiol.* **126**:811–825.
- Kliebenstein, D.J., Lambrix, V.M., Reichelt, M., Gershenzon, J., and Mitchell-Olds, T.** (2001b). Gene duplication in the diversification of secondary metabolism: tandem 2-oxoglutarate-dependent dioxygenases control glucosinolate biosynthesis in *Arabidopsis*. *Plant Cell* **13**:681–694.
- Kliebenstein, D.J., Figuth, A., and Mitchell-Olds, T.** (2002). Genetic architecture of plastic methyl jasmonate responses in *Arabidopsis thaliana*. *Genetics* **161**:1685–1696.
- Koo, A.J.K., Cooke, T.F., and Howe, G.A.** (2011). Cytochrome P450 CYP94B3 mediates catabolism and inactivation of the plant hormone jasmonoyl-L-isoleucine. *Proc. Natl. Acad. Sci. USA* **108**:9298–9303.
- Kumar, R., Ichihashi, Y., Kimura, S., Chitwood, D.H., Headland, L.R., Peng, J., Maloof, J.N., and Sinha, N.R.** (2012). A high-throughput method for illumina RNA-Seq library preparation. *Front. Plant Sci.* **3**:202.
- Lankau, R.A.** (2007). Specialist and generalist herbivores exert opposing selection on a chemical defense. *New Phytol.* **175**:176–184.
- Li, G., and Quiros, C.F.** (2003). In planta side-chain glucosinolate modification in *Arabidopsis* by introduction of dioxygenase Brassica homolog BoGSL-ALK. *Theor. Appl. Genet.* **106**:1116–1121.
- Li, J., Hansen, B.G., Ober, J.A., Kliebenstein, D.J., and Halkier, B.A.** (2008). Subclade of flavin-monooxygenases involved in aliphatic glucosinolate biosynthesis. *Plant Physiol.* **148**:1721–1733.
- Li, B., Gaudinier, A., Tang, M., Taylor-Teeples, M., Nham, N.T., Ghaffari, C., Benson, D.S., Steinmann, M., Gray, J.A., Brady, S.M., et al.** (2014). Promoter-based integration in plant defense regulation. *Plant Physiol.* **166**:1803–1820.
- Lobet, G., Pagès, L., and Draye, X.** (2011). A novel image-analysis toolbox enabling quantitative analysis of root system architecture. *Plant Physiol.* **157**:29–39.



## Molecular Plant

- Lorenzo, O., Chico, J.M., Sanchez-Serrano, J.J., and Solano, R. (2004). JASMONATE-INSENSITIVE1 encodes a MYC transcription factor essential to discriminate between different jasmonate-regulated defense responses in *Arabidopsis*. *Plant Cell* **16**:1938–1950.
- Mauricio, R. (1998). Costs of resistance to natural enemies in field populations of the annual plant *Arabidopsis thaliana*. *Am. Nat.* **151**:20–28.
- Mewis, I., Appel, H.M., Hom, A., Raina, R., and Schultz, J.C. (2005). Major signaling pathways modulate *Arabidopsis* glucosinolate accumulation and response to both phloem-feeding and chewing insects. *Plant Physiol.* **138**:1149–1162.
- Mewis, I., Tokuhisa, J.G., Schultz, J.C., Appel, H.M., Ulrichs, C., and Gershenzon, J. (2006). Gene expression and glucosinolate accumulation in *Arabidopsis thaliana* in response to generalist and specialist herbivores of different feeding guilds and the role of defense signaling pathways. *Phytochemistry* **67**:2450–2462.
- Mikkelsen, M.D., Petersen, B.L., Glawischnig, E., Jensen, A.B., Andreasson, E., and Halkier, B.A. (2003). Modulation of CYP79 genes and glucosinolate profiles in *Arabidopsis* by defense signaling pathways. *Plant Physiol.* **131**:298–308.
- Naur, P., Petersen, B.L., Mikkelsen, M.D., Bak, S., Rasmussen, H., Olsen, C.E., and Halkier, B.A. (2003). CYP83A1 and CYP83B1, two nonredundant cytochrome P450 enzymes metabolizing oximes in the biosynthesis of glucosinolates in *Arabidopsis*. *Plant Physiol.* **133**:63–72.
- Orrell, D., Ramsey, S., Marelli, M., Smith, J.J., Petersen, T.W., Atauri, P.d., Aitchison, J.D., and Bolouri, H. (2006). Feedback control of stochastic noise in the yeast galactose utilization pathway. *Physica D* **217**:64–76.
- Petersen, B., Chen, S., Hansen, C., Olsen, C., and Halkier, B. (2002). Composition and content of glucosinolates in developing *Arabidopsis thaliana*. *Planta* **214**:562–571.
- Pfalz, M., Mikkelsen, M.D., Bednarek, P., Olsen, C.E., Halkier, B.A., and Kroymann, J. (2011). Metabolic engineering in *Nicotiana benthamiana* reveals key enzyme functions in *Arabidopsis* indole glucosinolate modification. *Plant Cell* **23**:716–729.
- Piotrowski, M., Schemenewitz, A., Lopukhina, A., Muller, A., Janowitz, T., Weiler, E.W., and Oecking, C. (2004). Desulfoglucosinolate sulfotransferases from *Arabidopsis thaliana* catalyze the final step in the biosynthesis of the glucosinolate core structure. *J. Biol. Chem.* **279**:50717–50725.
- Reichelt, M., Brown, P.D., Schneider, B., Oldham, N.J., Stauber, E., Tokuhisa, J., Kliebenstein, D.J., Mitchell-Olds, T., and Gershenzon, J. (2002). Benzoic acid glucosinolate esters and other glucosinolates from *Arabidopsis thaliana*. *Phytochemistry* **59**:663–671.
- Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139–140.
- Rowe, H.C., Hansen, B.G., Halkier, B.A., and Kliebenstein, D.J. (2008). Biochemical networks and epistasis shape the *Arabidopsis thaliana* metabolome. *Plant Cell* **20**:1199–1216.
- Sanders, P.M., Lee, P.Y., Biesgen, C., Boone, J.D., Beals, T.P., Weiler, E.W., and Goldberg, R.B. (2000). The arabidopsis DELAYED DEHISCENCE1 gene encodes an enzyme in the jasmonic acid synthesis pathway. *Plant Cell* **12**:1041–1061.
- Schweizer, F., Fernandez-Calvo, P., Zander, M., Diez-Diaz, M., Fonseca, S., Glauser, G., Lewsey, M.G., Ecker, J.R., Solano, R., and Reymond, P. (2013). *Arabidopsis* basic helix-loop-helix transcription factors MYC2, MYC3, and MYC4 regulate glucosinolate biosynthesis, insect performance, and feeding behavior. *Plant Cell* **25**:3117–3132.

## Feed-back Regulation of Glucosinolate Biosynthesis

- Smith, A.M., and Stitt, M. (2007). Coordination of carbon supply and plant growth. *Plant Cell Environ.* **30**:1126–1149.
- Sønderby, I.E., Hansen, B.G., Bjarnholt, N., Ticconi, C., Halkier, B.A., and Kliebenstein, D.J. (2007). A systems biology approach identifies a R2R3 MYB gene subfamily with distinct and overlapping functions in regulation of aliphatic glucosinolates. *PLoS One* **2**:e1322.
- Sønderby, I.E., Burow, M., Rowe, H.C., Kliebenstein, D.J., and Halkier, B.A. (2010a). A complex interplay of three R2R3 MYB transcription factors determines the profile of aliphatic glucosinolates in *Arabidopsis*. *Plant Physiol.* **153**:348–363.
- Sønderby, I.E., Geu-Flores, F., and Halkier, B.A. (2010b). Biosynthesis of glucosinolates—gene discovery and beyond. *Trends Plant Sci.* **15**:283–290.
- Stenzel, I., Hause, B., Miersch, O., Kurz, T., Maucher, H., Weichert, H., Ziegler, J., Feussner, I., and Wasternack, C. (2003). Jasmonate biosynthesis and the allene oxide cyclase family of *Arabidopsis thaliana*. *Plant Mol. Biol.* **51**:895–911.
- Stingl, N., Krischke, M., Fekete, A., and Mueller, M.J. (2013). Analysis of defense signals in *Arabidopsis thaliana* leaves by ultra-performance liquid chromatography/tandem mass spectrometry: jasmonates, salicylic acid, abscisic acid. *Methods Mol. Biol.* **1009**:103–113.
- Stintzi, A., and Browse, J. (2000). The *Arabidopsis* male-sterile mutant, opr3, lacks the 12-oxophytodieneic acid reductase required for jasmonate synthesis. *Proc. Natl. Acad. Sci. USA* **97**:10625–10630.
- Stotz, H.U., Pittendrigh, B.R., Kroymann, J., Weniger, K., Fritsche, J., Bauke, A., and Mitchell-Olds, T. (2001). Induced plant defense responses against chewing insects. Ethylene signaling reduces resistance of *Arabidopsis* against Egyptian cotton worm but not diamondback moth. *Plant Physiol.* **124**:1007–1018, Erratum in *Plant Physiol.* **125**:2203–2203.
- Thies, W. (1979). Detection and utilization of a glucosinolate sulfohydrolase in the edible snail, *Helix pomatia*. *Naturwissenschaften* **66**:364–365.
- Thines, B., Katsir, L., Melotto, M., Niu, Y., Mandaokar, A., Liu, G., Nomura, K., He, S.Y., Howe, G.A., and Browse, J. (2007). JAZ repressor proteins are targets of the SCF(CO1) complex during jasmonate signalling. *Nature* **448**:661–665.
- Tierens, K.F., Thomma, B.P., Brouwer, M., Schmidt, J., Kistner, K., Porzel, A., Mauch-Mani, B., Cammue, B.P., and Broekaert, W.F. (2001). Study of the role of antimicrobial glucosinolate-derived isothiocyanates in resistance of *Arabidopsis* to microbial pathogens. *Plant Physiol.* **125**:1688–1699.
- Trapnell, C., Pachter, L., and Salzberg, S.L. (2009). TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* **25**:1105–1111.
- Wentzell, A.M., Rowe, H.C., Hansen, B.G., Ticconi, C., Halkier, B.A., and Kliebenstein, D.J. (2007). Linking metabolic QTLs with network and cis-eQTLs controlling biosynthetic pathways. *PLoS Genet.* **3**:1687–1701.
- Yi, T.-M., Huang, Y., Simon, M.I., and Doyle, J. (2000). Robust perfect adaptation in bacterial chemotaxis through integral feedback control. *Proc. Natl. Acad. Sci. USA* **97**:4649–4653.
- Zrybko, C.L., Fukuda, E.K., and Rosen, R.T. (1997). Determination of glucosinolates in domestic and wild mustard by high-performance liquid chromatography with confirmation by electrospray mass spectrometry and photodiode-array detection. *J. Chromatogr. A* **767**:43–52.
- Zust, T., Joseph, B., Shimizu, K.K., Kliebenstein, D.J., and Turnbull, L.A. (2011). Using knockout mutants to reveal the growth costs of defensive traits. *Proc. Biol. Sci.* **278**:2598–2603.