

UC Davis

UC Davis Previously Published Works

Title

Body Site Staphylococcus aureus Colonization among Maintenance Hemodialysis Patients

Permalink

<https://escholarship.org/uc/item/60g0p042>

Journal

Nephron, 129(2)

ISSN

1660-8151

Authors

Eells, Samantha J
Kalantar-Zadeh, Kamyar
Bolaris, Michael A
[et al.](#)

Publication Date

2015

DOI

10.1159/000369348

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Body Site *Staphylococcus aureus* Colonization among Maintenance Hemodialysis Patients

Samantha J. Eells^{a–c} Kamyar Kalantar-Zadeh^{c, d} Michael A. Bolaris^{a, b}
Larissa May^{e, f} Loren G. Miller^{a, b}

^aDivision of Infectious Diseases, Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center, and ^bDavid Geffen School of Medicine at UCLA and Harbor-UCLA Medical Center, Torrance, Calif., ^cUCLA Fielding School of Public Health, Los Angeles, Calif., ^dDivision of Nephrology and Hypertension, University of California Irvine Medical Center, Orange, Calif., and ^eDepartment of Emergency Medicine, School of Medicine, and ^fDepartment of Epidemiology and Biostatistics, Milken Institute of Public Health, The George Washington University, Washington, D.C., USA

Key Words

Staphylococcus aureus · Colonization · Maintenance hemodialysis

Abstract

Background: Patients on maintenance hemodialysis therapy are at high risk for health care-associated infections. *Staphylococcus aureus* is a common cause of health care-associated infections among maintenance hemodialysis patients. It is established that *S. aureus* colonization is associated with an increased risk for subsequent infection in this population. There is an increasing number of reports that extranasal *S. aureus* colonization is more common than previously believed and in certain body sites even more common than nasal colonization. There are few data describing extranasal colonization among maintenance hemodialysis patients. **Methods:** We surveyed 100 patients at 3 body sites (anterior nares, oropharynx, and inguinal region) for *S. aureus* colonization. Participants were also administered a standardized survey to assess risk factors for *S. aureus* colonization. **Results:** We found that 42% (95% CI 32–52) of patients were *S. aureus* colonized in >1 body site. Extranasal colonization was found among 32% (95% CI 23–41). There

were trends suggestive of an association between *S. aureus* colonization and younger age (OR 0.97, 95% CI 0.94–1.001, $p = 0.06$) and not having been hospitalized in the previous 12 months (OR 0.44, 95% CI 0.19–1.06, $p = 0.14$). **Conclusion:** Extranasal *S. aureus* colonization is common among maintenance hemodialysis patients with a prevalence of approximately one third. Future *S. aureus* decolonization efforts may need to consider not just nasal decolonization but also decolonization of the skin and oropharynx.

© 2015 S. Karger AG, Basel

Background

Patients on maintenance hemodialysis are at high risk of health care-associated infections, especially those caused by *Staphylococcus aureus*. *S. aureus*, including methicillin-resistant *S. aureus* (MRSA), causes a wide range of community- and health care-associated diseases, ranging from skin and soft-tissue infections to severe sepsis [1]. *S. aureus* is one of the most common pathogens identified in bloodstream infections among hemodialysis patients [2]. Persons receiving maintenance hemodialysis are at very high risk for infection with invasive disease from *S. aureus*

and MRSA infection compared to the general population. While annual invasive MRSA rates in the general population are 0.2–0.4 infections per 1,000 persons [3], rates among patients on maintenance hemodialysis have been estimated at 37 cases per 1,000 persons [4], a roughly 100-fold higher risk. Of critical concern is that invasive MRSA infection among patients on maintenance hemodialysis is associated with a 17% mortality rate [4].

Asymptomatic *S. aureus* carriage has been an area of great interest due to the risk of subsequent *S. aureus* infection in the colonized individual [5–7]. The ecologic niche for *S. aureus* has traditionally been thought to be the anterior nares [8, 9]. Population-based surveys found that 27–30% of the US population is nasally colonized with *S. aureus* and 1.2–1.8% is nasally colonized with MRSA [10]. The prevalence of nasal *S. aureus* colonization is 30–57% among patients on maintenance hemodialysis [11–13]. MRSA nasal colonization ranges from 5.6 to 12% among maintenance hemodialysis patients [13, 14].

Nasal carriage of *S. aureus* is known to play an important role as an endogenous source for *S. aureus* and MRSA infections that contribute significantly to morbidity, mortality, and cost of end-stage renal disease management [2]. However, recent data suggest that extranasal colonization is more common than previously recognized. Among inpatients with an *S. aureus* infection, MRSA colonization in the nares, axilla, inguinal area, and rectum was 25, 6, 11, and 13%, respectively, and 37% overall were MRSA colonized [15]. A study of households with a history of a recent *S. aureus* skin infection found that up to 50% of household members were *S. aureus* colonized and that a nares-only survey would miss 48% of *S. aureus* colonization and 51% of MRSA colonization [16]. The magnitude of pharyngeal and inguinal colonization has led to a paradigm shift in the understanding of body colonization with *S. aureus* [17]. Older investigations conducted prior to recent increases in MRSA prevalence in patients on maintenance hemodialysis found that 10% of patients (6/59 patients) had *S. aureus* umbilical colonization [18], 8% (3/37 patients) had inguinal *S. aureus* [19], and 7% (2/28 patients) had oropharyngeal *S. aureus* colonization [12]. Because body colonization, especially of extranasal sites, is more widespread than initially believed, interventions to prevent *S. aureus* and MRSA infection may need to consider extranasal decolonization as well as traditional nasal decolonization with agents such as mupirocin [20]. The objective of the present investigation is to identify the frequency and factors associated with extranasal *S. aureus* colonization among maintenance hemodialysis patients.

Methods

This cross-sectional investigation took place at Harbor-UCLA Medical Center-associated outpatient dialysis centers in Torrance, Long Beach, and Hawthorne, Calif., USA, from July 2010 through November 2010. Potential subjects were identified via daily screening of the dialysis centers. All adults receiving dialysis at participating centers were eligible for participation. Interested patients underwent the informed consent process and signed consent forms prior to any research-related samples or surveys being collected. This investigation and the associated consent form were approved by the Los Angeles Biomedical Research Institute at Harbor-UCLA Medical Center Institutional Review Board.

A standardized survey instrument from a published investigation on risk factors for skin infections was modified and used to assess risk factors for *S. aureus* colonization in this investigation [21]. Subjects were surveyed about past medical history and behavioral risk factors for *S. aureus* colonization. Based on data suggesting that nasal colonization surveillance alone is insensitive for detecting *S. aureus* body colonization [15], we collected nasal, oropharyngeal, and inguinal culture swabs to test for the presence of *S. aureus*.

Culture swabs were transported immediately to the microbiology laboratory and were enriched in trypticase soy broth with 7% sodium chloride overnight at 35°C. The broth was subcultured to BBL CHROMagar *S. aureus* and MRSA plates (BD, Franklin Lakes, N.J., USA) and incubated aerobically for 24 h at 35°C. Isolates were confirmed as *S. aureus* by the catalase test and StaphAureux tests (Remel, Lenexa, Kans., USA). MRSA isolates, confirmed using CHROMagar MRSA plates, were subcultured twice for purity. DNA was extracted, digested with *Sma*I, and subjected to pulsed-field gel electrophoresis (PFGE) as previously described [22]. DNA profiles were analyzed using GelCompar software (Applied Maths, Austin, Tex., USA), and a reference database from the Centers for Disease Control and Prevention (CDC) containing the MRSA USA types strain patterns was used to assign the pulsed-field type (PFT) and determine strain relatedness.

Bivariate analysis was used to compare variables from the risk factor survey hypothesized to be associated with extranasal *S. aureus* colonization. Secondary post hoc bivariate analyses were conducted using different outcomes of (1) any *S. aureus* colonization and (2) extranasal *S. aureus* colonization only. Bivariate analyses were assessed using odds ratios (OR) adjusted for 95% confidence intervals (CI) and the associated *p* values. All variables were considered significant at the $\alpha = 0.05$ level. Data analyses were performed using SAS (version 9.3; SAS Institute, Cary, N.C., USA).

Results

Among the 100 patients enrolled, the mean age of subjects was 51 years (median 42, range 20–88), 68% were male, 55% were Hispanic, 14% of cases were African-American, 7% were Caucasian, and 24% were of mixed or other race/ethnicity. The mean duration of maintenance hemodialysis was 5 years (median 3.5). Fifty-one percent had been hospitalized in the previous 12 months and 18 had a history of ever having a skin or soft-tissue infection.

Table 1. Baseline characteristics associated with extranasal *S. aureus* colonization among maintenance hemodialysis patients

	All patients (n = 100)	Extranasal colonization (n = 32)	No extranasal colonization (n = 68)	OR	95% CI	p value
Age, years						
Mean ± SD	51±15	47±15	53±15	0.97	0.94, 1.001	0.06
Median (range)	52 (20–88)	46 (23–84)	54 (20–88)			
Male gender	66 (68)	25 (78)	41 (63)	2.09	0.79, 5.56	0.14
Ethnicity						
African-American	14 (14)	5 (16)	9 (13)	0.72	0.10, 5.17	0.74
Hispanic	55 (55)	15 (47)	40 (59)	ref.		
Caucasian	7 (7)	2 (6)	5 (7)	0.68	0.20, 2.34	0.54
Other/unknown	24 (24)	10 (31)	14 (21)	1.29	0.33, 5.02	0.72
History of an <i>S. aureus</i> infection	3 (3)	0 (0)	3 (5)	–	–	0.24
History of a skin or soft-tissue infection	18 (19)	7 (23)	11 (16)	1.54	0.53, 4.49	0.42
Close contact with someone who had an <i>S. aureus</i> infection in the previous 12 months	2 (2)	0 (0)	2 (3)	–	–	0.47
Method of receipt of dialysis						
Venous catheter	9 (9)	1 (3)	8 (12)	ref.		
Arteriovenous graft or fistula	90 (91)	31 (97)	59 (88)	4.20	0.50, 35.2	0.26
Hospitalization in the previous 12 months	51 (51)	12 (38)	39 (57)	0.44	0.19, 1.06	0.06
Admitted to a long-term care facility in the previous 12 months	6 (6)	2 (6)	4 (6)	1.09	0.19, 6.27	0.99
Years of dialysis						
Mean ± SD	5.0±5.2	5.9±5.2	4.7±5.3	1.04	0.97, 1.13	0.29
Median (range)	3.5 (0.04–31)	4.8 (0.06–25)	3.0 (0.04–31)			
Charlson comorbidity score						
Mean ± SD	3.3±1.7	2.9±1.4	3.4±1.9	0.82	0.60, 1.11	0.19
Median (range)	2.5 (2.0–14.0)	2.0 (2.0–7.0)	3.0 (2.0–14.0)			

Values are n (%) unless otherwise indicated. ref. = Reference group; SD = standard deviation; – = cannot be calculated due to zero cells.

The baseline characteristics of the patients surveyed are summarized in table 1. *S. aureus* colonization results are summarized in table 2. Overall, 42% (95% CI 32–52) of patients were *S. aureus* colonized at >1 body site. Six percent (95% CI 1–11) were colonized with MRSA. Extranasal *S. aureus* colonization was found among 32% (95% CI 23–41) of patients and 4% of patients had extranasal MRSA colonization (95% CI 0–5). The overlap of *S. aureus* colonization among the three body sites surveyed can be found in figure 1.

In our primary bivariate analysis, there were trends of associations between extranasal *S. aureus* colonization and younger age (OR 0.97, 95% CI 0.94–1.001) and not having been hospitalized in the previous 12 months (OR 0.44, 95% CI 0.19–1.06; table 1). In our secondary analysis of factors associated with extranasal *S. aureus* colonization only, we found trends suggestive of an association between extranasal colonization only and more years of dialysis (OR 1.08, 95% CI 0.99–1.18) and lower Charlson comorbidity score

(OR 0.57, 95% CI 0.32–1.03). In our secondary analysis of factors associated with overall *S. aureus* colonization, there was an association between any *S. aureus* colonization and younger age (OR 0.97, 95% CI 0.94–0.99).

We recovered 9 MRSA isolates and 64 methicillin-susceptible *S. aureus* (MSSA) isolates for PFGE typing. Of the 20 patients with *S. aureus* cultured from >1 body site, all recovered isolates were indistinguishable from each other, suggesting clonality. The 9 MRSA isolates recovered were all the USA300-MRSA PFT.

Discussion

S. aureus is a common clinical problem resulting in hospitalization, morbidity, and mortality among hemodialysis patients. Our prevalence study found that 42% of maintenance hemodialysis patients were colonized with *S. aureus* when surveyed at the anterior nares, orophar-

Table 2. Body site *S. aureus* colonization results among maintenance hemodialysis patients (n = 100)

	Percentage of colonized patients (95% CI)
Colonization of any body site	
Any <i>S. aureus</i>	42 (32, 52)
MSSA	36 (27, 45)
MRSA	6.0 (1.0, 11)
Nasal colonization	
Any <i>S. aureus</i>	28 (19, 37)
MSSA	24 (16, 32)
MRSA	4.0 (0.2, 7.8)
Throat colonization	
Any <i>S. aureus</i>	27 (18, 35)
MSSA	24 (16, 32)
MRSA	3.0 (0.0, 6.0)
Inguinal-region colonization	
Any <i>S. aureus</i>	17 (10, 24)
MSSA	15 (8.0, 22)
MRSA	2.0 (0.0, 5.0)
Extranasal colonization	
Any <i>S. aureus</i>	32 (23, 41)
MSSA	28 (19, 37)
MRSA	4.0 (0.2, 7.8)
Colonization at >1 body site	
Any <i>S. aureus</i>	19 (11, 27)
MSSA	16 (9, 23)
MRSA	3.0 (0.0, 6.0)
Extranasal colonization only	
Any <i>S. aureus</i>	14 (7.0, 21)
MSSA	12 (6.0, 18)
MRSA	2.0 (0.0, 5.0)
Nasal colonization only	
Any <i>S. aureus</i>	10 (4.0, 16)
MSSA	9.0 (3.0, 15)
MRSA	1.0 (0.2, 5.0)
Throat colonization only	
Any <i>S. aureus</i>	10 (4.0, 16)
MSSA	8.0 (3.0, 13)
MRSA	2.0 (0.0, 5.0)
Inguinal-region colonization only	
Any <i>S. aureus</i>	3.0 (0.0, 6.0)
MSSA	3.0 (0.0, 6.0)
MRSA	0.0 (-)

- = Cannot be calculated.

ynx, and inguinal region with 6% of patients being colonized with MRSA. Extranasal surveillance resulted in 33% additional patients identified with *S. aureus* and MRSA colonization. This is similar to other investigations examining the utility of extranasal *S. aureus* colonization surveillance in populations at high risk for *S. aureus* infection [15, 16, 23].




Fig. 1. Overlap of *S. aureus* colonization at the anterior nares, oropharynx, and inguinal region. Each circle size is proportional to the amount of *S. aureus* detected at that given anatomic site.

Trends of association seen between *S. aureus* extranasal colonization and the factors of younger age and not having been hospitalized in the previous 12 months may be a marker that younger and healthier hemodialysis patients are more likely to have *S. aureus* extranasal colonization. The association between younger age and overall *S. aureus* colonization was significant in our post hoc analyses. In an era where community-associated *S. aureus* is common and endemic among younger persons [21], this relationship is unsurprising.

There are limitations to our investigation. First, the patients were enrolled from a single metropolitan area and findings may not be generalizable to other hemodialysis populations. However, the patient population at our institution is similar to that of other large urban US populations and is ethnically diverse. Second, we relied on patients to self-report factors associated with colonization. Patients may not recall health care exposures over the previous 12 months. Third, we did not examine all possible body sites for colonization. However, the increased yield from determining other *S. aureus* colonization sites in addition to nasal, oropharyngeal, and inguinal colonization is not well understood and the sites chosen for surveillance appear to be those of highest yield for nonnasal sites [15, 16, 23]. Fourth, the sample size of our investigation was relatively limited making it difficult to determine factors

associated with colonization. Nevertheless, our investigation is larger than previous surveys of nonnasal *S. aureus* colonization by nearly 2- to 3-fold [12, 18, 19].

In conclusion, we found that extranasal *S. aureus* colonization was common among maintenance hemodialysis patients. Although it is unclear if decolonization efforts targeting MRSA or *S. aureus* in patients on maintenance hemodialysis will reduce subsequent infections [24], future interventional efforts in this field should consider decolonization of skin sites and the throat in addition to nasal decolonization.

References

- 1 Lowy FD: *Staphylococcus aureus* infections. *N Engl J Med* 1998;339:520–532.
- 2 Patel PR, Kallen AJ, Arduino MJ: Epidemiology, surveillance, and prevention of bloodstream infections in hemodialysis patients. *Am J Kidney Dis* 2010;56:566–577.
- 3 Fridkin SK, Hageman JC, Morrison M, Sanza LT, Como-Sabetti K, Jernigan JA, Harriman K, Harrison LH, Lynfield R, Farley MM; Active Bacterial Core Surveillance Program of the Emerging Infections Program Network: Methicillin-resistant *Staphylococcus aureus* disease in three communities. *N Engl J Med* 2005;352:1436–1444.
- 4 Nguyen DB, Lessa FC, Belflower R, Mu Y, Wise M, Nadle J, Bamberg WM, Petit S, Ray SM, Harrison LH, Lynfield R, Dumyati G, Thompson J, Schaffner W, Patel PR; Active Bacterial Core Surveillance (ABCs) MRSA Investigators of the Emerging Infections Program: Invasive methicillin-resistant *Staphylococcus aureus* infections among patients on chronic dialysis in the United States, 2005–2011. *Clin Infect Dis* 2013;57:1393–1400.
- 5 Casewell MW: The nose: an underestimated source of *Staphylococcus aureus* causing wound infection. *J Hosp Infect* 1998;40(suppl B):S3–S11.
- 6 von Eiff C, Becker K, Machka K, Stammer H, Peters G: Nasal carriage as a source of *Staphylococcus aureus* bacteremia. *N Engl J Med* 2001;344:11–16.
- 7 Wertheim HF, Vos MC, Ott A, van Belkum A, Voss A, Kluytmans JA, van Keulen PH, Vandenbroucke-Grauls CM, Meester MH, Verbrugh HA: Risk and outcome of nosocomial *Staphylococcus aureus* bacteraemia in nasal carriers versus non-carriers. *Lancet* 2004;364:703–705.
- 8 Kluytmans J, van Belkum A, Verbrugh H: Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev* 1997;10:505–520.
- 9 Williams RE: Healthy carriage of *Staphylococcus aureus*: its prevalence and importance. *Bacteriol Rev* 1963;27:56–71.
- 10 Gorwitz RJ, Kruszon-Moran D, McAllister SK, McQuillan G, McDougal LK, Fosheim GE, Jensen BJ, Killgore G, Tenover FC, Kuehnert MJ: Changes in the prevalence of nasal colonization with *Staphylococcus aureus* in the United States, 2001–2004. *J Infect Dis* 2008;197:1226–1234.
- 11 Johnson LB, Jose J, Yousif F, Pawlak J, Saravolatz LD: Prevalence of colonization with community-associated methicillin-resistant *Staphylococcus aureus* among end-stage renal disease patients and healthcare workers. *Infect Control Hosp Epidemiol* 2009;30:4–8.
- 12 Koziol-Montewka M, Chudnicka A, Ksiazek A, Majdan M: Rate of *Staphylococcus aureus* nasal carriage in immunocompromised patients receiving haemodialysis treatment. *Int J Antimicrob Agents* 2001;18:193–196.
- 13 Lederer SR, Riedelsdorf G, Schiffel H: Nasal carriage of methicillin resistant *Staphylococcus aureus*: the prevalence, patients at risk and the effect of elimination on outcomes among out-clinic haemodialysis patients. *Eur J Med Res* 2007;12:284–288.
- 14 Hadley AC, Karchmer TB, Russell GB, McBride DG, Freedman BI: The prevalence of resistant bacterial colonization in chronic hemodialysis patients. *Am J Nephrol* 2007;27:352–359.
- 15 Yang ES, Tan J, Eells S, Rieg G, Tagudar G, Miller LG: Body site colonization in patients with community-associated methicillin-resistant *Staphylococcus aureus* and other types of *S. aureus* skin infections. *Clin Microbiol Infect* 2010;16:425–431.
- 16 Miller LG, Eells SJ, Taylor AR, David MZ, Ortiz N, Zychowski D, Kumar N, Cruz D, Boyle-Vavra S, Daum RS: *Staphylococcus aureus* colonization among household contacts of patients with skin infections: risk factors, strain discordance, and complex ecology. *Clin Infect Dis* 2012;54:1523–1535.
- 17 Miller LG, Diep BA: Clinical practice: colonization, fomites, and virulence: rethinking the pathogenesis of community-associated methicillin-resistant *Staphylococcus aureus* infection. *Clin Infect Dis* 2008;46:752–760.
- 18 Swartz R, Messana J, Starman B, Weber M, Reynolds J: Preventing *Staphylococcus aureus* infection during chronic peritoneal dialysis. *J Am Soc Nephrol* 1991;2:1085–1091.
- 19 Ludlam HA, Young AE, Berry AJ, Phillips I: The prevention of infection with *Staphylococcus aureus* in continuous ambulatory peritoneal dialysis. *J Hosp Infect* 1989;14:293–301.
- 20 Kallen AJ, Jernigan JA, Patel PR: Decolonization to prevent infections with *Staphylococcus aureus* in patients undergoing hemodialysis: a review of current evidence. *Semin Dial* 2011;24:533–539.
- 21 Miller LG, Perdreau-Remington F, Bayer AS, Diep B, Tan N, Bharadwa K, Tsui J, Perlroth J, Shay A, Tagudar G, Ibebuogu U, Spellberg B: Clinical and epidemiologic characteristics cannot distinguish community-associated methicillin-resistant *Staphylococcus aureus* infection from methicillin-susceptible *S. aureus* infection: a prospective investigation. *Clin Infect Dis* 2007;44:471–482.
- 22 McDougal LK, Steward CD, Killgore GE, Chaitram JM, McAllister SK, Tenover FC: Pulsed-field gel electrophoresis typing of oxacillin-resistant *Staphylococcus aureus* isolates from the United States: establishing a national database. *J Clin Microbiol* 2003;41:5113–5120.
- 23 McKinnell JA, Huang SS, Eells SJ, Cui E, Miller LG: Quantifying the impact of extranasal testing of body sites for methicillin-resistant *Staphylococcus aureus* colonization at the time of hospital or intensive care unit admission. *Infect Control Hosp Epidemiol* 2013;34:161–170.
- 24 Rebmann T, Barnes SA; Association for Professionals in Infection Control and Epidemiology: Preventing infections in hemodialysis: an executive summary of the APIC Elimination Guide. *Am J Infect Control* 2011;39:72–75.

Acknowledgements

We would like to thank Reginald Winston and Diana Romero for assistance with collection of study data. This work was supported in part by the National Institutes of Health-National Institute for Diabetes and Digestive and Kidney Diseases STEP-UP program.

Disclosure Statement

The authors declare that they have no conflicts of interest.