

UC Irvine

UC Irvine Previously Published Works

Title

The Aizenman-Sims-Starr scheme and Parisi formula for mixed p-spin spherical models

Permalink

<https://escholarship.org/uc/item/60s9h8mw>

Journal

Electronic Journal of Probability, 18(0)

ISSN

1083-6489

Author

Chen, Wei-Kuo

Publication Date

2013-10-29

DOI

10.1214/EJP.v18-2580

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

The Aizenman-Sims-Starr scheme and Parisi formula for mixed p -spin spherical models

Wei-Kuo Chen*

Abstract

The Parisi formula for the free energy in the spherical models with mixed even p -spin interactions was proven in Michel Talagrand [16]. In this paper we study the general mixed p -spin spherical models including p -spin interactions for odd p . We establish the Aizenman-Sims-Starr scheme and from this together with many well-known results and Dmitry Panchenko's recent proof on the Parisi ultrametricity conjecture [11], we prove the Parisi formula.

Keywords: Parisi formula; ultrametricity.

AMS MSC 2010: 60K35; 82B44.

Submitted to EJP on January 28, 2013, final version accepted on October 14, 2013.

1 Introduction and main results.

In the past decades, the Sherrington-Kirkpatrick model [13] of Ising spin glass has been intensively studied with the aim of understanding the strange magnetic behavior of certain alloys. Despite only being a mean-field model, the formula for its free energy has an intricate and beautiful variational formula, first discovered by G. Parisi [8, 9], based on a hierarchical replica symmetry breaking scheme. As it is hard to compute the free energy from the Parisi formula, the closely related spherical model was studied by A. Crisanti and H. J. Sommers [4]; this model is widely believed to retain the main features of the Ising case, and the analogous Parisi formula admits a more explicit representation for the free energy.

Mathematically, the Parisi formula for the Ising and spherical spin glass models was proven rigorously and generalized to mixtures of p -spin interactions in the seminal works of M. Talagrand [16, 17] following the discovery of the replica symmetry breaking interpolation scheme by F. Guerra [7]. For technical reasons, only the mixture of even p -spin interactions was considered. The main missing ingredient in proving the general case was the validity of the ultrametric property for the Gibbs measure, which was recently verified by D. Panchenko [11]. This together with a series of known results led to the Parisi formula for the Ising spin glass model with general mixed p -spin interactions [12].

*University of California at Irvine, USA. E-mail: weikuoc@uci.edu

In view of the approach in [12], the most crucial step is the derivation of the inequality that the liminf of the free energy is bounded from below by the Parisi formula. It is well-known, in the case of mixed even p -spin models, that the limiting free energy can be represented by a new variational formula through the Aizenman-Sims-Starr (A.S.S.) scheme [1]. An important implication of this scheme is that it provides a natural lower bound for the liminf of the free energy for *general* mixed p -spin models. Since under a small perturbation of the Hamiltonian the asymptotic Gibbs measures in the A.S.S. scheme satisfy the Ghirlanda-Guerra (G.G.) identities, the main result in [11] implies that the support of these measures are asymptotically ultrametric. This property then allows us to characterize the asymptotic Gibbs measure via the Poisson-Dirichlet cascades and consequently recover the Parisi formula in the general mixed p -spin case.

In the present paper, we will establish the Aizenman-Sims-Starr (A.S.S.) scheme and prove the Parisi formula in the spherical model with general mixed p -spin interactions taking the approach of [12]. However, the task turned out to be much more difficult, mostly, due to the fact that uniform measure on the sphere is not a product measure. In particular, as we shall see in the proof of Theorem 2.1 below, the establishment of the A.S.S. scheme in this case becomes much more intricate since in general its construction requires a certain decoupling of the configuration space. Also, the approximation of the Parisi formula using the ultrametricity of the Gibbs measure will be more involved. These obstacles are the main concerns of the paper and will be studied in the next two sections.

Let us now state the mixed p -spin spherical model. For given $N \geq 1$, we denote by $S_N = \{\boldsymbol{\sigma} \in \mathbb{R}^N : \|\boldsymbol{\sigma}\| = \sqrt{N}\}$ the configuration space and by λ_N its normalized surface measure. Consider the pure p -spin Hamiltonians $H_{N,p}(\boldsymbol{\sigma})$ for $p \geq 1$ indexed by $\boldsymbol{\sigma} \in S_N$,

$$H_{N,p}(\boldsymbol{\sigma}) = \frac{1}{N^{(p-1)/2}} \sum_{1 \leq i_1, \dots, i_p \leq N} g_{i_1, \dots, i_p} \sigma_{i_1} \cdots \sigma_{i_p}, \tag{1.1}$$

where the random variables g_{i_1, \dots, i_p} for all $p \geq 1$ and all (i_1, \dots, i_p) are i.i.d. standard Gaussian. The Hamiltonian of the mixed p -spin spherical model is given by

$$H_N(\boldsymbol{\sigma}) = \sum_{p \geq 1} \beta_p H_{N,p}(\boldsymbol{\sigma}) \tag{1.2}$$

with coefficients (β_p) that satisfy $\beta_p \geq 0$ and decrease fast enough, $\sum_{p \geq 1} 2^p \beta_p^2 < \infty$. This technical assumption ensures that on S_N the series (1.2) converges a.s. and the covariance of the Gaussian process is given by a function of the normalized scalar product, called the overlap, $R_{1,2} = N^{-1} \sum_{i \leq N} \sigma_i^1 \sigma_i^2$ of spin configurations $\boldsymbol{\sigma}^1$ and $\boldsymbol{\sigma}^2$,

$$\mathbb{E} H_N(\boldsymbol{\sigma}^1) H_N(\boldsymbol{\sigma}^2) = N \xi(R_{1,2}),$$

where \mathbb{E} is the expectation with respect to all Gaussian randomness and $\xi(x) = \sum_{p \geq 1} \beta_p^2 x^p$.

Now we formulate the Parisi formula. Given $k \geq 1$, let $\mathbf{m} = (m_\ell)_{0 \leq \ell \leq k}$ and $\mathbf{q} = (q_\ell)_{0 \leq \ell \leq k+1}$ be two sequences satisfying

$$\begin{aligned} 0 &= m_0 \leq m_1 \leq \dots \leq m_{k-1} \leq m_k = 1, \\ 0 &= q_0 \leq q_1 \leq \dots \leq q_k \leq q_{k+1} = 1. \end{aligned}$$

One may think of the triplet $(k, \mathbf{m}, \mathbf{q})$ as a distribution function \mathbf{x} on $[0, 1]$ with $x(q) = m_\ell$ if $q_\ell \leq q < q_{\ell+1}$ for $0 \leq \ell \leq k$ and $x(1) = 1$. We define for $1 \leq \ell \leq k$,

$$d_\ell = \sum_{\ell \leq p \leq k} m_p (\xi'(q_{p+1}) - \xi'(q_p)).$$

For a given parameter $b > d_1$, we define $D_\ell = b - d_\ell$ for $1 \leq \ell \leq k$ and $D_{k+1} = b$. Let $\theta(x) = x\xi'(x) - \xi(x)$. The formulation of the Parisi formula involves a functional \mathcal{P} of the distribution function \mathbf{x} ,

$$\mathcal{P}(\mathbf{x}) = \inf_{b > d_1} \frac{1}{2} \left(b - 1 - \log b + \frac{\xi'(q_1)}{D_1} + \sum_{1 \leq \ell \leq k} \frac{1}{m_\ell} \log \frac{D_{\ell+1}}{D_\ell} - \sum_{1 \leq \ell \leq k} m_\ell (\theta(q_{\ell+1}) - \theta(q_\ell)) \right),$$

which is the limit of the Parisi functional described below. For every $M \geq 1$, let $\mathbf{z}_p = (z_{p,1}, \dots, z_{p,M})$ for $0 \leq p \leq k$ be centered Gaussian random vectors with $\mathbb{E}z_{p,i}z_{p,i'} = \delta_{i,i'}(\xi'(q_{p+1}) - \xi'(q_p))$ for any $i \neq i'$ and let $(\mathbf{z}_p)_{0 \leq p \leq k}$ be independent of each other. Starting from

$$X_{k+1}^M = \log \int_{S_M} \exp(\boldsymbol{\varepsilon} \cdot (\mathbf{z}_0 + \dots + \mathbf{z}_k)) d\lambda_M(\boldsymbol{\varepsilon}),$$

we define decreasingly for $0 \leq p \leq k$,

$$X_p^M = \frac{1}{m_p} \log \mathbb{E}_p \exp m_p X_{p+1}^M,$$

where \mathbb{E}_p denotes the expectation in the random vectors $(z_i)_{i \geq p}$. When $m_p = 0$, this means $X_p^M = \mathbb{E}_p X_{p+1}^M$. Set the Parisi functional

$$\mathcal{P}_M(x) = X_0^M - \frac{1}{2} \sum_{1 \leq p \leq k} m_p (\theta(q_{p+1}) - \theta(q_p)).$$

A series of nontrivial applications of elementary large deviation principle implies $\mathcal{P}(\mathbf{x}) = \lim_{M \rightarrow \infty} \mathcal{P}_M(\mathbf{x})$ (see Proposition 3.1 [16]). Let us denote the partition function of the model by

$$Z_N = \int_{S_N} \exp H_N(\boldsymbol{\sigma}) d\lambda_N(\boldsymbol{\sigma}).$$

Below is our main result.

Theorem 1.1 (The Parisi formula). *We have*

$$\lim_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log Z_N = \inf_{\mathbf{x}} \mathcal{P}(\mathbf{x}), \tag{1.3}$$

where the infimum is taken over all possible choices of \mathbf{x} .

The quantity in the limit on the left-hand side of (1.3) is usually called the free energy of this model and the formula on the right-hand side of (1.3) is the famous Parisi formula in the spherical form, which is also known to have the Crisanti-Sommers representation [4] (see Theorem 4.1 in [16]). Of course, one may also consider the model in the present of external field and prove the equation (1.3), but for the clarity of our discussion, we will only consider the Hamiltonian of the form (1.2).

2 The A.S.S. scheme for spherical models.

The main result of this section is the construction of the A.S.S. scheme for spherical models. For $M, N \geq 1$, let us denote by $(z(\boldsymbol{\sigma}) = (z_1(\boldsymbol{\sigma}), \dots, z_M(\boldsymbol{\sigma})) : \boldsymbol{\sigma} \in S_N)$ and $(y(\boldsymbol{\sigma}) : \boldsymbol{\sigma} \in S_N)$ two independent families of centered Gaussian random variables with covariances

$$\mathbb{E}z_i(\boldsymbol{\sigma}^1)z_{i'}(\boldsymbol{\sigma}^2) = \delta_{i,i'}\xi'(R_{1,2}) \tag{2.1}$$

$$\mathbb{E}y(\boldsymbol{\sigma}^1)y(\boldsymbol{\sigma}^2) = \theta(R_{1,2}) \tag{2.2}$$

for every $\sigma^1, \sigma^2 \in S_N$. These random variables are independent of all the other randomness. Let $G_{M,N}$ be the Gibbs measure corresponding to the Hamiltonian

$$H_{M,N}(\sigma) = \sum_{p \geq 1} \frac{\beta_p}{(M+N)^{(p-1)/2}} \sum_{1 \leq i_1, \dots, i_p \leq N} g_{i_1, \dots, i_p} \sigma_{i_1} \cdots \sigma_{i_p} \tag{2.3}$$

with reference measure λ_N on S_N and let $\langle \cdot \rangle_{M,N}$ be the Gibbs average of $G_{M,N}$. The following is our main result.

Theorem 2.1 (The A.S.S. scheme). *For any $\delta > 0$ and $M \geq 1$, the liminf of $N^{-1} \mathbb{E} \log Z_N$ is bounded from below by*

$$\begin{aligned} \frac{1}{M} \liminf_{N \rightarrow \infty} & \left(\mathbb{E} \log \int_{S_M} \langle \exp(\varepsilon \cdot z(\sigma)) \rangle_{M,N} d\lambda_M(\varepsilon) - \mathbb{E} \log \langle \exp(\sqrt{M}y(\sigma)) \rangle_{M,N} \right) \\ & - \delta \xi'(1) + \frac{1}{M} \log \nu_M(A_\delta), \end{aligned} \tag{2.4}$$

where ν_M is the probability measure of M independent standard Gaussian r.v.s and

$$A_\delta = \{ \varepsilon \in \mathbb{R}^M : M \leq \|\varepsilon\|^2 \leq M(1 + \delta) \}.$$

Note that a standard large deviation result yields $\lim_{M \rightarrow \infty} M^{-1} \log \nu_M(A_\delta) = 0$ for every $\delta > 0$. Thus (2.4) gives a lower bound for the liminf of the free energy that will play an essential role in proving the Parisi formula in Section 3. Suppose that μ_N is the uniform probability measure on $\Sigma_N = \{-1, +1\}^N$. If we replace the measure space (S_N, λ_N) by (Σ_N, μ_N) and ignore the error term $-\delta \xi'(1) + M^{-1} \log \nu_M(A_\delta)$ in (2.4), then Theorem 2.1 gives the A.S.S. scheme for the Ising spin glass model with general mixed p -spin interactions (see e.g. Section 15.8 [18]).

We will follow the spirit in [1] to establish the A.S.S. scheme. However, since uniform measure on the spherical configuration space in the spherical models is not a product measure, we have to seek a proper way to approximate this measure by another one that possesses a certain product structure. To deal with this obstacle, we will use the idea of Poincaré’s lemma (see page 77 [14]) that the projection on any fixed M coordinates of the uniform probability measure λ_{M+N} on the sphere S_{M+N} converge weakly to M -dimensional standard Gaussian measure as N tends to infinity. The argument of our proof proceeds as follows. Recall the definition of H_{M+N} from (1.2). For $\rho \in S_{M+N}$, let us write $\rho = (\sigma, \varepsilon)$ for the first N coordinates $\sigma = (\sigma_1, \dots, \sigma_N)$ and the last M cavity coordinates $\varepsilon = (\varepsilon_1, \dots, \varepsilon_M)$ and write

$$H_{M+N}(\rho) = H_{M,N}(\sigma) + \sum_{i \leq M} \varepsilon_i Z_i(\sigma) + \gamma(\rho),$$

where $H_{M,N}$ is given in (2.3), the term $\varepsilon_i Z_i(\sigma)$ consists of all terms in H_{M+N} with only one factor ε_i from ε present, and the last term is the sum of terms with at least two factors in ε . From now on, without causing any ambiguity, for any given vectors $\omega, \omega' \in \mathbb{R}^K$ for some $K \geq 1$, we define $R(\omega, \omega') = K^{-1} \sum_{i \leq K} \omega_i \omega'_i$. One can check easily that

$$\mathbb{E} H_{M,N}(\sigma^1) H_{M,N}(\sigma^2) = (M+N) \xi \left(\frac{N}{M+N} R(\sigma^1, \sigma^2) \right), \tag{2.5}$$

$$\mathbb{E} Z_i(\sigma^1) Z_i(\sigma^2) = \xi' \left(\frac{N}{M+N} R(\sigma^1, \sigma^2) \right), \tag{2.6}$$

$$\mathbb{E} \gamma(\rho)^2 \leq \frac{M^2}{M+N} \xi''(1). \tag{2.7}$$

Let us notice an elementary fact that for a real-valued sequence (c_N) , if $\liminf_{N \rightarrow \infty} c_N/N = r \in \mathbb{R}$, then $M^{-1} \liminf_{N \rightarrow \infty} (c_{M+N} - c_N) \leq r$ for every $M \geq 1$. This implies

$$\liminf_{N \rightarrow \infty} \frac{\mathbb{E} \log Z_N}{N} \geq \frac{1}{M} \liminf_{N \rightarrow \infty} \mathbb{E} \log \frac{Z_{M+N}}{Z_N} \tag{2.8}$$

and thus to obtain Theorem 2.1, it suffices to prove that (2.4) is the lower bound of the right-hand side of (2.8). Let us write $\mathbb{E} \log Z_{M+N}/Z_N$ as

$$\begin{aligned} & \mathbb{E} \log \int_{A_\delta} \langle \exp(\varepsilon \cdot z(\sigma)) \rangle_{M,N} d\nu_M(\varepsilon) - \mathbb{E} \log \frac{\int_{S_N} \exp H_N d\lambda_N(\sigma)}{\int_{S_N} \exp H_{M,N} d\lambda_N(\sigma)} \\ & + \mathbb{E} \log \frac{Z_{M+N}}{\int_{A_\delta} J_{M,N} d\nu_M(\varepsilon)}, \end{aligned} \tag{2.9}$$

where

$$J_{M,N}(\varepsilon) = \int_{S_N} \exp(H_{M,N}(\sigma) + \varepsilon \cdot z(\sigma)) d\lambda_N(\sigma), \quad \varepsilon \in \mathbb{R}^M.$$

In what follows, we prove several lemmas that will be used to bound these three terms in (2.9) as N tends to infinity.

Lemma 2.2. For $z \in \mathbb{R}^M$, we have

$$\int_{A_\delta} \exp(\varepsilon \cdot z) d\nu_M(\varepsilon) \geq \nu_M(A_\delta) \int_{S_M} \exp(\varepsilon \cdot z) d\lambda_M(\varepsilon). \tag{2.10}$$

Proof. Using the rotational invariance of ν_M , there exists a probability measure γ_M on \mathbb{R}^+ such that ν_M is the image of $\gamma_M \times \lambda_M$ under the map $(s, \varepsilon) \mapsto s\varepsilon$. For $1 \leq s \leq (1 + \delta)^{1/2}$, using symmetry,

$$\begin{aligned} \int \exp(s\varepsilon \cdot z) d\lambda_M(\varepsilon) &= \int \cosh(s\varepsilon \cdot z) d\lambda_M(\varepsilon) \\ &\geq \int \cosh(\varepsilon \cdot z) d\lambda_M(\varepsilon) = \int \exp(\varepsilon \cdot z) d\lambda_M(\varepsilon). \end{aligned}$$

Therefore, (2.10) holds by

$$\begin{aligned} \int_{A_\delta} \exp(\varepsilon \cdot z) d\nu_M(\varepsilon) &= \int_1^{(1+\delta)^{1/2}} \int_{S_M} \exp(s\varepsilon \cdot z) d\lambda_M(\varepsilon) d\gamma_M(s) \\ &\geq \int_1^{(1+\delta)^{1/2}} \int_{S_M} \exp(\varepsilon \cdot z) d\lambda_M(\varepsilon) d\gamma_M(s) \\ &= \gamma_M(\{1 \leq s \leq (1 + \delta)^{1/2}\}) \int_{S_M} \exp(\varepsilon \cdot z) d\lambda_M(\varepsilon). \end{aligned}$$

□

From this lemma and Fubini's theorem, the first term in (2.9) is bounded from below by

$$\mathbb{E} \log \int_{A_\delta} \langle \exp(\varepsilon \cdot z(\sigma)) \rangle_{M,N} d\lambda_M(\varepsilon) + \log \nu_M(A_\delta) \tag{2.11}$$

Using the standard Gaussian interpolation technique, Lemma 2.3 below takes care of the second term in (2.9).

Lemma 2.3. *We have*

$$\lim_{N \rightarrow \infty} \left| \mathbb{E} \log \frac{\int_{S_N} \exp H_N(\boldsymbol{\sigma}) d\lambda_N(\boldsymbol{\sigma})}{\int_{S_N} \exp H_{M,N}(\boldsymbol{\sigma}) d\lambda_N(\boldsymbol{\sigma})} - \mathbb{E} \log \left\langle \exp(\sqrt{M}y(\boldsymbol{\sigma})) \right\rangle_{M,N} \right| = 0. \quad (2.12)$$

Proof. Suppose $(y'(\boldsymbol{\sigma}) : \boldsymbol{\sigma} \in S_N)$ is a family of centered Gaussian random variables independent of all the other randomness such that in distribution

$$H_N(\boldsymbol{\sigma}) = H_{M,N}(\boldsymbol{\sigma}) + \sqrt{M}y'(\boldsymbol{\sigma}).$$

It is easy to check that $\mathbb{E}y'(\boldsymbol{\sigma}^1)y'(\boldsymbol{\sigma}^2) = \sum_{p \geq 1} \beta_p^2 D_p N^p R(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2)^p$, where

$$D_p = \frac{1}{M} \left(\frac{1}{N^{p-1}} - \frac{1}{(M+N)^{p-1}} \right)$$

satisfies $N^p D_p \rightarrow p - 1$. Recall $y(\boldsymbol{\sigma})$ from (2.2). Let us consider the interpolating free energy

$$\psi(t) = \mathbb{E} \log \left\langle \exp \left(\sqrt{M}(\sqrt{t}y'(\boldsymbol{\sigma}) + \sqrt{1-t}y(\boldsymbol{\sigma})) \right) \right\rangle_{M,N}.$$

Then $|\psi(1) - \psi(0)|$ gives the quantity in the limit on the left-hand side of (2.12). Thus it suffices to prove that $\psi'(t) \rightarrow 0$ uniformly in $t \in (0, 1)$ as $N \rightarrow \infty$, which can be easily verified by using Gaussian integration by parts, as $N \rightarrow \infty$,

$$\begin{aligned} \sup_{0 < t < 1} |\psi'(t)| &\leq M \sup_{\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \in S_N} |\mathbb{E}y'(\boldsymbol{\sigma}^1)y'(\boldsymbol{\sigma}^2) - \mathbb{E}y(\boldsymbol{\sigma}^1)y(\boldsymbol{\sigma}^2)| \\ &= M \sup_{\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2 \in S_N} \left| \sum_{p \geq 1} \beta_p^2 (D_p N^p - (p-1)) R(\boldsymbol{\sigma}^1, \boldsymbol{\sigma}^2)^p \right| \rightarrow 0. \end{aligned}$$

□

Let us continue to find a lower bound for the liminf of the third term in (2.9). It relies on an elementary lemma that allows us to decouple the surface measure (S_{M+N}, λ_{M+N}) as the product measure $(S_N \times A_\delta, \lambda_N \times F_M)$ asymptotically. For $K \geq 1$, we denote by S_K^1 the unit sphere in \mathbb{R}^K and by $|S_K^1|$ the area of S_K^1 . Set

$$\begin{aligned} A_{M,N} &= \prod_{j=1}^M \left[-\sqrt{M+N+1-j}, \sqrt{M+N+1-j} \right], \\ b_{M,N} &= \prod_{j=1}^M \frac{|S_{M+N-j}^1|}{|S_{M+N+1-j}^1| \sqrt{M+N+1-j}}. \end{aligned}$$

Define

$$\begin{aligned} F_{M,N}(\boldsymbol{\varepsilon}) &= b_{M,N} \prod_{j=1}^M \left(1 - \frac{\varepsilon_j^2}{M+N+1-j} \right)^{\frac{M+N-j-2}{2}}, \quad \boldsymbol{\varepsilon} \in A_{M,N}, \\ F_M(\boldsymbol{\varepsilon}) &= \left(\frac{1}{2\pi} \right)^{M/2} \exp \left(-\frac{\|\boldsymbol{\varepsilon}\|^2}{2} \right), \quad \boldsymbol{\varepsilon} \in \mathbb{R}^M. \end{aligned}$$

Define $a_1 = 1$ and for $2 \leq \ell \leq M+1$,

$$a_\ell(\boldsymbol{\varepsilon}) = \prod_{j=1}^{\ell-1} \sqrt{1 + \frac{1 - \varepsilon_j^2}{M+N-j}}, \quad \boldsymbol{\varepsilon} \in A_{M,N}.$$

Lemma 2.4. *Suppose that f is a nonnegative function defined on S_{M+N} . Then we have*

$$\int_{S_{M+N}} f(\boldsymbol{\rho}) d\lambda_{M+N}(\boldsymbol{\rho}) = \int_{A_{M,N}} F_{M,N}(\boldsymbol{\varepsilon}) d\boldsymbol{\varepsilon} \int_{S_N} f(\boldsymbol{\sigma} a_{M+1, \varepsilon_1} a_1, \dots, \varepsilon_M a_M) d\lambda_N(\boldsymbol{\sigma}). \tag{2.13}$$

Proof. This is an elementary calculus formula that follows by an induction argument starting from

$$\int_{S_K} f(\boldsymbol{\rho}) d\lambda_K(\boldsymbol{\rho}) = \frac{|S_{K-1}^1|}{|S_K^1| \sqrt{K}} \int_{-\sqrt{K}}^{\sqrt{K}} \left(1 - \frac{\varepsilon^2}{K}\right)^{\frac{K-3}{2}} d\varepsilon \int_{S_{K-1}} f\left(\boldsymbol{\sigma} \sqrt{\frac{K - \varepsilon^2}{K-1}}, \varepsilon\right) d\lambda_{K-1}(\boldsymbol{\sigma})$$

for every $K \geq 2$. □

As an immediate consequence of Lemma 2.4, the cavity coordinates converge weakly to the M -dimensional standard Gaussian random vector as N tends to infinity. We write $\mathbb{E} \log Z_{M+N} / \int_{A_\delta} J_{M,N} d\nu_M(\boldsymbol{\varepsilon})$ as

$$\mathbb{E} \log \frac{Z_{M+N}}{\int_{A_\delta} F_{M,N} J_{M,N} d\boldsymbol{\varepsilon}} + \mathbb{E} \log \frac{\int_{A_\delta} F_{M,N} J_{M,N} d\boldsymbol{\varepsilon}}{\int_{A_\delta} F_M J_{M,N} d\boldsymbol{\varepsilon}}. \tag{2.14}$$

Then the first term of (2.14) can be controlled by the following lemma.

Lemma 2.5. *We have*

$$\liminf_{N \rightarrow \infty} \mathbb{E} \log \frac{Z_{M+N}}{\int_{A_\delta} F_{M,N} J_{M,N} d\boldsymbol{\varepsilon}} \geq -M\delta\xi'(1). \tag{2.15}$$

Proof. Suppose that $(H'_{M,N}(\boldsymbol{\sigma}) : \boldsymbol{\sigma} \in S_N)$ is an independent copy of $(H_{M,N}(\boldsymbol{\sigma}) : \boldsymbol{\sigma} \in S_N)$ and is independent of all the other randomness. We consider the following interpolating Hamiltonian

$$\varphi(t) = \mathbb{E} \log \int_{A_\delta} F_{M,N}(\boldsymbol{\varepsilon}) d\boldsymbol{\varepsilon} \int_{S_N} \exp(H_{1,t}(\boldsymbol{\rho}) + H_{2,t}(\boldsymbol{\rho}) + H_{3,t}(\boldsymbol{\rho})) d\lambda_N(\boldsymbol{\sigma}),$$

where for $\boldsymbol{\rho} = (\boldsymbol{\sigma}, \boldsymbol{\varepsilon})$ with $\boldsymbol{\sigma} \in S_N$ and $\boldsymbol{\varepsilon} \in A_\delta$,

$$\begin{aligned} H_{1,t}(\boldsymbol{\rho}) &= \sqrt{t} H_{M,N}(\boldsymbol{\sigma} a_{M+1}) + \sqrt{1-t} H'_{M,N}(\boldsymbol{\sigma}) \\ H_{2,t}(\boldsymbol{\rho}) &= \sum_{i \leq M} \varepsilon_i \left(\sqrt{t} a_i Z_i(\boldsymbol{\sigma} a_{M+1}) + \sqrt{1-t} z_i(\boldsymbol{\sigma}) \right) \\ H_{3,t}(\boldsymbol{\rho}) &= \sqrt{t} \gamma(\boldsymbol{\sigma} a_{M+1}, \varepsilon_1 a_1, \dots, \varepsilon_M a_M). \end{aligned}$$

From (2.13), obviously $\varphi(1) \leq \mathbb{E} \log Z_{M+N}$ when N is large enough. On the other hand, using Gaussian integration by parts and replicas $\boldsymbol{\rho}^1 = (\boldsymbol{\sigma}^1, \boldsymbol{\varepsilon}^1)$ and $\boldsymbol{\rho}^2 = (\boldsymbol{\sigma}^2, \boldsymbol{\varepsilon}^2)$, it follows that

$$\varphi'(t) = \frac{1}{2} \sum_{j \leq 3} (\mathbb{E} \langle U_j(\boldsymbol{\rho}, \boldsymbol{\rho}) \rangle_t - \mathbb{E} \langle U_j(\boldsymbol{\rho}^1, \boldsymbol{\rho}^2) \rangle_t),$$

where $\langle \cdot \rangle_t$ is the Gibbs average corresponding to random weights $F_{M,N} \exp(H_{1,t} + H_{2,t} + H_{3,t})$ with reference measure $\lambda_N \times d\boldsymbol{\varepsilon}$ on $S_N \times A_\delta$ and

$$\begin{aligned} U_1(\boldsymbol{\rho}^1, \boldsymbol{\rho}^2) &= \mathbb{E} H_{M,N}(\boldsymbol{\sigma}^1 a_{M+1}(\boldsymbol{\varepsilon}^1)) H_{M,N}(\boldsymbol{\sigma}^2 a_{M+1}(\boldsymbol{\varepsilon}^2)) - \mathbb{E} H'_{M,N}(\boldsymbol{\sigma}^1) H'_{M,N}(\boldsymbol{\sigma}^2) \\ U_2(\boldsymbol{\rho}^1, \boldsymbol{\rho}^2) &= \sum_{i \leq M} \varepsilon_i^1 \varepsilon_i^2 (a_i(\boldsymbol{\varepsilon}^1) a_i(\boldsymbol{\varepsilon}^2) \mathbb{E} Z_i(\boldsymbol{\sigma}^1 a_{M+1}(\boldsymbol{\varepsilon}^1)) Z_i(\boldsymbol{\sigma}^2 a_{M+1}(\boldsymbol{\varepsilon}^2)) - \mathbb{E} z_i(\boldsymbol{\sigma}^1) z_i(\boldsymbol{\sigma}^2)) \\ U_3(\boldsymbol{\rho}^1, \boldsymbol{\rho}^2) &= \mathbb{E} \gamma(\boldsymbol{\sigma}^1 a_{M+1}(\boldsymbol{\varepsilon}^1), \varepsilon_1^1 a_1(\boldsymbol{\varepsilon}^1), \dots, \varepsilon_M^1 a_M(\boldsymbol{\varepsilon}^1)) \\ &\quad \cdot \gamma(\boldsymbol{\sigma}^2 a_{M+1}(\boldsymbol{\varepsilon}^2), \varepsilon_1^2 a_1(\boldsymbol{\varepsilon}^2), \dots, \varepsilon_M^2 a_M(\boldsymbol{\varepsilon}^2)). \end{aligned}$$

From (2.1), (2.6), and (2.7), it is easy to see that for $j = 2, 3$,

$$\lim_{N \rightarrow \infty} \sup_{\rho^1, \rho^2 \in \mathcal{S}_N \times A_\delta} |U_j(\rho^1, \rho^2)| \rightarrow 0$$

and from (2.5) and using mean value theorem,

$$\begin{aligned} & \lim_{N \rightarrow \infty} \sup_{\rho^1, \rho^2 \in \mathcal{S}_N \times A_\delta} |U_1(\rho^1, \rho^2)| \\ & \leq \lim_{N \rightarrow \infty} \xi'(1)(M+N) \sup_{\varepsilon^1, \varepsilon^2 \in A_\delta} \left| \frac{N}{M+N} a_{M+1}(\varepsilon^1) a_{M+1}(\varepsilon^2) - 1 \right| \\ & = \xi'(1) \sup_{\varepsilon^1, \varepsilon^2 \in A_\delta} \left| M - \frac{\|\varepsilon^1\|^2 + \|\varepsilon^2\|^2}{2} \right| \leq M\delta\xi'(1). \end{aligned}$$

Thus we conclude that $\limsup_{N \rightarrow \infty} \sup_{0 < t < 1} |\varphi'(t)| \leq M\delta\xi'(1)$ and so (2.15) follows by

$$\begin{aligned} \liminf_{N \rightarrow \infty} \mathbb{E} \log \frac{Z_{M+N}}{\int_{A_\delta} F_{M,N} J_{M,N} d\varepsilon} & \geq \liminf_{N \rightarrow \infty} (\varphi(1) - \varphi(0)) \\ & \geq -\limsup_{N \rightarrow \infty} \sup_{0 < t < 1} |\varphi'(t)| \geq -M\delta\xi'(1). \end{aligned}$$

□

Finally, let us deal with the second term of (2.14) below.

Lemma 2.6. *We have*

$$\liminf_{N \rightarrow \infty} \mathbb{E} \log \frac{\int_{A_\delta} F_{M,N} J_{M,N} d\varepsilon}{\int_{A_\delta} F_M J_{M,N} d\varepsilon} \geq 0. \tag{2.16}$$

Proof. Using the inequality $\log(1-x) \geq -2x$ for $0 < x < 1/2$, when N is sufficiently large, we have for every $\varepsilon \in A_\delta$,

$$\begin{aligned} & \log \frac{F_{M,N}(\varepsilon)}{F_M(\varepsilon)} \\ & = \log b_{M,N} - \frac{M}{2} \log \frac{1}{2\pi} + \sum_{j=1}^M \frac{M+N-j-2}{2} \log \left(1 - \frac{\varepsilon_j^2}{M+N+1-j} \right) + \frac{\|\varepsilon\|^2}{2} \\ & \geq \log b_{M,N} - \frac{M}{2} \log \frac{1}{2\pi} + \frac{3\|\varepsilon\|^2}{M+N}. \end{aligned}$$

Note that $\lim_{N \rightarrow \infty} b_{M,N} = 1/(2\pi)^{M/2}$ as can be seen from (2.13) with $f = 1$. By Jensen's inequality, we obtain

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \mathbb{E} \log \frac{\int_{A_\delta} F_M J_{M,N} \exp(\log F_{M,N}/F_M) d\varepsilon}{\int_{A_\delta} F_M J_{M,N} d\varepsilon} \\ & \geq \liminf_{N \rightarrow \infty} \mathbb{E} \frac{\int_{A_\delta} F_M J_{M,N} \log F_{M,N}/F_M d\varepsilon}{\int_{A_\delta} F_M J_{M,N} d\varepsilon} \\ & \geq \liminf_{N \rightarrow \infty} \left(\log b_{M,N} - \frac{M}{2} \log \frac{1}{2\pi} + \frac{3M}{M+N} \right) = 0 \end{aligned}$$

and this gives (2.16).

□

Proof of Theorem 2.1. From (2.8), (2.9), and (2.14), using Lemmas 2.3, 2.5, and 2.6, the \liminf of $N^{-1}\mathbb{E} \log Z_N$ is bounded from below by

$$\frac{1}{M} \liminf_{N \rightarrow \infty} \left(\mathbb{E} \log \int_{A_\delta} \langle \exp(\varepsilon \cdot \mathbf{z}(\boldsymbol{\sigma})) \rangle_{M,N} d\nu_M(\varepsilon) - \mathbb{E} \log \langle \exp(\sqrt{M}y(\boldsymbol{\sigma})) \rangle_{M,N} \right) - \delta\xi'(1)$$

and then (2.11) yields (2.4). □

3 Proof of the Parisi formula for spherical models.

This section is devoted to proving Theorem 1.1. Our argument is started with Guerra’s replica symmetry breaking interpolation scheme in the spherical models (see Section 3 in [16]). In the case when p -spin interactions for odd $p \geq 3$ are not present in (1.2), this scheme implies that the upper limit of the free energy is bounded from above by $\inf_x \mathcal{P}(x)$. The fact that this bound still holds in the general mixed p -spin case relies on Talagrand’s positivity principle on the overlap (see [15] and Section 12.3 in [18]) that is deduced from the validity of the G.G. identities [6] derived by a perturbation trick on the Hamiltonian. This part of the argument is well-known and has been used in several places. For example, the readers are referred to Theorem 14.4.4 [18] for a great detailed discussion.

Our main concern in the proof will be concentrated on obtaining the lower bound, that is, the \liminf of the free energy is bounded from below by $\inf_x \mathcal{P}(x)$, that will be established in several steps and is based on the A.S.S. scheme. Again due to the non-product structure of the uniform measure on the sphere, additional technical issues occur. For instance, in the Ising spin glass model, the argument for the lower bound only uses the A.S.S. scheme with a single cavity coordinate, that is, $M = 1$ (see Theorem 3.5 [2], Lemma 11 [10], or Proposition 2 [12]), but this is not the case in our model. Indeed, recalling the formulation of the Parisi formula in the spherical model, one may figure out immediately that it is necessary to deal with the A.S.S. scheme (2.4) with all $M \geq 1$ and compute their lower limit in M . Under this circumstance, we find the fact that the Parisi functionals $(\mathcal{P}_M)_{M \geq 1}$ have the same Lipschitz constant with respect to the L_1 -norm as stated in Lemma 3.5 below is of great use.

The A.S.S. scheme for the perturbed Hamiltonian. For each $u = (u_p)_{p \geq 1}$ with $1 \leq u_p \leq 2$, let us consider a perturbation Hamiltonian

$$H_N^{pert}(\boldsymbol{\sigma}) = \frac{1}{N^{1/8}} \sum_{p \geq 1} \frac{u_p}{2^p} H'_{N,p}(\boldsymbol{\sigma}),$$

where $H'_{N,p}(\boldsymbol{\sigma})$ are independent copies of the p -spin Hamiltonian in (1.1). Since the perturbation term is of a small order, replacing H_N with $H_N + H_N^{pert}$ in (1.3) obviously does not affect the limit and in addition, the following A.S.S. scheme also holds. Consider another perturbation Hamiltonian

$$H_{M,N}^{pert}(\boldsymbol{\sigma}) = \frac{1}{(M+N)^{1/8}} \sum_{p \geq 1} \frac{u_p}{2^p} \left(\frac{N}{M+N} \right)^{(p-1)/2} H'_{N,p}(\boldsymbol{\sigma}).$$

Let $G_{M,N}^-$ be the Gibbs measure corresponding to the Hamiltonian $H_{M,N} + H_{M,N}^{pert}$ with reference measure λ_N and $\langle \cdot \rangle_{M,N}^-$ be the average with respect to the product Gibbs measure $G_{M,N}^{-\otimes \infty}$.

Proposition 3.1 (The A.S.S. scheme). *Recall A_δ and ν_M from Theorem 2.1. For every $\delta > 0$ and $M \geq 1$, we have*

$$\begin{aligned} & \liminf_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log Z_N \\ & \geq \frac{1}{M} \liminf_{N \rightarrow \infty} \sup_u \left(\mathbb{E} \log \int_{S_M} \langle \exp(\varepsilon \cdot \mathbf{z}(\boldsymbol{\sigma})) \rangle_{M,N}^- d\lambda_M(\varepsilon) - \mathbb{E} \log \langle \exp(\sqrt{M}y(\boldsymbol{\sigma})) \rangle_{M,N}^- \right) \\ & \quad - \delta \xi'(1) + \frac{1}{M} \log \nu_M(A_\delta), \end{aligned} \tag{3.1}$$

where the supremum is taken over all $u = (u_p)_{p \geq 1}$ with $1 \leq u_p \leq 2$.

Starting with the Hamiltonian $H_{M+N} + H_{M+N}^{pert}$ and using the fact that H_{M+N}^{pert} is of a small order, one may easily find that the same arguments in the proof of Theorem 2.1 also work in Proposition 3.1, so we will leave this part of the proof to the readers. In what follows, for each $M \geq 1$, we will construct a sequence $(u^{M,N})_{N \geq 1}$ for $u^{M,N} = (u_p^{M,N})_{p \geq 1}$ with $1 \leq u_p^{M,N} \leq 2$ along which the G.G. identities for the Gibbs measure $G_{M,N}^-$ and the A.S.S. scheme for the Hamiltonian $H_{M,N} + H_{M,N}^{pert}$ hold simultaneously. Let $(\boldsymbol{\sigma}^\ell)_{\ell \geq 1}$ be an i.i.d. sample from $G_{M,N}^-$ and let $R^{M,N} = (R_{\ell,\ell'}^{M,N})_{\ell,\ell' \geq 1}$ be the normalized Gram matrix, or matrix of overlaps, of this sample. Let \mathcal{C} be the collection of all (p, n, f) with $p \geq 1$, $n \geq 2$, and f being a monomial of $(R_{\ell,\ell'}^{M,N})_{\ell,\ell' \leq n}$. We define

$$\begin{aligned} & \Phi_{M,N}(p, n, f) \\ & = \left| \mathbb{E} \langle f(R_{1,n+1}^{M,N})^p \rangle_{M,N}^- - \frac{1}{n} \mathbb{E} \langle f \rangle_{M,N}^- \mathbb{E} \langle (R_{1,2}^{M,N})^p \rangle_{M,N}^- - \frac{1}{n} \sum_{\ell=2}^n \mathbb{E} \langle f(R_{1,\ell}^{M,N})^p \rangle_{M,N}^- \right| \end{aligned}$$

for $(p, n, f) \in \mathcal{C}$. Then the G.G. identities can be stated as follows (see Theorem 12.3.1 [18]).

Proposition 3.2. *For $M \geq 1$, we have*

$$\lim_{N \rightarrow \infty} \mathbb{E}_u \Phi_{M,N}(p, n, f) = 0 \tag{3.2}$$

for all $(p, n, f) \in \mathcal{C}$, where we view $u = (u_p)_{p \geq 1}$ as a sequence of i.i.d. uniform random variables on $[1, 2]$ and \mathbb{E}_u is the expectation with respect to the randomness of u .

Since \mathcal{C} is a countable set, we can enumerate it as $\{(p_j, n_j, f_j)\}_{j \geq 1}$. Let us define

$$\begin{aligned} \Phi_{M,N}(u) &= \sum_{j \geq 1} \frac{1}{2^j} \Phi_{M,N}(p_j, n_j, f_j), \\ \Lambda_{M,N}(u) &= \frac{1}{M} \left(\mathbb{E} \log \int_{S_M} \langle \exp(\varepsilon \cdot \mathbf{z}(\boldsymbol{\sigma})) \rangle_{M,N}^- d\lambda_M(\varepsilon) - \mathbb{E} \log \langle \exp(\sqrt{M}y(\boldsymbol{\sigma})) \rangle_{M,N}^- \right). \end{aligned}$$

By Chebyshev's inequality, $\mathbb{P}_u(\Phi_{M,N} \leq \eta) \geq 1 - \mathbb{E}_u \Phi_{M,N} / \eta$ for every $\eta > 0$. If we pick $\eta = 2\mathbb{E}_u \Phi_{M,N}$, it implies that for any M and N we can always find $u^{M,N} = (u_p^{M,N})_{p \geq 1}$ such that $\Phi_{M,N}(u^{M,N}) \leq 2\mathbb{E}_u \Phi_{M,N}$. Note that using $\Phi_{M,N}(n, p, f) \leq 2$ and the G.G. identities (3.2) implies $\lim_{N \rightarrow \infty} \mathbb{E}_u \Phi_{M,N} = 0$. Consequently, we have

$$\lim_{N \rightarrow \infty} \Phi_{M,N}(u^{M,N}) = 0 \tag{3.3}$$

and from (3.1), $\liminf_{N \rightarrow \infty} N^{-1} \mathbb{E} \log Z_N$ is bounded from below by

$$\liminf_{N \rightarrow \infty} \Lambda_{M,N}(u^{M,N}) - \delta \xi'(1) + \frac{1}{M} \log \nu_M(A_\delta). \tag{3.4}$$

In other words, for each $M \geq 1$, both the G.G. identities and the A.S.S. scheme hold along $(u^{M,N})_{N \geq 1}$. Let us redefine the Hamiltonian $H_{M,N}^{pert}$ and the Gibbs measure $G_{M,N}^-$ by fixing parameters $u = u^{M,N}$.

Asymptotic Gibbs' measures. Next, we will find a sequence of asymptotic Gibbs' measures that satisfy the G.G. identities and represent the lower limit in (3.4). For each $M \geq 1$, let us pick a subsequence (N_n) along which the limits in (3.3) and (3.4) are achieved and the distribution $R^{M,N}$ under $\mathbb{E}G_{M,N}^{-\otimes \infty}$ converges in the sense of convergence of finite dimensional distributions to the distribution of some array R^M . To lighten the notation, we assume without loss of generality that the sequence (N_n) coincides with the natural numbers. Under $\mathbb{E}G_{M,N}^{-\otimes \infty}$, the array $R^{M,N}$ is weakly exchangeable, that is,

$$\left(R_{\pi(\ell), \pi(\ell')}^{M,N} \right) \stackrel{d}{=} \left(R_{\ell, \ell'}^{M,N} \right)$$

for any permutation π of finitely many indices. This property will still be preserved in the limit such that R^M is a weakly exchangeable symmetric non-negative definite array. Such an array is called a Gram-de Finetti array and is known to have the Dobysh-Sudakov representation [5].

Proposition 3.3 (The Dobysh-Sudakov representation). *If $(R_{\ell, \ell'})_{\ell, \ell' \geq 1}$ is a Gram-de Finetti array such that $R_{\ell, \ell} = 1$, then there exists a random measure G on the unit ball of a separable Hilbert space such that*

$$(R_{\ell, \ell'})_{\ell, \ell' \geq 1} \stackrel{d}{=} \left(\tau^\ell \cdot \tau^{\ell'} + \delta_{\ell, \ell'} (1 - \|\tau^\ell\|^2) \right)_{\ell, \ell' \geq 1},$$

where (τ^ℓ) is an i.i.d. sample from G .

Obviously the array R^M satisfies $R_{\ell, \ell}^M = 1$. Using Proposition 3.3, let us denote by G_M the random measure generating the array R^M and by (τ^ℓ) an i.i.d. sample from G_M . We also denote by $\langle \cdot \rangle_M$ the average with respect to G_M . From (3.3) and an approximation argument, the measure G_M satisfies the G.G. identities,

$$\mathbb{E} \langle f \psi(R_{1, n+1}^M) \rangle_M = \frac{1}{n} \mathbb{E} \langle f \rangle_M \mathbb{E} \langle \psi(R_{1, 2}^M) \rangle_M + \frac{1}{n} \sum_{\ell=2}^n \mathbb{E} \langle f \psi(R_{1, \ell}^M) \rangle_M \quad (3.5)$$

for all $n \geq 2$, all bounded measurable function f of $(R_{\ell, \ell'})_{\ell, \ell' \leq n}$, and measurable function ψ . These identities will be used to identify the asymptotic Gibbs measures G_M in next subsection.

Let us continue to check that the limit of $\Lambda_{M,N}(u^{M,N})$ in N can be expressed in terms of G_M , which relies on the following observation. Suppose for the moment that $R = (R_{\ell, \ell'})_{\ell, \ell' \geq 1}$ is an arbitrary Gram-de Finetti array such that $R_{\ell, \ell} = 1$. Let \mathcal{L} be its distribution and G be any random measure generating R as Proposition 3.3. Suppose that $\mathbf{z}(\boldsymbol{\tau}) = (z_i(\boldsymbol{\tau}))_{i \leq M}$ and $y(\boldsymbol{\tau})$ are two Gaussian processes on the unit ball of some Hilbert space generating R with covariances

$$\mathbb{E} z_i(\boldsymbol{\tau}^1) z_{i'}(\boldsymbol{\tau}^2) = \delta_{i, i'} \xi'(\boldsymbol{\tau}^1 \cdot \boldsymbol{\tau}^2), \quad \mathbb{E} y(\boldsymbol{\tau}^1) y(\boldsymbol{\tau}^2) = \theta(\boldsymbol{\tau}^1 \cdot \boldsymbol{\tau}^2),$$

let $\boldsymbol{\eta}$ be a M -dimensional standard Gaussian random vector, and let η be a standard Gaussian random variable independent of everything else. Denote by $\langle \cdot \rangle$ the Gibbs average corresponding to G and define

$$\begin{aligned} \Psi_M(\mathcal{L}) &= \frac{1}{M} \mathbb{E} \log \int_{S_M} \mathbb{E}_\eta \left\langle \exp(\boldsymbol{\varepsilon} \cdot (\mathbf{z}(\boldsymbol{\tau}) + \boldsymbol{\eta}(\xi'(1) - \xi'(\|\boldsymbol{\tau}\|^2))^{1/2})) \right\rangle d\lambda_M(\boldsymbol{\varepsilon}) \\ &\quad - \frac{1}{M} \mathbb{E} \log \mathbb{E}_\eta \left\langle \exp(\sqrt{M}(y(\boldsymbol{\tau}) + \eta(\xi'(1) - \xi'(\|\boldsymbol{\tau}\|^2))^{1/2})) \right\rangle. \end{aligned}$$

Lemma 3.4. For each $M \geq 1$, $\mathcal{L} \rightarrow \Psi_M(\mathcal{L})$ is a well-defined continuous function with respect to the weak convergence of the distribution \mathcal{L} .

This is a generalization of Lemma 3 in [12] and can be verified exactly in the same way. Let $\mathcal{L}^{M,N}$ and \mathcal{L}^M be the distributions generating $R^{M,N}$ and R^M , respectively. Then Lemma 3.4 leads to

$$\lim_{N \rightarrow \infty} \Lambda_{M,N}(u^{M,N}) = \lim_{N \rightarrow \infty} \Psi_M(\mathcal{L}^{M,N}) = \Psi_M(\mathcal{L}^M). \tag{3.6}$$

Identifying the asymptotic Gibbs' measures using ultrametricity. Since the asymptotic Gibbs measure G_M satisfies the G.G. identities (3.5), the main result in [11] implies that the support of G_M is ultrametric with probability one, that is,

$$\mathbb{E} \langle I(R_{1,2}^M \geq \min(R_{1,3}^M, R_{2,3}^M)) \rangle_M = 1.$$

This implies that for any q , the inequality $q \leq \tau^\ell \cdot \tau^{\ell'}$ defines an equivalent relation and therefore, the array $(I(q \leq R_{\ell,\ell'}^M))_{\ell,\ell' \geq 1}$ is nonnegative definite since it is block-diagonal with blocks consisting of all elements equal to one. Consider a sequence of functions $(H_k)_{k \geq 1}$ on $[0, 1]$ by defining $H_k(q) = j/k$ if $j/(k+1) \leq q < (j+1)/(k+1)$ for $0 \leq j \leq k$ and $H_k(1) = 1$. Then we see that $R_k^M = (H_k(R_{\ell,\ell'}^M))_{\ell,\ell' \geq 1}$ is symmetric, weakly exchangeable, and satisfies the G.G. identities. In addition, it is non-negative definite since it can be written as a convex combination of nonnegative definite arrays,

$$H_k(R_{\ell,\ell'}^M) = \sum_{j=1}^k \frac{1}{k} I\left(\frac{j}{k+1} \leq R_{\ell,\ell'}^M\right).$$

Thus by the Dovbysh-Sudakov representation, R_k^M can be generated by a sample from a random measure G_M^k on the unit ball of some Hilbert space. Since $\sup_{0 \leq q \leq 1} |H_k(q) - q| \leq 1/k$, $(R_k^M)_{k \geq 1}$ converges weakly to R^M . From Lemma 3.4, we then obtain

$$\lim_{k \rightarrow \infty} \Psi_M(\mathcal{L}_k^M) = \Psi_M(\mathcal{L}^M), \tag{3.7}$$

where \mathcal{L}_k^M is the distribution for R_k^M . On the other hand, from the Baffioni-Rosati theorem [3], an ultrametric measure, such as G_M^k , that satisfies the G.G. identities and under which the overlaps R_k^M take finitely many values $(p/k)_{0 \leq p \leq k}$ can be identified as the Poisson-Dirichlet cascade with parameters

$$\begin{aligned} \mathbf{m}^{M,k} &= (m_p^{M,k})_{0 \leq p \leq k} \text{ with } m_p^{M,k} = \mathbb{E} \langle I((R_k^M)_{1,2} \leq p/k) \rangle_M, \\ \mathbf{q}^{M,k} &= (q_p^{M,k})_{0 \leq p \leq k+1} \text{ with } q_p^{M,k} = p/(k+1). \end{aligned}$$

For details of this result, one may consult the proof of Theorem 15.3.6 in [18]. Let \mathbf{x}_k^M be the distribution function corresponding to this triplet $(k, \mathbf{m}^{M,k}, \mathbf{q}^{M,k})$. Using the same argument as Theorem 14.2.1 in [18], we obtain

$$\Psi_M(\mathcal{L}_k^M) = \mathcal{P}_M(\mathbf{x}_k^M). \tag{3.8}$$

Lemma 3.5. For any two triplets $(k, \mathbf{m}, \mathbf{q})$ and $(k', \mathbf{m}', \mathbf{q}')$ associated with the distribution functions x and x' respectively, we have

$$|\mathcal{P}_M(x) - \mathcal{P}_M(x')| \leq \frac{\xi'(1)}{2} \int_0^1 |x(q) - x'(q)| dq. \tag{3.9}$$

It is well-known that the same Lipschitz property as (3.9) for the Parisi functional in the Ising spin glass model was established by F. Guerra [7]. In the spherical models, despite of the non-product structure on the configuration space, one may find the same approach as Theorem 14.11.2 in [18] can also be applied to obtain (3.9). Let \mathcal{F} be the collection of all distribution functions \mathbf{x} on $[0, 1]$. Define a metric d on \mathcal{F} by letting $d(\mathbf{x}, \mathbf{x}') = \int_0^1 |\mathbf{x}(q) - \mathbf{x}'(q)| dq$ for every $\mathbf{x}, \mathbf{x}' \in \mathcal{F}$. Using the Lipschitz inequality (3.9), we can extend the functional \mathcal{P}_M continuously to \mathcal{F} .

Proof of Theorem 1.1. If we denote by \mathbf{x}^M the distribution of $R_{1,2}^M$, it follows that by using (3.7) and (3.8),

$$\Psi_M(\mathcal{L}^M) = \mathcal{P}_M(\mathbf{x}^M). \quad (3.10)$$

From Helly's selection theorem, $(\mathbf{x}^M)_{M \geq 1}$ contains a weakly convergent subsequence with limit \mathbf{x}^∞ . For clarity, we simply use (\mathbf{x}^M) to denote this subsequence. Using (3.9) and (3.10), if we set $\mathbf{x}_k^\infty = H_k(\mathbf{x}^\infty)$, then

$$\begin{aligned} \limsup_{M \rightarrow \infty} |\Psi_M(\mathcal{L}^M) - \mathcal{P}_M(\mathbf{x}_k^\infty)| &= \limsup_{M \rightarrow \infty} |\mathcal{P}_M(\mathbf{x}^M) - \mathcal{P}_M(\mathbf{x}_k^\infty)| \\ &\leq \frac{\xi'(1)}{2} \left(\limsup_{M \rightarrow \infty} d(\mathbf{x}^M, \mathbf{x}^\infty) + d(\mathbf{x}^\infty, \mathbf{x}_k^\infty) \right) \\ &= \frac{\xi'(1)}{2} d(\mathbf{x}^\infty, \mathbf{x}_k^\infty) \end{aligned}$$

for every $k \geq 1$. To sum up, from (3.4), (3.6), and this inequality, we conclude that

$$\begin{aligned} \liminf_{N \rightarrow \infty} \frac{1}{N} \mathbb{E} \log Z_N &\geq \limsup_{M \rightarrow \infty} \left(\Psi_M(\mathcal{L}^M) + \frac{1}{M} \log \nu_M(A_\delta) \right) - \delta \xi'(1) \\ &\geq \limsup_{M \rightarrow \infty} \mathcal{P}_M(\mathbf{x}_k^\infty) - \limsup_{M \rightarrow \infty} |\Psi_M(\mathcal{L}^M) - \mathcal{P}_M(\mathbf{x}_k^\infty)| - \delta \xi'(1) \\ &\geq \inf_{\mathbf{x}} \mathcal{P}(\mathbf{x}) - \frac{\xi'(1)}{2} d(\mathbf{x}^\infty, \mathbf{x}_k^\infty) - \delta \xi'(1) \end{aligned}$$

for every $k \geq 1$ and $\delta > 0$ and this completes our proof by letting $\delta \rightarrow 0$ and $k \rightarrow \infty$. □

References

- [1] Aizenman, M., Sims, R., Starr, S. (2003) An extended variational principle for the SK spin-glass model. *Phys. Rev. B*, **68**, 214403.
- [2] Arguin, L.-P., Chatterjee, S. (2013) Random overlap structures: properties and applications to spin glasses. *Probab. Theory Related Fields*, **156**, no. 1-2, 375-413. MR-3055263
- [3] Baffioni, F., Rosati, F. (2000) Some exact results on the ultrametric overlap distribution in mean field spin glass models. *Eur. Phys. J. B*, **17**, 439-447.
- [4] Crisanti, A., Sommers, H. J. (1982) The spherical p -spin interaction spin glass model: The statics. *Z. Phys. B. Condensed Matter*, **83** 341-354.
- [5] Dovbysh, L. N., Sudakov, V. N. (1982) Gram-de Finetti matrices. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov.*, **119**, 77-86. MR-0666087
- [6] Ghirlanda, S., Guerra, F. (1998) General properties of overlap probability distributions in disordered spin systems. Towards Parisi ultrametricity. *J. Phys. A*, **31**, no. 46, 9149-9155. MR-1662161
- [7] Guerra, F. (2003) Broken replica symmetry bounds in the mean field spin glass model. *Comm. Math. Phys.*, **233**, no. 1, 1-12. MR-1957729
- [8] Parisi, G. (1979) Infinite number of order parameters for spin-glasses. *Phys. Rev. Lett.*, **43**, 1754-1756. MR-0702601

- [9] Parisi, G. (1980) A sequence of approximate solutions to the S-K model for spin glasses. *J. Phys. A*, **13**, L-115.
- [10] Panchenko, D. (2013) Spin glass models from the point of view of spin distributions. *Ann. of Probab.*, **41**, no. 3A, 1315-1361. MR-3098679
- [11] Panchenko, D. (2013) The Parisi ultrametricity conjecture. *Ann. of Math. (2)*, **177**, no. 1, 383-393. MR-2999044
- [12] Panchenko, D. (2011) The Parisi formula for mixed p -spin models. To appear in the *Ann. of Probab.*
- [13] Sherrington, D., Kirkpatrick, S. (1972) Solvable model of a spin glass. *Phys. Rev. Lett.*, **35**, 1792-1796.
- [14] Stroock, D. (1993) Probability theory. An analytic view, first edition. Cambridge University Press, Cambridge. MR-1267569
- [15] Talagrand, M. (2003) On Guerra's broken replica-symmetry bound. *C. R. Math. Acad. Sci. Paris*, **337**, no. 7, 477-480. MR-2023757
- [16] Talagrand, M. (2006) Free energy of the spherical mean field model. *Probab. Theory Related Fields*, **134**, no. 3, 339-382. MR-2226885
- [17] Talagrand, M. (2006) The Parisi formula. *Ann. of Math. (2)*, **163**, no. 1, 221-263. MR-2195134
- [18] Talagrand, M. (2011) Mean-Field Models for Spin Glasses. *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge A Series of Modern Surveys in Mathematics*, Vol. 54, 55. Springer-Verlag.

Acknowledgments. The author would like to thank Dmitry Panchenko for motivating this work and his advisor, Michael Cranston, for many helpful discussions. Special thanks are due to Alexander Vandenberg-Rodes for several suggestions regarding the presentation of the paper.