

# UC San Diego

## UC San Diego Previously Published Works

### Title

Spinal 12-lipoxygenase-derived hepoxilin A3 contributes to inflammatory hyperalgesia via activation of TRPV1 and TRPA1 receptors

### Permalink

<https://escholarship.org/uc/item/62w3b0x3>

### Journal

Proceedings of the National Academy of Sciences of the United States of America, 109(17)

### ISSN

0027-8424

### Authors

Gregus, Ann M  
Doolen, Suzanne  
Dumlao, Darren S  
et al.

### Publication Date

2012-04-24

### DOI

10.1073/pnas.1110460109

Peer reviewed

# Spinal 12-lipoxygenase-derived hepoxilin A<sub>3</sub> contributes to inflammatory hyperalgesia via activation of TRPV1 and TRPA1 receptors

Ann M. Gregus<sup>a,1</sup>, Suzanne Doolen<sup>b,1</sup>, Darren S. Dumlao<sup>c,1</sup>, Matthew W. Buczynski<sup>c</sup>, Toshifumi Takasusuki<sup>a</sup>, Bethany L. Fitzsimmons<sup>a</sup>, Xiao-Ying Hua<sup>a</sup>, Bradley K. Taylor<sup>b</sup>, Edward A. Dennis<sup>c,2</sup>, and Tony L. Yaksh<sup>a,2</sup>

<sup>a</sup>Department of Anesthesiology and <sup>c</sup>Department of Pharmacology, Chemistry, and Biochemistry, University of California at San Diego, La Jolla, CA 92093-0818; and <sup>b</sup>Department of Physiology, University of Kentucky, Lexington, KY 40536

Edited by David Julius, University of California, San Francisco, CA, and approved March 15, 2012 (received for review June 28, 2011)

Peripheral inflammation initiates changes in spinal nociceptive processing leading to hyperalgesia. Previously, we demonstrated that among 102 lipid species detected by LC-MS/MS analysis in rat spinal cord, the most notable increases that occur after intraplantar carrageenan are metabolites of 12-lipoxygenases (12-LOX), particularly hepoxilins (HXA<sub>3</sub> and HXB<sub>3</sub>). Thus, we examined involvement of spinal LOX enzymes in inflammatory hyperalgesia. In the current work, we found that intrathecal (IT) delivery of the LOX inhibitor nordihydroguaiaretic acid prevented the carrageenan-evoked increase in spinal HXB<sub>3</sub> at doses that attenuated the associated hyperalgesia. Furthermore, IT delivery of inhibitors targeting 12-LOX (CDC, Baicalein), but not 5-LOX (Zileuton) dose-dependently attenuated tactile allodynia. Similarly, IT delivery of 12-LOX metabolites of arachidonic acid 12(S)-HpETE, 12(S)-HETE, HXA<sub>3</sub>, or HXB<sub>3</sub> evoked profound, persistent tactile allodynia, but 12(S)-HpETE and HXA<sub>3</sub> produced relatively modest, transient heat hyperalgesia. The pronociceptive effect of HXA<sub>3</sub> correlated with enhanced release of Substance P from primary sensory afferents. Importantly, HXA<sub>3</sub> triggered sustained mobilization of calcium in cells stably overexpressing TRPV1 or TRPA1 receptors and in acutely dissociated rodent sensory neurons. Constitutive deletion or antagonists of TRPV1 (AMG9810) or TRPA1 (HC030031) attenuated this action. Furthermore, pretreatment with antihyperalgesic doses of AMG9810 or HC030031 reduced spinal HXA<sub>3</sub>-evoked allodynia. These data indicate that spinal HXA<sub>3</sub> is increased by peripheral inflammation and promotes initiation of facilitated nociceptive processing through direct activation of TRPV1 and TRPA1 at central terminals.

eicosanoid | pain | central sensitization

Tissue injury and inflammation are associated with hyperalgesia mediated by facilitated spinal nociceptive processing that can be modulated by lipids derived from arachidonic acid (AA) and other polyunsaturated fatty acids (PUFA), including eicosanoids synthesized via three enzymatic pathways: (i) cyclooxygenase (COX)–prostaglandins (PG); (ii) 5-, 12-, and 12/15-lipoxygenases (LOX)–leukotrienes, hydroxyeicosatetraenoic acids (HETEs), hepoxilins (HXA<sub>3</sub> and HXB<sub>3</sub>), lipoxins, resolvins, and protectins; and (iii) cytochrome P450–epoxyeicosatrienoic acids and HETEs (1). Substantial evidence indicates that peripheral injury or direct activation of spinal dorsal horn receptors [Neurokinin 1 (NK1), AMPA, and NMDA] increases eicosanoid formation and that spinal delivery of COX inhibitors reduces the associated hyperalgesia (2, 3). Recently, we reported that paw carrageenan increases spinal production of both COX and 12-LOX metabolites of AA, including 12(S)-HETE in cerebrospinal fluid (CSF) and hepoxilins in the lumbar spinal cord (4).

Several groups point to a peripheral role for 5- and 12-LOX in nociception, as shown by antihyperalgesic actions of LOX inhibitors administered via systemic routes (5–9). It has been suggested that spinal 12-LOX may play a role in morphine physical dependence (10), yet there is little work defining its involvement in spinal nociception. The chemosensors Transient Receptor Potential Vanilloid 1

(TRPV1) and Ankyrin 1 (TRPA1) have been implicated in peripheral inflammation-evoked hyperpathic states (11). Irritants, such as formalin and oxidized lipids, can cause hyperalgesia via activation of spinal TRPA1 (12–14). For example, 4-hydroxy-2-nonenal (HNE) elicits spinal release of Substance P (SP) and triggers nociceptive behaviors via activation of TRPA1 receptors (15). Both 12(S)-HpETE and its reduction product 12(S)-HETE directly stimulate TRPV1 in cultured dorsal root ganglion (DRG) neurons (16). Noxious heat or depolarization releases 12-LOX metabolites of linoleic acid that activate spinal TRPV1 and contribute to complete Freund's adjuvant (CFA)-induced mechanical allodynia (17, 18). Given that linoleic acid is used in vivo for biosynthesis of AA and that HNE is generated by peroxidation of 12/15-LOX products of *n*-6 PUFA, we hypothesized that spinal 12-LOX metabolites of AA, particularly hepoxilins, also contribute to inflammatory hyperalgesia through stimulation of nociceptive afferents. Although hepoxilins increase intracellular Ca<sup>2+</sup> in neutrophils (19, 20) and neurons (21) and enhance vascular permeability in rat skin (22, 23), its receptors remain undefined. Our findings demonstrate that peripheral inflammation increases spinal HXA<sub>3</sub>, which produces hyperesthesia via activation of TRPV1 and TRPA1 and spinal SP release.

## Results

### Inhibition of Spinal 12-LOX Attenuates Inflammatory Hyperalgesia.

First, we examined involvement of spinal LOX in the development of hyperalgesia following intraplantar (IPLT) carrageenan. Intrathecal (IT) pretreatment with the LOX inhibitor nordihydroguaiaretic acid (NDGA) dose-dependently attenuated tactile allodynia as depicted by timecourse (Fig. 1A) and area under the curve (AUC), or percent hyperalgesic index (Fig. 1C), without overt toxicity. An antihyperalgesic dose of IT NDGA at the upper limit of solubility (60 μg) also prevented the development of thermal hyperalgesia (Fig. S1A and B) and partially reversed established tactile allodynia (Fig. S1C). IT NDGA did not alter paw volume at any of the doses tested (Fig. S1D), indicating that spinal lipoxygenases contribute centrally to inflammatory hyperesthesia.

Recently, we demonstrated that IPLT carrageenan increases spinal levels of AA metabolites of 12-LOX but not of 5-LOX (4). Therefore, we reasoned that spinal 12-LOX likely contributes to inflammatory hyperesthesia. To address this question, we examined the effect of IT pretreatment with 5- or 12-LOX inhibitors on

Author contributions: A.M.G., S.D., D.S.D., M.W.B., X.-Y.H., B.K.T., E.A.D., and T.L.Y. designed research; A.M.G., S.D., D.S.D., T.T., and B.L.F. performed research; A.M.G., S.D., D.S.D., and T.T. analyzed data; and A.M.G., S.D., D.S.D., M.W.B., X.-Y.H., B.K.T., E.A.D., and T.L.Y. wrote the paper.

The authors declare no conflict of interest.


This article is a PNAS Direct Submission.

<sup>1</sup>A.M.G., S.D., and D.S.D. contributed equally to this work.

<sup>2</sup>To whom correspondence may be addressed. E-mail: edennis@ucsd.edu or tyaksh@ucsd.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1110460109/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1110460109/-DCSupplemental).





**Fig. 2.** Bioactive 12-LOX metabolites produce robust tactile allodynia after spinal delivery. (A–D) Dose–response depicting timecourses (Left) and AUC values (Right) reveal sustained reduction of tactile thresholds after a single IT administration of nanogram to microgram amounts of (A and B) HXA<sub>3</sub> or (C and D) 12(S)-HpETE. \**P* < 0.05, \*\**P* < 0.01, \*\*\**P* < 0.001 vs. saline vehicle; *n* = 5–6.

(Fig. S5 A and B). This response was attenuated by treatment with 10 μM HC030031 (83%) (Fig. S5C). Application of the TRPV1-selective antagonist AMG9810 at 1 μM, a concentration that exceeds its IC<sub>50</sub> of ~0.1 μM (28), reduced HXA<sub>3</sub>-evoked Ca<sup>2+</sup>-responses by 61% and 12(S)-HpETE-evoked Ca<sup>2+</sup>-responses by 55% (Fig. 3C and Fig. S5D). HXA<sub>3</sub> (1 μM) also elicited intracellular Ca<sup>2+</sup> accumulation in cells heterologously overexpressing TRPA1 (HEK-TRPA1-tet) or TRPV1 (CHO-TRPV1) compared with their respective controls (Fig. 3D). Similarly, superfusion of HXA<sub>3</sub> (1 μM) on acutely dissociated adult mouse DRG cells increased free-Ca<sup>2+</sup> levels, a response that was dampened in both TRPA1 and TRPV1 knockout mice (Fig. 3E and F). It is important to note that although the majority of cells recorded from either TRPA1 or TRPV1 KO mice were not considered HXA<sub>3</sub>-sensitive according to our pre-established criteria (see *SI Materials and Methods*), some cells were very weakly responsive. Of this population, the percent sensitivity to capsaicin was as follows: WT, 36%; TRPA1 KO, 20%; TRPV1 KO, 0%. Thus, HXA<sub>3</sub>-mediated calcium mobilization was drastically reduced but not completely abolished in the absence of either TRPA1 or TRPV1. These data indicate that HXA<sub>3</sub>, as well as its parent compound 12(S)-HpETE, produce potent activation of both TRPA1 and TRPV1 receptors on sensory neurons.

We then examined whether the hyperalgesic activity of HXA<sub>3</sub> is related to its activation of primary sensory afferents and subsequent release of pronociceptive neurotransmitters. We investigated in rat spinal cord *in vivo* if IT HXA<sub>3</sub> increases SP release from peptidergic primary afferents by measuring NK1 receptor internalization in L4, L5, and L6 levels of lumbar spinal dorsal horn. We found no difference in the percentage of internalized NK1 receptors at 10 min after IT HXA<sub>3</sub> (1 μg), before the onset of allodynia (Fig. S6A). In contrast, NK1 receptor internalization is significantly increased in lamina I of L4, L5, and L6 at 30 min after spinal administration of HXA<sub>3</sub> (Fig. S6B), concurrent with the onset of tactile allodynia. Thus, IT HXA<sub>3</sub> evokes spinal release of SP from nociceptive primary sensory afferents in direct temporal correlation with hyperesthesia.

**HXA<sub>3</sub>-Induced Activation of TRPV1 and TRPA1 Contributes to Tactile Allodynia.** Next, we sought to determine if HXA<sub>3</sub>-evoked tactile allodynia is mediated via stimulation of TRPV1 or TRPA1 receptors on nociceptive primary sensory afferents. IT pretreatment with the maximum soluble dose (10 μg) for antagonists


of TRPA1 (HC030031) or of TRPV1 (AMG9810) significantly attenuated IPLT carrageenan-evoked tactile allodynia and thermal hyperalgesia (Fig. S7A and B). These observations corroborate other studies that use the same doses of these antagonists in different rodent models of inflammatory pain (14, 29, 30). Similarly, IT HXA<sub>3</sub>-induced tactile allodynia is significantly attenuated by spinal pretreatment with antihyperalgesic doses of HC030031 or AMG9810 (Fig. 4). Taken together, the present study provides strong evidence that HXA<sub>3</sub> produces hyperesthesia at the spinal level by activating TRPA1 and TRPV1 receptors in a subpopulation of primary sensory afferents and by facilitating release of nociceptive mediators in dorsal horn (Fig. 5).

## Discussion

Rat lipoxygenases are classified as 5-, 12-, and 12/15-LOX, named according to the stereospecific insertion of O<sub>2</sub> into PUFA (1). To date, most reports that examine the role of LOX in hyperalgesia focus on 5-LOX, which catalyzes the rate-limiting step in the synthesis of leukotrienes (LT), potent regulators of innate immune cell functions and chemoattractants for neutrophils (31). Systemic treatment with 5-LOX inhibitors or LT receptor antagonists attenuated inflammatory hyperalgesia (5, 8, 32). However, spinal 5-LOX inhibition did not prevent carrageenan-induced allodynia, which supports our previous observation that spinal 5-LOX metabolites are either not present or are at levels below the assay limit of detection (4). Nonetheless, our data do not preclude the possibility of inflammation-induced activation of 5-LOX in brain, DRG, or peripherally at the site of injury. For example, intradermal administration of LTB<sub>4</sub> produces hyperalgesia that is attenuated by depletion of neutrophils (33). In addition, systemic administration of inhibitors targeting 5-LOX or LT receptors reduces mechanical allodynia and, respectively, prevents increased levels of LTB<sub>4</sub> (8) and neutrophil infiltration (32) in paw tissue during inflammation. Thus, it appears that induction of 5-LOX activity occurs largely outside of the lumbar spinal cord.

In contrast, we found that spinal inhibition of 12-LOX significantly attenuated hyperesthesia. Conversely, spinal delivery of either HXA<sub>3</sub> or HXB<sub>3</sub>, the parent molecule 12(S)-HpETE, or its reduction product 12(S)-HETE, produced persistent tactile allodynia. Our findings regarding thermal responses to 12(S)-HpETE are consistent with Trang et al., who found that IT delivery at doses up to 1 μg has no effect on thermal (heat) nociceptive thresholds in rats (34). Similar responses were observed following IT administration of COX metabolites PGE<sub>2</sub> or PGF<sub>2α</sub>, which elicit robust mechanical but weak thermal hyperalgesia (35). Others have illustrated that activation of 12-LOX in the paw produces mechanical allodynia (36) as well as thermal hyperalgesia (6), which is attenuated by local administration of LOX inhibitors (7). Furthermore, several groups have shown that during inflammation, spinal TRPV1 receptors contribute significantly to both tactile allodynia and thermal hyperalgesia, but TRPV1 on peripheral terminals preferentially mediates the latter (7, 37–39). Peripheral activation of TRPA1 produces some thermal hyperalgesia, but the resulting tactile allodynia is the more robust response (40). Thus, the current work demonstrates that spinal HXA<sub>3</sub> contributes to the development of inflammatory hyperalgesia, preferentially initiating tactile allodynia via activation of TRPV1 and TRPA1 on central terminals. Although mechanisms underlying thermal and mechanical hypersensitivity remain to be elucidated, differential expression of these phenomena have been described in other animal models of pain syndromes (41, 42).

An important question relates to the mechanisms by which hepxilins cause inflammatory hyperalgesia. Both HXA<sub>3</sub> and its precursor 12(S)-HpETE stimulated TRPV1 and TRPA1 on nociceptive sensory neurons. However, in an aqueous environment, lipid hydrogen peroxides (such as 12-HpETE) have a short T<sub>1/2</sub> on the order of minutes (43); in contrast, HXA<sub>3</sub> and structurally similar compounds created by 15-LOX have a half-life on the order of hours (20, 44). Moreover, the *in vivo* degradation of each of these metabolites is handled by two distinct enzyme classes, glutathione peroxidase for 12-HpETE and soluble




**Fig. 3.** HXA<sub>3</sub> produces direct activation of TRPV1 and TRPA1 in sensory neurons. (A) Representative micrographs of cultured rat sensory neurons before (Upper) and after (Lower) application of 1 μM HXA<sub>3</sub>. Arrowheads, Fura-2-loaded cells exhibiting Ca<sup>2+</sup> mobilization in response to HXA<sub>3</sub>. (Scale bars, 50 μm.) (B) Representative traces of HXA<sub>3</sub>-evoked calcium responses in rat sensory neurons and attenuation by the TRPA1 antagonist HC030031 (Upper) or the TRPV1 antagonist AMG9810 (Lower). Peak Ca<sup>2+</sup> responses represent mean change in fluorescence ratio (Δ340/380) ± SEM. (C) HXA<sub>3</sub>-evoked Ca<sup>2+</sup>-responses were reduced by HC030031 (10 μM) (VEH, 0.117 ± 0.01 vs. HC030031, 0.001 ± 0.02; \*\*\*P < 0.001, n = 16 cells, 4 rats) and AMG 9810 (1 μM) (VEH, 0.159 ± 0.03 vs. AMG9810 0.061 ± 0.03; \*P < 0.05, n = 5 cells, 3 rats). (D) HXA<sub>3</sub> (1 μM) also increased calcium mobilization in HEK-TRPA1-tet [TRPA1(+)] Induced, 0.077 ± 0.008 vs. TRPA1(-) Uninduced control, 0.020 ± 0.004; \*\*\*P < 0.001, n = 4 cells] and in CHO-TRPV1 cells [TRPV1(+)] 0.070 ± 0.001 vs. TRPV1(-) CHO control, 0.012 ± 0.001; \*P < 0.05, n = 3–5 cells]. (E) Representative traces of HXA<sub>3</sub>-evoked calcium responses in mouse sensory neurons from WT (Top) and attenuation in TRPA1 (Middle), or TRPV1 knockout (KO) (Bottom). (F) Percent of WT, TRPA1, or TRPV1 KO neurons in which 1 μM HXA<sub>3</sub> increased free intracellular Ca<sup>2+</sup> levels. (WT, 15.2 ± 4.9% n = 42 cells; TRPA1 KO, 0.0 ± 0.0%, n = 34 cells; TRPV1 KO, 1.1 ± 1.1%; n = 51 cells; \*P < 0.05, 3 mice per group). Cell viability was confirmed using 50 mM K<sup>+</sup>; functional TRPV1 or TRPA1 receptors were verified with 500 nM capsaicin or 5 μM icilin, respectively. VEH, artificial CSF.

epoxide hydrolase for HXA<sub>3</sub> (1). Thus, although 12-HpETE and HXA<sub>3</sub> may interact with these channels via a similar mechanism, their bioavailability in vivo may differ.


Collectively, our findings are intriguing given the differential effect of spinally delivered 12-LOX products on thermal and tactile sensitivity. Several groups have demonstrated that TRPA1 modulates nociceptive tactile thresholds, triggers release of the neuropeptides from primary sensory afferents, and mediates the mechanical allodynia produced by inflammatory agents and chemical irritants (13, 15, 25). Similarly, we observed a time-dependent increase in spinal SP release following IT administration of HXA<sub>3</sub> at a dose that produces hyperalgesia. Although others have demonstrated that systemic or IT delivery of the TRPA1 antagonist HC030031 attenuates mechanical allodynia induced by formalin, mustard oil, or CFA (14, 25, 26), the spinal actions of this agent have not yet been examined following carrageenan or after IT delivery of 12-LOX metabolites. In this study, we found that TRPA1 receptors are required for the observed effects of HXA<sub>3</sub> on calcium mobilization in sensory neurons and tactile allodynia.

TRPA1 receptors are expressed by a subpopulation of peptidergic somatosensory neurons that often coexpress TRPV1 (45). It has been suggested that 12-LOX-mediated activation of TRPV1 in hippocampal neurons mediates long-term depression (46) and in primary afferent nociceptors contributes to inflammatory pain hypersensitivity (6). Until recently, TRPV1 receptors were viewed primarily as sensors of noxious heat, the

activation of which underlies thermal hyperalgesia (47). However, data are now emerging in support of TRPV1-mediated involvement in tactile allodynia during persistent inflammation by CFA. For example, it has been reported that 9-HODE and 13-HODE contribute to mechanical hypersensitivity via stimulation of TRPV1 (17, 18). Similarly, we found that TRPV1 receptors also are required for the observed effects of HXA<sub>3</sub> on calcium mobilization in sensory neurons and tactile allodynia.



**Fig. 4.** HXA<sub>3</sub>-evoked tactile allodynia is mediated by spinal TRPV1 and TRPA1. (A) Timecourse and (B) corresponding AUC reveals significant attenuation of tactile allodynia following IT HXA<sub>3</sub> (1 μg) by IT pretreatment with antihyperalgesic doses (10 μg) of antagonists of TRPA1 (HC030031) or TRPV1 (AMG9810). \*P < 0.05, \*\*P < 0.01 vs. VEH 3; n = 5–7.



**Fig. 5.** Model of HXA<sub>3</sub>-mediated hyperalgesic effects at the spinal level. HXA<sub>3</sub> is produced through 12-LOX either from arachidonic acid or via 12-HpETE. Cellular sources of HXA<sub>3</sub> may represent DRG neurons or satellite cells, spinal neurons, or glia, and circulating leukocytes or platelets. Spinally generated HXA<sub>3</sub> activates TRPV1 and TRPA1, resulting in calcium mobilization and release of SP from nociceptive afferents, internalization of NK1 receptors in dorsal horn, and ultimately tactile allodynia.

We demonstrate here that low micromolar concentrations of HXA<sub>3</sub> produce potent activation of TRPV1 and TRPA1 receptors on sensory neurons and increased spinal SP release concurrent with the development of tactile allodynia. Our observations of HXA<sub>3</sub>-mediated activation of TRPV1 or TRPA1 in heterologous overexpression systems indicate a direct interaction with these channels. However, HXA<sub>3</sub> may also exert effects via direct binding and gating of TRPV1/TRPA1 multimeric complexes, binding to TRPV1 and transactivation of TRPA1, or binding to a G protein-coupled receptor and signaling via second messenger systems to increase membrane trafficking of TRPV1/TRPA1. Recent studies demonstrate functional interaction of these receptors in sensory neurons (48, 49). Similar to HXA<sub>3</sub>, bradykinin, nitric oxide, and BH4 all require both TRPV1 and TRPA1 receptor function for hyperalgesia (12, 50). Phospholipase C-dependent sensitization and increased plasma membrane trafficking of TRPA1 has been described for bradykinin and mustard oil, respectively (51, 52). Accordingly, it has been shown in neutrophils that HXA<sub>3</sub> produces a biphasic increase in levels of intracellular calcium through G protein-coupled receptor-mediated signaling via phospholipase C (20, 53).

Ultimately, although the local concentration of individual 12-LOX products may be subthreshold for activation of TRP channels, it is possible that several metabolites could function in concert to produce hyperalgesia, as suggested previously (17). The identities, temporal profiles, concentrations, and targets of these mediators also may vary with different pain states. In summary, this work is unique in providing evidence of hepxilin receptors in the CNS and that HXA<sub>3</sub>-mediated activation of TRPA1 and TRPV1 contributes to hyperesthesia at the spinal level. These findings highlight the importance of developing novel anti-inflammatory therapeutics targeting this pathway. Furthermore, these findings advance our understanding of distinct mechanisms contributing to tactile allodynia versus heat hyperalgesia.

## Materials and Methods

**Animals.** Male Sprague-Dawley rats (300–350 g; Harlan) or mice (25–30 g; C57BL/6 and TRPV1 knockout, Jackson Laboratories, or TRPA1 knockout, University of Kentucky) were used in accordance with protocols approved by the Institutional Animal Care and Use Committees of the University of

California at San Diego and the University of Kentucky. All testing was performed by a blinded observer. IT catheter implantation of rats and drug delivery was performed as described previously (54).

**Drugs.** NDGA, Baicalein, 12(S)-HpETE, and 12(S)-HETE (Cayman); CDC, hepxilins HXA<sub>3</sub> and HXB<sub>3</sub> (Enzo); Zileuton, HC-030031, and AMG9810 (Tocris) were prepared in vehicles (VEH): NDGA in 20% (wt/vol)  $\beta$ -cyclodextrin/saline; Baicalein, CDC, and Zileuton in 3% (vol/vol) DMSO/3% Cremaphor-EL/saline; HC030031 and AMG9810 in 3% (vol/vol) DMSO/3% Tween-80/saline.

**Behavioral Testing.** Peripheral inflammation and hyperalgesia were assessed as previously described (4) after IT delivery of drugs 30 min before or 90 min after 2% (wt/vol) carrageenan injected IPLT into the left hind paw. The 12-LOX metabolites were prepared fresh before use (16) and IT injections were performed immediately after dilution followed by measurement of nociceptive behaviors. Vehicle or antagonists of TRPA1 or TRPV1 were given IT 30 min before HXA<sub>3</sub>. Data were expressed as response latency vs. time and as AUC, or hyperalgesic index (percent change from baseline  $\times$  minute).

**Immunohistochemistry.** At 10 or 30 min after IT delivery of saline or HXA<sub>3</sub> (1  $\mu$ g), rats were perfused and the lumbar spinal cord was processed for NK1 receptor internalization (55, 56).

**Heterologous Overexpression Systems, DRG Cell Preparation, and Ca<sup>2+</sup> Imaging.** Inducible HEK293 cells stably overexpressing TRPA1 (HEK-TRPA1-tet) and CHO cells stably overexpressing TRPV1 (CHO-TRPV1) were maintained as previously described (15, 17). L4 and L5 DRGs were dissected from male Sprague-Dawley rats at 4–5 wk old or mice (WT, TRPV1, or TRPA1 knockout) at 8 wk old, dissociated and used within 24 h of culture, as previously described (57), with modifications by S.D. and B.K.T. (*SI Materials and Methods*).

**LC-MS/MS.** LC-MS/MS of L4/L5 rat spinal cord was conducted using a tandem quadrupole mass spectrometer (ABI 4000 Q-Trap) as previously described (4, 58, 59).

**Statistics.** Normalized raw data of spinal eicosanoids were filtered using the Grubbs' test (4). Data were expressed as the mean  $\pm$  SEM, with  $P < 0.05$  as significant.  $P$  values were determined using SPSS software (PASW version 18, SPSS) as follows:  $t$  test for two-group analysis, standard or repeated-measures ANOVA for multiple group analysis with Bonferroni or Dunnett's post hoc for behavioral or biochemical data, respectively.

**ACKNOWLEDGMENTS.** We thank Drs. David Julius and Ardem Patapoutian for the generous gifts of HEK-TRPA1-tet cells and CHO-TRPV1 cells; Alex Chesler and Adrienne Dubin for excellent technical expertise; Gregory Frolenkov for

1. Buczynski MW, Dumlao DS, Dennis EA (2009) Thematic Review Series: Proteomics. An integrated omics analysis of eicosanoid biology. *J Lipid Res* 50:1015–1038.
2. Dirig DM, Isakson PC, Yaksh TL (1998) Effect of COX-1 and COX-2 inhibition on induction and maintenance of carrageenan-evoked thermal hyperalgesia in rats. *J Pharmacol Exp Ther* 285:1031–1038.
3. Malmberg AB, Yaksh TL (1992) Hyperalgesia mediated by spinal glutamate or substance P receptor blocked by spinal cyclooxygenase inhibition. *Science* 257:1276–1279.
4. Buczynski MW, et al. (2010) Inflammatory hyperalgesia induces essential bioactive lipid production in the spinal cord. *J Neurochem* 114:981–993.
5. Jain NK, Kulkarni SK, Singh A (2001) Role of cysteinyl leukotrienes in nociceptive and inflammatory conditions in experimental animals. *Eur J Pharmacol* 423:85–92.
6. Shin J, et al. (2002) Bradykinin-12-lipoxygenase-VR1 signaling pathway for inflammatory hyperalgesia. *Proc Natl Acad Sci USA* 99:10150–10155.
7. Yoo S, et al. (2009) Lipoxygenase inhibitors suppressed carrageenan-induced Fos-expression and inflammatory pain responses in the rat. *Mol Cells* 27:417–422.
8. Cortes-Burgos LA, et al. (2009) CJ-13610, an orally active inhibitor of 5-lipoxygenase is efficacious in preclinical models of pain. *Eur J Pharmacol* 617:59–67.
9. Szabó A, et al. (2005) Role of transient receptor potential vanilloid 1 receptors in adjuvant-induced chronic arthritis: In vivo study using gene-deficient mice. *J Pharmacol Exp Ther* 314:111–119.
10. Trang T, Sutak M, Quirion R, Jhamandas K (2003) Spinal administration of lipoxygenase inhibitors suppresses behavioural and neurochemical manifestations of naloxone-precipitated opioid withdrawal. *Br J Pharmacol* 140:295–304.
11. Patapoutian A, Tate S, Woolf CJ (2009) Transient receptor potential channels: Targeting pain at the source. *Nat Rev Drug Discov* 8:55–68.
12. Bautista DM, et al. (2006) TRPA1 mediates the inflammatory actions of environmental irritants and proalgesic agents. *Cell* 124:1269–1282.
13. Kwan KY, et al. (2006) TRPA1 contributes to cold, mechanical, and chemical nociception but is not essential for hair-cell transduction. *Neuron* 50:277–289.
14. da Costa DS, et al. (2010) The involvement of the transient receptor potential A1 (TRPA1) in the maintenance of mechanical and cold hyperalgesia in persistent inflammation. *Pain* 148:431–437.
15. Trevisani M, et al. (2007) 4-Hydroxynonenal, an endogenous aldehyde, causes pain and neurogenic inflammation through activation of the irritant receptor TRPA1. *Proc Natl Acad Sci USA* 104:13519–13524.
16. Hwang SW, et al. (2000) Direct activation of capsaicin receptors by products of lipoxygenases: Endogenous capsaicin-like substances. *Proc Natl Acad Sci USA* 97:6155–6160.
17. Patwardhan AM, Scotland PE, Akopian AN, Hargreaves KM (2009) Activation of TRPV1 in the spinal cord by oxidized linoleic acid metabolites contributes to inflammatory hyperalgesia. *Proc Natl Acad Sci USA* 106:18820–18824.
18. Patwardhan AM, et al. (2010) Heat generates oxidized linoleic acid metabolites that activate TRPV1 and produce pain in rodents. *J Clin Invest* 120:1617–1626.
19. Dho S, Gristein S, Corey EJ, Su WG, Pace-Asciak CR (1990) Hexoxilin A3 induces changes in cytosolic calcium, intracellular pH and membrane potential in human neutrophils. *Biochem J* 266:63–68.
20. Mrsny RJ, et al. (2004) Identification of hexoxilin A3 in inflammatory events: A required role in neutrophil migration across intestinal epithelia. *Proc Natl Acad Sci USA* 101:7421–7426.
21. Amer RK, Pace-Asciak CR, Mills LR (2003) A lipoxygenase product, hexoxilin A(3), enhances nerve growth factor-dependent neurite regeneration post-axotomy in rat superior cervical ganglion neurons in vitro. *Neuroscience* 116:935–946.
22. Wang MM, Reynaud D, Pace-Asciak CR (1999) In vivo stimulation of 12(S)-lipoxygenase in the rat skin by bradykinin and platelet activating factor: Formation of 12(S)-HETE and hexoxilins, and actions on vascular permeability. *Biochim Biophys Acta* 1436:354–362.
23. Wang MM, Demin PM, Pace-Asciak CR (1999) Stereoselective actions of hexoxilins A3 and B3 and their cyclopropane analogs (HxdeltaA3 and HxdeltaB3) on bradykinin and PAF-evoked potentiation of vascular leakage in rat skin. *Gen Pharmacol* 33:377–382.
24. Sutherland M, Schewe T, Nigam S (2000) Biological actions of the free acid of hexoxilin A3 on human neutrophils. *Biochem Pharmacol* 59:435–440.
25. McNamara CR, et al. (2007) TRPA1 mediates formalin-induced pain. *Proc Natl Acad Sci USA* 104:13525–13530.
26. Eid SR, et al. (2008) HC-030031, a TRPA1 selective antagonist, attenuates inflammatory- and neuropathy-induced mechanical hypersensitivity. *Mol Pain* 4:48.
27. Vilceanu D, Stucky CL (2010) TRPA1 mediates mechanical currents in the plasma membrane of mouse sensory neurons. *PLoS ONE* 5:e12177.
28. Gava NR, et al. (2005) AMG 9810 [(E)-3-(4-t-butylphenyl)-N-(2,3-dihydrobenzo[b][1,4]dioxin-6-yl)acrylamide], a novel vanilloid receptor 1 (TRPV1) antagonist with anti-hyperalgesic properties. *J Pharmacol Exp Ther* 313:474–484.
29. Yu L, et al. (2008) The role of TRPV1 in different subtypes of dorsal root ganglion neurons in rat chronic inflammatory nociception induced by complete Freund's adjuvant. *Mol Pain* 4:61.
30. Wei H, et al. (2010) Roles of cutaneous versus spinal TRPA1 channels in mechanical hypersensitivity in the diabetic or mustard oil-treated non-diabetic rat. *Neuropharmacology* 58:578–584.
31. Murphy RC, Gijón MA (2007) Biosynthesis and metabolism of leukotrienes. *Biochem J* 405:379–395.
32. Singh VP, Patil CS, Kulkarni SK (2005) Differential effect of zileuton, a 5-lipoxygenase inhibitor, against nociceptive paradigms in mice and rats. *Pharmacol Biochem Behav* 81:433–439.
33. Levine JD, Lau W, Kwiat G, Goetzl EJ (1984) Leukotriene B4 produces hyperalgesia that is dependent on polymorphonuclear leukocytes. *Science* 225:743–745.
34. Trang T, McNaull B, Quirion R, Jhamandas K (2004) Involvement of spinal lipoxygenase metabolites in hyperalgesia and opioid tolerance. *Eur J Pharmacol* 491:21–30.
35. Turnbach ME, Spraggins DS, Randich A (2002) Spinal administration of prostaglandin E(2) or prostaglandin F(2alpha) primarily produces mechanical hyperalgesia that is mediated by nociceptive specific spinal dorsal horn neurons. *Pain* 97:33–45.
36. Aley O, Levine JD (2003) Contribution of 5- and 12-lipoxygenase products to mechanical hyperalgesia induced by prostaglandin E(2) and epinephrine in the rat. *Exp Brain Res* 148:482–487.
37. Bölskei K, et al. (2005) Investigation of the role of TRPV1 receptors in acute and chronic nociceptive processes using gene-deficient mice. *Pain* 117:368–376.
38. McGaraughty S, et al. (2008) Contributions of central and peripheral TRPV1 receptors to mechanically evoked and spontaneous firing of spinal neurons in inflamed rats. *J Neurophysiol* 100:3158–3166.
39. Kanai Y, Hara T, Imai A, Sakakibara A (2007) Differential involvement of TRPV1 receptors at the central and peripheral nerves in CFA-induced mechanical and thermal hyperalgesia. *J Pharm Pharmacol* 59:733–738.
40. Tsgareli MG, et al. (2010) Behavioral evidence of thermal hyperalgesia and mechanical allodynia induced by intradermal cinnamaldehyde in rats. *Neurosci Lett* 473:233–236.
41. Christianson CA, et al. (2010) Characterization of the acute and persistent pain state present in K/BxN serum transfer arthritis. *Pain* 151:394–403.
42. Scherrer G, et al. (2010) VGLUT2 expression in primary afferent neurons is essential for normal acute pain and injury-induced heat hypersensitivity. *Proc Natl Acad Sci USA* 107:22296–22301.
43. Pace-Asciak CR, Asotra S (1989) Biosynthesis, catabolism, and biological properties of HPETEs, hydroperoxide derivatives of arachidonic acid. *Free Radic Biol Med* 7:409–433.
44. Hamburger M, Herman CA, Herman RP (1986) Novel biological transformations of 15-Ls-hydroperoxy-5,8,11,13-eicosatetraenoic acid. *Biochim Biophys Acta* 877:447–457.
45. Bautista DM, et al. (2005) Pungent products from garlic activate the sensory ion channel TRPA1. *Proc Natl Acad Sci USA* 102:12248–12252.
46. Gibson HE, Edwards JG, Page RS, Van Hook MJ, Kauer JA (2008) TRPV1 channels mediate long-term depression at synapses on hippocampal interneurons. *Neuron* 57:746–759.
47. Caterina MJ, et al. (2000) Impaired nociception and pain sensation in mice lacking the capsaicin receptor. *Science* 288:306–313.
48. Salas MM, Hargreaves KM, Akopian AN (2009) TRPA1-mediated responses in trigeminal sensory neurons: interaction between TRPA1 and TRPV1. *Eur J Neurosci* 29:1568–1578.
49. Staruschenko A, Jeske NA, Akopian AN (2010) Contribution of TRPV1-TRPA1 interaction to the single channel properties of the TRPA1 channel. *J Biol Chem* 285:15167–15177.
50. Miyamoto T, Dubin AE, Petrus MJ, Patapoutian A (2009) TRPV1 and TRPA1 mediate peripheral nitric oxide-induced nociception in mice. *PLoS ONE* 4:e7596.
51. Bandell M, et al. (2004) Noxious cold ion channel TRPA1 is activated by pungent compounds and bradykinin. *Neuron* 41:849–857.
52. Schmidt M, Dubin AE, Petrus MJ, Earley TJ, Patapoutian A (2009) Nociceptive signals induce trafficking of TRPA1 to the plasma membrane. *Neuron* 64:498–509.
53. Nigam S, Nodes S, Cichon G, Corey EJ, Pace-Asciak CR (1990) Receptor-mediated action of hexoxilin A3 releases diacylglycerol and arachidonic acid from human neutrophils. *Biochem Biophys Res Commun* 171:944–948.
54. Malkmus SA, Yaksh TL (2004) Intrathecal catheterization and drug delivery in the rat. *Methods Mol Med* 99:109–121.
55. Abbadie C, Trafton J, Liu H, Mantyh PW, Basbaum AI (1997) Inflammation increases the distribution of dorsal horn neurons that internalize the neurokinin-1 receptor in response to noxious and non-noxious stimulation. *J Neurosci* 17:8049–8060.
56. Mantyh PW, et al. (1995) Rapid endocytosis of a G protein-coupled receptor: substance P evoked internalization of its receptor in the rat striatum in vivo. *Proc Natl Acad Sci USA* 92:2622–2626.
57. Lu SG, Zhang X, Gold MS (2006) Intracellular calcium regulation among subpopulations of rat dorsal root ganglion neurons. *J Physiol* 577:169–190.
58. Harmon GS, et al. (2010) Pharmacological correction of a defect in PPAR-gamma signaling ameliorates disease severity in Ctrf-deficient mice. *Nat Med* 16:313–318.
59. Blahó VA, Buczynski MW, Brown CR, Dennis EA (2009) Lipidomic analysis of dynamic eicosanoid responses during the induction and resolution of Lyme arthritis. *J Biol Chem* 284:21599–21612.