

UC Davis

UC Davis Previously Published Works

Title

A Syst-OMICS Approach to Ensuring Food Safety and Reducing the Economic Burden of Salmonellosis.

Permalink

<https://escholarship.org/uc/item/62x8k47v>

Journal

Frontiers in microbiology, 8(JUN)

ISSN

1664-302X

Authors

Emond-Rheault, Jean-Guillaume

Jeukens, Julie

Freschi, Luca

et al.

Publication Date

2017

DOI

10.3389/fmicb.2017.00996

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



OPEN ACCESS

Edited by:

Sabah Bidawid,
Health Canada, Canada

Reviewed by:

Young Min Kwon,
University of Arkansas, United States
Sheng Chen,
Hong Kong Polytechnic University,
Hong Kong

***Correspondence:**

Roger C. Levesque
rclevesq@ibis.ulaval.ca
Lawrence Goodridge
lawrence.goodridge@mcmcgill.ca

† These authors have contributed
equally to this work.

Specialty section:

This article was submitted to
Food Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 29 March 2017

Accepted: 17 May 2017

Published: 02 June 2017

Citation:

Emond-Rheault J-G, Jeukens J,
Freschi L, Kukavica-Ibrulj I, Boyle B,
Dupont M-J, Colavecchio A,
Barrere V, Cadieux B, Arya G,
Bekal S, Berry C, Burnett E,
Cavestri C, Chapin TK, Crouse A,
Daigle F, Danyluk MD, Delaquis P,
Dewar K, Doualla-Bell F, Fliss I,
Fong K, Fournier E, Franz E,
Garduno R, Gill A, Gruenheid S,
Harris L, Huang CB, Huang H,
Johnson R, Joly Y, Kerhoas M,
Kong N, Lapointe G, Larivière L,
Loignon S, Malo D, Moineau S,
Mottawea W, Mukhopadhyay K,
Nadon C, Nash J, Ngueng Feze I,
Ogunremi D, Perets A, Pilar AV,
Reimer AR, Robertson J, Rohde J,
Sanderson KE, Song L, Stephan R,
Tamber S, Thomassin P, Tremblay S,
Usongo V, Vincent C, Wang S,
Weadge JT, Wiedmann M,
Wijnands L, Wilson ED, Wittum T,
Yoshida C, Youfsi K, Zhu L,
Weimer BC, Goodridge L and
Levesque RC (2017) A Syst-OMICS
Approach to Ensuring Food Safety
and Reducing the Economic Burden
of Salmonellosis.
Front. Microbiol. 8:996.
doi: 10.3389/fmicb.2017.00996

A Syst-OMICS Approach to Ensuring Food Safety and Reducing the Economic Burden of Salmonellosis

Jean-Guillaume Emond-Rheault^{1†}, Julie Jeukens^{1†}, Luca Freschi^{1†}, Irena Kukavica-Ibrulj¹, Brian Boyle¹, Marie-Josée Dupont¹, Anna Colavecchio², Virginie Barrere², Brigitte Cadieux², Gitanjali Arya³, Sadjia Bekal⁴, Chrystal Berry³, Elton Burnett², Camille Cavestri⁵, Travis K. Chapin⁶, Alanna Crouse², France Daigle⁷, Michelle D. Danyluk⁶, Pascal Delaquis⁸, Ken Dewar^{2,9}, Florence Doualla-Bell⁴, Ismail Fliss⁵, Karen Fong¹⁰, Eric Fournier⁴, Eelco Franz¹¹, Rafael Garduno¹², Alexander Gill¹³, Samantha Gruenheid², Linda Harris¹⁴, Carol B. Huang¹⁵, Hongsheng Huang¹⁶, Roger Johnson³, Yann Joly², Maud Kerhoas⁷, Nguyet Kong¹⁵, Gisèle Lapointe¹⁷, Line Larivière², Stéphanie Loignon⁵, Danielle Malo², Sylvain Moineau⁵, Walid Mottawea^{2,18}, Kakali Mukhopadhyay², Céline Nadon³, John Nash³, Ida Ngueng Feze², Dele Ogunremi¹⁶, Ann Perets³, Ana V. Pilar², Aleisha R. Reimer³, James Robertson³, John Rohde¹⁹, Kenneth E. Sanderson²⁰, Lingqiao Song², Roger Stephan²¹, Sandeep Tamber¹³, Paul Thomassin², Denise Tremblay⁵, Valentine Usongo⁴, Caroline Vincent⁴, Siyun Wang¹⁰, Joel T. Weadge²², Martin Wiedmann²³, Lucas Wijnands¹¹, Emily D. Wilson²², Thomas Wittum²⁴, Catherine Yoshida³, Khadija Youfsi⁴, Lei Zhu², Bart C. Weimer¹⁵, Lawrence Goodridge^{2*} and Roger C. Levesque^{1*}

¹ Institute for Integrative and Systems Biology, Université Laval, Québec City, QC, Canada, ² McGill University, Montréal, QC, Canada, ³ National Microbiology Laboratory, Public Health Agency of Canada, Ottawa, ON, Canada, ⁴ Laboratoire de Santé Publique du Québec, Sainte-Anne-de-Bellevue, QC, Canada, ⁵ Université Laval, Québec City, QC, Canada, ⁶ Institute of Food and Agricultural Sciences, University of Florida, Gainesville, FL, United States, ⁷ Département de Microbiologie, Infectiologie et Immunologie, Université de Montréal, Montréal, QC, Canada, ⁸ Agriculture and Agri-Food Canada, Summerland, BC, Canada, ⁹ Génome Québec Innovation Center, Montréal, QC, Canada, ¹⁰ Food Safety Engineering, Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC, Canada, ¹¹ National Institute for Public Health and the Environment, Bilthoven, Netherlands, ¹² Canadian Food Inspection Agency, Halifax, NS, Canada, ¹³ Bureau of Microbial Hazards, Health Canada, Ottawa, ON, Canada, ¹⁴ UC Davis Food Science and Technology, Davis, CA, United States, ¹⁵ UC Davis School of Veterinary Medicine, Davis, CA, United States, ¹⁶ Canadian Food Inspection Agency, Ottawa, ON, Canada, ¹⁷ Food Science, University of Guelph, Guelph, ON, Canada, ¹⁸ Department of Microbiology and Immunology, Faculty of Pharmacy, Mansoura University, Mansoura, Egypt, ¹⁹ Department of Microbiology and Immunology, Dalhousie University, Halifax, NS, Canada, ²⁰ Department of Biological Sciences, University of Calgary, Calgary, AB, Canada, ²¹ Institute for Food Safety and Hygiene, University of Zurich, Zurich, Switzerland, ²² Biological and Chemical Sciences, Wilfrid Laurier University, Waterloo, ON, Canada, ²³ Department of Food Science, Cornell University, Ithaca, NY, United States, ²⁴ College of Veterinary Medicine, The Ohio State University, Columbus, OH, United States

The *Salmonella* Syst-OMICS consortium is sequencing 4,500 *Salmonella* genomes and building an analysis pipeline for the study of *Salmonella* genome evolution, antibiotic resistance and virulence genes. Metadata, including phenotypic as well as genomic data, for isolates of the collection are provided through the *Salmonella* Foodborne Syst-OMICS database (SalFoS), at <https://salfos.ibis.ulaval.ca/>. Here, we present our strategy and the analysis of the first 3,377 genomes. Our data will be used to draw potential links between strains found in fresh produce, humans, animals and the environment. The ultimate goals are to understand how *Salmonella* evolves over time, improve the accuracy of diagnostic methods, develop control methods in the field, and identify prognostic markers for evidence-based decisions in epidemiology and surveillance.

Keywords: *Salmonella*, foodborne pathogen, next-generation sequencing, bacterial genomics, phylogeny, antibiotic resistance, database

IMPORTANCE OF FOODBORNE *Salmonella* AS A MODEL IN LARGE-SCALE BACTERIAL GENOMICS

Salmonella enterica is a foodborne bacterial pathogen having at least 2,600 serotypes (Gal-Mor et al., 2014)¹ that contaminates a diversity of foods and is a leading cause of foodborne illnesses and mortality globally. In fact, there are an estimated 93.3 million cases of gastroenteritis due to non-typhoidal *Salmonella* infections each year, resulting in approximately 155,000 deaths (Majowicz et al., 2010). In Canada, non-typhoidal salmonellosis accounts for more than 88,000 cases of foodborne illness each year, and has among the highest incidence rate of any bacterial foodborne pathogen (Thomas et al., 2015). *S. enterica* is responsible for more than 50% of fresh produce-borne outbreaks, the highest number of foodborne outbreaks of any inspected food commodity in North America (Kozak et al., 2013). Because of its remarkable genomic diversity, *Salmonella* is found in complex environmental and ecological niches and survives in harsh environments for long periods (Podolak et al., 2010; Fatica and Schneider, 2011). Several research groups have identified relationships between some of the 2,557 *S. enterica* serotypes and specific foods, which suggests, that some food commodities act as reservoirs for particular serotypes (Kim, 2010; Jackson et al., 2013; Nuesch-Inderbinen et al., 2015).

Salmonella outbreaks are monitored with support from the PulseNet surveillance system in 86 countries² (Ribot and Hise, 2016; Scharff et al., 2016). PulseNet Canada³ is a national surveillance system used to quickly identify and respond to foodborne disease outbreaks, centralized at the National Microbiology Laboratory in Winnipeg, MB, and working in close collaboration with a network of federal and provincial public health laboratories and epidemiologists. Still, despite the availability of thousands of sequenced genomes, knowledge of genome evolution integrated with transmission and epidemiology is limited for produce-related outbreaks.

Studies of *S. enterica* population structure in humans, animals, food and the environment are central to understand the biodiversity, evolution, ecology and epidemiology of this pathogen. However, studies describing the genetic structure of *Salmonella* populations are commonly based on isolates drawn overwhelmingly from clinical collections (Hoffmann et al., 2014). This approach has resulted in a limited view of *Salmonella*'s evolutionary history (D'costa et al., 2006; Perry and Wright, 2014). In *Salmonella* as in many other bacterial pathogens, there is limited knowledge on how genome content, rearrangements and the complement of genes including those acquired by horizontal gene transfer (HGT) contribute to strain-specific phenotypes, including virulence (Casadevall, 2017). Various studies have sought to resolve the population structure of *Salmonella* using complementary subtyping methods including pulsed-field gel electrophoresis (PFGE), multiple

loci VNTR analysis (MLVA), 7-gene housekeeping schemes, whole-genome multi-locus sequence typing (wgMLST) profiles, pan- and core genome studies, and CRISPR analysis to define molecular signatures, pathogen subtypes and the potential for pathogenicity (Shariat and Dudley, 2014; Rouli et al., 2015; Liu et al., 2016). Next-generation sequencing (NGS) coupled with whole-genome comparison is well-positioned to become the gold standard subtyping method, as it offers previously unmatched resolution for phylogenetic analysis and rapid subtyping during investigation of food contamination and outbreaks (Ashton et al., 2016; Bekal et al., 2016).

THE Syst-OMICS Strategy


The application of genomics to infectious pathogens via WGS is transforming the practice of *Salmonella* diagnostics, epidemiology and surveillance. Genomic data are increasingly used to understand infectious disease epidemiology (Didelot et al., 2017). With rapidly falling costs and turnaround time, microbial WGS and analysis is becoming a viable strategy to identify the geographic origin of bacterial pathogens (Weedmark et al., 2015; Hoffmann et al., 2016). The objective of the Canadian-based international Syst-OMICS consortium is to sequence a minimum of 4,500 genomes, include the data in the *Salmonella* Foodborne Syst-OMICS database (SalFoS) at <https://SalFoS.ibis.ulaval.ca/>, share this information plus available metadata with Canadian federal and provincial regulators and the food industry, and develop pipelines to study these genomes. Genomics data will support molecular epidemiology and source attribution of outbreaks and has the potential for future genotypic antimicrobial susceptibility testing, as well as the identification of novel therapeutic targets and prognostic markers. Moreover, the large-scale genomics and evolutionary biology tools developed may lead to new strategies for countering not only *Salmonella* infections, but other pathogens as well (Little et al., 2012).

The Syst-OMICS project is based upon a systems approach (flowchart and screening method available in Supplementary File 1). First, the genome diversity of 4,500 isolates will be assessed using high-quality WGS, assembly, annotation and phylogeny. This data will be used for *in silico* serotyping (Yoshida et al., 2016), as well as analysis of virulence (Chen et al., 2012), antibiotic resistance (Jia et al., 2016) and mobilome gene content (Lanza et al., 2014). Based on this genomic data, a funnel-type model will be applied such that 300 isolates will be selected for *in vitro* high-throughput screening (HTS) in cell lines to determine attachment, adhesion, invasion and replication of each isolate (protocol adapted to 96-well plates from Forest et al., 2007). From the results, isolates will be categorized as being of high, medium, or low virulence. A limited number of those isolates will then be selected for further screening *in vivo* using a mouse model (Roy et al., 2007) and *in vitro* using gastrointestinal fermenter models (Kheadr et al., 2010; Le Blay et al., 2012). These data will identify isolates to represent the different levels of virulence that will be used to develop novel diagnostic and control tools. We propose to enhance food safety and lower the economic burden of salmonellosis through a farm-to-table

¹<https://www.cdc.gov/salmonella/reportspubs/salmonella-atlas/serotype-snapshots.html>

²<http://www.pulsenetinternational.org/networks/usa/>

³<https://www.nml-lnm.gc.ca/Pulsenet/index-eng.htm>



systematic approach to control *Salmonella*, with a focus on new control methods in agricultural production, more specific diagnostics and improved bacterial subtyping methods to support investigation of foodborne outbreaks, as no single intervention is likely to produce meaningful and lasting effects.

THE *Salmonella* FOODBORNE Syst-OMICS DATABASE (SalFoS)


Salmonella Foodborne Syst-OMICS database is an online web application that relies on a MySQL 5 database. It was designed not only to store data for the *Salmonella* strain collection but also to provide access to each isolate's phenotypic, genomic, virulence, serotype, mobilome and epidemiological data. Different levels of access may be granted, but data modification is strictly reserved to the curators. It includes isolate identification, host, provider, date of isolation, geographical origin, phenotypic data, DNA extraction details, NGS information and genome assembly statistics. SalFoS currently contains NGS data and unpublished draft genomes from produce, human, animal and environmental isolates. Upon publication, draft genomes of SalFoS will become available at NCBI and EnteroBase⁴.

⁴<http://enterobase.warwick.ac.uk/>

The SalFoS collection currently contains 2,498 entries for *Salmonella*, as well as for *Citrobacter*, *Hafnia* and *Proteus*, three genera often identified as false-positives by a number of *Salmonella* detection schemes. It includes previously described collections such as the unique *Salmonella* Genetic Stock Centre strains, described at <http://people.ucalgary.ca/~kesander/>. This collection was assembled with the aim of representing maximal genomic diversity.

SEQUENCING 4,500 *Salmonella* GENOMES: OBJECTIVES AND STRATEGY

Our working hypothesis is that a very high-quality, large-scale bacterial genome database available through a user-friendly pipeline will have a major impact for epidemiology, diagnosis, prevention and treatment. By generating a comprehensive genome sequence database truly representative of the foodborne *Salmonella* population, we will: (1) assemble a large and representative strain collection, with associated genome data, useful for antimicrobial testing, identification of resistance markers, data mining for new therapeutic targets and development of machine learning strategies; (2) develop



platforms and pipelines to manage and analyze this information, which will allow identification of prognostic markers, fast epidemiological tracking and reduction of socio-economic costs. We seek to develop user-friendly tools that will enable epidemiologists, microbiologists, clinicians and others to interpret genomic data, thus leading to informed decisions in cases of food contamination and outbreaks. The contamination of fresh produce by *Salmonella* will be addressed through the development of natural solutions to control the presence of *Salmonella* on fruits and vegetables as they are growing in the fields. New tests will also be developed so that fresh produce can be quickly and efficiently tested for the presence of *Salmonella* before being sold to consumers. In the context of outbreak investigation, the genomic data will be used to assess high-quality SNPs and core/whole genome MLST for their usefulness in genetic discrimination in addition to other emerging methods such as CRISPR and prophage sequence typing. As for outbreak investigation software, the National Microbiology Laboratory-Public Health Agency of Canada group has implemented the Integrated Rapid Infectious Disease Analysis project (IRIDA)⁵ and developed the SNVPhyl phylogenomics pipeline that is in use by PulseNet Canada for microbial genomic epidemiology (Petkau et al., 2016). A complementary system called the Metagenomics Computation and Analytics Workbench (MCAW) is being implemented as a computing service for food safety (Edlund et al., 2016; Weimer et al., 2016).

Sequencing for this project is performed on an Illumina MiSeq instrument (at the Plateforme d'Analyses Génomiques of the IBIS, Université Laval, Quebec City, QC, Canada), at a rate of 120 genomes per week, using 300 bp paired-end libraries, and with a median coverage of 45×. In order to perform core genome phylogenetic analysis, the pan-genome, i.e., the complete repertoire of genes of a species, is determined using a recently developed software capable of handling high-quality NGS data from thousands of genomes: Saturn V version 1.0⁶ (Jeukens et al., 2017). Additional analyses focus on genes implicated in virulence using comparative genomics predictions of confirmed and predicted virulence factors (Yang et al., 2008), and resistance identification based on the comprehensive antibiotic resistance database (CARD) (Mcarthur et al., 2013; Jia et al., 2016). A set of new reference *Salmonella* genomes representing maximal genomic diversity among foodborne pathogens will then be selected for PacBio Sequel sequencing to become fully assembled and annotated as a single circular chromosome.

THE IBIS BIOINFORMATICS PIPELINE FOR GENOME ASSEMBLY

When working with hundreds or thousands of genomes, analysis software for assembly, annotation, statistics for quality control and selection of additional reference genomes is required to extract relevant information in an automated and reliable fashion

⁵<http://dev.irida.ca/>

⁶<https://github.com/efresch/saturnV>

with minimal human intervention. Ideally, this software should be platform independent and able to analyze sequence data directly without being tied to proprietary data formats. This insures maximal flexibility and reduces lag time to a minimum. We are currently using an integrated pipeline for *de novo* assembly of microbial genomes based on the A5 pipeline (Tritt et al., 2012). It was parallelized on a Silicon Graphics UV 300 using up to 120 cores to accommodate raw data from 120 genomes and provide assembly statistics as well as reference genome alignment metrics in as little as 2 h. This automated approach currently results in a median of 35 scaffolds per genome (median N50 = 462 kb).

PHYLOGENY OF *Salmonella*

Once isolates from a given outbreak are sequenced, patterns of shared variations can be used to infer which isolates within the outbreak are most closely related to each other (e.g., Didelot et al., 2017). As a future strategy for the Syst-OMICS project, this could be applied to partially sampled and on-going *Salmonella* outbreaks. Here, as a first step in the study of *S. enterica* diversity and epidemiology, we used 3,377 genomes; 1,627 were from a collaboration with UC Davis (Bart C. Weimer), and 1,750 were part of SalFoS. All genomes with >100 scaffolds were eliminated; this filter typically removes the vast majority of low coverage (i.e., low quality) assemblies and mixed cultures. As our assembly pipeline also includes alignment on a suite of reference genomes, it is also possible to ensure that genomes used belong to *S. enterica*. The core (conserved) genome was identified with Saturn V, and consisted of 839 genes, which were used for phylogenetic analysis. This number of core genes, which seems small compared to other studies (2,882 core genes for 73 genomes from 2 subspecies, Leekitcharoenphon et al., 2012), is due to both the extensive diversity and the high number of genomes used. As depicted in **Figure 1**, this population of *S. enterica* strains could be divided into seven major groups. They correspond to *S. enterica* subspecies *enterica* clades A and B and a collection of branching subspecies previously defined as *salamae*, *arizonae*, *diarizonae*, *houtenae* and *indica*. The significant number of strains (3,377) included in our analysis and their wide-ranging sources (including environmental, human, animal and food) is essential to understand the diversity of *Salmonella* as a foodborne pathogen and in defining levels of virulence. The remarkable genomic diversity exhibited in **Figure 1** is thought to enable the colonization of a wide range of ecological niches. The *Salmonella* Syst-OMICS consortium will provide fine-scale analysis of this diversity via virulence factors, antibiotic resistance genes as well as complete core and accessory genomes.

LINKING SalFoS WITH THE COMPREHENSIVE ANTIBIOTIC RESISTANCE DATABASE

The SalFoS database is intended to become an established platform for searching and comparing multiple genome

sequences for *Salmonella* isolates. The database will also incorporate genome annotation and serotype prediction based on SISTR (Yoshida et al., 2016). Close attention to the links between specific genomic islands and patterns of SNPs in the core genome will help identify diagnostic sequences and SNP combinations for the development of new *Salmonella* subtyping methods with the highest resolution to date. This will be done using *de novo* island prediction with IslandViewer (Langille and Brinkman, 2009; Dhillon et al., 2015) as well as with gene presence-absence from SaturnV.

As an additional feature, we routinely determine the resistome of the genomes in SalFoS, i.e., the genes and variants likely involved in antibiotic resistance. This is done using the Resistance Gene Identifier (RGI) available with the CARD (Mcarthur et al., 2013; Jia et al., 2016), at <http://arpcard.mcmaster.ca/>. **Figure 2** summarizes the resistomes of 3,377 genomes. In fact, the original dataset contained 1,003 unique resistomes, composed of various combinations of 195 different genes and variants. Despite this impressive diversity, the most striking feature shown in **Figure 2** is that the two most frequently observed resistomes, which are extremely similar, account for 23% of the strains. They are therefore highly conserved and warrant further investigation. These results will be exploited to study and understand the pool of resistance genes present in *Salmonella* strains, with a focus on strains found in fresh produce, to understand the links between foodborne *Salmonella* and environmental strains with respect to resistance genes.

LINKING GENOMIC AND CLINICAL DATA

It will be essential to match phenotypic, epidemiological and available clinical *Salmonella* data (antibiotic resistance, virulence, and anonymized clinical observations) to the genomic data produced. We will categorize metadata in SalFoS so that isolates can be sorted by phenotype, allowing rapid identification of linked genomic signatures and the development of prognostic approaches for diagnostic, epidemiology and surveillance. We will develop tools to rapidly collate data for a given strain type and produce a concise phenotypic and clinical profile that provides users with an evidence-based decision-making platform. The Canadian Food Inspection Agency, Health Canada, Agriculture Canada, provincial public health laboratories and the National Microbiology Laboratory-Public Health Agency of Canada group are expected to be end-users of the projects outcomes.

REFERENCES

- Ashton, P. M., Nair, S., Peters, T. M., Bale, J. A., Powell, D. G., Painset, A., et al. (2016). Identification of *Salmonella* for public health surveillance using whole genome sequencing. *PeerJ* 4:e1752. doi: 10.7717/peerj.1752
- Bekal, S., Berry, C., Reimer, A. R., Van Domselaar, G., Beaudry, G., Fournier, E., et al. (2016). Usefulness of high-quality core genome single-nucleotide variant analysis for subtyping the highly clonal and the most prevalent *Salmonella enterica* serovar heidelberg clone in the context of outbreak investigations. *J. Clin. Microbiol.* 54, 289–295. doi: 10.1128/JCM.02200-15

FUTURE GENOMIC AND BIOLOGICAL STUDIES OF *Salmonella*

We will continuously improve SalFoS by adding *Salmonella* strains, NGS data and analysis as well as experimental results. Another aim of the Syst-OMICS consortium is to avoid duplication of efforts in *Salmonella* genomics and enhance interest from researchers having common goals. Additional members are welcome to join in and expand on our original Genome Canada project. We also intend to seek collaboration with other groups to connect our database with those developed for other *Salmonella* genomes. Finally, the *Salmonella* Syst-OMICS project could be a model for other groups interested in the bacterial genomics of infectious diseases, a strategy that we are also pursuing for *Pseudomonas aeruginosa* (Freschi et al., 2015).

AUTHOR CONTRIBUTIONS

J-GER, JJ, LF, IK-I and RL collected strains, performed the analyses and drafted the manuscript. BB provided support for sequencing and analysis. MD contributed to the development of SalFoS. All other authors handled strains and collected metadata. All authors revised the manuscript.

ACKNOWLEDGMENTS

We express our gratitude to members of the genomics analysis and bioinformatics platforms at IBIS. We also acknowledge Betty Wilkie, Ketna Mistry, Robert Holtslander and Shaun Kenaghan from the NML *Salmonella* reference laboratory for their assistance with serotyping. RL, LG, AG, ST, PD, DM, SG, SB, FD, SW, SM, GL, INF, YJ, PT, CN, RG, JoR, JW are funded by Genome Canada, provincial genome centers Génome Québec and Genome BC, and the Ontario Ministry of Research and Innovation. SM holds a Tier 1 Canada Research Chair in Bacteriophages.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <http://journal.frontiersin.org/article/10.3389/fmicb.2017.00996/full#supplementary-material>

- Casadevall, A. (2017). The pathogenic potential of a microbe. *mSphere* 2:e00015-17. doi: 10.1128/mSphere.00015-17
- Chen, L., Xiong, Z., Sun, L., Yang, J., and Jin, Q. (2012). VFDB 2012 update: toward the genetic diversity and molecular evolution of bacterial virulence factors. *Nucleic Acids Res.* 40, D641–D645. doi: 10.1093/nar/gkr989
- D'costa, V. M., Mcgrann, K. M., Hughes, D. W., and Wright, G. D. (2006). Sampling the antibiotic resistome. *Science* 311, 374–377. doi: 10.1126/science.1120800
- Dhillon, B. K., Laird, M. R., Shay, J. A., Winsor, G. L., Lo, R., Nizam, F., et al. (2015). IslandViewer 3: more flexible, interactive genomic island discovery, visualization and analysis. *Nucleic Acids Res.* 43, 27. doi: 10.1093/nar/gkv401

- Didelot, X., Fraser, C., Gardy, J., and Colijn, C. (2017). Genomic infectious disease epidemiology in partially sampled and ongoing outbreaks. *Mol. Biol. Evol.* 34, 997–1007. doi: 10.1093/molbev/msw275
- Edlund, S. B., Beck, K. L., Haiminen, N., Parida, L. P., Storey, D. B., Weimer, B. C., et al. (2016). Design of the MCAW compute service for food safety bioinformatics. *IBM J. Res. Dev.* 60:12. doi: 10.1147/JRD.2016.2584798
- Fatica, M. K., and Schneider, K. R. (2011). *Salmonella* and produce: survival in the plant environment and implications in food safety. *Virulence* 2, 573–579. doi: 10.4161/viru.2.6.17880
- Forest, C., Faucher, S. P., Poirier, K., Houle, S., Dozois, C. M., and Daigle, F. (2007). Contribution of the stg fimbrial operon of *Salmonella enterica* serovar typhi during interaction with human cells. *Infect. Immun.* 75, 5264–5271. doi: 10.1128/iai.00674-07
- Freschi, L., Jeukens, J., Kukavica-Ibrulj, I., Boyle, B., Dupont, M. J., Laroche, J., et al. (2015). Clinical utilization of genomics data produced by the international *Pseudomonas aeruginosa* consortium. *Front. Microbiol.* 6:1036. doi: 10.3389/fmicb.2015.01036
- Gal-Mor, O., Boyle, E. C., and Grassl, G. A. (2014). Same species, different diseases: how and why typhoidal and non-typhoidal *Salmonella enterica* serovars differ. *Front. Microbiol.* 5:391. doi: 10.3389/fmicb.2014.00391
- Hoffmann, M., Luo, Y., Monday, S. R., Gonzalez-Escalona, N., Ottesen, A. R., Muruvanda, T., et al. (2016). Tracing origins of the *Salmonella* bareilly strain causing a food-borne outbreak in the United States. *J. Infect. Dis.* 213, 502–508. doi: 10.1093/infdis/jiv297
- Hoffmann, M., Zhao, S., Pettengill, J., Luo, Y., Monday, S. R., Abbott, J., et al. (2014). Comparative genomic analysis and virulence differences in closely related *Salmonella enterica* serotype Heidelberg isolates from humans, retail meats, and animals. *Genome Biol. Evol.* 6, 1046–1068. doi: 10.1093/gbe/evu079
- Jackson, B. R., Griffin, P. M., Cole, D., Walsh, K. A., and Chai, S. J. (2013). Outbreak-associated *Salmonella enterica* serotypes and food commodities, United States, 1998–2008. *Emerg. Infect. Dis.* 19, 1239–1244. doi: 10.3201/eid1908.121511
- Jeukens, J., Freschi, L., Vincent, A. T., Emond-Rheault, J. G., Kukavica-Ibrulj, I., Charette, S. J., et al. (2017). A pan-genomic approach to understand the basis of host adaptation in *Achromobacter*. *Genome Biol. Evol.* doi: 10.1093/gbe/evx061 [Epub ahead of print].
- Jia, B., Raphenya, A. R., Alcock, B., Waglechner, N., Guo, P., Tsang, K. K., et al. (2016). CARD 2017: expansion and model-centric curation of the comprehensive antibiotic resistance database. *Nucleic Acids Res.* 45, D566–D573. doi: 10.1093/nar/gkw1004
- Kheadr, E., Zihler, A., Dabour, N., Lacroix, C., Le Blay, G., and Fliss, I. (2010). Study of the physicochemical and biological stability of pediocin PA-1 in the upper gastrointestinal tract conditions using a dynamic in vitro model. *J. Appl. Microbiol.* 109, 54–64. doi: 10.1111/j.1365-2672.2009.04644.x
- Kim, S. (2010). *Salmonella* serovars from foodborne and waterborne diseases in Korea, 1998–2007: total isolates decreasing versus rare serovars emerging. *J. Kor. Med. Sci.* 25, 1693–1699. doi: 10.3346/jkms.2010.25.12.1693
- Kozak, G. K., Macdonald, D., Landry, L., and Farber, J. M. (2013). Foodborne outbreaks in Canada linked to produce: 2001 through 2009. *J. Food Prot.* 76, 173–183. doi: 10.4315/0362-028X.JFP-12-126
- Langille, M. G. I., and Brinkman, F. S. L. (2009). IslandViewer: an integrated interface for computational identification and visualization of genomic islands. *Bioinformatics* 25, 664–665. doi: 10.1093/bioinformatics/btp030
- Lanza, V. F., De Toro, M., Garcillan-Barcia, M. P., Mora, A., Blanco, J., Coque, T. M., et al. (2014). Plasmid flux in *Escherichia coli* ST131 sublineages, analyzed by plasmid constellation network (PLACNET), a new method for plasmid reconstruction from whole genome sequences. *PLoS Genet.* 10:e1004766. doi: 10.1371/journal.pgen.1004766
- Le Blay, G., Hammami, R., Lacroix, C., and Fliss, I. (2012). Stability and inhibitory activity of pediocin PA-1 against *Listeria* sp. in simulated physiological conditions of the human terminal ileum. *Probiotics Antimicrob. Proteins* 4, 250–258. doi: 10.1007/s12602-012-9111-1
- Leekitcharoenphon, P., Lukjancenko, O., Friis, C., Aarestrup, F. M., and Ussery, D. W. (2012). Genomic variation in *Salmonella enterica* core genes for epidemiological typing. *BMC Genomics* 13:88. doi: 10.1186/1471-2164-13-88
- Little, T. J., Allen, J. E., Babayan, S. A., Matthews, K. R., and Colegrave, N. (2012). Harnessing evolutionary biology to combat infectious disease. *Nat. Med.* 18, 217–220. doi: 10.1038/nm.2572
- Liu, Y. Y., Chen, C. C., and Chiou, C. S. (2016). Construction of a pan-genome allele database of *Salmonella enterica* serovar enteritidis for molecular subtyping and disease cluster identification. *Front. Microbiol.* 7:2010. doi: 10.3389/fmicb.2016.02010
- Majowicz, S. E., Musto, J., Scallan, E., Angulo, F. J., Kirk, M., O'Brien, S. J., et al. (2010). The global burden of nontyphoidal *Salmonella* gastroenteritis. *Clin. Infect. Dis.* 50, 882–889. doi: 10.1086/650733
- Mcarthur, A. G., Waglechner, N., Nizam, F., Yan, A., Azad, M. A., Baylay, A. J., et al. (2013). The comprehensive antibiotic resistance database. *Antimicrob. Agents Chemother.* 57, 3348–3357. doi: 10.1128/AAC.00419-13
- Nuesch-Inderbinen, M., Cernela, N., Althaus, D., Hachler, H., and Stephan, R. (2015). *Salmonella enterica* serovar szentes, a rare serotype causing a 9-month outbreak in 2013 and 2014 in Switzerland. *Foodborne Pathog. Dis.* 12, 887–890. doi: 10.1089/fpd.2015.1996
- Perry, J. A., and Wright, G. D. (2014). Forces shaping the antibiotic resistome. *Bioessays* 36, 1179–1184. doi: 10.1002/bies.201400128
- Petkau, A., Mabon, P., Sieffert, C., Knox, N., Cabral, J., Iskander, M., et al. (2016). SNVPhyl: a single nucleotide variant phylogenomics pipeline for microbial genomic epidemiology. *bioRxiv* 092940. doi: 10.1101/092940
- Podolak, R., Enache, E., Stone, W., Black, D. G., and Elliott, P. H. (2010). Sources and risk factors for contamination, survival, persistence, and heat resistance of *Salmonella* in low-moisture foods. *J. Food Prot.* 73, 1919–1936.
- Ribot, E. M., and Hise, K. B. (2016). Future challenges for tracking foodborne diseases: pulseNet, a 20-year-old US surveillance system for foodborne diseases, is expanding both globally and technologically. *EMBO Rep.* 17, 1499–1505. doi: 10.15252/embr.201643128
- Rouli, L., Merhej, V., Fournier, P. E., and Raoult, D. (2015). The bacterial pangenome as a new tool for analysing pathogenic bacteria. *New Microbes New Infect* 7, 72–85. doi: 10.1016/j.nmni.2015.06.005
- Roy, M.-F., Riendeau, N., Bédard, C., Hélie, P., Min-Oo, G., Turcotte, K., et al. (2007). Pyruvate kinase deficiency confers susceptibility to *Salmonella* Typhimurium infection in mice. *J. Exp. Med.* 204, 2949–2961. doi: 10.1084/jem.20062606
- Scharff, R. L., Besser, J., Sharp, D. J., Jones, T. F., Peter, G.-S., and Hedberg, C. W. (2016). An Economic Evaluation of PulseNet. *Am. J. Prev. Med.* 50, S66–S73. doi: 10.1016/j.amepre.2015.09.018
- Shariat, N., and Dudley, E. G. (2014). CRISPRs: molecular signatures used for pathogen subtyping. *Appl. Environ. Microbiol.* 80, 430–439. doi: 10.1128/AEM.02790-13
- Thomas, M. K., Murray, R., Flockhart, L., Pintar, K., Fazil, A., Nesbitt, A., et al. (2015). Estimates of foodborne illness-related hospitalizations and deaths in Canada for 30 specified pathogens and unspecified agents. *Foodborne Pathog. Dis.* 12, 820–827. doi: 10.1089/fpd.2015.1966
- Timme, R. E., Pettengill, J. B., Allard, M. W., Strain, E., Barrangou, R., Wehnes, C., et al. (2013). Phylogenetic diversity of the enteric pathogen *Salmonella enterica* subsp. *enterica* inferred from genome-wide reference-free SNP characters. *Genome Biol. Evol.* 5, 2109–2123. doi: 10.1093/gbe/evt159
- Tritt, A., Eisen, J. A., Facciotti, M. T., and Darling, A. E. (2012). An integrated pipeline for de novo assembly of microbial genomes. *PLoS ONE* 7:e42304. doi: 10.1371/journal.pone.0042304
- Weedmark, K. A., Mabon, P., Hayden, K. L., Lambert, D., Van Domselaar, G., Austin, J. W., et al. (2015). Clostridium botulinum Group II isolate phylogenomic profiling using whole-genome sequence data. *Appl. Environ. Microbiol.* 81, 5938–5948. doi: 10.1128/aem.01155-15
- Weimer, B. C., Storey, D. B., Elkins, C. A., Baker, R. C., Markwell, P., Chambliss, D. D., et al. (2016). Defining the food microbiome for authentication, safety, and process management. *IBM J. Res. Dev.* 60:13. doi: 10.1147/JRD.2016.2582598
- Yang, J., Chen, L., Sun, L., Yu, J., and Jin, Q. (2008). VFDB 2008 release: an enhanced web-based resource for comparative pathogenomics. *Nucleic Acids Res.* 36, D539–D542. doi: 10.1093/nar/gkm951
- Yoshida, C. E., Kruczkiewicz, P., Laing, C. R., Lingohr, E. J., Gannon, V. P., Nash, J. H., et al. (2016). The *Salmonella* in silico typing resource (SISTR): an open web-accessible tool for rapidly typing and subtyping draft *Salmonella* genome assemblies. *PLoS ONE* 11:e0147101. doi: 10.1371/journal.pone.0147101

Conflict of Interest Statement: The handling Editor declared a shared affiliation, though no other collaboration, with the authors ST and AG, and the handling Editor states that the process met the standards of a fair and objective review.

The other authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Emond-Rheault, Jeukens, Freschi, Kukavica-Ibrulj, Boyle, Dupont, Colavecchio, Barrere, Cadieux, Arya, Bekal, Berry, Burnett, Cavestri, Chapin, Crouse, Daigle, Danyluk, Delaquis, Dewar, Doualla-Bell, Fliss, Fong, Fournier,

Franz, Garduno, Gill, Gruenheid, Harris, Huang, Huang, Johnson, Joly, Kerhoas, Kong, Lapointe, Larivière, Loignon, Malo, Moineau, Mottawea, Mukhopadhyay, Nadon, Nash, Ngueng Feze, Ogunremi, Perets, Pilar, Reimer, Robertson, Rohde, Sanderson, Song, Stephan, Tamber, Thomassin, Tremblay, Usongo, Vincent, Wang, Weadge, Wiedmann, Wijnands, Wilson, Wittum, Yoshida, Youfsi, Zhu, Weimer, Goodridge and Levesque. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.