

# UC Irvine

## UC Irvine Previously Published Works

### Title

Developmental and cultural perspectives on children's postoperative pain management at home

### Permalink

<https://escholarship.org/uc/item/64c8s83x>

### Journal

Pain Management, 4(6)

### ISSN

1758-1869

### Authors

Jenkins, Brooke N  
Fortier, Michelle A

### Publication Date

2014-11-01

### DOI

10.2217/pmt.14.44


### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Developmental and cultural perspectives on children's postoperative pain management at home

Brooke N Jenkins<sup>1</sup> & Michelle A Fortier<sup>\*1</sup>



### Practice Points

#### Background

- Postoperative pain in children is extremely common and is undertreated in the home setting.

#### Parent management of child pain

- Interventions based on empirical findings may be an avenue to improve the management of children's postoperative pain in the home setting.

#### Cultural factors in pain

- Pain management is impacted by parent beliefs regarding analgesic use for children, developmental and language status of children and cultural and language factors.

#### Best practice

- Proper pain assessment, around-the-clock dosing and a multimodal approach are essential for the optimal management of children's postoperative pain.

**SUMMARY** Outpatient surgery is extremely common in children, and approximately 4 million children experience significant pain after surgery in the USA each year. Management of children's postoperative pain in the home setting is suboptimal and is impacted by characteristics of children and parents, as well as the larger family and cultural context. In particular, developmental status of the child, parental beliefs regarding pain expression and analgesic use in children, cultural values and language barriers can affect management of children's postoperative pain. Targeting the myriad barriers to children's pain management by capitalizing upon the use of tailored interventions may help bridge the gap between the translation of pain management guidelines to the home setting.

### KEYWORDS

- cultural • developmental
- parent management
- pediatric • postoperative pain

Pain is a universal experience that is affected by biological and psychosocial factors. As defined by the International Association for the Study of Pain, "pain is ... an unpleasant sensory and emotional experience associated with actual or potential tissue damage, or described in terms of such damage" [1]. Included in the experience of pain are myriad psychosocial factors that are widely accepted to influence the experience and expression of pain, including race/ethnicity, culture, development and the family environment [2,3]. Accordingly, pain often presents a dilemma to the treating healthcare provider given the complexity of its experience.

Postoperative pain is extremely common in children. Each year millions of children in the USA undergo surgery [4], many of whom experience significant amounts of postoperative pain [5]. Pain continues to be prevalent upon discharge home; in fact, our group and others have documented that approximately 80% of children report immediate postoperative pain at home and up to half continue to experience pain 1 week following surgery [6,7]. The undertreatment of pain in the postoperative setting is particularly problematic given the impact of children's pain experiences. Specifically, children may experience maladaptive behavioral changes, increased analgesic requirements and

<sup>1</sup>Department of Anesthesiology & Perioperative Care, University of California, Irvine, CA 92697, USA

\*Author for correspondence: Tel.: +1 714 456 2833; Fax: +1 714 480 0733; mfortier@uci.edu

delayed postoperative recovery due to pain [8]. Furthermore, postoperative pain is a strong predictor of unanticipated hospitalization [9], the development of chronic pain [10] and greater sensitivity toward pain in the future [11]. Moreover, early exposure to pain has been associated years later to adults' reports of increased pain and anxiety during medical events [12]. Recent findings have even demonstrated alterations in pain neuropathways as a result of activation of the nociceptive system early in development [13]. Accordingly, it is vital to provide optimal pain management in children after surgery in an effort to prevent negative sequelae.

### Parent management of child pain

Changes in healthcare have led to a transition of children's surgeries to be conducted primarily on an outpatient basis [4]. Accordingly, parents are becoming increasingly responsible for management of children's postoperative pain. Unfortunately, a growing body of literature indicates that parents tend to provide suboptimal postoperative pain management. For example, research by our group confirms that children experience high levels of pain following one of the most common pediatric surgeries, tonsillectomy and adenoidectomy (T&A), yet parents provide very few doses of analgesics in the home setting [6]. Additional research illustrates that families may fail to adhere to recommendations for administration of analgesia from their healthcare providers, but not necessarily because their children are not experiencing pain [14–18]. For example, the majority of parents provide fewer than the prescribed number of analgesic doses to children [14,15,18]. Furthermore, out of the doses that parents do administer to children, the overwhelming majority (70%) are subtherapeutic. Parents may also substitute weaker medications than those prescribed, or stretch the time interval between analgesic doses [14–18].

The reasons behind parental undertreatment of children's postoperative pain are not entirely clear. Managing child postoperative pain may be difficult due to a number of reasons including child factors, such as fatigue from sleep disturbances due to surgery [19], medication factors, such as inadequate medication strength, system factors, such as lack of instruction on how to administer medication, and parental factors, such as attitudes toward medication [20]. Examining parental factors, our group and others have established a link between parental

misconceptions regarding analgesic use and pain expression in children and administration of pain medication in the home setting. Data from our research center have documented that parents endorse many misconceptions about using analgesia with children, such as beliefs that pain medication works best the less often it is used and analgesia should only be used when the pain is severe [21]. Parents also report fears of side effects and addiction potential and therefore, may withhold medication from their children. These misconceptions have also been empirically connected to the undertreatment of children's pain by parents. Specifically, our group has shown that the more parents endorse misconceptions regarding analgesic use for children, the fewer doses administered to children after surgery [22]. Moreover, we have also documented that parents may misunderstand the myriad ways that children can express pain. For example, many parents report that "Children always express pain by crying or whining," "Children complain about pain to get attention" and "Children who are quiet are not in pain" [21]. Thus, parents who are not able to detect pain will likely be unable to optimally treat pain. These misconceptions about medication and the lack of pain detection by parents are important as medication is only effective when used.

The Parents Postoperative Pain Measure (PPPM) [23] is one tool that may be helpful in allowing parents to detect child pain. The PPPM is a 15-item checklist of behaviors indicative of postoperative pain (e.g., whining, refusing to eat, holding the sore part of his/her body, etc.). This measure can be used by parents to assess their child's pain by looking for the occurrence of these behaviors. Furthermore, this measure also reflects behavioral indices of recovery. Because of the utility of the measure and because it is the only validated measure of parent report of child postoperative pain, the PPPM has been used in many studies [6–8].

In addition to using tools to assess pain, parents play a vital role in reducing children's pain through behavioral strategies. For example, Taddio and colleagues have reviewed and graded several guidelines for pain management based on empirical evidence [24]. In particular, such guidelines highlight parental use of distraction and coaching (e.g., encouraging child use of coping skills) as effective parent behavioral pain management techniques for acute pain in children.

---

### Developmental factors in pain

Developmental stages play a pivotal role in how a child's pain is managed. Frequencies of vocal expressions of pain such as crying, wailing, ingressive vocalization appear in children experiencing postoperative pain but specific rates of occurrence of these expressions change as children age and develop [25]. Verbal expressions often make assessing child pain easier compared with other vocalizations. When children are unable to express their pain verbally, healthcare provider and parental assessment of child pain is more challenging as adults cannot simply ask children about their discomfort. Given that self-report is considered the gold standard of pain management, lack of ability to verbally report, either due to age or developmental delay/cognitive impairment can provide challenges for the optimal management of postoperative pain.

Although pain assessment in nonverbal children can be difficult, validated observational tools have been developed in order to assess pain in this population. For young children, the Neonatal Infant Pain Scale (used for infants) [26] and the Face, Legs, Activity, Cry and Consolability (FLACC) scale (used for children under 4 years of age) [27] each provide observers with a checklist to assess pain-related behaviors. For children slightly older but who are still not verbally fluent, self-assessment pain measures through the use of scales such as the Faces Pain Scale-Revised (used for children aged 4 to 12 years) [28] and Wong-Baker FACES (used for children 3 years and older) [29] are helpful. These scales allow children to point to their level of pain depicted through faces. Additionally, the Visual Analog Scale allows children over the age of 7 years to point to their level of pain on a line 100 mm long with one end representing 'no pain' and the other representing 'worst pain' [30].

Observational measures can also be adapted for children with cognitive disabilities. If a child's verbal ability is impaired, self-report nonverbal measures or observational measures can be used. For example, the Noncommunicating Children's Pain Checklist, an observational measure and the FLACC have both been utilized to assess pain in children with disabilities [31–33].

In addition to determining verbal ability, the age of a child may impact the expression and management of pain. For example, older children have more resources in terms of coping strategies to handle their pain compared with

younger children [34]. Specifically, older children are better at describing their pain and using cognitive pain management skills. Older children are also better at distinguishing between their distressful states by identifying differences between fear and physical pain [35]. It is therefore not surprising that older children develop fewer behavioral problems after surgery compared with younger children [36]. For these reasons, the age of a child plays a large role in the course of their postoperative pain and implies that observational tools should be used as complements to self-report measures of pain in younger children.

---

### Cultural factors in pain

It is widely accepted that cultural variables can influence the expression and treatment of pain and, in fact, there is a growing body of ethnic and racial disparities in pain management [37,38]. Such disparities have been documented in the experience of postoperative pain in children. For example, examining differences in postoperative recovery, African-American children were shown to have higher pain scores and require more pain medication compared with white children [39]. In addition, white children experienced more analgesic side effects compared with African-American children. Our group has a growing body of evidence on cultural factors that impact management of children's postoperative pain. Specifically, findings from our laboratory suggest ethnic and language differences in parental beliefs about children's pain expression, suggesting that Spanish speaking parents report misconceptions to a greater degree than English speaking parents, even after controlling for group differences in socioeconomic status [40].

Language is another factor that can influence management of children's postoperative pain. For example, language barriers may lead to parents not fully understanding pain management instructions of healthcare providers and such barriers have been shown to be associated with low treatment adherence [41]. Moreover, there may be cultural preferences for use of complementary and alternative medicine (CAM) in addition to or in place of traditional pain management strategies [42], and data do support differences in CAM use based upon language. Our laboratory has documented that both English speaking mothers are more likely to use complementary and alternative medicine to treat child pain compared with Spanish

speaking mothers [43]. These differences were not associated with mothers' beliefs about CAM, nonetheless, such cultural beliefs should be explored further with more culturally sensitive measures of CAM attitudes.

Although ethnic disparities in children's pain management exist, we cannot fully tease apart the role of cultural values and socioeconomic factors. It is possible that values and beliefs are shaped by culture which in turn affects outcomes in pain management [44]. It is also possible that these differences could relate to socioeconomic status. For example, education may impact parent understanding of pain management and willingness to administer medication to children. It is likely that the disparities are a function of the more complex interplay of culture and SES and future research to more clearly identify the relationship among these variables is needed.

---

### Best practice

Assessment is key in adequately treating pain; therefore, ensuring that parents and caregivers can conduct proper assessment of children's postoperative pain at home is imperative. Educating parents on the use of validated pain scales and ensuring parents understand the various ways pain can be expressed by children (e.g., verbal distress, withdrawal, behavioral changes) can help parents detect when children may be in pain but are unable to directly express their discomfort. Avoiding assumptions regarding parent understanding of appropriate analgesic administration is vital. Given that analgesics are weight based and require proper measurement, teaching parents how to administer analgesics is crucial. The media recently reported on the issue of parental confusion regarding administration of liquid analgesics to children and how the 'old school' method of using a teaspoon can lead to errors in dosing. Moreover, parents may not know the differences or similarities in the variety of brand and generic medication names – we frequently encounter parents in our setting who are unaware that ibuprofen, Advil and Motrin are all the same analgesic, for example. In terms of dosing, it has long been proposed that optimal management of children's postoperative pain ought to include around-the-clock dosing of analgesics to prevent pain. We teach parents that it takes more medication to treat pain that has become severe than it does to prevent pain. Finally, pain is best treated using a multimodal approach – in other words, combining

pharmacological and nonpharmacological pain management strategies. Behavioral strategies, such as distraction, imagery and relaxation can positively impact children's acute pain when used in conjunction with analgesics.

---

### Conclusion & future perspective

Postoperative pain in children remains as a significant problem, particularly given that the majority of postoperative pain management occurs at home under the responsibility of parents and caregivers. A large body of research indicates parents provide suboptimal pain management in the home setting, which may relate to parental beliefs regarding analgesic use and pain expression in children. Developmental and cultural issues can also contribute to challenges in children's pain management. A child's developmental state due to age or ability affects pain expression. Similarly, a child's ability to cope with pain depends on age and maturity level. Cultural issues including ethnicity, language and socioeconomic status play a role in pain management as well. Accordingly, research in the areas of parental, developmental and cultural issues in children's postoperative pain management suggest that a tailored, rather than a 'one size fits all' approach is needed. Tailored interventions target specific populations and eliminate extraneous information so that information provided is directly relevant, and therefore, more likely to be effective [45]. Such interventions can be modified accordingly to children's age, parental beliefs, cultural values and any other factors empirically associated with children's pain management after surgery. Development of tailored interventions for children's postoperative pain may provide a fruitful avenue for improving children's pain management in the home setting.

Children's pain management has long been neglected in both clinical and research settings; however, we have seen a dramatic increase in the focus on this topic in recent decades. Children's pain management in the medical setting has greatly improved, particularly given the increased focus on pain as the 'fifth vital sign'. We also have available a wealth of knowledge on children's pain management via well-established and evidence-based guidelines, such as Practice Guidelines for Acute Pain Management in the Perioperative Setting published by the American Society of Anesthesiologists Task Force on Acute Pain Management [46]. However, there is a huge gap in the translation of knowledge

into practice, particularly with pain management in the home setting. Given the increased focus on children's pain management in the home setting and the incorporation of health information technology into the treatment of pain, it is expected that we would see significant improvements in the management of postoperative pain in children in the home setting in the next 5–10 years. In the next several years, we anticipate a growing number of studies reflecting development of tailored interventions to target pain in this area that capitalize on the Internet to provide ongoing access to strategies by parents to provide optimal treatment to children. In addition, given the growing use of ecological momentary assessment strategies such as pain diaries, healthcare providers

are increasingly provided with real-time data on pain, which allows for provision of real-time intervention to prevent increases in pain severity. Therefore, we hope to see a decrease in unnecessary suffering in children following surgery in the next decade.

**Financial & competing interests disclosure**

*The authors have no relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript. This includes employment, consultancies, honoraria, stock ownership or options, expert testimony, grants or patents received or pending or royalties.*

*No writing assistance was utilized in the production of this manuscript.*

**References**

Papers of special note have been highlighted as:  
 • of interest; •• of considerable interest

- 1 IASP Task Force on Taxonomy, *Classification of Chronic Pain (2nd Edition)*, IASP Press, Seattle, WA, USA (2004).
- 2 Free MM. Cross-cultural conceptions of pain and pain control. *Proc. (Bayl. Univ. Med. Cent.)* 15(2), 143–145 (2000).
- 3 Good MD, Brodwin PE, Good BJ. *Pain as Human Experience: An Anthropological Perspective*. University of California Press, CA, USA, 143–145 (1994).
- 4 Cullen KA, Hall MJ, Golosinskiy A. Ambulatory surgery in the United States, 2006. *Natl Heal. Stat. Rep.* 11, 1–25 (2009).
- 5 Fortier MA, Chou J, Maurer EL, Kain ZN. Acute to chronic postoperative pain in children: preliminary findings. *J. Pediatr. Surg.* 46(9), 1700–1705 (2011).
- 6 Fortier MA, MacLaren JE, Martin SR, Perret-Karimi D, Kain ZN. Pediatric pain after ambulatory surgery: where's the medication?. *Pediatrics* 124(4), e588–e595 (2009).
- **This study examined postoperative pain and medication consumption for children after surgery.**
- 7 Stewart DW, Ragg PG, Sheppard S, Chalkiadis GA. The severity and duration of postoperative pain and analgesia requirements in children after tonsillectomy, orchidopexy, or inguinal hernia repair. *Paediatr. Anaesth.* 22(2), 136–143 (2012).
- 8 Kain ZN, Mayes LC, Caldwell-Andrews AA, Karas DE, McClain BC. Preoperative anxiety, postoperative pain, and behavioral recovery in young children undergoing surgery. *Pediatrics* 118(2), 651–658 (2006).
- **By examining children before and after surgery, this study found that presurgery anxiety was related to greater levels of postoperative pain.**
- 9 Coley KC, Williams BA, Dapos SV, Chen C, Smith RB. Retrospective evaluation of readmissions after same day surgery and associated costs. *8180(2)*, 349–353 (2002).
- 10 Perkins FM, Kehlet H. Chronic pain as an outcome of surgery. A review of predictive factors. *Anesthesiology* 93(4), 1123–1133 (2000).
- 11 Taddio A, Katz A, Ilersich AL, Koren G. Effect of neonatal circumcision on pain response during subsequent routine vaccination. *Lancet* 349(9052), 599–603 (1997).
- 12 Pate J, Blount R, Cohen L, Smith A. Childhood medical experience and temperament as predictors of adult functioning in medical situations. *Child. Health Care* 25(4), 281–298 (1996).
- 13 Woolf CJ, Salter MW. Neuronal plasticity: increasing the gain in pain. *Science* 288(5472), 1765–1769 (2000).
- 14 Finley GA, McGrath PJ, Forward SP, McNeill G, Fitzgerald P. Parents' management of children's pain following 'minor' surgery. *Pain* 64(1), 83–87 (1996).
- **This investigation documents child pain and the amount of medication given by parents at home after surgery.**
- 15 Hamers JP, Abu-Saad HH. Children's pain at home following (adeno) tonsillectomy. *Eur. J. Pain* 6(3), 213–219 (2002).
- 16 Kankkunen P, Vehvilainen-Julkunen K, Pietila AM, Kokki H, Halonen P. Parents' perceptions and use of analgesics at home after children's day surgery. *Paediatr. Anaesth.* 13(2), 132–140 (2003).
- **This study found that at least some parents had inaccurate perceptions of children's analgesics.**
- 17 Munro HM, Malviya S, Lauder GR, Voepel-Lewis T, Tait A. Pain relief in children following outpatient surgery. *J. Clin. Anesth.* 11(3), 187–191 (1999).
- 18 Warnock FF, Lander J. Pain progression, intensity and outcomes following tonsillectomy. *Pain* 75(1), 37–45 (1998).
- 19 MacLaren JE, Kain ZN. Prevalence and predictors of significant sleep disturbances in children undergoing ambulatory tonsillectomy and adenoidectomy. *J. Pediatr. Psychol.* 33(3), 248–257 (2008).
- 20 Dorkham MC, Chalkiadis GA, von Ungern Sternberg BS, Davidson AJ. Effective postoperative pain management in children after ambulatory surgery, with a focus on tonsillectomy: barriers and possible solutions. *Paediatr. Anaesth.* 24(3), 239–248 (2014).
- 21 Zisk RY, Grey M, MacLaren JE, Kain ZN. Exploring sociodemographic and personality characteristic predictors of parental pain perceptions. *Anesth. Analg.* 104(4), 790–798 (2007).
- 22 Rony RYZ, Fortier MA, Chorney JM, Perret D, Kain ZN. Parental postoperative pain management: attitudes, assessment, and management. *Pediatrics* 125(6), e1372–e1378 (2010).
- 23 Chambers CT, Reid GJ, McGrath PJ, Finley GA. Development and preliminary validation

- of a postoperative pain measure for parents. *Pain* 68(2), 307–313 (1996).
- 24 Taddio A, Appleton M, Bortolussi R *et al.* Reducing the pain of childhood vaccination: an evidence-based clinical practice guideline. *CMAJ* 182(18), e843–e855 (2010).
- 25 Dubois A, Bringuier S, Capdevilla X, Pry R. Vocal and verbal expression of postoperative pain in preschoolers. *Pain Manag. Nurs.* 9(4), 160–165 (2008).
- 26 Lawrence J, Alcock D, McGrawth P, Kay J, MacMurray S, Dulberg C. The development of a tool to assess neonatal pain. *Neonatal Netw.* 12(6), 59–66 (1993).
- 27 Manworren R, Hynan L. Clinical validation of FLACC: preverbal patient pain scale. *Pediatr. Nurs.* 29(2), 140–146 (2003).
- 28 Bieri D, Reeve RA, Champion GD, Addicoat L, Ziegler JB. The Faces Pain Scale for the self-assessment of the severity of pain experienced by children: development, initial validation, and preliminary investigation for ratio scale properties. *Pain* 41(2), 139–150 (1990).
- 29 Hockenberry MJ, Wilson D. *Wong's Essentials of Pediatric Nursing (8th Edition)*. Mosby, St Louis, MO, USA (2009).
- 30 Todd KH, Funk KG, Funk JP, Bonacci R. Clinical significance of reported changes in pain severity. *Ann. Emerg. Med.* 27(4), 485–489 (1996).
- 31 Nader R, Oberlander TF, Chambers CT, Craig KD. Expression of pain in children with autism. *Clin. J. Pain* 20(2), 88–97 (2004).
- 32 Hadden KL, Von Baeyer CL. Pain in children with cerebral palsy: common triggers and expressive behaviors. *Pain* 99(1), 281–288 (2002).
- 33 McGrath PJ, Romus C, Canfield C, Campbell MA, Hennigar A. Behaviours caregivers use to determine pain in non-verbal, cognitively impaired individuals. *Dev. Med. Child Neurol.* 40(5), 340–343 (1998).
- 34 Bennett-Branson SM, Craig KD. Postoperative pain in children: developmental and family influences on spontaneous coping strategies. *Can. J. Behav. Sci.* 25(3), 355–383 (1993).
- 35 Young KD. Pediatric procedural pain. *Ann. Emerg. Med.* 45(2), 160–171 (2005).
- 36 Kain ZN, Linda MC, O'Conner TZ, Cicchetti DV. Preoperative anxiety in children. *Arch. Pediatr. Adolesc. Med.* 150(12), 1238 (1996).
- 37 Cintron A, Morrison RS. Pain and ethnicity in the United States: a systematic review. *J. Palliat. Med.* 9(6), 1454–1473 (2006).
- 38 Anderson KO, Green CR, Payne R. Racial and ethnic disparities in pain: causes and consequences of unequal care. *J. Pain* 10(12), 1187–1204 (2009).
- Examines the relevant literature surrounding racial disparities in child pain and finds that such disparities do exist.
- 39 Sadhasivam S, Chidambaran V, Ngamprasertwong P *et al.* Race and unequal burden of perioperative pain and opioid related adverse effects in children. *Pediatrics* 129(5), 832–838 (2012).
- 40 Batista ML, Fortier MA, Maurer EL, Tan E, Huszti HC, Kain ZN. Exploring the impact of cultural background on parental perceptions of children's pain. *Child. Health Care* 41(2), 97–110 (2012).
- 41 Hsu Y, Mao C, Wey M. Antihypertensive medication adherence among elderly Chinese Americans. *J. Transcult. Nurs.* 21(4), 297–305 (2010).
- 42 Birdee GS, Phillips RS, Davis RB, Gardiner P. Factors associated with pediatric use of complementary and alternative medicine. *Pediatrics* 125(2), 249–256 (2010).
- 43 Fortier MA, Gillis S, Gomez SH, Wang SM, Tan ET, Kain ZN. Attitudes toward and use of complementary and alternative medicine among Hispanic and white mothers. *Altern. Ther. Health Med.* 20(1), 13–19 (2014).
- 44 Zisk RY, Grey M, MacLaren JE, Kain ZN. Exploring sociodemographic and personality characteristic predictors of parental pain perceptions. *Anesth. Analg.* 104(4), 790–798 (2007).
- 45 Rimer BK, Lyna CMP, Glassman B, Yarnall K, Lipkus I, Barber L. The impact of tailored interventions on a community health center population. *Pat. Ed. Couns.* 37(2), 125–140 (1999).
- 46 American Society of Anesthesiologists Task Force on Acute Pain Management. Practice guidelines for acute pain management in the perioperative setting: an updated report. *Anesthesiology* 100(6), 1573–1581 (2004).