

Lawrence Berkeley National Laboratory

Recent Work

Title

DOES THE f_0 PARTICLE LIE ON THE POMERANCHUK TRAJECTORY?

Permalink

<https://escholarship.org/uc/item/6674q70v>

Author

Pignotti, Alberto.

Publication Date

1963-01-15

University of California
Ernest O. Lawrence
Radiation Laboratory

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

DOES THE f^0 PARTICLE LIE ON THE
POMERANCHUK TRAJECTORY?

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review Letters

UCRL-10600 Rev.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

DOES THE f^0 PARTICLE LIE ON THE POMERANCHUK TRAJECTORY?

Alberto Pignotti

January 15, 1963

fit into the Regge-pole scheme as possibly belonging to the trajectory proposed by Igi.⁵

We assume that a boson Regge trajectory as a function of the energy-squared satisfies the dispersion relation⁶

$$\alpha(t) = \alpha(\infty) + \frac{1}{\pi} \int_{t_0}^{\infty} \frac{\text{Im } \alpha(t')}{t' - t} dt' \quad \text{for } t_0 > 0, \quad (\text{H1})$$

with

$$\text{Im } \alpha(t) \geq 0 \quad \text{for } t_0 \leq t < \infty. \quad (\text{H2})$$

As a consequence, all the derivatives of $\alpha(t)$ for $t < t_0$ are positive, and this implies a severe restriction on the behavior of α in that region. In particular, we expect to be able to put a lower bound to the slope of α at the origin, if $\alpha(\infty)$, $\alpha(0)$, and $\alpha(t_1)$ for any negative value of t_1 are given. In some instances, this bound will be twice the slope of a straight line through $\alpha(t_1)$ and $\alpha(0)$. In order to derive this and other related results, it will be useful to prove the following lemma: Let $\xi(t)$ satisfy the dispersion relation

$$\xi(t) = \frac{1}{\pi} \int_a^{\infty} \frac{\text{Im } \xi(t')}{t' - t} dt', \quad (1)$$

with $\text{Im } \xi(t) \geq 0$ for $a \leq t < \infty$. It follows that

$$\xi(t_3) \geq \frac{\xi(t_1) \xi(t_2) (t_2 - t_1)}{(t_2 - t_3) \xi(t_2) + (t_3 - t_1) \xi(t_1)}, \quad (2)$$

for $t_1 < t_2$, t_1 , t_2 , and $t_3 \leq a$, where the \geq sign holds for $t_3 \leq t_1$ or $t_3 \geq t_2$, and the \leq sign for $t_1 \leq t_3 \leq t_2$. Proof: From Eq. (1) we have

-3-

$$\operatorname{Im} \xi(t) > 0 \quad \text{for} \quad \operatorname{Im} t > 0 ,$$

and

$$\left. \begin{array}{l} \operatorname{Re} \xi(t) > 0 \\ \operatorname{Im} \xi(t) = 0 \end{array} \right\} \quad \text{for} \quad t \text{ real and } -\infty < t < a .$$

Therefore the function

$$\eta(t) = -1/\xi(t) \quad (3)$$

has no singularities except for a right-hand cut and a pole at infinity, and can be represented as⁷

$$\eta(t) = C + At + \int_a^{\infty} \frac{1 + t't}{t' - t} \phi(t') dt' \quad (4)$$

with C , A , and $\phi(t)$ real, and $\phi(t) \geq 0$. From Eq. (4) it follows for t real and $-\infty < t < a$ that

$$\eta''(t) \geq 0 . \quad (5)$$

Therefore, because $\eta(t)$ is a concave function for $t < a$, we have

$$\eta(t_3) \geq \eta(t_2) \frac{t_3 - t_1}{t_2 - t_1} + \eta(t_1) \frac{t_3 - t_2}{t_1 - t_2} \quad (6)$$

for $t_1 < t_2$, t_1 , t_2 and $t_3 \leq a$. Here again the \geq sign holds for $t_3 \leq t_1$ or $t_3 \geq t_2$, and the \leq sign for $t_1 \leq t_3 \leq t_2$. Equation (2) follows trivially from Eqs. (6) and (3). This completes the lemma.

-4-

We now apply the above lemma to the function $\alpha(t) - \alpha(\infty)$. In particular, if we let $t_1 \rightarrow t_2 = 0$ and $t_3 = t$ in Eq. (2), we obtain for $-\infty < t < 0$

$$\alpha(t) > \alpha(\infty) + \frac{[\alpha(0) - \alpha(\infty)]^2}{\alpha(0) - \alpha(\infty) + \alpha'(0) |t|} = L(t) \quad (7)$$

or

$$\alpha'(0) > \frac{1}{|t|} \left\{ \frac{[\alpha(0) - \alpha(\infty)]^2}{\alpha(t) - \alpha(\infty)} - \alpha(0) + \alpha(\infty) \right\} . \quad (8)$$

In the case of the Pomeranchuk trajectory, relations (7) or (8) plus the values of $\alpha(t)$ obtained in the high-energy experiments provide a lower bound for $\alpha'(0)$.⁸

Our aim is now to study what restrictions on the energy and width of a resonance follow from a given value of $\alpha'(0)$. The idea is that the trajectory, which has a positive curvature up to threshold, has to stay above its tangent line at $t = 0$ until its imaginary part has become appreciably large, and that when this happens, the Regge pole is so far from physical values of l that it is likely to give rise either to a very broad resonance or to no resonance at all. The detailed argument is as follows: We want to find under what restrictions it is possible to have a resonance at a value t_R of the energy-squared such that

$$\text{Re } \alpha(t_R) < \alpha(0) + t_R \alpha'(0) , \quad (9)$$

i.e.,

$$t_R > [\text{Re } \alpha(t_R) - \alpha(0)]/\alpha'(0) . \quad (10)$$

We know that $\text{Re } \alpha(t_R)$ has to be equal to the spin of the resonance in question, but that this condition is not sufficient to have a resonance. In addition, our

-5-

experience with the nonrelativistic case shows that at $t = t_R$, $\text{Re } \alpha$ must still be increasing steeply as a function of t , and $\text{Im } \alpha$ must still be small. The width of the resonance in the energy variable is related to these two properties and is given approximately by the function⁶

$$\Gamma(t_R) = \frac{1}{\sqrt{t}} \left. \frac{\text{Im } \alpha(t)}{\frac{d}{dt} \text{Re } \alpha(t)} \right|_{t=t_R} \quad (11)$$

We require that the trajectory considered present these two characteristics at $t = t_R$ by assuming

$$\text{Re } \alpha(t) \text{ has at most one inflection point for } t < t_R \quad (H3)$$

$$\text{Im } \alpha(t) \leq \text{Im } \alpha(t_R) \frac{t - t_0}{t_R - t_0} \quad \text{for } t_0 < t < t_R \quad (H4)$$

From assumption (H3) and the fact that $\alpha''(t) > 0$ for $t \leq t_0$ it follows that

$$\left. \frac{d}{dt} \text{Re } \alpha(t) \right|_{t=t_R} < [\text{Re } \alpha(t_R) - \alpha(0)]/t_R \quad (12)$$

for the values of t_R satisfying relation (10). Our next step is to put a lower bound to $\text{Im } \alpha(t_R)$ such that, combined with Eqs. (11) and (12), it will provide a restriction on the permissible values for $\Gamma(t_R)$. It is clear that such a bound for $\text{Im } \alpha(t_R)$ exists and is larger than zero, because if we had $\text{Im } \alpha(t_R) = 0$, from assumptions (H1), (H2), and (H4) all the derivatives of $\text{Re } \alpha$ would be positive up to $t = t_R$, and therefore $\text{Re } \alpha$ would not satisfy relation (9). This suggests splitting α into two parts, β and γ , one of which has a zero imaginary part for $t < t_R$. Therefore, we define

-6-

$$\left. \begin{aligned} \operatorname{Im} \beta(t) &= \operatorname{Im} \alpha(t) && \text{for } t_0 < t < t_R \\ &= \operatorname{Im} \alpha(2t_R - t) && \text{for } t_R < t < 2t_R - t_0 \end{aligned} \right\} \quad (13a)$$

$$\beta(t) = \frac{1}{\pi} \int_{t_0}^{2t_R - t_0} \frac{\operatorname{Im} \beta(t')}{t' - t} dt' , \quad (13b)$$

and

$$\gamma(t) = \alpha(t) - \beta(t) . \quad (14)$$

From the above definitions it follows that $\beta(\infty) = 0$, $\beta(0) > 0$, $\beta(t_R) = 0$, $\gamma(\infty) = \alpha(\infty)$, $\gamma(0) < \alpha(0)$, $\gamma(t_R) = \operatorname{Re} \alpha(t_R)$. In order to have $\operatorname{Im} \gamma(t) \geq 0$ on the real axis we assume further¹⁰

$$\operatorname{Im} \alpha(t) \geq \operatorname{Im} \alpha(2t_R - t) \quad \text{for } t_R < t < 2t_R - t_0 , \quad (H5)$$

which is consistent with the properties discussed above. Thus we can apply our lemma to the function $\gamma(t) - \gamma(\infty)$. We let $t_1 \rightarrow t_2 = 0$ and $t_3 = t \leq a = t_R$ in Eq. (2) and obtain

$$\gamma(t) - \gamma(\infty) \geq \frac{[\gamma(0) - \gamma(\infty)]^2}{\gamma(0) - \gamma(\infty) - t\gamma'(0)} .$$

The denominator in this expression is positive because it is so at $t = 0$ and $\gamma(t)$ is finite for $t \leq t_R$. We can therefore write for $t = t_R$

$$\gamma'(0) \leq \frac{[\gamma(0) - \gamma(\infty)][\gamma(t_R) - \gamma(0)]}{t_R[\gamma(t_R) - \gamma(\infty)]} < \frac{[\alpha(0) - \alpha(\infty)][\operatorname{Re} \alpha(t_R) - \gamma(0)]}{t_R[\operatorname{Re} \alpha(t_R) - \alpha(\infty)]} . \quad (15)$$

-7-

Now, from assumption (H4) we have

$$\beta(0) \leq B_0(t_R) \operatorname{Im} \alpha(t_R)$$

and

$$\beta'(0) \leq B_1(t_R) \operatorname{Im} \alpha(t_R) ,$$

where

$$B_0(t_R) = \frac{1}{\pi(t_R - t_0)} \left[-t_0 \ln \left(\frac{2t_R - t_0}{t_0} \right) + 2t_R \ln \left(\frac{2t_R - t_0}{t_R} \right) \right]$$

and

$$B_1(t_R) = \frac{1}{\pi(t_R - t_0)} \ln \left[\frac{t_R^2}{t_0(2t_R - t_0)} \right] .$$

It then follows that

$$\left. \begin{aligned} \gamma(0) &\geq \alpha(0) - \operatorname{Im} \alpha(t_R) B_0(t_R) \\ \text{and} \\ \gamma'(0) &\geq \alpha'(0) - \operatorname{Im} \alpha(t_R) B_1(t_R) \end{aligned} \right\} . \quad (16)$$

From relations (15) and (16) we obtain a lower bound for $\operatorname{Im} \alpha(t_R)$. This result combined with relations (11) and (12) gives the final bound

$$\Gamma(t_R) > \frac{\sqrt{t_R} \left\{ \alpha(\infty) - \alpha(0) + \alpha'(0)t_R [\operatorname{Re} \alpha(t_R) - \alpha(\infty)] / [\operatorname{Re} \alpha(t_R) - \alpha(0)] \right\}}{[\alpha(0) - \alpha(\infty)] B_0(t_R) + [\operatorname{Re} \alpha(t_R) - \alpha(\infty)] B_1(t_R) t_R} . \quad (17)$$

We want to apply the above results to the Pomeranchuk trajectory, which controls the high-energy behavior of total cross sections and therefore satisfies the condition^{3,11}

$$\alpha(0) = 1 . \quad (H6)$$

Submitted to Physical Review Letters

UCRL-10600 Rev.

UNIVERSITY OF CALIFORNIA
Lawrence Radiation Laboratory
Berkeley, California

Contract No. W-7405-eng-48

DOES THE f^0 PARTICLE LIE ON THE POMERANCHUK TRAJECTORY?

Alberto Pignotti

January 15, 1963

DOES THE f^0 PARTICLE LIE ON THE POMERANCHUK TRAJECTORY?*

Alberto Pignotti†

Lawrence Radiation Laboratory
University of California
Berkeley, California

January 15, 1963

Two experimental groups have recently reported on the discovery of a $\pi\pi$ resonance at about 1250 MeV that is probably an $I = 0$, $J = 2$ state.^{1,2} It was conjectured that this resonance, called f^0 , is the "particle" predicted by Chew and Frautschi on the basis of the Regge-pole scheme.³ In that case, it should lie on the Pomeranchuk trajectory. A. Ahmadzadeh and I. Sakmar have tried to test this conjecture, starting from a four-parameter expression for the imaginary part of the Pomeranchuk trajectory $\alpha(t)$.⁴ They assume the usual properties for this trajectory and conclude that the f^0 cannot lie on it, if at the same time one accepts the published analysis of the high-energy pp scattering in terms of the Pomeranchuk trajectory. The purpose of this letter is to prove that, as conjectured by Ahmadzadeh and Sakmar, this result is independent of the choice of an Ansatz for $\text{Im } \alpha(t)$, provided some current ideas on Regge trajectories are correct. In the first part, some general bounds derived for boson Regge trajectories are used to relate values of $\alpha(t)$ for negative t to the slope of the trajectory at $t = 0$. Then it is shown how this slope restricts the possible values for the energy and width of a resonance that lies on the same trajectory. When these results are applied to the Pomeranchuk particle, it is concluded that either the slope of the Pomeranchuk trajectory has been seriously overestimated from the pp data, or the Pomeranchuk particle is likely to have an energy less than or approximately equal to the ρ -meson energy. Experimental work directed towards testing the latter possibility is encouraged. In that case, the f^0 resonance still could

fit into the Regge-pole scheme as possibly belonging to the trajectory proposed by Igi.⁵

We assume that a boson Regge trajectory as a function of the energy-squared satisfies the dispersion relation⁶

$$\alpha(t) = \alpha(\infty) + \frac{1}{\pi} \int_{t_0}^{\infty} \frac{\text{Im } \alpha(t')}{t' - t} dt' \quad \text{for } t_0 > 0, \quad (\text{H1})$$

with

$$\text{Im } \alpha(t) \geq 0 \quad \text{for } t_0 \leq t < \infty. \quad (\text{H2})$$

As a consequence, all the derivatives of $\alpha(t)$ for $t < t_0$ are positive, and this implies a severe restriction on the behavior of α in that region. In particular, we expect to be able to put a lower bound to the slope of α at the origin, if $\alpha(\infty)$, $\alpha(0)$, and $\alpha(t_1)$ for any negative value of t_1 are given. In some instances, this bound will be twice the slope of a straight line through $\alpha(t_1)$ and $\alpha(0)$. In order to derive this and other related results, it will be useful to prove the following lemma: Let $\xi(t)$ satisfy the dispersion relation

$$\xi(t) = \frac{1}{\pi} \int_a^{\infty} \frac{\text{Im } \xi(t')}{t' - t} dt', \quad (1)$$

with $\text{Im } \xi(t) \geq 0$ for $a \leq t < \infty$. It follows that

$$\xi(t_3) \begin{cases} \geq \\ \leq \end{cases} \frac{\xi(t_1) \xi(t_2) (t_2 - t_1)}{(t_2 - t_3) \xi(t_2) + (t_3 - t_1) \xi(t_1)}, \quad (2)$$

for $t_1 < t_2$, t_1 , t_2 , and $t_3 \leq a$, where the \geq sign holds for $t_3 \leq t_1$ or $t_3 \geq t_2$, and the \leq sign for $t_1 \leq t_3 \leq t_2$. Proof: From Eq. (1) we have

-3-

$$\operatorname{Im} \xi(t) > 0 \quad \text{for} \quad \operatorname{Im} t > 0 ,$$

and

$$\left. \begin{array}{l} \operatorname{Re} \xi(t) > 0 \\ \operatorname{Im} \xi(t) = 0 \end{array} \right\} \quad \text{for} \quad t \text{ real and } -\infty < t < a .$$

Therefore the function

$$\eta(t) = -1/\xi(t) \quad (3)$$

has no singularities except for a right-hand cut and a pole at infinity, and can be represented as⁷

$$\eta(t) = C + At + \int_a^{\infty} \frac{1 + t't}{t' - t} \phi(t') dt' \quad (4)$$

with C , A , and $\phi(t)$ real, and $\phi(t) \geq 0$. From Eq. (4) it follows for t real and $-\infty < t < a$ that

$$\eta''(t) \geq 0 . \quad (5)$$

Therefore, because $\eta(t)$ is a concave function for $t < a$, we have

$$\eta(t_3) \geq \eta(t_2) \frac{t_3 - t_1}{t_2 - t_1} + \eta(t_1) \frac{t_3 - t_2}{t_1 - t_2} \quad (6)$$

for $t_1 < t_2$, t_1 , t_2 and $t_3 \leq a$. Here again the \geq sign holds for $t_3 \leq t_1$ or $t_3 \geq t_2$, and the \leq sign for $t_1 \leq t_3 \leq t_2$. Equation (2) follows trivially from Eqs. (6) and (3). This completes the lemma.

-4-

We now apply the above lemma to the function $\alpha(t) - \alpha(\infty)$. In particular, if we let $t_1 \rightarrow t_2 = 0$ and $t_3 = t$ in Eq. (2), we obtain for $-\infty < t < 0$

$$\alpha(t) > \alpha(\infty) + \frac{[\alpha(0) - \alpha(\infty)]^2}{\alpha(0) - \alpha(\infty) + \alpha'(0) |t|} = L(t) \quad (7)$$

or

$$\alpha'(0) > \frac{1}{|t|} \left\{ \frac{[\alpha(0) - \alpha(\infty)]^2}{\alpha(t) - \alpha(\infty)} - \alpha(0) + \alpha(\infty) \right\} . \quad (8)$$

In the case of the Pomeranchuk trajectory, relations (7) or (8) plus the values of $\alpha(t)$ obtained in the high-energy experiments provide a lower bound for $\alpha'(0)$.⁸

Our aim is now to study what restrictions on the energy and width of a resonance follow from a given value of $\alpha'(0)$. The idea is that the trajectory, which has a positive curvature up to threshold, has to stay above its tangent line at $t = 0$ until its imaginary part has become appreciably large, and that when this happens, the Regge pole is so far from physical values of ℓ that it is likely to give rise either to a very broad resonance or to no resonance at all. The detailed argument is as follows: We want to find under what restrictions it is possible to have a resonance at a value t_R of the energy-squared such that

$$\text{Re } \alpha(t_R) < \alpha(0) + t_R \alpha'(0) , \quad (9)$$

i.e.,

$$t_R > [\text{Re } \alpha(t_R) - \alpha(0)] / \alpha'(0) . \quad (10)$$

We know that $\text{Re } \alpha(t_R)$ has to be equal to the spin of the resonance in question, but that this condition is not sufficient to have a resonance. In addition, our

experience with the nonrelativistic case shows that at $t = t_R$, $\text{Re } \alpha$ must still be increasing steeply as a function of t , and $\text{Im } \alpha$ must still be small. The width of the resonance in the energy variable is related to these two properties and is given approximately by the function⁶

$$\Gamma(t_R) = \frac{1}{\sqrt{t}} \left. \frac{\text{Im } \alpha(t)}{\frac{d}{dt} \text{Re } \alpha(t)} \right|_{t=t_R} \quad (11)$$

We require that the trajectory considered present these two characteristics at $t = t_R$ by assuming

$$\text{Re } \alpha(t) \text{ has at most one inflection point for } t < t_R \quad (H3)$$

$$\text{Im } \alpha(t) \leq \text{Im } \alpha(t_R) \frac{t - t_0}{t_R - t_0} \quad \text{for } t_0 < t < t_R \quad (H4)$$

From assumption (H3) and the fact that $\alpha''(t) > 0$ for $t \leq t_0$ it follows that

$$\left. \frac{d}{dt} \text{Re } \alpha(t) \right|_{t=t_R} < [\text{Re } \alpha(t_R) - \alpha(0)]/t_R \quad (12)$$

for the values of t_R satisfying relation (10). Our next step is to put a lower bound to $\text{Im } \alpha(t_R)$ such that, combined with Eqs. (11) and (12), it will provide a restriction on the permissible values for $\Gamma(t_R)$. It is clear that such a bound for $\text{Im } \alpha(t_R)$ exists and is larger than zero, because if we had $\text{Im } \alpha(t_R) = 0$, from assumptions (H1), (H2), and (H4) all the derivatives of $\text{Re } \alpha$ would be positive up to $t = t_R$, and therefore $\text{Re } \alpha$ would not satisfy relation (9). This suggests splitting α into two parts, β and γ , one of which has a zero imaginary part for $t < t_R$. Therefore, we define

-6-

$$\left. \begin{aligned} \operatorname{Im} \beta(t) &= \operatorname{Im} \alpha(t) && \text{for } t_0 < t < t_R \\ &= \operatorname{Im} \alpha(2t_R - t) && \text{for } t_R < t < 2t_R - t_0 \end{aligned} \right\} \quad (13a)$$

$$\beta(t) = \frac{1}{\pi} \int_{t_0}^{2t_R - t_0} \frac{\operatorname{Im} \beta(t')}{t' - t} dt' , \quad (13b)$$

and

$$\gamma(t) = \alpha(t) - \beta(t) . \quad (14)$$

From the above definitions it follows that $\beta(\infty) = 0$, $\beta(0) > 0$, $\beta(t_R) = 0$, $\gamma(\infty) = \alpha(\infty)$, $\gamma(0) < \alpha(0)$, $\gamma(t_R) = \operatorname{Re} \alpha(t_R)$. In order to have $\operatorname{Im} \gamma(t) \geq 0$ on the real axis we assume further¹⁰

$$\operatorname{Im} \alpha(t) \geq \operatorname{Im} \alpha(2t_R - t) \quad \text{for } t_R < t < 2t_R - t_0 , \quad (H5)$$

which is consistent with the properties discussed above. Thus we can apply our lemma to the function $\gamma(t) - \gamma(\infty)$. We let $t_1 \rightarrow t_2 = 0$ and $t_3 = t \leq a = t_R$ in Eq. (2) and obtain

$$\gamma(t) - \gamma(\infty) \geq \frac{[\gamma(0) - \gamma(\infty)]^2}{\gamma(0) - \gamma(\infty) - t\gamma'(0)} .$$

The denominator in this expression is positive because it is so at $t = 0$ and $\gamma(t)$ is finite for $t \leq t_R$. We can therefore write for $t = t_R$

$$\gamma'(0) \leq \frac{[\gamma(0) - \gamma(\infty)][\gamma(t_R) - \gamma(0)]}{t_R[\gamma(t_R) - \gamma(\infty)]} < \frac{[\alpha(0) - \alpha(\infty)][\operatorname{Re} \alpha(t_R) - \gamma(0)]}{t_R[\operatorname{Re} \alpha(t_R) - \alpha(\infty)]} .$$

(15)

-7-

Now, from assumption (H4) we have

$$\beta(0) \leq B_0(t_R) \operatorname{Im} \alpha(t_R)$$

and

$$\beta'(0) \leq B_1(t_R) \operatorname{Im} \alpha(t_R),$$

where

$$B_0(t_R) = \frac{1}{\pi(t_R - t_0)} \left[-t_0 \ln \left(\frac{2t_R - t_0}{t_0} \right) + 2t_R \ln \left(\frac{2t_R - t_0}{t_R} \right) \right]$$

and

$$B_1(t_R) = \frac{1}{\pi(t_R - t_0)} \ln \left[\frac{t_R^2}{t_0(2t_R - t_0)} \right].$$

It then follows that

$$\left. \begin{aligned} \gamma(0) &\geq \alpha(0) - \operatorname{Im} \alpha(t_R) B_0(t_R) \\ \text{and} \\ \gamma'(0) &\geq \alpha'(0) - \operatorname{Im} \alpha(t_R) B_1(t_R) \end{aligned} \right\} \quad (16)$$

From relations (15) and (16) we obtain a lower bound for $\operatorname{Im} \alpha(t_R)$. This result combined with relations (11) and (12) gives the final bound

$$\Gamma(t_R) > \frac{\sqrt{t_R} \left\{ \alpha(\infty) - \alpha(0) + \alpha'(0) t_R [\operatorname{Re} \alpha(t_R) - \alpha(\infty)] / [\operatorname{Re} \alpha(t_R) - \alpha(0)] \right\}}{[\alpha(0) - \alpha(\infty)] B_0(t_R) + [\operatorname{Re} \alpha(t_R) - \alpha(\infty)] B_1(t_R) t_R} \quad (17)$$

We want to apply the above results to the Pomeranchuk trajectory, which controls the high-energy behavior of total cross sections and therefore satisfies the condition^{3,11}

$$\alpha(0) = 1. \quad (\text{H6})$$

We also take

$$\alpha(\infty) \geq -1 \quad (\text{H7})$$

as suggested by the work of Gribov and Pomeranchuk.¹² It can be easily verified that the bounds given by relations (7), (8), and (17) are increasing functions of $\alpha(\infty)$. Therefore we replace in these relations $\alpha(\infty) = -1$ and our results are valid a fortiori if $\alpha(\infty) > -1$. Finally we put $t_0 = 4m_\pi^2$ and $\text{Re } \alpha(t_R) = 2$ as is the case for the Pomeranchuk particle. We then obtain from Eq. (17)

$$\Gamma(t_R) > \frac{\sqrt{t_R} [3t_R\alpha'(0) - 2]}{2B_0(t_R) + 3t_R B_1(t_R)} = \Phi(t_R) \quad \text{for} \quad t_R > 1/\alpha'(0). \quad (\text{18})$$

In Fig. 1 the function $L(t)$ from relations (7), (H6), and (H7) is plotted for $\alpha'(0) = 1/80$, $1/50$, and $1/35$. The values of $\alpha(t)$ obtained by the Brookhaven and Cornell groups from the analysis of the high-energy data are also indicated.^{13,14} If we apply the inequality (8) to the three values for lower $|t|$, we obtain $\alpha'(0) > 1/29$ with 90% confidence. Eventually one may expect this bound to be lowered by the presence of other than statistical errors in these data. However, the values of $\Phi(t_R)$ obtained from Eq. (18) and plotted in Fig. 2 show that unless $\alpha'(0)$ is less than $1/80$, then $\Gamma(80)$ has to be much larger than the experimental width of the f^0 resonance. Therefore, within the scheme presented here, and provided the analysis of the high-energy pp data is accepted, this resonance cannot lie on the Pomeranchuk trajectory. The Pomeranchuk particle might be found at a value of t smaller than or approximately equal to $30m_\pi^2$.^{15,16} Alternatively one may conclude that other l -plane singularities for pp scattering (e.g., the P' and ω Regge poles) have conspired to simulate a falsely large slope for the Pomeranchuk trajectory.

-9-

I am indebted to Professors Geoffrey F. Chew, Gyo Takeda, and Charles Zemach for encouragement and many helpful discussions and Messrs. A. Ahmadzadeh and I. Sakmar for day-to-day communication of their results. I also want to thank Dr. David L. Judd for his kind hospitality at the Lawrence Radiation Laboratory.

FOOTNOTES AND REFERENCES

- * Word done under the auspices of the U. S. Atomic Energy Commission during a fellowship sponsored by the Consejo Nacional de Investigaciones Científicas y Técnicas of Argentina. This does not imply that this institution either approves or assumes any liabilities for the information contained in this report.
- † On leave from Facultad de Ciencias Exactas y Naturales, Buenos Aires, Argentina.
1. W. Selove, V. Hagopian, H. Brody, A. Baker, and E. Leboy, Phys. Rev. Letters 9, 272 (1962).
 2. J. J. Veillet, J. Hennessy, H. Bingham, M. Block, D. Dryard, A. Lagarrigue, P. Mittner, and A. Rousset, Ecole Polytechnique, Paris, and G. Bellini, M. di Corato, E. Fiorini, and P. Negri, Istituto Nazionale de Fisica Nucleare, Milano, preprint (n. d.),.
 3. G. F. Chew and S. Frautschi, Phys. Rev. Letters 7, 394 (1961). We will call this particle the Pomeranchuk particle.
 4. A. Ahmadzadeh and I. Sakmar, Lawrence Radiation Laboratory Report UCRL-10584, December 1962.
 5. K. Igi, Phys. Rev. Letters 9, 76 (1962). We call P' the trajectory proposed in this paper.
 6. This and other properties are discussed by G. F. Chew in "The Self-Consistent S-Matrix with the Regge Asymptotic Behaviour," CERN preprint TH/292, August 16, 1962, to be published in the Physical Review; further references are given here.
 7. J. A. Shohat and J. D. Tamarkin, The Problem of Moments, (American Mathematical Society, New York, N. Y., 1943) p. 23.

-11-

8. We will present here some other results that follow from our lemma applied to a boson Regge trajectory. We call t_g the value of t at which α vanishes. In Eq. (2) let $t_1 = t_g$, $t_2 = 0$, and $t_3 = t$; then for $t_g < t < 0$ we have

$$\alpha(t) < \alpha(\infty) + \frac{t_g \alpha(\infty) [\alpha(0) - \alpha(\infty)]}{-t\alpha(0) + t_g \alpha(\infty)} . \quad (F1)$$

Let $t_1 \rightarrow t_2 = t$, $t_3 = 0$; it follows that, for $-\infty < t < 0$,

$$\alpha'(t) < [\alpha(t) - \alpha(\infty)][\alpha(t) - \alpha(0)]/t[\alpha(0) - \alpha(\infty)] .$$

Finally, from relations (3) and (5) we obtain for $t < t_0$

$$\alpha''(t) > \frac{2[\alpha'(t)]^2}{\alpha(t) - \alpha(\infty)} .$$


9. This last hypothesis is generous near threshold for the Pomeranchuk trajectory because there we know $\text{Im } \alpha(t) \sim (t - t_0)^{\text{Re } \alpha(t_0) + 1/2}$ and $\text{Re } \alpha(t_0) > 1$ [see A. O. Barut and D. E. Zwanziger, Phys. Rev. 127, 974 (1962)]. Moreover, the maximum for $\text{Re } \alpha(t)$ is expected to occur at approximately the same energy as the inflection point for $\text{Im } \alpha(t)$. Therefore, as $\text{Re } \alpha(t)$ has not yet reached its maximum at the resonance energy, $\text{Im } \alpha(t_R)$ is presumably still concave, which is stronger than required by the linear assumption (H4). Obviously this hypothesis has to be modified for trajectories such that $\text{Re } \alpha(t_0) < 0.5$.
10. A weaker bound may be chosen without modifying the essence of our results. However such a change is hardly expected to be necessary, and it would make our proof more complicated.
11. G. F. Chew and S. C. Frautschi, Phys. Rev. Letters 8, 41 (1962).

12. V. N. Gribov and I. Ya. Pomeranchuk, in 1962 Annual International Conference on High Energy Physics at CERN (CERN Scientific Information Service, Geneva, 1962).
13. W. F. Baker, E. W. Jenkins, A. L. Read, G. Cocconi, V. T. Cocconi, and J. Orear, Phys. Rev. Letters 9, 221 (1962).
14. The reader is referred also to the data of A. N. Diddens, E. Lillethun, G. Manning, A. E. Taylor, T. G. Walker, and A. M. Wetherell, Phys. Rev. Letters 9, 111 (1962). If one considers all their data, the result $\alpha(-70) < 0$ is definitely established. However, their results for $t > -50$ are too linear and therefore not consistent with Eq. (F1) in footnote 8, and the values of $\alpha(\infty)$ and $\alpha(0)$ given by relations (H6) and (H7). If one takes only their data for $(s/2M^2) > 10.5$, their results agree with those of Baker et al.,¹³ although with larger errors.
15. Both groups reporting on the f^0 have observed some anomalies in the angular distribution around the ρ peak that can be attributed to an $I = 0$ state because they were not present in the analogous experiment with $\pi^+\pi^0$ or $\pi^-\pi^0$. On the other hand, Dr. Takeda has pointed out to me that if the Pomeranchuk particle has about the same energy as the ρ meson, its width is likely to be less than the experimental resolution, on the basis of the measured cross sections for ρ^+ and ρ^0 .
- A. Ahmadzadeh and I. Sakmar have also suggested the possible connection of the Pomeranchuk particle with other experimental information for $t < 30m_\pi^2$.
16. The possibility of the Pomeranchuk trajectory "bending down" soon and never reaching 2 cannot be excluded. However, this is not likely to occur if the quantum numbers of the f^0 have been assigned correctly, which would imply that another trajectory with the same quantum numbers as the Pomeranchuk, but below it at $t = 0$, is able to reach $\text{Re } \alpha = 2$.

FIGURE LEGENDS


Fig. 1. Curves a, b, and c are lower bounds for $\alpha(t)$ given by Eqs. (7), (H6), and (H7) for $\alpha'(0) \leq 1/80$, $1/50$, and $1/35$, respectively. The experimental data are from Ref. 13.

Fig. 2. Curves a, b, and c show lower bounds for $\Gamma(t_R)$ given by Eq. (18) for $\alpha'(0) \geq 1/80$, $1/50$, and $1/35$, respectively. The experimental values for the energy squared and width of the f^0 are indicated (Refs. 1 and 2).



MU-28927

Fig. 1



MU-28928-A

Fig. 2

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

