

UC San Diego

UC San Diego Previously Published Works

Title

Targeting G1/S phase cell-cycle genomic alterations and accompanying co-alterations with individualized CDK4/6 inhibitor–based regimens

Permalink

<https://escholarship.org/uc/item/66p2z8d2>

Journal

JCI Insight, 6(1)

ISSN

2379-3708

Authors

Kato, Shumei

Okamura, Ryosuke

Adashek, Jacob J

et al.

Publication Date

2021-01-11

DOI

10.1172/jci.insight.142547

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Targeting G₁/S phase cell-cycle genomic alterations and accompanying co-alterations with individualized CDK4/6 inhibitor-based regimens

Shumei Kato,¹ Ryosuke Okamura,¹ Jacob J. Adashek,² Noor Khalid,¹ Suzanna Lee,¹ Van Nguyen,¹ Jason K. Sicklick,³ and Razelle Kurzrock¹

¹Center for Personalized Cancer Therapy and Division of Hematology and Oncology, Department of Medicine, Moores Cancer Center at UC San Diego Health, La Jolla, California, USA. ²Department of Internal Medicine, University of South Florida, H. Lee Moffitt Cancer Center & Research Institute, Tampa, Florida, USA. ³Center for Personalized Cancer Therapy and Division of Surgical Oncology, Department of Surgery, Moores Cancer Center at UC San Diego Health, La Jolla, California, USA.

BACKGROUND. Although CDK4/6 inhibitors are an established treatment for hormone receptor-positive, HER2-negative metastatic breast cancers, their benefit in other malignancies remains limited.

METHODS. We investigated factors associated with clinical outcomes from CDK4/6 inhibitor-based therapy among patients with G₁/S phase cell-cycle alterations (CDK4/6 amplifications, CCND1/2/3 amplifications, or CDKN2A/B alterations).

RESULTS. Overall, 2457 patients with diverse solid tumors that underwent clinical-grade, next-generation sequencing (182–465 genes) and therapy outcome of (non-breast cancer) patients treated with matched CDK4/6 inhibitors were analyzed. G₁/S phase cell-cycle alterations occurred in 20.6% (507 of 2457) of patients; 99% of those patients (*n* = 501) harbored ≥1 characterized co-alteration (median, 4; range, 0–24). In 40 patients with G₁/S phase cell-cycle alterations given CDK4/6 inhibitors as part of their regimen, significantly longer median progression-free survival (PFS) was observed when CDK4/6 inhibitor-based therapies matched a larger proportion of tumor alterations, often because CDK4/6 inhibitors were administered together with other drugs that were matched to genomic co-alterations, hence achieving a high matching score (high vs. low [≥50% vs. <50%] matching score, PFS, 6.2 vs. 2.0 months, *P* < 0.001 [*n* = 40] [multivariate]) and higher rate of stable disease ≥6 months or an objective response (57% vs. 21%, *P* = 0.048).

CONCLUSION. In summary, in cell-cycle-altered cancers, matched CDK4/6 inhibitors, as part of an individualized regimen targeting a majority of genomic alterations, was independently associated with longer PFS.

TRIAL REGISTRATION. ClinicalTrials.gov NCT02478931.

FUNDING. Joan and Irwin Jacobs Fund, National Cancer Institute (P30 CA023100, R01 CA226803), and the FDA (R01 FD006334).

Conflict of interest: SK serves as a consultant for Foundation Medicine; and receives speakers' fees from Roche. JKS receives research funding from Novartis Pharmaceuticals, Amgen Pharmaceuticals, and Foundation Medicine; consultant fees from Grand Rounds, Loxo, and Deciphera; and speakers' fees from Roche. RK receives research funding from Incyte, Genentech, Merck Serono, Pfizer, Sequenom, Foundation Medicine, Guardant Health, Grifols, and Konica Minolta as well as consultant fees from Loxo, X-Biotech, Actuate Therapeutics, Genentech, and NeoMed. She receives speakers' fees from Roche and has an equity interest in IDbyDNA and CureMatch Inc.

Copyright: © 2021, Kato et al. This is an open access article published under the terms of the Creative Commons Attribution 4.0 International License.

Submitted: July 22, 2020

Accepted: November 18, 2020

Published: January 11, 2021

Reference information: *JCI Insight*. 2021;6(1):e142547.
<https://doi.org/10.1172/jci.insight.142547>.

Introduction

The cell cycle is tightly regulated by various checkpoints, which are populated by signaling molecules that need to be at threshold levels and appropriately phosphorylated by various kinases (1). Certain cancers have the ability to hijack various players within these cycle cascades, triggering uncontrolled growth and cell division (2). Specifically, cyclin-dependent kinases 4 and 6 (CDK4/6) require cyclin D1 to function and phosphorylate the retinoblastoma (RB) protein, which inactivates this tumor suppressor, allowing for the progression of cell cycle during the G₁/S phase. Without cyclin D1 or appropriately functioning CDK4/6 enzymes, the cells will not adequately

divide and proliferate. This cell-cycle feedback loop permits regulated growth and assures that cells only divide when necessary. However, several genomic alterations inappropriately fuel cell-cycle progression. Well-defined cyclin-related mechanisms that drive tumorigenesis include increases in expression/amplification of *CDK4/6*, upregulation of cyclin D, or deleterious alterations in the *Rb*, *CCNE1*, *CDKN2A*, or *CDKN2B* genes (3–5). The aforementioned G_1/S phase cell-cycle modulator alterations exist in anywhere from 9.5% to 73.8% of a variety of tumor types, making this pathway an attractive therapeutic target (6).

There are currently 3 CDK4/6 inhibitors — palbociclib, ribociclib, and abemaciclib — that are FDA approved for the treatment of hormone receptor–positive (HR-positive), HER2-negative metastatic breast cancer in conjunction with an aromatase inhibitor (7–10). While these agents result in improved progression-free survival (PFS) and overall survival (OS) in this patient population, there remains no clear biomarker that predicts response to CDK4/6 inhibitors (11–13). Theoretically, amplification of *CDK4/6* and cyclin D1, D2, and/or D3 (*CCND1/2/3*) and alterations in *CDKN2A/B* are putative markers to predict the response from CDK4/6 inhibitors; however, there have been mixed reports in this regard. For instance, in the American Society of Clinical Oncology's TAPUR study, participants with *CDKN2A* alterations (expected to increase CDK4/6 expression) or *CDK4* amplifications were assigned to receive palbociclib. Patients with head and neck cancer, soft tissue sarcoma, and bronchus/lung cancers did demonstrate benefit and continued on to the second portion of the trial as part of Simon's optimal 2-stage design (14). However, patients with pancreatic and gallbladder cancers with *CDKN2A* alterations did not derive significant benefit from CDK4/6 inhibition. The discrepancy in findings between tumor histologies confounds the ability to identify a biomarker of responsiveness. Furthermore, no cassette of markers has proved important in patients with breast cancer treated with CDK4/6 inhibitors (15). It is therefore still unclear, despite the pharmacologically driven properties of these agents supporting their effect on the G_1/S phase cell-cycle pathway, how to best ascertain in advance if there is a subset of non–breast cancer patients who may respond to CDK4/6 inhibitors.

One hypothesis for why certain G_1/S phase cell-cycle–associated genes have not been reliable markers to predict sensitivity to CDK4/6 inhibitors (11, 16) relates to the frequent finding of important genomic co-alterations (2). On average, patients with metastatic cancer have approximately 2–5 deleterious genomic alterations when assessed with a fixed panel derived from next-generation sequencing (NGS) (17–19). Although targeting the cell-cycle pathway may be appealing, it may also be less rewarding than anticipated due to this phenomenon. Indeed, although certain drivers, such as *EGFR* or *ALK* or *NTRK* aberrations, may be effectively targeted by matched monotherapy, not all patients respond and resistance often develops (20–23). It is plausible, therefore, that, even in these cases, primary or secondary resistance could be driven by co-alterations or driver feedback loops. For instance, in colorectal cancer with *BRAF* mutations, *BRAF* inhibitors alone are ineffective. Meanwhile, the *BRAF* inhibitor encorafenib, together with the *EGFR* antibody cetuximab, targets both *BRAF* and the feedback *EGFR* driver pathway; this efficacious combination was recently approved by the FDA (24). Indeed, targeting one specific signal in a complicated network of genomic drivers may be ineffective (25), and recent studies demonstrate that the greater the proportion of signals targeted, the better the outcome (26–28).

Herein, we used NGS to interrogate the complex genomic landscape of 2457 patients with diverse cancers, of whom 507 patients harbored specific, potentially sensitizing G_1/S phase cell-cycle (*CDK4/6*, *CCND1/2/3*, or *CDKN2A/B*) gene alterations. In a subset of patients with cancer (not breast cancer) with sensitizing cell-cycle gene alterations treated with CDK4/6 inhibitors, we show examples of responders. Additionally, we show that, overall, there was a significantly improved PFS and higher rate of stable disease ≥ 6 months or having a response observed when a majority of genomic alterations/co-alterations were targeted, as compared with administration of matched CDK4/6 inhibitors alone in the face of complex molecular portfolios.

Results

*Characteristics of patients with alterations in the potentially sensitizing G_1/S phase cell-cycle signaling pathway (*CDK4/6* amplifications, *CCND1/2/3* amplifications, or *CDKN2A/B* alterations).* Among 2457 patients with diverse solid tumors, 507 patients (21%) had at least 1 characterized genomic alteration in sensitizing G_1/S phase cell-cycle signaling genes — *CDK4/6*, *CCND1/2/3*, or *CDKN2A/B* — on tissue NGS (Figure 1). Among the 507 patients, the most common diagnosis was brain tumors (16%, $n = 83$), non–small cell lung cancers (15%, $n = 77$), and skin cancers, including melanoma (13%, $n = 67$). Among the G_1/S phase cell-cycle alterations of interest, *CDKN2A/B* alterations (71%, $n = 359$) were the most commonly observed in this series, followed by *CCND1* amplification (15%, $n = 75$) and *CDK4* amplification (12%, $n = 61$) (Table 1).




Figure 1. Consort diagram of patients with alterations in the G₁/S phase cell-cycle signaling pathway (n = 507).

Most patients with alterations in G₁/S phase cell-cycle signaling pathway had genomic co-alterations. Among 507 patients with diverse tumors harboring CDK4/6 amplifications, CCND1/2/3 amplifications, or CDKN2A/B alterations, 99% (n = 501) had at least 1 deleterious co-alteration (median, 4 co-alterations [excludes the cyclin alteration]; range, 0–24) in tissue NGS and the remaining 6 patients whose tumors did not have a co-alteration only had a CDKN2A/B alteration. The most common co-alterations were seen in the TP53 (48% of the 507 patients, n = 241), EGFR (17%, n = 87), TERT (16%, n = 82), and KRAS (16%, n = 81) genes (Figure 2). Co-alterations in cell-cycle resistant genes (RB and CCNE1) were rare (each occurring in less than 2% of cases).

In patients with cell-cycle–altered tumors, CDK4/6 inhibitor–based therapy with high matching score was associated with significantly longer PFS and a higher rate of clinical benefit (stable disease ≥6 months or objective response). Among 507 patients with CDK4/6 amplifications, CCND1/2/3 amplifications, or CDKN2A/B alterations, 40 patients with diverse cancers (excluding patients with breast cancer) were treated with CDK4/6 inhibitor–containing regimens and evaluated for PFS (Figure 1 and Supplemental Table 1; supplemental material available online with this article; <https://doi.org/10.1172/jci.insight.142547DS1>). (The breast cancer cohort was excluded from the analysis because the combination of CDK4/6 inhibitors with antihormone agents is already FDA approved for patients with breast cancer [ref. 29]). None of the treated patients had a coexisting cell-cycle gene alteration in RB or CCNE1.

Among those 40 patients with diverse cancers, PFS was not associated with age, sex, types of cancer, line of therapy, or treatment regimen in univariate analysis (Table 2 and Figure 3A). Among patients with CDKN2A/B alterations, PFS was worse, but the difference was not significant (median PFS between CDKN2A/B alteration vs. not, 4.0 vs. 6.8 months; P = 0.10 [univariate]). PFS was significantly longer among patients with Eastern Cooperative Oncology Group Performance Status (ECOG PS) of 0–1 (median PFS between ECOG PS 0–1 vs. 2–3, 6.1 vs. 1.6 months; P = 0.04) and in patients who had a higher matching score (i.e., matching score roughly equivalent to the number of alterations targeted divided by total number of deleterious alterations) (median PFS between matching score ≥50% vs. <50%, 6.2 vs. 2.0 months; P = 0.001 [univariate]) (Table 2 and Figure 3B).

After the multivariate analysis, CDKN2A/B alteration remained a factor independently associated with poor PFS (HR, 2.76; 95% CI, 1.10–6.93; P = 0.03) and high matching score was an independent factor for longer PFS (≥ 50% vs. <50%) (HR, 0.24; 95% CI, 0.11–0.51; P < 0.001) (Table 2).

In line with the favorable longer PFS seen, patients with a matching score of ≥50% achieved stable disease for ≥6 months or a stable objective response rate at higher rates (matching score of ≥ 50% vs. < 50%, 57% vs. 21%, P = 0.048) (Figure 3C). Improved PFS with a higher matching score also translated into numerically longer median OS, which, however, was not statistically significant (median OS between

Table 1. Characteristics of 507 patients with alterations in the G₁/S phase cell-cycle signaling pathway

Characteristics	Total patients (n = 507)
Median age ^a , yr (range)	62.1 (20.8–92.6)
Sex, n (%)	
Men	265 (52.3%)
Women	242 (47.7%)
Race or ethnicity, n (%)	
White	354 (69.8%)
Asian	57 (11.2%)
Hispanic	55 (10.8%)
African American	12 (2.4%)
Other/unknown	29 (5.8%)
Type of cancer, n (%)	
Brain	83 (16.4%)
Lung, non-small cell	77 (15.2%)
Skin/melanoma	67 (13.2%)
Hepato-pancreato-biliary	58 (11.4%)
Breast	51 (10.1%)
Gastrointestinal, noncolorectal	46 (9.1%)
Head and neck	37 (7.3%)
Genitourinary/prostate	15 (3.0%)
Gynecologic	14 (2.8%)
Colorectal	12 (2.4%)
Thyroid	7 (1.4%)
Other	40 (7.9%)
Median no. of characterized co-alterations (range)	4.0 (0–24)
Alterations in <i>CCND1/2/3</i> , <i>CDK4/6</i> , and/or <i>CDKN2A/B</i> genes as part of the G ₁ /S cell-cycle signaling pathways, n (%) ^b	
<i>CCND1</i> amplification	75 (14.8%)
<i>CCND2</i> amplification	17 (3.4%)
<i>CCND3</i> amplification	21 (4.1%)
<i>CDK4</i> amplification	61 (12.0%)
<i>CDK6</i> amplification	26 (5.1%)
<i>CDKN2A/B</i> alteration	359 (70.8%)

^aAge at the time of metastatic/locally advanced disease. ^bSome patients had more than one genomic alteration in the *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B* genes; each alteration is counted separately.

matching score $\geq 50\%$ vs. $< 50\%$, 8.3 vs. 5.3 months; $P = 0.15$ [univariate]) (Figure 3D). Similar clinical outcomes were observed among the 33 patients who were managed with regimens that did not contain immunotherapy (matching score $\geq 50\%$ vs. $< 50\%$, PFS, 8.8 vs. 3.2 months [$P = 0.001$], OS, 13.0 vs. 8.0 months [$P = 0.08$] [univariate]) (Supplemental Table 1).

Examples of responding patients treated with CDK4/6 inhibitory therapy. Case 1 (patient ID 269) is a 43-year-old woman with metastatic high-grade ovarian carcinoma with neuroendocrine features and 2 prior lines of therapy. The patient's tumor harbored a sole alteration in *CDKN2A/B* and demonstrated a response with single-agent palbociclib (30% regression; partial response by RECIST 1.1; tumor marker, CA 125, 328 [baseline] down to 50 U/ml [reference range, 0–34 U/ml], PFS, 8.0 months) (Figure 4A and Supplemental Table 1).

Case 2 (patient ID 501) is a 68-year-old man with metastatic gastrointestinal stromal tumor with *BRAF* V600E and *CDKN2A* alterations (30), who presented after the tumor progressed on BRAF/MEK-targeted therapy. Addition of palbociclib led to resolution of ¹⁸F-fluorodeoxyglucose-avid diseases per PET/CT scan and PFS of 11.3 months without significant toxicities (Figure 4B and Supplemental Table 1).

Discussion

The cell cycle allows normal cellular growth and proliferation and is highly regulated by a series of cyclin molecules and their dependent constellation of kinases, whose signals must be integrated to




Figure 2. Summary of co-alterations observed in tumors harboring *CDK4/6* amplifications, *CCND1/2/3* amplifications, or *CDKN2A/B* alterations (*n* = 507). Among 507 patients with diverse tumors harboring *CDK4/6* amplifications, *CCND1/2/3* amplifications, or *CDKN2A/B* alterations, most patients (99%, *n* = 501) had ≥ 1 characterized co-alteration (median, 4; range, 0–24) in tissue NGS. The most common co-alterations were in *TP53* (approximately 48% of patients, *n* = 241), *EGFR* (17% of patients, *n* = 87), *TERT* (16% of patients, *n* = 82), and *KRAS* genes (16% of patients, *n* = 81). Genomic alterations with frequency of $\geq 1.0\%$ were included.

determine if it is appropriate for cells to divide. The activity of these kinases relies on the production of their cognate cyclin partners, represented by the D-type cyclins *CCND1/2/3*. In parallel, *CDK4/6* kinases are also regulated by phosphorylation events and the presence of physiological kinase inhibitor proteins. These inhibitors are encoded by the *CDKN2* gene family — *CDKN2A* and *CDKN2B*, which yield selective CDK inhibitors for *CDK4/6* (e.g., p16INK4a and p15INK4b) (31). The dysregulation through genomic alteration of the aforementioned major players, including *CDK4/6*, *CCND1/2/3*, and *CDKN2A/B* genes, has been implicated in the pathogenesis of diverse malignancies. Currently, *CDK4/6* inhibitors are approved for patients with HR-positive, HER2-negative metastatic breast cancer, as these inhibitors improve PFS and OS when given with hormone modulators (7–10). Even so, investigators have failed to identify a reliable biomarker for *CDK4/6* inhibitors, despite attempts made in several studies of breast cancer (11–13, 15). Moreover, *CDK4/6* inhibitors given as monotherapy matched to cognate alterations in a variety of cancers have mostly fared poorly (14, 15).

We hypothesized that the lack of association between *CDK4/6* amplifications, *CCND1/2/3* amplifications, and/or *CDKN2A/B* alterations and outcome after administration of *CDK4/6* inhibitors may be due to intratumoral heterogeneity and complexity, resulting in a large proportion of metastatic tumors with cyclin alterations also carrying genomic co-alterations that differ from patient to patient. Consistent with our hypothesis, in our cohort of 507 patients with a variety of cancers harboring *G₁/S* phase cell-cycle gene alterations (*CDK4/6* amplifications, *CCND1/2/3* amplifications, or *CDKN2A/B* alterations), 99% of patients (*n* = 501) had at least one genomic co-alteration (median, 4; range, 0–24). These alterations were heterogeneous and affected multiple oncogenic signaling pathways, including those regulated by mitogen-activated protein kinase, phosphoinositide 3-kinase, and β -catenin/Wnt; other kinase families and *BRCA*-associated genes were also affected (Figure 2).

Table 2. Analysis of factors associated with progression-free survival among patients with alterations in *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B* G₁/S phase cell-cycle genes, who received *CDK4/6* inhibitor-based therapies (n = 40)

Patient characteristics (n = 40)	Univariate analysis		Multivariate analysis ^a	
	Median PFS months	P value	Hazard ratio (95%CI)	P value
Age (yr), ≥62 (n = 18) vs. <62 (n = 22)	4.0 vs. 4.5	0.62	-	-
Sex, men (n = 21) vs. women (n = 19)	4.6 vs. 4.5	0.33	-	-
ECOG PS, 0-1 (n = 30) vs. 2-3 (n = 10)	6.1 vs. 1.6	0.04	0.65 (0.29-1.44)	0.29
Types of cancer				
Hepato-pancreato-biliary (n = 10) vs. other (n = 30)	1.9 vs. 4.6	0.80	-	-
Gastrointestinal (n = 6) vs. other (n = 34)	6.8 vs. 4.0	0.68	-	-
Alterations in cell-cycle signaling pathways ^b				
<i>CCND1</i> amplification (n = 3) vs. not (n = 37)	2.6 vs. 4.6	0.41	-	-
<i>CCND2</i> amplification (n = 2) vs. not (n = 38)	0.7 vs. 4.5	0.18	-	-
<i>CCND3</i> amplification (n = 1) vs. not (n = 39)	8.0 vs. 4.5	0.67	-	-
<i>CDK4</i> amplification (n = 7) vs. not (n = 33)	3.8 vs. 4.6	0.61	-	-
<i>CDK6</i> amplification (n = 3) vs. not (n = 37)	2.0 vs. 4.6	0.79	-	-
<i>CDKN2A/B</i> alteration (n = 31) vs. not (n = 9)	4.0 vs. 6.8	0.10	2.76 (1.10-6.93)	0.03
Line of therapy				
As first line (n = 8) vs. ≥ second line (n = 32)	6.1 vs. 4.0	0.25	-	-
CDK4/6 inhibitory therapy ^c				
Combination (n = 31) vs. single agent (n = 9)	4.6 vs. 2.8	0.26	-	-
Matching score ≥ 50% (n = 25) vs. < 50% (n = 15)	6.2 vs. 2.0	0.001	0.24 (0.11-0.51)	<0.001

^aVariables with $P \leq 0.10$ in univariate analysis (the log-rank test) were included in the multivariate analysis. ^bNo patient had co-alteration in *RB1* or *CCNE1*.

^cCombination indicates that patient received one or more additional therapy along with *CDK4/6* inhibitor. The *CDK4/6* inhibitor was matched in these cases with an alteration in *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B*, but any additional drugs were not necessarily matched to a co-alteration in that patient. See definition of "matching score" in Methods. In patients with a matching score of higher than ≥50%, higher degrees of matching were achieved by matching the genomic co-alterations to cognate drugs, in addition to matching the *CDK4/6* inhibitors to *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B* alterations. ECOG PS, Eastern Cooperative Oncology Group Performance Status; PFS, progression-free survival. Significant values are shown in bold.

The presence of coexisting disrupted oncogenic pathways could potentially lead to resistance to *CDK4/6* inhibitors. Consistent with this notion, in the current study, even combination therapy that matched *CDK4/6* inhibitors to *CDK4/6* amplifications, *CCND1/2/3* amplifications, or *CDKN2A/B* alterations, but without necessarily matching to genomic co-alterations, did not achieve a better clinical outcome when compared with patients who received matched *CDK4/6* inhibitors alone (combination approach [implying matched *CDK4/6* inhibitor and at least one other drug] vs. *CDK4/6* inhibitor alone, PFS, 4.6 months vs. 2.8 months, $P = 0.26$) (Table 2). However, when the *CDK4/6* inhibitor-based regimens were given together with customized additional drugs matched to genomic co-alterations in that patient's tumor (resulting in a high matching score (≥50%), overall longer PFS was observed when compared with that of patients who were treated with a *CDK4/6*-matched regimen with a low matching score (<50%) (PFS 6.2 vs. 2.0 months, $P < 0.001$ [P values were calculated after multivariate analysis]) (Table 2). The clinical benefit rate (stable disease ≥6 months as well as a higher objective response rate) was also improved (57% vs. 21%; $P = 0.048$) (Figure 3C).

In the era of precision oncology, the majority of cancer clinical trials are aimed at a prespecified genomic target of interest, and many patients are being treated with single-matched drugs. Salutary effects have been observed by targeting certain genomic alterations, such as *NTRK*, *RET*, and *ALK* fusions or *BRAF* V600 as well as *EGFR* mutations (22, 32-35). However, resistance is inevitable and there is likely a limitation of benefit with single agents in the setting of genomically complex advanced cancers. To overcome these limitations, future directions for the development of cancer clinical trials may require a more flexible, individualized treatment strategy that is tailored to each patient's tumor genomic profile. To this end, we have recently reported the outcome of the I-PREDICT and WINTHER trials, wherein we investigated personalized approaches based on genomic and/or transcriptomic profiling among patients with treatment-refractory solid tumors (26, 28). We demonstrated improvement in PFS and OS when targeting a larger fraction of identified molecular alterations, reflecting a high matching score, consistent with the observations in the current study. Further prospective trials with this tactic focused on patients with cell-cycle alterations are required.




Figure 3. Progression-free survival among patients with alterations in *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B G₁/S* phase cell-cycle genes, who received *CDK4/6* inhibitor-based therapy (n = 40). (A) Progression-free survival (PFS) comparison between patients who received *CDK4/6* inhibitor-based therapy as part of the combination therapies (n = 31) and patients who received *CDK4/6* inhibitor as a single agent (n = 9). Among patients with diverse cancers (n = 40) who received *CDK4/6* inhibitor-based therapy, there was no significant difference in PFS between patients who received combination therapy and those who received single agents (combination vs. single agent, 4.6 vs. 2.8 months, P = 0.26). (B) PFS among patients who received *CDK4/6* inhibitor-based therapy with a matching score of ≥50% (n = 25) versus those with a matching score of <50% (n = 15). Among patients with diverse cancers (n = 40) who received *CDK4/6* inhibitor-based therapy, patients who were treated with a combination of agents with higher matching scores had significantly longer PFS (median PFS for matching score ≥50% vs. <50%, 6.2 vs. 2.0 months, P = 0.001). (C) Response to *CDK4/6* inhibitor-based therapies among patients with *CCND1/2/3*, *CDK4/6*, and/or *CDKN2A/B G₁/S* phase cell-cycle gene alterations. Comparison between patients who received *CDK4/6* inhibitor-based therapy with matching score of ≥50% (n = 23) and patients with matching score of <50% (n = 14). There was a significant difference in achieving stable disease ≥6 months/partial response among patients who received therapy with matching score of ≥50% as compared with those with a matching score of <50 (57% vs. 21%, P = 0.048). (Among 40 patients treated with matched *CDK4/6* inhibitor-based therapies, 37 patients were assessable for response.) (D) Overall survival (OS) comparison (n = 40) between patients who received *CDK4/6* inhibitor-based therapy with matching score of ≥50% (n = 25) and patients with matching score of <50% (n = 15). Among patients who received *CDK4/6* inhibitor-based therapy (n = 40), there was no significant difference in OS between those with a matching score of ≥50% vs. <50% (median OS between matching score ≥ 50% vs. < 50%, 8.3 vs. 5.3 months; P = 0.15). Reverse Kaplan-Meier calculation for A, B, and D revealed no difference between groups, indicating that the median follow-up between groups was similar. MS, matching score; PD, progressive disease; PR, partial response; SD, stable disease.

There are several important limitations to the current report. First, the study has a small sample size. Second, while PFS and clinical benefit rate were improved with a greater degree of matching, survival changes did not reach statistical significance. A larger prospective trial that is controlled and randomized is needed, especially to mitigate the effect of confounders that may not be known despite the multivariate analysis. Third, molecular characteristics of tumors can have dynamic changes, especially with therapeutic pressure. Future studies may require serial profiling, such as with circulating tumor cell-free DNA analysis. Fourth, our study assessed only pathogenic somatic alterations, and not germ-line anomalies. Finally, this study included heterogeneous cancer diagnoses, and the number of patients in specific histologies was small (and patients were treated at various time points in their disease), which precluded the ability to interpret the results for specific disease types.

In conclusion, we have evaluated 2457 patients with diverse solid tumors and shown that potentially sensitizing *G₁/S* phase cell-cycle molecular aberrations, such as *CDK4/6* amplifications, *CCND1/2/3*





Figure 4. Examples of responders treated with CDK4/6 inhibitory therapy. (A) Case 1 (patient ID 269): Forty-three-year-old woman with metastatic high-grade ovarian carcinoma with neuroendocrine features that harbored *CDKN2A/B* alteration without any genomic co-alterations on the NGS panel of 315 genes and 2 lines of prior therapy demonstrated partial response with single-agent palbociclib lasting 8 months. NGS of tumor showed a single alteration in *CDKN2A/B*, for which the patient was started on palbociclib. Restaging scan with CT overall showed 30% regression, indicating partial response at the 4-month time point (response by RECIST 1.1). Along with the radiographic response, reduction of tumor marker, CA 125 was seen (CA 125: 328 U/ml down to 50 U/ml [reference range 0–34 U/ml]). **(B)** Case 2 (patient ID 501): Sixty-eight-year-old man with metastatic gastrointestinal stromal tumor (GIST) with alterations in *BRAF* V600E, *CDKN2A* p16INK4a splice site 150+1G > A and *LRP1B* deletion exon 23 presented after progressing treatment with dabrafenib (BRAF inhibitor) and trametinib (MEK inhibitor) based on underlying *BRAF* V600E mutation (30). Addition of palbociclib led to partial response lasting 11.3 months. Although progression was seen with a new pulmonary nodule and worsening rectal lesion (left to middle, circle), one of the right lower lung masses appeared to be stable (left to middle, arrow), and thus the decision was made to continue on dabrafenib/trametinib and to add palbociclib based on additional alteration in *CDKN2A*. Two months after the addition of palbociclib, restaging scan with PET/CT scan showed resolution of ¹⁸F-fluorodeoxyglucose-avid lung nodules as well as improvement in rectal lesion (middle to right). PFS was 11.3 months without significant toxicities.



amplifications, and/or *CDKN2A/B* alterations, were observed in 507 patients (21%). Most participants with these alterations (99% of patients, 501 of 507) had at least one genomic co-alteration. Among patients with *CDK4/6* amplifications, *CCND1/2/3* amplifications, and/or *CDKN2A/B* abnormalities, adding additional drugs to the *CDK4/6* inhibitor-based regimen without consideration of genomic co-alterations did not improve clinical outcome. However, significant improvement in PFS and in clinical benefit rate were observed when matched *CDK4/6* inhibitors were given as part of a tailored regimen that affected a larger proportion of genomic alterations, with achievement of a high matching score.

Because the genomic co-alterations differed from patient to patient, individualized combination therapies were often required. These results imply that, in the case of CDK4/6 inhibitors given to patients whose tumors harbor potentially sensitizing cyclin alterations, personalized consideration of important molecular co-alterations warrants further investigation as a direction for achieving benefit.

Methods

Study population. Patients were generally matched after presentation to a Molecular Tumor Board (36–38). Some patients were navigated to prospective precision studies, such as I-PREDICT (28). A total of 2457 patients with solid tumors who underwent tissue NGS were analyzed from January 2013 to April 2018. All patients were at UCSD. Among them, 507 patients with genomic alterations in *CDK4/6*, *CCND1/2/3*, or *CDKN2A/B* were included for more in-depth assessment ($n = 507$) (Figure 1). These genes were chosen because they are potentially sensitizing to CDK4/6 inhibitors.

Tissue NGS. All tissue DNA analyses were performed by a clinical laboratory improvement amendments–certified lab, Foundation Medicine Inc., as per methods previously described in detail (39) (<https://www.foundationmedicine.com>), except for 2 patients, with analyses performed at UCSD NGS and HLIQ Oncology (182–465 cancer-related genes). Briefly, 50–200 ng genomic DNA was extracted and purified from the submitted FFPE tumor samples. DNA was adaptor ligated, and hybrid capture was performed for all coding exons of 182–406 cancer-related genes plus selected introns from 14–31 genes frequently rearranged in cancer (Illumina HiSeq platform). Sequencing was performed with an average sequencing depth of coverage of $>250\times$, with $>100\times$ at $>99\%$ of exons. Somatic mutations were identified with $>99\%$ sensitivity and 99% specificity for base substitutions at $>95\%$ sensitivity for copy number alterations, and $\geq 5\%$ mutant allele frequency. Gene amplification was reported at ≥ 8 copies above ploidy, with ≥ 6 copies considered equivocal (with the exception of *ERBB2*, for which ≥ 5 copies is considered equivocal amplification). Tumor-mutation burden was classified into 3 categories: low (<6 mutations/mb), intermediate (6–19 mutations/mb), and high (≥ 20 mutations/mb). Variants of unknown significance were excluded from all analyses.

Molecular matching score. The molecular matching score was developed in an attempt to assess the association between coverage of deleterious genomic alterations by targeted therapy that patients may have received based on these molecular alterations and the clinical outcome, as previously described (27, 28, 41, 42). The score is roughly equal to the total number of deleterious alterations affected divided by the total number of deleterious alterations in each patient. Under this system, the higher the molecular matching score, the better the match. See Supplemental Methods for further description. Matching scores were determined while blinded to outcome.

Statistics. Patient characteristics; prevalence of alterations in *CDK4/6*, *CCND1/2/3*, or *CDKN2A/B*; and genomic co-alterations were summarized by descriptive statistics such as Kaplan-Meier and Log-rank test were used (as stated below). Among 40 patients with cancer (but not breast cancer) who underwent CDK4/6 inhibitor–based therapies, we assessed PFS, which was defined as time between start of the treatment and disease progression confirmed by imaging or clinical findings. OS was defined as time between start of therapy until the last follow-up. Patients with ongoing therapy without progression at the last follow-up date were censored for PFS at that date. Patients alive at last follow-up were censored for OS. PFS and OS were assessed by the Kaplan-Meier method. Reverse Kaplan-Meier was also done to determine whether there were differences in median follow-up times between groups (40). Log-rank test and Cox regression analysis were used to compare subgroups of patients. All tests were 2 sided, and variables with $P \leq 0.1$ were included for multivariate analysis. $P \leq 0.05$ was considered significant. Statistical analyses were performed using SPSS version 24 software (IBM Corporation).

Study approval. All investigations in this study were approved by and analyzed according to the guidelines of Moores Cancer Center at UC San Diego Health Internal Review Board under the Profile-Related Evidence Determining Individualized Cancer Therapy study (PREDICT study, NCT02478931). Patients gave informed consent before participating in investigational therapies.

Author contributions

SK, JJA, JKS, and RK drafted the manuscript; SK and RK designed the study; SK and RO analyzed the data; and RO, NK, SL, and VN collected the data. All authors read and approved the final manuscript.

Acknowledgments

This work was supported in part by the Joan and Irwin Jacobs Fund (to RK), the National Cancer Institute/NIH (P30 CA023100, to RK and JKS), the NIH (R01 CA226803, to JKS), and the FDA (R01 FD006334, to JKS) as well as Jon Strong (to JKS).

Address correspondence to: Shumei Kato, Center for Personalized Cancer Therapy and Division of Hematology and Oncology, Department of Medicine, UC San Diego Moores Cancer Center, 3855 Health Sciences Drive, La Jolla, California 92093, USA. Phone: 858.822.2372; Email: smkato@ucsd.edu.

1. Barnum KJ, O'Connell MJ. Cell cycle regulation by checkpoints. *Methods Mol Biol.* 2014;1170:29–40.
2. Kato S, et al. Cyclin-dependent kinase pathway aberrations in diverse malignancies: clinical and molecular characteristics. *Cell Cycle.* 2015;14(8):1252–1259.
3. Schwaederle M, et al. Cyclin alterations in diverse cancers: Outcome and co-amplification network. *Oncotarget.* 2015;6(5):3033–3042.
4. Shapiro GI. Cyclin-dependent kinase pathways as targets for cancer treatment. *J Clin Oncol.* 2006;24(11):1770–1783.
5. Sheppard KE, McArthur GA. The cell-cycle regulator CDK4: an emerging therapeutic target in melanoma. *Clin Cancer Res.* 2013;19(19):5320–5328.
6. Helsten T, et al. Cell-cycle gene alterations in 4,864 tumors analyzed by next-generation sequencing: implications for targeted therapeutics. *Mol Cancer Ther.* 2016;15(7):1682–1690.
7. Turner NC, et al. Overall survival with palbociclib and fulvestrant in advanced breast cancer. *N Engl J Med.* 2018;379(20):1926–1936.
8. Im SA, et al. Overall survival with ribociclib plus endocrine therapy in breast cancer. *N Engl J Med.* 2019;381(4):307–316.
9. Hortobagyi GN, et al. Ribociclib as first-line therapy for HR-Positive, advanced breast cancer. *N Engl J Med.* 2016;375(18):1738–1748.
10. Goetz MP, et al. MONARCH 3: abemaciclib as initial therapy for advanced breast cancer. *J Clin Oncol.* 2017;35(32):3638–3646.
11. Finn RS, et al. The cyclin-dependent kinase 4/6 inhibitor palbociclib in combination with letrozole versus letrozole alone as first-line treatment of oestrogen receptor-positive, HER2-negative, advanced breast cancer (PALOMA-1/TRIO-18): a randomised phase 2 study. *Lancet Oncol.* 2015;16(1):25–35.
12. Shapiro GI. Genomic biomarkers predicting response to selective CDK4/6 inhibition: progress in an elusive search. *Cancer Cell.* 2017;32(6):721–723.
13. Turner NC, et al. Cyclin E1 expression and palbociclib efficacy in previously treated hormone receptor-positive metastatic breast cancer. *J Clin Oncol.* 2019;37(14):1169–1178.
14. Mangat PK, et al. Rationale and design of the targeted agent and profiling utilization registry study. *JCO Precision Oncology.* 2018;2:1–14.
15. Finn RS, et al. Biomarker analyses of response to cyclin-dependent kinase 4/6 inhibition and endocrine therapy in women with treatment-naïve metastatic breast cancer. *Clin Cancer Res.* 2020;26(1):110–121.
16. DeMichele A, et al. CDK 4/6 inhibitor palbociclib (PD0332991) in Rb+ advanced breast cancer: phase II activity, safety, and predictive biomarker assessment. *Clin Cancer Res.* 2015;21(5):995–1001.
17. Kato S, et al. Analysis of circulating tumor DNA and clinical correlates in patients with esophageal, gastroesophageal junction, and gastric adenocarcinoma. *Clin Cancer Res.* 2018;24(24):6248–6256.
18. Kato S, et al. Rare tumor clinic: The University of California San Diego Moores Cancer Center experience with a precision therapy approach. *Oncologist.* 2018;23(2):171–178.
19. Kato S, et al. Utility of genomic analysis in circulating tumor DNA from patients with carcinoma of unknown primary. *Cancer Res.* 2017;77(16):4238–4246.
20. Ramalingam SS, et al. Overall survival with osimertinib in untreated, EGFR-mutated advanced NSCLC. *N Engl J Med.* 2020;382(1):41–50.
21. Solomon BJ, et al. First-line crizotinib versus chemotherapy in ALK-positive lung cancer. *N Engl J Med.* 2014;371(23):2167–2177.
22. Drlon A, et al. Efficacy of larotrectinib in TRK fusion-positive cancers in adults and children. *N Engl J Med.* 2018;378(8):731–739.
23. Okamura R, et al. Analysis of NTRK alterations in pan-cancer adult and pediatric malignancies: implications for NTRK-targeted therapeutics. *JCO Precis Oncol.* 2018;2(1):1–20.
24. Kopetz S, et al. Encorafenib, binimetinib, and cetuximab in BRAF V600E-mutated colorectal cancer. *N Engl J Med.* 2019;381(17):1632–1643.
25. Le Tourneau C, et al. Molecularly targeted therapy based on tumour molecular profiling versus conventional therapy for advanced cancer (SHIVA): a multicentre, open-label, proof-of-concept, randomised, controlled phase 2 trial. *Lancet Oncol.* 2015;16(13):1324–34.
26. Rodon J, et al. Genomic and transcriptomic profiling expands precision cancer medicine: the WINETHER trial. *Nat Med.* 2019;25(5):751–758.
27. Schwaederle M, et al. Precision oncology: the UC San Diego Moores Cancer Center PREDICT experience. *Mol Cancer Ther.* 2016;15(4):743–752.
28. Sicklick JK, et al. Molecular profiling of cancer patients enables personalized combination therapy: the I-PREDICT study. *Nat Med.* 2019;25(5):744–750.
29. O'Leary B, et al. Treating cancer with selective CDK4/6 inhibitors. *Nat Rev Clin Oncol.* 2016;13(7):417–430.
30. Falchook GS, et al. BRAF mutant gastrointestinal stromal tumor: first report of regression with BRAF inhibitor dabrafenib (GSK2118436) and whole exomic sequencing for analysis of acquired resistance. *Oncotarget.* 2013;4(2):310–315.
31. Knudsen ES, Witkiewicz AK. The strange case of CDK4/6 inhibitors: mechanisms, resistance, and combination strategies. *Trends Cancer.* 2017;3(1):39–55.
32. Zhou C, et al. Erlotinib versus chemotherapy as first-line treatment for patients with advanced EGFR mutation-positive

- non-small-cell lung cancer (OPTIMAL, CTONG-0802): a multicentre, open-label, randomised, phase 3 study. *Lancet Oncol.* 2011;12(8):735–742.
33. Shaw AT, et al. Crizotinib versus chemotherapy in advanced ALK-positive lung cancer. *N Engl J Med.* 2013;368(25):2385–2394.
34. Hyman DM, et al. Vemurafenib in multiple nonmelanoma cancers with BRAF V600 mutations. *N Engl J Med.* 2015;373(8):726–736.
35. Kato S, et al. *RET* aberrations in diverse cancers: next-generation sequencing of 4,871 patients. *Clin Cancer Res.* 2017;23(8):1988–1997.
36. Schwaederle M, et al. Molecular tumor board: the University of California-San Diego Moores Cancer Center experience. *Oncologist.* 2014;19(6):631–6.
37. Patel M, et al. Molecular tumor boards: realizing precision oncology therapy. *Clin Pharmacol Ther.* 2018;103(2):206–209.
38. Kato S, et al. Real-world data from a molecular tumor board demonstrates improved outcomes with a precision N-of-One strategy. *Nat Commun.* 2020;11(1):4965.
39. Frampton GM, et al. Development and validation of a clinical cancer genomic profiling test based on massively parallel DNA sequencing. *Nat Biotechnol.* 2013;31(11):1023–1031.
40. Schemper M, Smith TL. A note on quantifying follow-up in studies of failure time. *Control Clin Trials.* 1996;17(4):343–346.
41. Mazumdar M, Glassman JR. Categorizing a prognostic variable: review of methods, code for easy implementation and applications to decision-making about cancer treatments. *Stat Med.* 2000;19(1):113–132.
42. Wheler JJ, et al. Cancer therapy directed by comprehensive genomic profiling: a single center study. *Cancer Res.* 2016;76(13):3690–3701.