

UC Irvine

UC Irvine Previously Published Works

Title

Maternal caregiving ameliorates the consequences of prenatal maternal psychological distress on child development

Permalink

<https://escholarship.org/uc/item/6829460d>

Journal

Development and Psychopathology, 34(4)

ISSN

0954-5794

Authors

Grande, Leah A
Swales, Danielle A
Sandman, Curt A
[et al.](#)

Publication Date

2022-10-01

DOI

10.1017/s0954579421000286

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Published in final edited form as:

Dev Psychopathol. 2022 October ; 34(4): 1376–1385. doi:10.1017/S0954579421000286.

Maternal Caregiving Ameliorates the Consequences of Prenatal Maternal Psychological Distress on Child Development

Leah A. Grande¹, Danielle A. Swales¹, Curt A. Sandman², Laura M. Glynn³, Elysia Poggi Davis^{1,2}

¹Department of Psychology, University of Denver, Denver, CO 80210

²Department of Psychiatry and Human Behavior, University of California-Irvine, Irvine, CA 92697

³Department of Psychology, Chapman University, Orange, CA 92866

Abstract

Children exposed to prenatal maternal psychological distress are at elevated risk for a range of adverse outcomes; however, it remains poorly understood whether postnatal influences can ameliorate impairments related to prenatal distress. The current study evaluated if maternal care could mitigate child cognitive and emotional impairments associated with prenatal psychological distress. Prenatal maternal psychological distress was assessed (anxiety, depression, and perceived stress) for 136 mothers at 5 prenatal and 4 postpartum time points. Quality of maternal care (sensitivity to nondistress, positive regard and intrusiveness reverse-scored) were assessed during a mother-child play interaction at 6 and 12 months. Child cognitive function and negative emotionality were assessed at 2 years, using The Bayley Scales and the Early Childhood Behavior Questionnaire. Elevated prenatal distress was associated with poorer child cognitive function and elevated negative emotionality. Children exposed to elevated prenatal maternal distress did not, however, display these outcomes if they received high quality caregiving. Specifically, maternal care moderated the relation between prenatal psychological distress and child cognitive function and negative emotionality. This association remained after consideration of postnatal maternal psychological distress and relevant covariates. Sensitive maternal care was associated with altered offspring developmental trajectories, supporting child resilience following prenatal distress exposure.

Keywords

Fetal programming; maternal care; prenatal stress; resilience; cognitive function; depression; parenting

Fetal life is an exceptionally rapid period of neurological development, and is a time when the fetus is highly susceptible to both beneficial and harmful environmental influences (Barker, 1998). Prenatal maternal psychological distress is linked to profound and lasting consequences for developmental trajectories and increases risk for subsequent mental health

Correspondence to: Elysia Davis, 2155 South Race Street, Denver, CO 80210, elysia.davis@du.edu, 303-871-3790.

Conflicts of Interest: None

problems (Davis et al., 2007; Davis & Sandman, 2010, 2012; Demers et al., 2020; Glynn et al., 2018; Van den Bergh et al., 2017). The evidence that prenatal maternal stress and mental health has long reaching implications for offspring psychopathology raises the need to investigate factors that can reduce or eliminate the consequences of prenatal adversity. High quality parental caregiving is a likely factor that may increase resiliency in the face of prenatal adversity (Davis et al., 2019; Kok et al., 2015; NICHD Early Childcare Research Network, 1999). The current study addresses an important knowledge gap by investigating whether high quality maternal caregiving during infancy ameliorates the consequences of prenatal maternal distress on child cognitive and emotional development, thereby shifting trajectories of risk towards more optimal mental health.

Fetal programming research illustrates that children exposed to maternal psychological distress (e.g. anxiety, depression, and stress) during pregnancy have poorer cognitive function and elevated negative emotionality during infancy, childhood, and adolescence (Blair et al., 2011; Glynn et al., 2018; Korja et al., 2017; Madigan et al., 2018; Mahrer et al., 2019; Sandman & Davis, 2010; Van den Bergh et al., 2005, 2017). Importantly, these prospective and longitudinal studies document the predictive importance of prenatal distress on child outcomes, even after covarying potential confounders, including postnatal maternal psychological distress. Further, an independent line of research focusing on postnatal experiences reports that sensitive and responsive maternal care is associated with benefits for child cognitive and emotional development, as well as adult psychological health (Davis et al., 2017; Deans, 2018; Fan et al., 2014; Farrell et al., 2019; Malmberg et al., 2016; Spinrad & Stifter, 2002). Maternal caregiving during the first year postpartum may be especially important, because this is a sensitive window for mother-infant relationship development (Ainsworth, 1979; Feldman, 2007).

Although the opportunity to assess the joint contributions of prenatal and postnatal experiences are rare in human research, and often are limited by a lack of objective assessment of maternal behavior (Sharp et al., 2012, 2015), experimental research with animals shows that high quality maternal care can compensate for exposure to prenatal maternal stress. Specifically, the adverse effects of prenatal maternal stress on offspring cognition, emotion stress regulation, and brain development can be prevented with experimental manipulations of maternal quality of care (Bogoch et al., 2007; Lemaire et al., 2006; Rainekei et al., 2014; Wakshlak & Weinstock, 1990).

Several studies with clinical populations have tested the benefit of sensitive, responsive care on human infant development. Among mothers with an anxiety or a depression diagnosis during pregnancy, high quality of maternal care is associated with a reduction in the correlation between prenatal maternal mental health and offspring cortisol regulation at 4 months (Kaplan et al., 2008), on negative affect responses to still-face at 7 months (Grant et al., 2010a), and on Bayley cognitive development scores at 7 months (Grant et al., 2010b). In addition to the relatively small sample sizes of these studies (ranging from 47 to 84), limitations include reliance on assessment of a single diagnostic category (i.e., diagnoses of anxiety or depressive disorders; Grant et al., 2010a, 2010b; Kaplan et al., 2008). Reliance on a single timepoint assessment as well as diagnostic cut offs in a single disorder may not fully capture fetal exposure to maternal psychological distress for several

reasons. First, there is compelling evidence that maternal distress across the spectrum and including in the non-clinical range is linked to child outcomes (Davis & Sandman, 2012; Glynn et al., 2018; Kingston et al., 2012; Krueger & Markon, 2006; Lee et al., 2007). Second, women frequently experience comorbidity of stress and internalizing symptoms from multiple disorders all of which may impact the fetus (Falah-Hassani et al., 2016; Glynn et al., 2018). For these reasons, it is important to assess prenatal exposure to maternal anxiety, depression and stress multiple times during gestation to create a reliable index of exposure to multiple types of maternal distress over the entire course of gestation and then evaluate the compensatory benefit of maternal care.

A recent study, that addressed these limitations, reported that high levels of maternal positive engagement at ages 2.5 to 5 years alleviated the cognitive delays associated with prenatal maternal distress and suggested that parental care may be an important target for intervention (Schechter et al., 2017). The present study extends the findings by Schechter et al. (2017) in two important ways. First, by assessing maternal care during the first postnatal year (6 and 12 months), a sensitive window for attachment formation, this study assesses a potential target for early intervention. Second, the current study assesses both child cognitive function and negative emotionality, two important indices of early child development. Specifically, the current prospective research characterized maternal psychological distress (anxiety, depression, and perceived stress) across five gestational intervals and four postpartum time points. This approach was used to address the question: Does the quality of maternal care mitigate the link between prenatal exposure to maternal psychological distress and child cognitive function and negative emotionality?

Methods

Participants

Study participants included 136 mothers and their children (79 male, 57 female) participating in a longitudinal study evaluating the role of early experiences on development. Pregnant women were recruited from a large university medical center in Southern California. At recruitment, inclusion criteria were: (1) adult (> 18 years of age), (2) English-speaking, and (3) intrauterine, singleton pregnancy. Exclusion criteria at recruitment were: (1) presence of uterine or cervical abnormalities, (2) conditions such as endocrine, hepatic, or renal disorders, or use of corticosteroid medication, and (3) self-reported abuse of tobacco, alcohol, or recreational drugs in the pregnancy. An additional postnatal inclusion criterion for the current study was gestational age of 34 weeks or greater at birth. Descriptive information for the sample is provided in Table 1. All mothers provided informed, written consent for themselves and their child as approved by the Institutional Review Board for Protection of Human Subjects.

Procedures

Measures of maternal psychological distress were collected at five time points prenatally and four time points postnatally. Maternal sensitivity was assessed at 6 and 12 months postpartum and child cognitive and emotional outcomes were evaluated at 2 years (see

Figure 1 for study timeline). Gestational and child ages at each assessment are presented in the Appendix Table 1. Appendix Table 6 shows missing data for each measure.

Measures

Maternal Psychological Distress—Maternal anxiety was assessed using the 10-item State Anxiety subscale of the State-Trait Personality Inventory (STAI; Spielberger, 1983). Maternal depressive symptoms were assessed using the 9-item short form of the Center for Epidemiologic Studies Depression Inventory (CES-D; Santor & Coyne, 1997). Generalized or non-specific perceptions of stress were assessed using the 12-item version of Cohen's Perceived Stress Scale (PSS; Cohen, Kamarck, & Mermelstein, 1983). The internal consistencies of the three measures across time points were good (STAI $\alpha = .87 - .90$, CES-D $\alpha = .85 - .88$, PSS $\alpha = .87 - .91$ in the current sample). Mean levels of maternal depression, stress, and anxiety and distress composites at prenatal and postnatal time points are presented in the Appendix Table 2. The correlation between the three distress indicators and the pattern of association for anxiety, stress, and depression across pregnancy are presented in Appendix Tables 3-5.

Mean depression, perceived stress, and anxiety scores were standardized and averaged to create a composite prenatal distress and a composite postnatal distress score. This distress composite score was used as an index of prenatal exposure to multiple distress indicators, in order to provide a test of the ability of postnatal maternal sensitivity to compensate for prenatal maternal distress exposure. Prior research has indicated that the composite of anxiety, stress, and depression throughout pregnancy consistently predicts child outcomes (Glynn et al., 2018; Howland et al., 2020).

Quality of Maternal Care—Maternal care was characterized in the present investigation using a maternal sensitivity index developed by the NICHD early Child Care Study comprised of three indicators (a composite of positive regard, sensitivity to nondistress, and intrusiveness reverse-scored) that is a potent predictor of numerous child development outcomes (NICHD Early Child Care Research Network, 1997, 1999, 2001).

Mothers were video-recorded interacting with their infants in a semi-structured 10-minute play episode in the lab at 6 and 12 months of age. During this play interaction, mothers were given a standard set of age-appropriate toys and told to play with their infants as they would at home. Maternal behavior was scored from video using a coding system developed for the NICHD Study of Early Child Care and Youth Development (Glynn et al., 2016; NICHD Early Child Care Research Network, 1999). Based on standard procedures, a composite rating of maternal sensitivity was created by summing 4-point ratings of sensitivity to nondistress, positive regard, and intrusiveness (reverse-scored). A composite maternal sensitivity score was calculated by averaging 6- and 12-month scores. All coders were blind to other data gathered on study participants. Twenty percent of sessions were selected at random, without coder knowledge, and coded again by a second independent coder to obtain an index of inter-rater reliability. Reliability for each of the subscales were: sensitivity to nondistress (90%), intrusiveness reverse-scored (90%), and positive regard (93%).

Child Cognitive Function—The Bayley Scales of Infant Development, 2nd edition (BSID-II) Mental Developmental Index (MDI) was administered to assess child cognitive function at 2 years of age (Bayley, 1993). Examiners were directly supervised by a clinical psychologist. Interrater reliability, calculated on 15% of the assessments at each age, was 93%.

Child Negative Emotionality—Child negative emotionality at 2 years of age was assessed via maternal report on the Early Childhood Behavior Questionnaire (ECBQ; Putnam, Gartstein, & Rothbart, 2006). Prior research demonstrates that the ECBQ Negative Affectivity scale at 2 years has demonstrated inter-rater reliability and longitudinal stability (Putnam et al., 2006). Within the current sample, the Negative Affectivity scale had excellent internal consistency ($\alpha = .95$).

Measurement of Covariates

Sociodemographic Characteristics—Maternal socioeconomic status, age, cohabitation with child's father, and race and ethnicity were collected via maternal interview. Maternal socioeconomic status was calculated as the sum of standardized numbers of year of maternal education and standardized household income.

Pregnancy and Birth Outcomes—Maternal obstetric complications, parity, infant sex, birth weight, and 5-minute Apgar score were abstracted from the medical record. Estimated date of delivery was calculated utilizing both early ultrasound measures and date of last menstrual period based on ACOG guidelines, and used to assess gestational age at birth (Committee on Obstetric Practice, the American Institute of Ultrasound in Medicine, and the Society for Maternal-Fetal Medicine, 2017). An obstetric complications score was calculated, indicating the presence or absence of any pregnancy-related complications, including: prenatal infection, pregnancy-induced hypertension, gestational diabetes, oligohydramnios, polyhydramnios, preterm labor, anemia, vaginal bleeding, or placenta previa (Hobel, 1982).

Maternal Intelligence—Maternal intelligence was assessed at a subsequent visit via the Perceptual Organization Index (POI) of the Wechsler Adult Intelligence Scale (WAIS-III) once postnatally. The POI was a proxy of general intelligence, because it is highly correlated with the general factor g of intelligence ($r = .94$) (Deary, 2001). The WAIS and its subscales are widely used and valid measures of intelligence. Previous studies demonstrate the validity of the WAIS, as well as the reliability of the POI score ($r = .93$) (Wechsler et al., 1997). The WAIS-III was administered at a postnatal visit ($M = 5.34$, $SD = 1.99$ years post-delivery).

Statistical Analyses

Sociodemographic and Obstetric Covariates—Potential covariates were selected based on the literature (Blair et al., 2011; Polanska et al., 2017) and included infant characteristics (biological sex at birth, gestational age at birth, birthweight percentile adjusted for sex, and 5-minute Apgar score), as well as obstetric (obstetric health related factors, parity) and maternal characteristics (socioeconomic status, age, cohabitation with infant's father, race and ethnicity, and the WAIS, an index of intelligence). All regression

models included covariates associated ($p < .10$) with the child outcome. Thus, the following covariates were included in cognitive function analyses: maternal age, socioeconomic status, parity, cohabitation with child's father, maternal index of intelligence (POI), sum of obstetric risk, ethnicity (Non-Hispanic White or Latina), gestational age at birth (GAB), and child sex. Covariates included in negative emotionality analyses were maternal age, socioeconomic status, cohabitation with child's father, maternal index of intelligence (POI), and ethnicity (Non-Hispanic White or Latina). The correlations between all potential covariates, prenatal and postnatal distress, maternal sensitivity, and child outcomes are presented in Appendix Table 7.

Psychological Distress, Maternal Care and Child Outcomes—Initial bivariate correlations were performed to test whether prenatal and postnatal maternal psychological distress were associated with child outcomes, cognitive function and negative emotionality. The relations between the maternal sensitivity composite and child outcomes were also evaluated via bivariate correlation.

Moderation of Prenatal Psychological Distress by Maternal Caregiving—Regression models tested the primary hypothesis that the maternal sensitivity composite moderates the relation between prenatal maternal psychological distress and child outcome. Continuous predictor variables were mean-centered in the regression model. Significant interactions were probed by calculating and plotting simple slopes. Finally, to test the potential contribution of postnatal maternal psychological distress, regression models included it as an additional covariate.

Secondary Analyses: Assessment of Sex Differences, Individual Distress Indicators, and Timing and Subscales of Maternal Care

Sex differences: Secondary analyses were conducted in order to explore sex differences as follows 1) whether there are sex differences in the association between prenatal distress and child outcomes, 2) sex differences in the association between the maternal sensitivity composite and child outcomes and 3) whether the interaction between prenatal psychological distress and maternal sensitivity was moderated by child sex (three-way interaction).

Individual distress indicators: To test whether the maternal sensitivity composite moderated the effect of each of the three distress indicators, regression models were conducted separately for prenatal anxiety, stress and depression (see Appendix Tables 7-8).

Timing and subscales of maternal care: To evaluate whether care at 6 or 12 months was a more important moderator of prenatal distress (timing) interaction models were conducted with maternal sensitivity separately at 6 and 12 months (see Appendix Tables 9-10). Finally, to test the role of the individual subscales that comprise the sensitivity composite (positive regard, sensitivity to nondistress, and intrusiveness reverse-scored) in moderating the effect of prenatal maternal distress, moderation analyses were conducted separately with each of the three subscales (Appendix Tables 11-12).

Results

Child Developmental Outcomes

Descriptive information for the cognitive and emotional outcomes is shown in Table 1. Poorer child cognitive function was associated with higher child negative emotionality ($r = -.32, p < .001$).

Maternal Distress, Maternal Care and Child Cognitive Function

Elevated prenatal psychological distress composite was associated with poorer cognitive function ($r = -.28, p = .001$) and as shown in Appendix Table 5, the pattern of association was similar across the 5 gestational timepoints. Further, elevated postnatal psychological distress was associated with poorer child cognitive function at 2 years of age ($r = -.26, p = .003$). In contrast, a higher maternal sensitivity composite score was associated with enhanced child cognitive function ($r = .47, p < .001$).

Does Postnatal Care Moderate the Relation Between Prenatal Psychological Distress and Child Cognitive Function?

The maternal sensitivity composite moderated the association between prenatal distress and child cognitive function even with the inclusion of covariates ($b = 2.80, t(119) = 2.25, p = .027$; see Table 2). As shown in Figure 2, children exposed to elevated prenatal maternal distress and low maternal sensitivity exhibited the poorest cognitive performance, but children exposed to higher prenatal maternal distress who then received sensitive maternal care did not display deficits in cognitive function. This association remained when *postnatal* distress was additionally included as a covariate ($b = 2.78, t(118) = 2.22, p = .028$; see Appendix Table 13a).

Cognitive Function Secondary Analyses: Assessment of Sex Differences, Individual Distress Indicators, and Timing of Maternal Care

Sex differences: Secondary analyses revealed that child sex did not moderate the effects of prenatal psychological distress ($\beta = 0.070$) or maternal sensitivity ($\beta = 0.002$) on child cognitive function; additionally, the three-way interaction of child sex, prenatal distress, and maternal sensitivity was nonsignificant ($\beta = 0.072$) ($ps > .52$).

Individual distress indicators: The moderating effect of sensitivity on prenatal anxiety ($\beta = 0.143$), stress ($\beta = 0.166$), and depression ($\beta = 0.114$) examined separately yielded similar effect sizes (see Appendix Table 7).

Timing and subscales of maternal care: The moderating role of maternal sensitivity on child cognitive function was similar when maternal sensitivity was examined separately at 6 months ($\beta = 0.181$) and 12 months ($\beta = 0.158$) (Appendix Table 9). Additionally, the moderating effect of the individual subscales that comprise the sensitivity composite, including positive regard ($\beta = 0.154$), sensitivity to nondistress ($\beta = 0.151$), and intrusiveness reverse-scored ($\beta = 0.022$), revealed similar effect sizes (Appendix Table 11).

Maternal Distress, Maternal Care, and Child Negative Emotionality

Elevated prenatal maternal psychological distress was associated with higher child negative emotionality at 2 years of age ($r = .28, p = .002$), and as shown in Appendix Table 5, the pattern of association was similar across the 5 gestational timepoints. Elevated postnatal maternal psychological distress ($r = .25, p = .005$) was associated with higher child negative emotionality at 2 years of age. A higher maternal sensitivity composite score was associated with lower child negative emotionality ($r = -.24, p = .008$).

Does Postnatal Care Moderate the Relation between Prenatal Psychological Distress and Child Negative Emotionality?

As shown in Figure 3, children exposed to elevated prenatal distress and who received high quality maternal caregiving exhibited low negative emotionality even after consideration of covariates ($b = -0.11, t(109) = -2.01, p = .047$; see Table 3). With the addition of postnatal maternal distress as a covariate this association remained statistically significant ($b = -0.11, t(108) = -2.03, p = .045$; see Appendix Table 13b).

Negative Emotionality Secondary Analyses: Assessment of Sex Differences, Individual Distress Indicators, and Timing of Maternal Care

Sex differences: Secondary analyses revealed that child sex did not moderate the effects of prenatal psychological distress ($\beta = 0.065$) or maternal sensitivity ($\beta = -0.133$) on child negative emotionality; additionally, the three-way interaction of child sex, prenatal distress, and maternal sensitivity was nonsignificant ($\beta = 0.104$) ($ps > .29$).

Individual distress indicators: The moderating effect of sensitivity on prenatal anxiety ($\beta = -0.135$), stress ($\beta = -0.136$), and depression ($\beta = -0.165$) examined separately yielded similar effect sizes (see Appendix Table 8).

Timing and subscales of maternal care: The moderating role of maternal sensitivity on child negative emotionality was similar when maternal sensitivity was examined separately at 6 ($\beta = -0.162$) and 12 months ($\beta = -0.176$) (Appendix Table 10). Additionally, the moderating effect of the individual subscales that comprise the sensitivity composite, including positive regard ($\beta = -0.083$), sensitivity to nondistress ($\beta = -0.151$), and intrusiveness reverse-scored ($\beta = -0.166$), revealed similar effect sizes (Appendix Table 12).

Discussion

Consistent with the fetal programming literature, we show that prenatal exposure to maternal psychological distress (anxiety, depression and perceived stress) is associated with child cognitive and emotional vulnerabilities (Buss et al., 2010; Davis et al., 2007, 2019; Davis & Sandman, 2010; Glynn et al., 2018; Kingston et al., 2012; Van den Bergh et al., 2017). Further, our findings are consistent with decades of research that have established the importance of parental care during sensitive periods, such as the first postnatal year, for promoting optimal developmental outcomes (Ainsworth, 1979). The present study provides new evidence that high quality maternal care during the first postnatal year

ameliorates the negative cognitive and emotional outcomes that follow exposure to prenatal maternal distress. These findings remained after considering potential confounding factors including SES, postnatal maternal distress, and maternal intelligence scores. These data indicate that consequences of fetal programming are malleable and that prenatal and postnatal experiences synergistically impact child development. Specifically, we find that high-quality caregiving can compensate for the impact of prenatal maternal distress by altering developmental trajectories and improving child mental health.

The present study addresses a key issue in the fetal programming literature by demonstrating that the consequences of prenatal exposures are modifiable by postnatal experiences. Our data are consistent with experimental rodent models that report manipulations of maternal care compensate for prenatal adversity (Bogoch et al., 2007; Lemaire et al., 2006; Raine et al., 2014; Wakshlak & Weinstock, 1990) as well as studies showing that infants with a secure attachment relationship to their mother do not show behavioral problems following prenatal stress (Ali et al., 2020; Bergman et al., 2008). There are several important contributions of our project to the relatively small human literature evaluating whether maternal behavior towards her infant mitigates the consequences of prenatal stress (Grant et al., 2010a, 2010b; Kaplan et al., 2008; Schechter et al., 2017; Sharp et al., 2012, 2015). First, maternal distress was assessed repeatedly throughout gestation and we have shown previously that this composite measure of anxiety, stress, and depression symptoms predicts both cognitive and emotional outcomes through childhood and adolescence (Glynn et al., 2018; Howland et al., 2020). Second, we assessed a composite of maternal sensitivity twice during the first postnatal year, a sensitive window for attachment formation (Ainsworth, 1979). Thus, our design is a rigorous test of the hypothesis that high-quality maternal care can offset the impact of prenatal distress exposure after covarying for confounding factors including postnatal maternal distress. Third, we characterize both child cognitive and emotional function. Finally, these data provide evidence that infancy may be a sensitive window when maternal care can mitigate the consequences of prenatal maternal distress, thereby identifying a plausible target for early intervention following prenatal adversity.

Maternal care that is warm, sensitive and responsive to infant's signals (Ainsworth et al., 1978), may play a particularly important role in child neurodevelopment and long-term developmental outcomes (Malmberg et al., 2016; Spinrad & Stifter, 2002; Wang et al., 2019). Thus, it is highly plausible that high quality maternal care during the first postnatal year may alter neurodevelopmental trajectories following prenatal exposures towards more optimal outcomes. We show that maternal care at 6 and 12 months similarly moderated the association between prenatal maternal distress and child outcomes, suggesting that care throughout the first postnatal year may be important for ameliorating the impact of prenatal distress (Appendix Tables 9-10). The maternal sensitivity composite in the current study measured positive regard, sensitivity to nondistress, and intrusiveness reverse-scored, key aspects of maternal behavior the influence development. Findings suggest that all of these components of maternal sensitivity captured by this measure contribute to positive child development following prenatal maternal distress (See Appendix Tables 11-12). Future research could consider other aspects of positive parenting and paternal parenting behaviors that may ameliorate the impact of prenatal maternal distress.

There are a number of potential mechanisms by which prenatal maternal psychological distress may have consequences for child cognitive and emotional development, including alterations to fetal neurodevelopment (Sandman, Class, et al., 2015; Sandman et al., 2016; Schuurmans & Kurrasch, 2013; Wu et al., 2020). Animal studies demonstrate changes in offspring brain structure following prenatal stress exposure, such as reduced hippocampal volume and neurogenesis (Bogoch et al., 2007; Charil et al., 2010). Children exposed to prenatal maternal psychological distress show reduced gray matter volume and thickness in frontal, temporal, and limbic areas, as well as reduced total gray matter density (Adamson et al., 2018; Buss et al., 2010; Davis et al., 2019; Demers et al., 2020; Sandman, Buss, et al., 2015). Cross-species studies provide mechanistic evidence that dendritic atrophy may be a pathway by which prenatal maternal distress disrupts offspring brain development (Curran et al., 2017; Sandman et al., 2018). These neural systems impacted by prenatal exposures may be modifiable by high-quality postnatal maternal care. The first year postpartum continues to be a sensitive window of heightened neuroplasticity and rapid neural growth (Gee, 2016; Gilmore et al., 2018; Knickmeyer et al., 2008) and maternal care may modify developmental trajectories. Consistent with this possibility, high quality maternal care is associated with enhanced child hippocampal volume growth; this growth trajectory is further associated with improved child emotion regulation (Luby et al., 2017) as well as with greater child gray matter volume at 8 years (Kok et al., 2015). Experimental animal and cross species research provide strong evidence that maternal care directly impacts neural circuits underlying cognitive and emotional vulnerabilities that are impacted by prenatal maternal distress (Gee, 2016; Granger et al., 2021; Liu et al., 2000; Rao et al., 2010).

The current study has several limitations. Although the use of a community sample highlights the importance of subclinical variation in maternal psychological symptoms of distress, the range of observed maternal psychological distress and maternal sensitivity is constrained. Because of this, the current study may underestimate the associations among prenatal distress, maternal sensitivity, and child outcomes. We also cannot completely disentangle the role of genetic influences. However, evidence from experimental animal research consistently reports the effects of maternal care on offspring outcomes are independent from effects of genetics, and provides experimental evidence that maternal care can compensate for prenatal exposures (Francis et al., 1999; Liu et al., 2000).

Decades of research have confirmed the prenatal period as a time of enhanced responsiveness to environmental input, when maternal psychological distress can have a profound influence on child development (Bush et al., 2017; Davis et al., 2007, 2018; Davis & Sandman, 2010, 2012; Doyle et al., 2015; Van den Bergh et al., 2017; Vehmeijer et al., 2019). Few studies have directly assessed processes that promote resilience following prenatal adversity (Atzl et al., 2019; D'Anna-Hernandez & Rivera, 2014; Davis & Narayan, in press; Rosand et al., 2011). Our findings, coupled now with those from Schechter et al. (2017), provide strong support for postnatal prevention and intervention efforts to reduce the consequences of prenatal adversity. Efficacious interventions exist to support the transition to parenthood, and the development of positive parenting skills (Bick & Dozier, 2010; Eshel et al., 2006; Nillni et al., 2018). Thus, interventions to promote maternal psychological health and sensitive, responsive caregiving has enduring benefits that could thus mitigate the life-long cognitive and emotional consequences of prenatal psychological distress.

Acknowledgements:

We wish to thank the families who participated in this project. The assistance of Megan Faulkner, Natalie Hernandez, and Kendra Leak of the Women and Children's Health and Well-Being Project, Department of Psychiatry & Human Behavior, University of California and the Department of Psychology, Chapman University is gratefully acknowledged. This research was supported by the National Institutes of Health [R01 HD065823; P50 MH096889; R03 MH86062; R01 HD51852; R01 NS041298].

Appendix

Appendix Table 1

Mean Fetal/Child Age at Assessment

	Prenatal (Gestational Weeks)					Postnatal (Months)				
	15	19	25	31	36	3	6	12	24	Composite
Mean Fetal/Child Age	15.63 (1.16)	19.74 (1.06)	25.74 (1.03)	31.07 (0.89)	36.83 (0.93)	3.10 (0.50)	6.06 (0.34)	12.12 (0.32)	24.17 (0.41)	

Values presented as Means (SD)

Appendix Table 2

Mean Levels of Maternal Depression, Anxiety, Stress, and Distress Composites Across Prenatal and Postnatal Time points.

	Prenatal (Gestational Weeks)						Postnatal (Months)				
	15	19	25	31	36	Composite	3	6	12	24	Composite
CES-D	5.06 (4.44)	4.44 (4.05)	4.91 (4.64)	5.53 (4.42)	6.50 (5.17)	5.21 (3.84)	4.02 (3.95)	4.05 (4.48)	3.77 (4.49)	4.29 (4.63)	4.02 (3.57)
STAI	17.07 (5.34)	16.84 (5.25)	16.46 (5.27)	16.86 (5.39)	17.35 (5.71)	16.95 (4.56)	16.81 (4.68)	16.12 (4.98)	16.65 (5.31)	16.73 (5.51)	16.60 (4.19)
PSS	11.39 (6.43)	10.45 (5.83)	10.53 (6.97)	10.31 (6.78)	10.96 (6.43)	10.72 (5.66)	10.83 (6.56)	9.93 (6.30)	9.86 (5.94)	10.81 (6.28)	10.40 (5.20)

Values presented as Means (SD)

Appendix Table 3

Correlations between Prenatal Stress, Depression, and Anxiety Composites and Child Cognitive Function and Negative Emotionality

Variable	1	2	3	4	5
1. Depression Composite	—				
2. Stress Composite	.792 ^{**}	—			
3. Anxiety Composite	.842 ^{**}	.792 ^{**}	—		
4. Cognitive Function	-.242 ^{**}	-.300 ^{**}	-.245 ^{**}	—	
5. Negative Emotionality	.259 ^{**}	.315 ^{**}	.210 [*]	-.323 ^{**}	—

* $p < .05$.

** $p < .01$.

Appendix Table 4a

Correlations Between Depression (CESD) at Each Prenatal Timepoint, Prenatal Psychological Distress, Child Cognitive Function, and Child Negative Emotionality

Variable	1	2	3	4	5	6	7	8
1. CESD 15 wks	—	.746**	.661**	.625**	.533**	.792**	-.234**	.289**
2. CESD 19 wks	.746**	—	.669**	.654**	.575**	.811**	-0.137	.241**
3. CESD 25 wks	.661**	.669**	—	.579**	.638**	.815**	-.252**	.274**
4. CESD 31 wks	.625**	.654**	.579**	—	.665**	.774**	-.214*	0.143
5. CESD 36 wks	.533**	.575**	.638**	.665**	—	.796**	-.207*	.227*
6. Prenatal Psychological Distress	.792**	.811**	.815**	.774**	.796**	—	-.281**	.281**
7. Cognitive Function 2 yrs	-.234**	-0.137	-.252**	-.214*	-.207*	-.281**	—	-.323**
8. Negative Emotionality 2yrs	.289**	.241**	.274**	0.143	.227*	.281**	-.323**	—

* $p < .05$.** $p < .01$.**Appendix Table 4b**

Correlations Between Anxiety (STAI) at Each Prenatal Timepoint, Prenatal Psychological Distress Composite, Child Cognitive Function, and Child Negative Emotionality

Variable	1	2	3	4	5	6	7	8
1. STAI 15 wks	—	.744**	.626**	.637**	.668**	.824**	-.244**	0.165
2. STAI 19 wks	.744**	—	.620**	.620**	.598**	.788**	-0.113	0.148
3. STAI 25 wks	.626**	.620**	—	.603**	.679**	.796**	-0.167	.189*
4. STAI 31 wks	.637**	.620**	.603**	—	.719**	.792**	-.241**	.257**
5. STAI 36 wks	.668**	.598**	.679**	.719**	—	.801**	-.266**	0.143
6. Prenatal Psychological Distress	.824**	.788**	.796**	.792**	.801**	—	-.281**	.281**
7. Cognitive Function 2 yrs	-.244**	-0.113	-0.167	-.241**	-.266**	-.281**	—	-.323**
8. Negative Emotionality 2 yrs	0.165	0.148	.189*	.257**	0.143	.281**	-.323**	—

* $p < .05$.** $p < .01$.**Appendix Table 4c**

Correlations Between Perceived Stress (PSS) at Each Prenatal Timepoint, Prenatal Psychological Distress Composite, Child Cognitive Function, and Child Negative Emotionality

Variable	1	2	3	4	5	6	7	8
1. PSS 15 wks	—	.747**	.660**	.706**	.630**	.802**	-.289**	.323**
2. PSS 19 wks	.747**	—	.696**	.707**	.625**	.796**	-.220*	.307**

Variable	1	2	3	4	5	6	7	8
3. PSS 25 wks	.660**	.696**	—	.725**	.690**	.816**	-.266**	.287**
4. PSS 31 wks	.706**	.707**	.725**	—	.739**	.820**	-.260**	.334**
5. PSS 36 wks	.630**	.625**	.690**	.739**	—	.786**	-.236**	0.134
6. Prenatal Psychological Distress	.802**	.796**	.816**	.820**	.786**	—	-.281**	.281**
7. Cognitive Function 2 yrs	-.289**	-.220*	-.266**	-.260**	-.236**	-.281**	—	-.323**
8. Negative Emotionality 2 yrs	.323**	.307**	.287**	.334**	0.134	.281**	-.323**	—

* $p < .05$.

** $p < .01$.

Appendix Table 5

Correlations for Psychological Distress at Each Timepoint

Variable	1	2	3	4	5	6	7
1. Distress 15 wks	—						
2. Distress 19 wks	.804**	—					
3. Distress 25 wks	.714**	.717**	—				
4. Distress 31 wks	.742**	.724**	.718**	—			
5. Distress 36 wks	.701**	.666**	.722**	.776**	—		
6. Cognitive Function 2yrs	-.295**	-.173*	-.251**	-.267**	-.263**	—	
7. Negative Emotionality 2yrs	.300**	.248**	.269**	.280**	0.161	-.323**	—

* $p < .05$.

** $p < .01$.

Appendix Table 6

Percent Missing Data for Each Timepoint

		Timepoint	N	% Missing Data
Depression (CESD)	Prenatal	15 weeks	125	11
		19 weeks	135	1
		25 weeks	135	1
		31 weeks	135	1
		36 weeks	98	38*
	Postnatal	3 months	128	8
		6 months	129	7
		1 year	128	8
		2 years	132	4
Perceived Stress (PSS)	Prenatal	15 weeks	131	5
		19 weeks	134	2
		25 weeks	135	1
		31 weeks	135	1

		Timepoint	N	% Missing Data
	Postnatal	36 weeks	129	7
		3 months	129	7
		6 months	131	5
		1 year	130	6
		2 years	135	1
Anxiety (STAI)	Prenatal	15 weeks	124	12
		19 weeks	135	1
		25 weeks	134	2
		31 weeks	135	1
		36 weeks	127	9
	Postnatal	3 months	129	7
		6 months	130	6
		1 year	130	6
		2 years	134	2
Maternal Sensitivity Composite		6 months	126	10
		1 year	125	11

* Due to an administrative/data collection error, some participants were not administered CESD at 36 weeks

Appendix Table 7

Correlations between Sociodemographic Covariates, Prenatal Distress, Maternal Care, and Child Outcomes

Variable	1	2	3	4	5	6	7	8	9	10	11	12
1. Gestational Age at Birth (GAB)	—	-0.082	0.156	-.192 [*]	-0.051	0.052	-0.107	-0.065	-0.080	0.036	0.009	0.063
2. Maternal Age	-0.082	—	.589 ^{**}	.190 [*]	.197 [*]	.316 ^{**}	-0.045	.239 ^{**}	-.279 ^{**}	0.145	0.160	-0.108
3. SES	0.156	.589 ^{**}	—	-.203 [*]	.353 ^{**}	.556 ^{**}	-0.075	.250 ^{**}	-.323 ^{**}	-0.033	0.107	-0.027
4. Parity	-.192 [*]	.190 [*]	-.203 [*]	—	0.123	-.297 ^{**}	0.061	-.184 [*]	.218 [*]	0.073	0.033	0.073
5. Cohabitation	-0.051	.197 [*]	.353 ^{**}	0.123	—	0.159	-0.010	-0.020	0.045	-0.023	.184 [*]	-0.097
6. Index of Intelligence (WAIS)	0.052	.316 ^{**}	.556 ^{**}	-.297 ^{**}	0.159	—	-.305 ^{**}	.364 ^{**}	-.450 ^{**}	-0.005	0.030	0.000
7. Obstetric Risk	-0.107	-0.045	-0.075	0.061	-0.010	-.305 ^{**}	—	-.235 ^{**}	.221 ^{**}	-0.040	-0.060	0.011
8. Non-Hispanic White	-0.065	.239 ^{**}	.250 ^{**}	-.184 [*]	-0.020	.364 ^{**}	-.235 ^{**}	—	-.655 ^{**}	0.027	0.054	-0.046
9. Latina	-0.080	-.279 ^{**}	-.323 ^{**}	.218 [*]	0.045	-.450 ^{**}	.221 ^{**}	-.655 ^{**}	—	-0.106	-0.060	0.105
10. Child Sex (male)	0.036	0.145	-0.033	0.073	-0.023	-0.005	-0.040	0.027	-0.106	—	0.110	-0.075

Variable	1	2	3	4	5	6	7	8	9	10	11	12
11. Apgar Score at 5 min	0.009	0.160	0.107	0.033	.184 [*]	0.030	-0.060	0.054	-0.060	0.110	—	-0.007
12. Birthweight Percentile by Sex	0.063	-0.108	-0.027	0.073	-0.097	0.000	0.011	-0.046	0.105	-0.075	-0.007	—
13. Prenatal Distress	-0.130	-.295 ^{**}	-.450 ^{**}	-0.037	-.422 ^{**}	-.317 ^{**}	-0.001	0.031	-0.069	0.069	0.009	-0.007
14. Postnatal Distress	-0.156	-.206 [*]	-.401 ^{**}	-0.010	-.301 ^{**}	-.191 [*]	-0.094	0.086	-0.093	0.100	0.009	-0.045
15. Maternal Sensitivity Composite	.196 [*]	.287 ^{**}	.496 ^{**}	-.352 ^{**}	0.145	.461 ^{**}	-0.167	.253 ^{**}	-.333 ^{**}	-0.015	0.017	-0.050
16. Cognitive Function	0.163	.238 ^{**}	.500 ^{**}	-.279 ^{**}	.202 [*]	.474 ^{**}	-0.165	.220 [*]	-.329 ^{**}	-.199 [*]	0.030	-0.019
17. Negative Emotionality	-0.101	-.298 ^{**}	-.367 ^{**}	0.098	-.197 [*]	-.352 ^{**}	0.105	-.355 ^{**}	.331 ^{**}	0.019	-0.016	-0.034

* $p < .05$.** $p < .01$.

Appendix Tables 7-8: Analyses using Individual Distress Subscales (Depressive Symptoms, Anxiety Symptoms, & Perceived Stress)

Appendix Table 7a

Regression Models Examining the Association Between Prenatal Maternal Depressive Symptoms and Child Cognitive Function, Including Postnatal Maternal Depressive Symptoms

Variable	Model 1: Prenatal Depression			Model 2: Prenatal Depression and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.084	0.336	0.025	-0.020	0.335	-0.006	0.003	0.333	0.001
SES	2.147 [†]	1.158	0.224	1.748	1.157	0.183	1.834	1.152	0.191
Parity	-2.381	1.742	-0.119	-1.494	1.768	-0.075	-1.568	1.758	-0.078
Cohabitation with Child's Father	4.975	4.785	0.090	4.349	4.726	0.079	2.723	4.816	0.050
Index of Intelligence (WAIS)	0.185 [†]	0.096	0.194	0.161 [†]	0.096	0.169	0.145	0.096	0.152
Obstetric Risk	-1.812	2.836	-0.050	-1.692	2.796	-0.046	-1.768	2.780	-0.049
Non-Hispanic White	-1.495	3.220	-0.046	-1.689	3.176	-0.052	-2.145	3.171	-0.066
Latina	-6.286 [†]	3.705	-0.177	-5.641	3.665	-0.158	-6.261 [†]	3.666	-0.176
Gestational Age at Birth (GAB)	0.871	1.023	0.067	0.644	1.014	0.049	0.540	1.011	0.041
Child Sex (male)	-6.490 [*]	2.485	-0.196	-6.253 [*]	2.452	-0.189	-6.360 [*]	2.439	-0.192
Postnatal Depression	-0.310	0.523	-0.068	-0.300	0.515	-0.066	-0.334	0.513	-0.073

Variable	Model 1: Prenatal Depression			Model 2: Prenatal Depression and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Prenatal Depression	0.042	0.551	0.010	0.035	0.543	0.008	0.033	0.540	0.008
Sensitivity Composite				2.615*	1.235	0.191	2.817*	1.235	0.205
Prenatal Depression * Sensitivity Composite							0.494	0.320	0.114
<i>R</i> ²	0.378			0.401			0.413		
<i>F</i>	6.085***			6.125***			5.923***		

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 7b

Regression Models Examining the Association Between Prenatal Maternal Anxiety Symptoms and Child Cognitive Function, including Postnatal Maternal Anxiety Symptoms

Variable	Model 1: Prenatal Anxiety			Model 2: Prenatal Anxiety and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.058	0.339	0.017	-0.051	0.339	-0.015	-0.050	0.335	-0.015
SES	2.205 [†]	1.146	0.230	1.835	1.144	0.192	1.879 [†]	1.130	0.196
Parity	-2.432	1.732	-0.121	-1.532	1.762	-0.076	-1.410	1.741	-0.070
Cohabitation with Child's Father	4.967	4.587	0.090	4.291	4.535	0.078	3.578	4.493	0.065
Index of Intelligence (WAIS)	0.179 [†]	0.096	0.188	0.155	0.095	0.163	0.150	0.094	0.158
Obstetric Risk	-1.812	2.844	-0.050	-1.700	2.805	-0.047	-1.731	2.771	-0.048
Non-Hispanic White	-1.532	3.254	-0.047	-1.820	3.212	-0.056	-2.132	3.176	-0.065
Latina	-6.353 [†]	3.716	-0.178	-5.688	3.679	-0.160	-5.428	3.635	-0.152
Gestational Age at Birth (GAB)	0.846	1.041	0.065	0.627	1.032	0.048	0.354	1.028	0.027
Child Sex (male)	-6.454*	2.487	-0.195	-6.263*	2.454	-0.189	-6.627**	2.431	-0.200
Postnatal Anxiety	-0.195	0.489	-0.050	-0.107	0.484	-0.027	-0.119	0.478	-0.031
Prenatal Anxiety	-0.018	0.467	-0.005	-0.084	0.462	-0.024	-0.097	0.456	-0.027

Variable	Model 1: Prenatal Anxiety			Model 2: Prenatal Anxiety and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Sensitivity Composite				2.601*	1.241	0.190	2.704*	1.226	0.197
Prenatal Anxiety * Sensitivity Composite							0.470*	0.235	0.143
<i>R</i> ²		0.377			0.400			0.419	
<i>F</i>		6.063***			6.094***			6.086***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 7c

Regression Models Examining the Association Between Prenatal Maternal Perceived Stress and Child Cognitive Function, Including Postnatal Maternal Perceived Stress

Variable	Model 1: Prenatal Perceived Stress			Model 2: Prenatal Perceived Stress and Sensitivity Composite			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.066	0.335	0.020	-0.038	0.336	-0.011	-0.012	0.330	-0.004
SES	1.929 [†]	1.140	0.201	1.632	1.138	0.170	1.713	1.119	0.179
Parity	-2.561	1.720	-0.128	-1.736	1.755	-0.087	-1.704	1.724	-0.085
Cohabitation with Child's Father	4.476	4.501	0.081	3.932	4.461	0.072	2.283	4.441	0.042
Index of Intelligence (WAIS)	0.144	0.099	0.151	0.125	0.098	0.132	0.126	0.097	0.133
Obstetric Risk	-2.455	2.843	-0.067	-2.209	2.815	-0.061	-2.248	2.766	-0.062
Non-Hispanic White	-2.132	3.195	-0.065	-2.277	3.161	-0.070	-2.956	3.120	-0.091
Latina	-7.454 [†]	3.860	-0.209	-6.802 [†]	3.833	-0.191	-7.292 [†]	3.772	-0.205
Gestational Age at Birth (GAB)	0.910	1.007	0.069	0.696	1.002	0.053	0.660	0.984	0.050
Child Sex (male)	-6.443*	2.452	-0.194	-6.257*	2.427	-0.189	-6.943**	2.404	-0.209
Postnatal Stress	-0.356	0.389	-0.114	-0.231	0.390	-0.074	-0.151	0.385	-0.049
Prenatal Stress	-0.094	0.395	-0.033	-0.155	0.392	-0.054	-0.152	0.386	-0.053
Sensitivity Composite				2.393 [†]	1.248	0.174	2.496*	1.227	0.182
Prenatal Stress * Sensitivity Composite							0.444*	0.193	0.166
<i>R</i> ²		0.389			0.408			0.433	
<i>F</i>		6.376***			6.299***			6.436***	

B = unstandardized coefficient. β = standardized coefficient.

[†]
 $p < .10$
 *
 $p < .05$.
 **
 $p < .01$.

 $p < .001$

Appendix Table 8a

Regression Models Examining the Association Between Prenatal Maternal Depressive Symptoms and Child Negative Emotionality, Including Postnatal Maternal Depressive Symptoms

Variable	Model 1: Prenatal Depression		Model 2: Prenatal Depression and Sensitivity Composite				Model 3: Interaction		
	B	SE B	B	B	SE B	β	B	SE B	B
Maternal Age	-0.016	0.011	-0.134	-0.016	0.011	-0.134	-0.018	0.011	-0.155
SES	-0.009	0.041	-0.027	-0.007	0.042	-0.020	-0.008	0.042	-0.023
Cohabitation with Child's Father	-0.160	0.194	-0.074	-0.162	0.195	-0.074	-0.104	0.195	-0.048
Index of Intelligence (WAIS)	-0.005	0.003	-0.136	-0.004	0.004	-0.132	-0.004	0.003	-0.112
Non-Hispanic White	-0.292*	0.127	-0.253	-0.293*	0.128	-0.253	-0.277*	0.126	-0.239
Latina	0.128	0.149	0.099	0.124	0.151	0.096	0.169	0.150	0.131
Postnatal Depression	0.028	0.019	0.174	0.028	0.019	0.175	0.030	0.019	0.184
Prenatal Depression	0.015	0.021	0.091	0.014	0.021	0.088	0.015	0.021	0.090
Sensitivity Composite				-0.011	0.047	-0.023	-0.013	0.047	-0.025
Prenatal Depression * Sensitivity Composite							-0.029*	0.015	-0.165
R^2		0.298			0.298			0.324	
F		5.834***			5.148***			5.168***	

B = unstandardized coefficient. β = standardized coefficient.

[†]
 $p < .10$
 *
 $p < .05$.
 **
 $p < .01$.

 $p < .001$

Appendix Table 8b

Regression Models Examining the Association Between Prenatal Maternal Anxiety Symptoms and Child Negative Emotionality, Including Postnatal Maternal Anxiety Symptoms

Variable	Model 1: Prenatal Anxiety		Model 2: Prenatal Anxiety and Sensitivity Composite				Model 3: Interaction		
	B	SE B	B	B	SE B	β	B	SE B	β
Maternal Age	-0.013	0.011	-0.114	-0.013	0.011	-0.114	-0.015	0.011	-0.126
SES	-0.023	0.041	-0.067	-0.021	0.042	-0.060	-0.021	0.042	-0.061
Cohabitation with Child's Father	-0.258	0.192	-0.119	-0.259	0.193	-0.119	-0.250	0.191	-0.115

Variable	Model 1: Prenatal Anxiety			Model 2: Prenatal Anxiety and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Index of Intelligence (WAIS)	-0.005	0.003	-0.145	-0.005	0.004	-0.140	-0.005	0.004	-0.138
Non-Hispanic White	-0.272*	0.129	-0.235	-0.273*	0.129	-0.236	-0.260*	0.129	-0.225
Latina	0.128	0.152	0.099	0.123	0.154	0.095	0.123	0.152	0.095
Postnatal Anxiety	0.028	0.018	0.201	0.028	0.018	0.200	0.031 [†]	0.018	0.217
Prenatal Anxiety	-0.006	0.017	-0.049	-0.007	0.017	-0.052	-0.009	0.017	-0.067
Sensitivity Composite				-0.013	0.048	-0.026	-0.009	0.048	-0.017
Prenatal Anxiety * Sensitivity Composite							-0.016	0.010	-0.135
<i>R</i> ²		0.276			0.276			0.294	
<i>F</i>		5.240***			4.626***			4.499***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 8c

Regression Models Examining the Association Between Prenatal Maternal Perceived Stress and Child Negative Emotionality, Including Postnatal Maternal Perceived Stress

Variable	Model 1: Prenatal Perceived Stress			Model 2: Prenatal Perceived Stress and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.013	0.011	-0.108	-0.013	0.011	-0.109	-0.014	0.011	-0.117
SES	-0.017	0.040	-0.050	-0.017	0.041	-0.049	-0.020	0.041	-0.056
Cohabitation with Child's Father	-0.184	0.189	-0.085	-0.184	0.190	-0.085	-0.136	0.191	-0.062
Index of Intelligence (WAIS)	-0.002	0.004	-0.072	-0.002	0.004	-0.072	-0.003	0.004	-0.075
Non-Hispanic White	-0.222 [†]	0.126	-0.192	-0.223 [†]	0.127	-0.192	-0.196	0.127	-0.170
Latina	0.216	0.157	0.167	0.215	0.159	0.166	0.247	0.159	0.191
Postnatal Stress	0.013	0.016	0.112	0.013	0.016	0.112	0.012	0.016	0.102
Prenatal Stress	0.017	0.016	0.155	0.017	0.016	0.155	0.016	0.016	0.149
Sensitivity Composite				-0.002	0.048	-0.005	0.004	0.047	0.008
Prenatal Stress * Sensitivity Composite							-0.014	0.009	-0.136

Variable	Model 1: Prenatal Perceived Stress			Model 2: Prenatal Perceived Stress and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
R^2		0.297			0.297			0.314	
<i>F</i>		5.810 ***			5.118 ***			4.946 ***	

B = unstandardized coefficient. β = standardized coefficient.

[†] $p < .10$

* $p < .05$.

** $p < .01$.

*** $p < .001$

Appendix Tables 9-10: Analyses of Maternal Care at 6 and 12 months

Appendix Table 9a

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Cognitive Function, with Maternal Care at 6 months

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite at 6 months			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.050	0.349	0.015	0.014	0.352	0.004	0.051	0.346	0.015
SES	2.102 [†]	1.160	0.221	1.973 [†]	1.172	0.207	2.024 [†]	1.150	0.212
Parity	-2.224	1.807	-0.110	-2.208	1.810	-0.110	-2.589	1.784	-0.128
Cohabitation	5.439	4.792	0.098	5.687	4.807	0.102	4.796	4.735	0.086
Index of Intelligence (WAIS)	0.175 [†]	0.099	0.186	0.158	0.101	0.167	0.138	0.100	0.146
Obstetric Risk	-2.557	2.995	-0.069	-2.561	2.999	-0.069	-4.123	3.023	0.111
Non-Hispanic White	-2.660	3.363	-0.081	-2.727	3.368	-0.083	-3.746	3.336	-0.114
Latina	-6.908 [†]	3.925	-0.191	-6.810 [†]	3.932	-0.188	-7.987*	3.894	-0.221
Gestational Age at Birth (GAB)	0.294	1.080	0.022	0.092	1.108	0.007	-0.229	1.097	-0.017
Child Sex (male)	-5.719*	2.586	-0.171	-5.376*	2.621	-0.161	-6.271*	2.602	-0.188
Prenatal Distress	-1.798	1.686	-0.102	-1.993	1.704	-0.113	-2.467	1.686	-0.140
Sensitivity Composite 6mo				1.135	1.346	0.074	1.579	1.336	0.102
Prenatal Distress * Sensitivity Composite 6mo							3.886*	1.710	0.181
R^2		0.383			0.387			0.414	
<i>F</i>		6.252 ***			5.775 ***			5.930 ***	

B = unstandardized coefficient. β = standardized coefficient.

[†] $p < .10$

*
 $p < .05$.
 **
 $p < .01$.

 $p < .001$

Appendix Table 9b

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Cognitive Function, with Maternal Care at 12 months

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite at 12 months			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.159	0.385	0.047	0.152	0.377	0.045	0.249	0.375	0.074
SES	1.913	1.221	0.198	1.419	1.216	0.147	1.411	1.198	0.146
Parity	-2.659	1.909	-0.130	-1.414	1.945	-0.069	-1.576	1.920	-0.077
Cohabitation	4.206	4.870	0.073	2.621	4.822	0.045	0.560	4.860	0.010
Index of Intelligence (WAIS)	0.193 [†]	0.103	0.195	0.185 [†]	0.101	0.186	0.171 [†]	0.100	0.172
Obstetric Risk	0.021	3.157	0.001	0.520	3.102	0.014	1.108	3.071	0.029
Non-Hispanic White	-2.254	3.396	-0.068	-2.449	3.330	-0.074	-2.528	3.283	-0.076
Latina	-7.350 [†]	3.901	-0.204	-6.623 [†]	3.837	-0.184	-6.660 [†]	3.783	-0.185
Gestational Age at Birth (GAB)	0.703	1.140	0.049	0.671	1.117	0.047	0.531	1.104	0.037
Child Sex (male)	-7.679 ^{**}	2.637	-0.227	-7.841 ^{**}	2.586	-0.232	-7.982 ^{**}	2.550	-0.236
Prenatal Distress	-1.551	1.757	-0.082	-0.913	1.744	-0.048	-0.738	1.721	-0.039
Sensitivity Composite 12mo				2.141	0.915 [*]	0.215	1.950 [*]	0.907	0.196
Prenatal Distress * Sensitivity Composite 12mo							1.844 [*]	0.905	0.158
<i>R</i> ²		0.376			0.406			0.428	
<i>F</i>		6.027 ^{***}			6.207 ^{***}			6.215 ^{***}	

B = unstandardized coefficient. β = standardized coefficient.

[†]
 $p < .10$
 *
 $p < .05$.
 **
 $p < .01$.

 $p < .001$

Appendix Table 10a

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Negative Emotionality, with Maternal Care at 6 months

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite at 6 months			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.019	0.012	-0.156	-0.018	0.012	-0.150	-0.017	0.012	-0.142
SES	-0.008	0.041	-0.023	-0.001	0.042	-0.002	-0.004	0.041	-0.010
Cohabitation	-0.225	0.202	-0.099	-0.259	0.205	-0.115	-0.270	0.202	-0.119
Index of Intelligence (WAIS)	-0.004	0.003	-0.117	-0.004	0.004	-0.103	-0.004	0.003	-0.107
Non-Hispanic White	-0.234 [†]	0.130	-0.199	-0.248 [†]	0.131	-0.211	-0.239 [†]	0.130	-0.203
Latina	0.140	0.157	0.105	0.112	0.160	0.084	0.132	0.158	0.099
Prenatal Distress	0.159 [*]	0.062	0.249	0.158 [*]	0.062	0.247	0.156 [*]	0.061	0.244
Sensitivity Composite 6mo				-0.052	0.053	-0.090	-0.076	0.054	-0.132
Prenatal Distress [*] Sensitivity Composite 6mo							-0.131 [†]	0.069	-0.162
<i>R</i> ²		0.308			0.314			0.338	
<i>F</i>		6.536 ^{***}			5.838 ^{***}			5.726 ^{***}	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

^{*} *p* < .05.

^{**} *p* < .01.

^{***} *p* < .001

Appendix Table 10b

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Negative Emotionality, with Maternal Care at 12 months

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite at 12 mo			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.009	0.012	-0.073	-0.007	0.012	-0.063	-0.012	0.012	-0.099
SES	-0.026	0.042	-0.074	-0.032	0.044	-0.089	-0.030	0.043	-0.085
Cohabitation	-0.180	0.198	-0.080	-0.189	0.199	-0.084	-0.120	0.199	-0.053
Index of Intelligence (WAIS)	-0.004	0.004	-0.103	-0.004	0.004	-0.109	-0.003	0.004	-0.093
Non-Hispanic White	-0.265 [*]	0.131	-0.226	-0.268 [*]	0.132	-0.228	-0.25 [†]	0.130	-0.214
Latina	0.203	0.156	0.155	0.208	0.157	0.159	0.242	0.156	0.185

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite at 12 mo			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Prenatal Distress	0.147*	0.065	0.217	0.153*	0.066	0.226	0.159*	0.066	0.234
Sensitivity Composite 12mo				0.018	0.035	0.049	0.034	0.035	0.093
Prenatal Distress * Sensitivity Composite 12mo							-0.076*	0.038	-0.176
<i>R</i> ²		0.305			0.307			0.334	
<i>F</i>		6.399***			5.593***			5.564***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Tables 11-12: Maternal Sensitivity Subscale Analyses (Sensitivity to Nondistress, Intrusiveness Reverse-Scored, Positive Regard)

Appendix Table 11a

Regression Model Examining Maternal Sensitivity to Nondistress as a Moderator of the Association Between Prenatal Distress and Child Cognitive Function

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity to Nondistress			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.038	0.334	0.011	-0.104	0.330	-0.031	-0.093	0.326	-0.028
SES	2.159 [†]	1.139	0.225	1.627	1.129	0.170	1.635	1.113	0.171
Parity	-2.503	1.728	-0.125	-1.459	1.730	-0.073	-1.481	1.706	-0.074
Cohabitation with Child's Father	4.281	4.612	0.078	3.678	4.505	0.067	1.888	4.522	0.034
Index of Intelligence (WAIS)	0.166 [†]	0.097	0.174	0.139	0.095	0.146	0.129	0.093	0.135
Obstetric Risk	-1.925	2.831	-0.053	-1.435	2.768	-0.039	-1.242	2.731	-0.034
Non-Hispanic White	-1.764	3.191	-0.054	-1.435	3.116	-0.044	-1.921	3.081	-0.059
Latina	-6.728 [†]	3.747	-0.189	-5.478	3.686	-0.154	-5.920	3.640	-0.166
Gestational Age at Birth (GAB)	0.863	1.017	0.066	0.632	0.996	0.048	0.464	0.985	0.035
Child Sex (male)	-6.473*	2.464	-0.195	-5.463*	2.434	-0.165	-5.965*	2.411	-0.180
Prenatal Distress	-1.376	1.616	-0.079	-1.324	1.577	-0.076	-1.403	1.555	-0.080

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity to Nondistress			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Sensitivity to Nondistress				7.622**	2.857	0.241	8.227**	2.831	0.260
Prenatal Distress* Sensitivity to Nondistress							5.485*	2.598	0.151
<i>R</i> ²		0.379			0.414			0.435	
<i>F</i>		6.715***		7	.060***			7.048***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 11b

Regression Model Examining Maternal Intrusiveness Reverse-Scored as a Moderator of the Association Between Prenatal Distress and Child Cognitive Function

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Intrusiveness Reverse-Scored			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.038	0.334	0.011	0.008	0.333	0.002	0.002	0.335	0.001
SES	2.159 [†]	1.139	0.225	1.916 [†]	1.145	0.200	1.956 [†]	1.158	0.204
Parity	-2.503	1.728	-0.125	-1.763	1.788	-0.088	-1.772	1.795	-0.088
Cohabitation with Child's Father	4.281	4.612	0.078	2.720	4.705	0.049	2.599	4.743	0.047
Index of Intelligence (WAIS)	0.166 [†]	0.097	0.174	0.148	0.097	0.156	0.149	0.097	0.156
Obstetric Risk	-1.925	2.831	-0.053	-2.226	2.824	-0.061	-2.210	2.835	-0.061
Non-Hispanic White	-1.764	3.191	-0.054	-2.295	3.195	-0.070	-2.277	3.208	-0.070
Latina	-6.728 [†]	3.747	-0.189	-7.197 [†]	3.741	-0.202	-7.238 [†]	3.759	-0.203
Gestational Age at Birth (GAB)	0.863	1.017	0.066	0.776	1.014	0.059	0.800	1.021	0.061
Child Sex (male)	-6.473*	2.464	-0.195	-6.717**	2.457	-0.203	-6.808**	2.488	-0.205
Prenatal Distress	-1.376	1.616	-0.079	-1.624	1.616	-0.093	-1.675	1.633	-0.096
Intrusiveness Reverse-Scored				4.403	2.937	0.128	4.174	3.059	0.121
Prenatal Distress* Intrusiveness							0.977	3.497	0.022

Variable	Model 1: Prenatal Distress		Model 2: Prenatal Distress and Intrusiveness Reverse-Scored				Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Reverse-Scored									
<i>R</i> ²		0.379			0.390			0.391	
<i>F</i>		6.715***			6.406***			5.874***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 11c

Regression Model Examining Maternal Positive Regard as a Moderator of the Association Between Prenatal Distress and Child Cognitive Function

Variable	Model 1: Prenatal Distress		Model 2: Prenatal Distress and Positive Regard				Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.038	0.334	0.011	0.011	0.336	0.003	0.092	0.334	0.028
SES	2.159 [†]	1.139	0.225	2.110 [†]	1.143	0.220	2.099 [†]	1.128	0.219
Parity	-2.503	1.728	-0.125	-2.425	1.734	-0.121	-2.627	1.713	-0.131
Cohabitation with Child's Father	4.281	4.612	0.078	4.728	4.659	0.086	3.989	4.609	0.073
Index of Intelligence (WAIS)	0.166 [†]	0.097	0.174	0.164 [†]	0.097	0.172	0.152	0.096	0.160
Obstetric Risk	-1.925	2.831	-0.053	-1.816	2.840	-0.050	-2.281	2.810	-0.063
Non-Hispanic White	-1.764	3.191	-0.054	-1.753	3.197	-0.054	-2.491	3.174	-0.076
Latina	-6.728 [†]	3.747	-0.189	-6.319 [†]	3.794	-0.177	-6.571 [†]	3.745	-0.185
Gestational Age at Birth (GAB)	0.863	1.017	0.066	0.792	1.023	0.060	0.503	1.019	0.038
Child Sex (male)	-6.473*	2.464	-0.195	-6.452*	2.469	-0.195	-6.201*	2.438	-0.187
Prenatal Distress	-1.376	1.616	-0.079	-1.219	1.633	-0.070	-1.059	1.613	-0.060
Positive Regard				2.006	2.691	0.059	1.486	2.666	0.044
Prenatal Distress* Positive Regard							5.768*	2.768	0.154
<i>R</i> ²		0.379			0.382			0.404	
<i>F</i>		6.715***			6.179***			6.197***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

p < .001

Appendix Table 12a

Regression Model Examining Maternal Sensitivity to Nondistress as a Moderator of the Association Between Prenatal Distress and Child Negative Emotionality

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity to Nondistress			Model 3: Interaction		
	B	SE B	β	B	SE B	β	B	SE B	β
Maternal Age	-0.012	0.011	-0.105	-0.012	0.011	-0.105	-0.014	0.011	-0.120
SES	-0.023	0.041	-0.065	-0.022	0.042	-0.062	-0.019	0.042	-0.054
Cohabitation with Child's Father	-0.172	0.193	-0.079	-0.174	0.194	-0.080	-0.141	0.193	-0.065
Index of Intelligence (WAIS)	-0.003	0.003	-0.100	-0.003	0.004	-0.097	-0.003	0.004	-0.090
Non-Hispanic White	-0.258*	0.127	-0.223	-0.259*	0.128	-0.224	-0.235 [†]	0.127	-0.203
Latina	0.152	0.152	0.117	0.149	0.154	0.115	0.187	0.154	0.145
Prenatal Distress	0.125*	0.060	0.201	0.124*	0.061	0.200	0.127*	0.060	0.205
Sensitivity to Nondistress				-0.013	0.109	-0.011	-0.026	0.108	-0.022
Prenatal Distress* Sensitivity to Nondistress							-0.214 [†]	0.116	-0.151
R ²		0.280			0.280			0.302	
F		6.173***			5.355***			5.246***	

B = unstandardized coefficient. β= standardized coefficient.

[†] p < .10

* p < .05.

** p < .01.

*** p < .001

Appendix Table 12b

Regression Model Examining Maternal Intrusiveness Reverse-Scored as a Moderator of the Association Between Prenatal Distress and Child Negative Emotionality

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Intrusiveness Reverse-Scored			Model 3: Interaction		
	B	SE B	β	B	SE B	β	B	SE B	β
Maternal Age	-0.012	0.011	-0.105	-0.013	0.011	-0.109	-0.010	0.011	-0.088
SES	-0.023	0.041	-0.065	-0.017	0.042	-0.050	-0.030	0.042	-0.087
Cohabitation with Child's Father	-0.172	0.193	-0.079	-0.154	0.196	-0.071	-0.076	0.198	-0.035
Index of Intelligence (WAIS)	-0.003	0.003	-0.100	-0.003	0.004	-0.088	-0.003	0.004	-0.085
Non-Hispanic White	-0.258*	0.127	-0.223	-0.255*	0.128	-0.221	-0.267*	0.126	-0.231

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Intrusiveness Reverse-Scored			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Latina	0.152	0.152	0.117	0.155	0.153	0.120	0.153	0.151	0.118
Prenatal Distress	0.125*	0.060	0.201	0.126*	0.060	0.203	0.142*	0.060	0.228
Intrusiveness Reverse-Scored				-0.073	0.128	-0.052	0.020	0.136	0.014
Prenatal Distress* Intrusiveness Reverse-Scored							-0.262 [†]	0.142	-0.166
<i>R</i> ²		0.280			0.282			0.304	
<i>F</i>		6.173***			5.409***			5.295***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

*** *p* < .001

Appendix Table 12c

Regression Model Examining Maternal Positive Regard as a Moderator of the Association Between Prenatal Distress and Child Negative Emotionality

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Positive Regard			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.012	0.011	-0.105	-0.012	0.011	-0.105	-0.014	0.012	-0.124
SES	-0.023	0.041	-0.065	-0.023	0.041	-0.067	-0.022	0.041	-0.062
Cohabitation with Child's Father	-0.172	0.193	-0.079	-0.166	0.196	-0.076	-0.174	0.196	-0.080
Index of Intelligence (WAIS)	-0.003	0.003	-0.100	-0.003	0.003	-0.100	-0.003	0.003	-0.099
Non-Hispanic White	-0.258*	0.127	-0.223	-0.258*	0.128	-0.223	-0.238 [†]	0.129	-0.206
Latina	0.152	0.152	0.117	0.158	0.155	0.122	0.171	0.156	0.132
Prenatal Distress	0.125*	0.060	0.201	0.127*	0.062	0.205	0.119 [†]	0.062	0.192
Positive Regard				0.022	0.108	0.018	0.022	0.108	0.018
Prenatal Distress* Positive Regard							-0.127	0.128	-0.083
<i>R</i> ²		0.280			0.280			0.287	
<i>F</i>		6.173***			5.360***			4.873***	

B = unstandardized coefficient. β = standardized coefficient.

[†] *p* < .10

* *p* < .05.

** *p* < .01.

 $p < .001$

Appendix Table 13: Main Analyses Including Postnatal Maternal Distress

Appendix Table 13a

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Cognitive Function, Including Postnatal Maternal Distress

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite			Model 3 : Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	<i>B</i>
Maternal Age	0.072	0.338	0.022	-0.035	0.338	-0.011	-0.018	0.332	-0.005
SES	2.036 [†]	1.154	0.213	1.692	1.151	0.177	1.769	1.133	0.185
Parity	-2.484	1.731	-0.124	-1.619	1.760	-0.081	-1.618	1.731	-0.081
Cohabitation	4.675	4.652	0.085	4.029	4.603	0.073	2.193	4.601	0.040
Index of Intelligence (WAIS)	0.172 [†]	0.097	0.180	0.148	0.096	0.156	0.135	0.095	0.141
Obstetric risk	-2.023	2.840	-0.056	-1.869	2.804	-0.051	-1.986	2.758	-0.054
Non-Hispanic White	-1.551	3.211	-0.048	-1.794	3.171	-0.055	-2.468	3.134	-0.076
Latina	-6.690 [†]	3.755	-0.188	-6.038	3.720	-0.170	-6.532 [†]	3.666	-0.183
Gestational Age at Birth (GAB)	0.820	1.021	0.063	0.616	1.012	0.047	0.437	0.999	0.033
Child sex (male)	-6.401 [*]	2.471	-0.193	-6.199 [*]	2.441	-0.187	-6.660 ^{**}	2.410	-0.201
Postnatal Distress	-1.701	2.334	-0.095	-1.224	2.316	-0.068	-1.080	2.279	-0.060
Prenatal Distress	0.014	2.503	0.001	-0.303	2.475	-0.017	-0.543	2.437	-0.031
Sensitivity Composite				2.533 [*]	1.239	0.185	2.700 [*]	1.221	0.197
Prenatal Distress [*] Sensitivity Composite							2.796 [*]	1.249	0.161
<i>R</i> ²		0.382			0.403			0.427	
<i>F</i>		6.176 ^{***}			6.173 ^{***}			6.283 ^{***}	

B = unstandardized coefficient. β = standardized coefficient.

[†] $p < .10$

^{*} $p < .05$.

^{**} $p < .01$.

^{***} $p < .001$

Appendix Table 13b

Regression Models Examining the Association Between Prenatal Maternal Psychological Distress and Child Negative Emotionality, Including Postnatal Maternal Distress

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.014	0.011	-0.121	-0.014	0.011	-0.122	-0.016	0.011	-0.138
SES	-0.013	0.041	-0.037	-0.012	0.042	-0.033	-0.013	0.042	-0.037
Cohabitation with Child's Father	-0.195	0.192	-0.090	-0.196	0.193	-0.090	-0.145	0.192	-0.067
Index of Intelligence (WAIS)	-0.004	0.003	-0.119	-0.004	0.004	-0.117	-0.004	0.004	-0.108
Non-Hispanic White	-0.265*	0.127	-0.230	-0.266*	0.127	-0.230	-0.240 [†]	0.126	-0.208
Latina	0.158	0.152	0.122	0.156	0.153	0.120	0.190	0.152	0.146
Postnatal Distress	0.122	0.084	0.194	0.122	0.084	0.194	0.123	0.083	0.196
Prenatal Distress	0.030	0.089	0.048	0.029	0.089	0.047	0.031	0.088	0.050
Sensitivity Composite				-0.007	0.047	-0.013	-0.002	0.047	-0.003
Prenatal Distress * Sensitivity Composite							-0.105*	0.052	-0.165
<i>R</i> ²		0.294			0.294			0.320	
<i>F</i>		5.718***			5.039***			5.076***	

B = unstandardized coefficient. β = standardized coefficient.

[†] $p < .10$

* $p < .05$.

** $p < .01$.

*** $p < .001$

References

- Adamson B, Letourneau N, & Lebel C (2018). Prenatal maternal anxiety and children's brain structure and function: A systematic review of neuroimaging studies. *Journal of Affective Disorders*, 241, 117–126. 10.1016/j.jad.2018.08.029 [PubMed: 30118945]
- Ainsworth M, Blehar M, Waters E, & Wall S (1978). Patterns of attachment: Assessed in the strange situation and at home.
- Ainsworth MS (1979). Infant–mother attachment. *American Psychologist*, 34(10), 932–937. [PubMed: 517843]
- Ali E, Letourneau N, Benzies K, Ntanda H, Dewey D, Campbell T, & Giesbrecht G (2020). Maternal prenatal anxiety and children's externalizing and internalizing behavioral problems: The moderating roles of maternal-child attachment security and child sex. *Canadian Journal of Nursing Research*, 52(2), 88–99. 10.1177/0844562119894184

- Atzl VM, Grande LA, Davis EP, & Narayan AJ (2019). Perinatal promotive and protective factors for women with histories of childhood abuse and neglect. *Child Abuse & Neglect*, 91, 63–77. 10.1016/j.chiabu.2019.02.008 [PubMed: 30831534]
- Barker D (1998). In utero programming of chronic disease. *Clinical Science*, 95(2), 115–128. [PubMed: 9680492]
- Bayley N (1993). Bayley Scales of Infant Development. Psychological Corporation.
- Bergman K, Sarkar P, Glover V, & O'Connor T. g. (2008). Quality of child–parent attachment moderates the impact of antenatal stress on child fearfulness. *Journal of Child Psychology and Psychiatry*, 49(10), 1089–1098. 10.1111/j.1469-7610.2008.01987.x [PubMed: 19017025]
- Bick J, & Dozier M (2010). Mothers' concentrations of oxytocin following close, physical interactions with biological and nonbiological children. *Developmental Psychobiology*, 52(1), 100–107. 10.1002/dev.20411 [PubMed: 20953313]
- Blair MM, Glynn LM, Sandman CA, & Davis EP (2011). Prenatal maternal anxiety and early childhood temperament. *Stress*, 14(6), 644–651. [PubMed: 21790468]
- Bogoch Y, Biala YN, Linial M, & Weinstock M (2007). Anxiety induced by prenatal stress is associated with suppression of hippocampal genes involved in synaptic function. *Journal of Neurochemistry*, 101(4), 1018–1030. [PubMed: 17254015]
- Bush NR, Jones-Mason K, Coccia M, Caron Z, Alkon A, Thomas M, Coleman-Phox K, Wadhwa PD, Laraia BA, Adler NE, & Epel ES (2017). Effects of pre- and postnatal maternal stress on infant temperament and autonomic nervous system reactivity and regulation in a diverse, low-income population. *Development and Psychopathology*, 29(5), 1553–1571. 10.1017/S0954579417001237 [PubMed: 29162167]
- Buss C, Davis EP, Muftuler LT, Head K, & Sandman CA (2010). High pregnancy anxiety during mid-gestation is associated with decreased gray matter density in 6–9-year-old children. *Psychoneuroendocrinology*, 35(1), 141–153. 10.1016/j.psyneuen.2009.07.010 [PubMed: 19674845]
- Charil A, Laplante DP, Vaillancourt C, & King S (2010). Prenatal stress and brain development. *Brain Research Reviews*, 65(1), 56–79. [PubMed: 20550950]
- Cohen S, Kamarck T, & Mermelstein R (1983). A global measure of perceived stress. *Journal of Health and Social Behavior*, 24(4), 385–396. [PubMed: 6668417]
- Committee on Obstetric Practice, the American Institute of Ultrasound in Medicine, and the Society for Maternal-Fetal Medicine. (2017). Committee Opinion No 700: Methods for estimating the due date. *Obstetrics and Gynecology*, 129(5), 150–154.
- Curran MM, Sandman CA, Davis EP, Glynn LM, & Baram TZ (2017). Abnormal dendritic maturation of developing cortical neurons exposed to corticotropin releasing hormone (CRH): Insights into effects of prenatal adversity? *PloS One*, 12(6), e0180311. 10.1371/journal.pone.0180311 [PubMed: 28658297]
- D'Anna-Hernandez K, & Rivera KD (2014). Understanding and alleviating cultural stressors and health disparities in the perinatal outcomes of Mexican-American women. *ZERO TO THREE*, 34(4), 37–45.
- Davis EP, Glynn LM, Schetter CD, Hobel CJ, Chicz-DeMet A, & Sandman CA (2007). Prenatal exposure to maternal depression and cortisol influences infant temperament. *Journal of the American Academy of Child Adolescent Psychiatry*, 46(6), 737–746. [PubMed: 17513986]
- Davis EP, Hankin BL, Glynn LM, Head K, Kim DJ, & Sandman CA (2019). Prenatal maternal stress, child cortical thickness, and adolescent depressive symptoms. *Child Development*. 10.1111/cdev.13252
- Davis EP, Hankin BL, Swales DA, & Hoffman MC (2018). An experimental test of the fetal programming hypothesis: Can we reduce child ontogenetic vulnerability to psychopathology by decreasing maternal depression? *Development and Psychopathology*, 30(3), 787–806. 10.1017/S0954579418000470 [PubMed: 30068416]
- Davis EP, & Narayan AJ (in press). Pregnancy as a period of risk, adaptation, and resilience for mothers and infants. *Developmental Psychopathology*.

- Davis EP, & Sandman CA (2010). The timing of prenatal exposure to maternal cortisol and psychosocial stress is associated with human infant cognitive development. *Child Development*, 81(1), 131–148. 10.1111/j.1467-8624.2009.01385.x [PubMed: 20331658]
- Davis EP, & Sandman CA (2012). Prenatal psychobiological predictors of anxiety risk in preadolescent children. *Psychoneuroendocrinology*, 37(8), 1224–1233. [PubMed: 22265195]
- Davis EP, Stout SA, Molet J, Vegetabile B, Glynn LM, Sandman CA, Heins K, Stern H, & Baram TZ (2017). Exposure to unpredictable maternal sensory signals influences cognitive development across species. *Proceedings of the National Academy of Sciences*, 114(39), 10390–10395. 10.1073/pnas.1703444114
- Deans CL (2018). Maternal sensitivity, its relationship with child outcomes, and interventions that address it: A systematic literature review. *Early Child Development and Care*, 1–24. 10.1080/03004430.2018.1465415
- Deary IJ (2001). Human intelligence differences: A recent history. *Trends in Cognitive Sciences*, 5(3), 127–130. 10.1016/S1364-6613(00)01621-1 [PubMed: 11239813]
- Demers CH, Aran O, Glynn LM, & Davis EP (2020). Prenatal programming of neurodevelopment: Imaging and structural changes. In Wazana A, Oberlander T, & Szekel E (Eds.), *Prenatal Stress and Child Development*.
- Doyle C, Werner E, Feng T, Lee S, Altemus M, Isler JR, & Monk C (2015). Pregnancy distress gets under fetal skin: Maternal ambulatory assessment & sex differences in prenatal development. *Developmental Psychobiology*, 57(5), 607–625. 10.1002/dev.21317 [PubMed: 25945698]
- Eshel N, Daelmans B, Mello MCD, & Martines J (2006). Responsive parenting: Interventions and outcomes. *Bulletin of the World Health Organization*, 84(12), 991–998. [PubMed: 17242836]
- Falah-Hassani K, Shiri R, & Dennis CL (2016). Prevalence and risk factors for comorbid postpartum depressive symptomatology and anxiety. *Journal of Affective Disorders*, 198, 142–147. 10.1016/j.jad.2016.03.010 [PubMed: 27016657]
- Fan A, Buka S, Kosik RO, Chen YS, Wang SJ, & Eaton WW (2014). Association between maternal behavior in infancy and adult mental health: A 30-year prospective study. *Comprehensive Psychiatry*, 55(2).
- Farrell AK, Waters TEA, Young ES, Englund MM, Carlson EE, Roisman GI, & Simpson JA (2019). Early maternal sensitivity, attachment security in young adulthood, and cardiometabolic risk at midlife. *Attachment & Human Development*, 21(1), 70–86. 10.1080/14616734.2018.1541517 [PubMed: 30428778]
- Feldman R (2007). Parent-infant synchrony and the construction of shared timing; physiological precursors, developmental outcomes, and risk conditions. *Journal of Child Psychology and Psychiatry*, 48(3–4), 329–354. 10.1111/j.1469-7610.2006.01701.x [PubMed: 17355401]
- Francis D, Diorio J, Liu D, & Meaney MJ (1999). Nongenomic transmission across generations of maternal behavior and stress responses in the rat. *Science*; Washington, 286(5442), 1155–1158. [PubMed: 10550053]
- Gee DG (2016). Sensitive periods of emotion regulation: Influences of parental care on frontoamygdala circuitry and plasticity. In Rutherford HJV & Mayes LC (Eds.), *Maternal brain plasticity: Preclinical and human research and implications for intervention*. *New Directions for Child and Adolescent Development* (Vol. 153, pp. 87–110). <http://doi.wiley.com/10.1002/cad.20166>
- Gilmore J, Knickmeyer R, & Gao W (2018). Imaging structural and functional brain development in early childhood. *Nature Reviews Neuroscience*, 19, 123–137. [PubMed: 29449712]
- Glynn LM, Davis EP, Sandman CA, & Goldberg WA (2016). Gestational hormone profiles predict human maternal behavior at 1-year postpartum. *Hormones and Behavior*, 85, 19–25. 10.1016/j.yhbeh.2016.07.002 [PubMed: 27427279]
- Glynn LM, Howland M, Sandman CA, Davis EP, Phelan M, Baram TZ, & Stern H (2018). Prenatal maternal mood patterns predict child temperament and adolescent mental health. *Journal of Affective Disorders*, 228, 83–90. [PubMed: 29241049]
- Granger SJ, Glynn LM, Sandman CA, Small SL, Obenaus A, Keator DB, Baram TZ, Stern H, Yassa MA, & Davis EP (2020). Aberrant maturation of the uncinatus fasciculus follows

exposure to unpredictable patterns of maternal signals. *Journal of Neuroscience*. 10.1523/JNEUROSCI.0374-20.2020

- Grant KA, McMahon C, Reilly N, & Austin MP (2010a). Maternal sensitivity moderates the impact of prenatal anxiety disorder on infant responses to the still-face procedure. *Infant Behavior and Development*, 33(4), 453–462. [PubMed: 20554325]
- Grant KA, McMahon C, Reilly N, & Austin MP (2010b). Maternal sensitivity moderates the impact of prenatal anxiety disorder on infant mental development. *Early Human Development*, 86(9), 551–556. 10.1016/j.earlhumdev.2010.07.004 [PubMed: 20709475]
- Hobel CJ (1982). Identification of the patient at risk. In *Perinatal medicine: Management of the high risk fetus and neonate*. Williams & Wilkins.
- Howland MA, Sandman CA, Davis EP, Stern HS, Phelan M, Baram TZ, & Glynn LM (2020). Prenatal maternal mood entropy is associated with child neurodevelopment. *Emotion*. 10.1037/emo0000726
- Kaplan LA, Evans L, & Monk C (2008). Effects of mothers' prenatal psychiatric status and postnatal caregiving on infant biobehavioral regulation: Can prenatal programming be modified? *Early Human Development*, 84(4), 249–256. 10.1016/j.earlhumdev.2007.06.004 [PubMed: 17761394]
- Kingston D, Tough S, & Whitfield H (2012). Prenatal and postpartum maternal psychological distress and infant development: A systematic review. *Child Psychiatry and Human Development*, 43(5), 683–714. [PubMed: 22407278]
- Knickmeyer R, Gouttard S, Kang C, Evans D, Wilber K, Smith JK, Hamer RM, Lin W, Gerig G, & Gilmore J (2008). A structural MRI study of human brain development from birth to 2 years. *Journal of Neuroscience*, 28(47), 12176–12181. [PubMed: 19020011]
- Kok R, Thijssen S, Bakermans-Kranenburg MJ, Jaddoe VWV, Verhulst FC, White T, van IJzendoorn MH, & Tiemeier H (2015). Normal variation in early parental sensitivity predicts child structural brain development. *Journal of the American Academy of Child & Adolescent Psychiatry*, 54(10), 824–831. 10.1016/j.jaac.2015.07.009 [PubMed: 26407492]
- Korja R, Nolvi S, Grant KA, & McMahon C (2017). The relations between maternal prenatal anxiety or stress and child's early negative reactivity or self-regulation: A systematic review. *Child Psychiatry & Human Development*, 48(6), 851–869. 10.1007/s10578-017-0709-0 [PubMed: 28124273]
- Krueger RF, & Markon KE (2006). Reinterpreting comorbidity: A model-based approach to understanding and classifying psychopathology. *Annual Review of Clinical Psychology*, 2(1), 111–133. 10.1146/annurev.clinpsy.2.022305.095213
- Lee AM, Lam SK, Sze Mun Lau S, Chong C, Chui H, & Fong D (2007). Prevalence, course, and risk factors for antenatal anxiety and depression. *Obstetrics and Gynecology*, 110(5), 1102–1112. [PubMed: 17978126]
- Lemaire V, Lamarque S, Le Moal M, Piazza P-V, & Abrous DN (2006). Postnatal stimulation of the pups counteracts prenatal stress-induced deficits in hippocampal neurogenesis. *Biological Psychiatry*, 59(9), 786–792. 10.1016/j.biopsych.2005.11.009 [PubMed: 16460692]
- Liu D, Diori J, Day JC, Francis DD, & Meaney MJ (2000). Maternal care, hippocampal synaptogenesis and cognitive development in rats. *Nature Neuroscience*, 3(8), 799. [PubMed: 10903573]
- Luby JL, Barch D, Whalen D, Tillman R, & Belden A (2017). Association between early life adversity and risk for poor emotional and physical health in adolescence: A putative mechanistic neurodevelopmental pathway. *JAMA Pediatrics*, 171(12), 1168–1175. [PubMed: 29084329]
- Madigan S, Oatley H, Racine N, Fearon RMP, Schumacher L, Akbari E, Cooke JE, & Tarabulsy GM (2018). A meta-analysis of maternal prenatal depression and anxiety on child socioemotional development. *Journal of the American Academy of Child & Adolescent Psychiatry*, 57(9), 645–657.e8. 10.1016/j.jaac.2018.06.012 [PubMed: 30196868]
- Mahrer N, Ramos I, Guardino C, Davis E, Ramey S, Shalowitz M, & Schetter C (2019). Pregnancy anxiety in expectant mothers predicts offspring negative affect: The moderating role of acculturation. *Early Human Development*, 141, 104932. 10.1016/j.earlhumdev.2019.104932 [PubMed: 31775096]

- Malmberg LE, Lewis S, West A, Murray E, Sylva K, & Stein A (2016). The influence of mothers' and fathers' sensitivity in the first year of life on children's cognitive outcomes at 18 and 36 months. *Child: Care, Health and Development*, 42(1), 1–7. 10.1111/cch.12294 [PubMed: 26538379]
- NICHD Early Child Care Research Network. (1997). The effects of infant child care on infant-mother attachment security: Results of the NICHD study of early child care. *Child Development*, 68(5), 860–879. 10.2307/1132038 [PubMed: 29106728]
- NICHD Early Child Care Research Network. (1999). Child care and mother-child interaction in the first three years of life. *Developmental Psychology*, 35(6), 1399–1413. [PubMed: 10563730]
- NICHD Early Child Care Research Network. (2001). Nonmaternal care and family factors in early development: An overview of the NICHD Study of Early Child Care. *Journal of Applied Developmental Psychology*, 22(5), 457–492. 10.1016/S0193-3973(01)00092-2
- Nillni YI, Mehralizade A, Mayer L, & Milanovic S (2018). Treatment of depression, anxiety, and trauma-related disorders during the perinatal period: A systematic review. *Clinical Psychology Review*, 66, 136–148. 10.1016/j.cpr.2018.06.004 [PubMed: 29935979]
- Polanska K, Krol A, Merecz-Kot D, Jurewicz J, Makowiec-Dabrowska T, Chiarotti F, Calamandrei G, & Hanke W (2017). Maternal stress during pregnancy and neurodevelopmental outcomes of children during the first 2 years of life. *Journal of Paediatrics and Child Health*, 53(3), 263–270. 10.1111/jpc.13422 [PubMed: 28168801]
- Putnam SP, Gartstein MA, & Rothbart MK (2006). Measurement of fine-grained aspects of toddler temperament: The early childhood behavior questionnaire. *Infant Behavior & Development*, 29(3), 386–401. 10.1016/j.infbeh.2006.01.004 [PubMed: 17138293]
- Raineki C, Lucion A, & Weinberg J (2014). Neonatal handling: An overview of the positive and negative effects. *Developmental Psychobiology*, 56(8), 1613–1625. [PubMed: 25132525]
- Rao H, Betancourt L, Giannetta JM, Brodsky NL, Korczykowski M, Avants BB, Gee JC, Wang J, Hurt H, Detre JA, & Farah MJ (2010). Early parental care is important for hippocampal maturation: Evidence from brain morphology in humans. *NeuroImage*, 49(1), 1144–1150. 10.1016/j.neuroimage.2009.07.003 [PubMed: 19595774]
- Røsand GB, Slinning K, Eberhard-Gran M, Røysamb E, & Tambs K (2011). Partner relationship satisfaction and maternal emotional distress in early pregnancy. *BMC Public Health*, 11(1), 161. 10.1186/1471-2458-11-161 [PubMed: 21401914]
- Sandman CA, Buss C, Head K, & Davis EP (2015). Fetal exposure to maternal depressive symptoms is associated with cortical thickness in late childhood. *Biological Psychiatry*, 77(4), 324–334. 10.1016/j.biopsych.2014.06.025 [PubMed: 25129235]
- Sandman CA, Class QA, Glynn LM, & Davis EP (2015). Neurobehavioral disorders and developmental origins of health and disease. In Rosenfeld CS (Ed.), *The Epigenome and Developmental Origins of Health and Disease* (pp. 235–266). Academic Press/Elsevier. <https://www.elsevier.com/books/the-epigenome-and-developmental-origins-of-health-and-disease/rosenfeld/978-0-12-801383-0>
- Sandman CA, Curran MM, Davis EP, Glynn LM, Head K, & Baram TZ (2018). Cortical thinning and neuropsychiatric outcomes in children exposed to prenatal adversity: A role for placental CRH? *American Journal of Psychiatry*, 175(5), 471–479. 10.1176/appi.ajp.2017.16121433 [PubMed: 29495899]
- Sandman CA, & Davis EP (2010). Gestational stress influences cognition and behavior. *Future Neurology*, 5(5), 675–690. 10.2217/fnl.10.35
- Sandman CA, Glynn LM, Davis EP, Kisilevsky BS, & Reissland N (2016). Neurobehavioral consequences of fetal exposure to gestational stress. In *Fetal Development: Research on Brain and Behavior, Environmental Influences and Emerging Technologies* (pp. 229–265). Springer. https://link.springer-com.du.idm.oclc.org/chapter/10.1007/978-3-319-22023-9_13
- Santor DA, & Coyne JC (1997). Shortening the CES–D to improve its ability to detect cases of depression. *Psychological Assessment*, 9(3), 233–243.
- Schechter JC, Brennan PA, Smith AK, Stowe ZN, Newport DJ, & Johnson KC (2017). Maternal prenatal psychological distress and preschool cognitive functioning: The protective role of positive parental engagement. *Journal of Abnormal Child Psychology*, 45(2), 249–260. [PubMed: 27150387]

- Schuermans C, & Kurrasch D (2013). Neurodevelopmental consequences of maternal distress: What do we really know? *Clinical Genetics*, 83(2), 108–117. 10.1111/cge.12049 [PubMed: 23140231]
- Sharp H, Hill J, Hellier J, & Pickles A (2015). Maternal antenatal anxiety, postnatal stroking and emotional problems in children: Outcomes predicted from pre- and postnatal programming hypotheses. *Psychological Medicine*, 45(2), 269–283. [PubMed: 25068652]
- Sharp H, Pickles A, Meaney M, Marshall K, Tibu F, & Hill J (2012). Frequency of infant stroking reported by mothers moderates the effect of prenatal depression on infant behavioural and physiological outcomes. *PLOS ONE*, 7(10).
- Spielberger C (1983). *Manual for the state-trait anxiety inventory*. Consulting Psychologists Press, Inc.
- Spinrad TL, & Stifter CA (2002). Maternal sensitivity and infant emotional reactivity: Concurrent and longitudinal relations. *Marriage & Family Review*, 34(3–4), 243–263. 10.1300/J002v34n03_03
- Van den Bergh BRH, Mulder EJH, Mennes M, & Glover V (2005). Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: Links and possible mechanisms. A review. *Neuroscience & Biobehavioral Reviews*, 29(2), 237–258. 10.1016/j.neubiorev.2004.10.007 [PubMed: 15811496]
- Van den Bergh BRH, Van Den Heuvel M, Lahti M, Braeken M, Rooij S, Entringer S, Hoyer D, Roseboom T, Raikkonen K, King S, & Schwab M (2017). Prenatal developmental origins of behavior and mental health: The influence of maternal stress in pregnancy. *Neuroscience & Biobehavioral Reviews*. <https://www.ncbi.nlm.nih.gov/pubmed/28757456>
- Vehmeijer FOL, Guxens M, Duijts L, & Marroun HE (2019). Maternal psychological distress during pregnancy and childhood health outcomes: A narrative review. *Journal of Developmental Origins of Health and Disease*, 10(3), 274–285. 10.1017/S2040174418000557 [PubMed: 30378522]
- Wakshlak A, & Weinstock W (1990). Neonatal handling reverses behavioral abnormalities induced in rats by prenatal stress. *Physiology & Behavior*, 48(2), 289–292. 10.1016/0031-9384(90)90315-U [PubMed: 2255733]
- Wang Q, Zhang H, Wee C-Y, Lee A, Poh JS, Chong Y-S, Tan KH, Gluckman PD, Yap F, Fortier MV, Rifkin-Graboi A, & Qiu A (2019). Maternal sensitivity predicts anterior hippocampal functional networks in early childhood. *Brain Structure and Function*, 224(5), 1885–1895. 10.1007/s00429-019-01882-0 [PubMed: 31055646]
- Wechsler D, Coalson DL, & Raiford SE (1997). *WAIS-III: Wechsler adult intelligence scale*. Psychological Corporation.
- Wu Y, Lu Y-C, Jacobs M, Pradhan S, Kapse K, Zhao L, Niforatos-Andescavage N, Vezina G, du Plessis AJ, & Limperopoulos C (2020). Association of prenatal maternal psychological distress with fetal brain growth, metabolism, and cortical maturation. *JAMA Network Open*, 3(1), e1919940–e1919940. 10.1001/jamanetworkopen.2019.19940 [PubMed: 31995213]




Figure 1.
Study timeline




Figure 2.

Maternal sensitivity composite was analyzed as a continuous variable using regression, but for illustrative purposes are depicted here as low (1 *SD* below the mean), average (at the mean), and high (1 *SD* above the mean) levels of maternal sensitivity. Prenatal maternal distress (on the x-axis) is the standardized composite of anxiety, depressive symptoms, and perceived stress scores. Children exposed to elevated prenatal maternal distress did not exhibit impaired cognitive function at age 2 if they received higher quality maternal caregiving.




Figure 3. Maternal sensitivity composite was analyzed as a continuous variable using regression, but for illustrative purposes are depicted here as low (1 *SD* below the mean), average (at the mean), and high (1 *SD* above the mean) levels of maternal sensitivity. Prenatal maternal distress (on the x-axis) is the standardized composite of anxiety, depressive symptoms, and perceived stress scores. Children exposed to elevated prenatal maternal distress did not exhibit high negative emotionality at age 2 if they received higher quality maternal caregiving.

Table 1

Sample Characteristics

Maternal Characteristics (N = 136)	(M ± SD) (range) or (%)*
Age at Delivery	30.28 ± 5.08 (19.06 – 44.63)
Cohabiting with Child's Father	91.0
Primiparous	43.4
Ethnicity	
Non-Hispanic White	50.7
Latina	29.4
African American or Black	2.9
Asian	8.1
Multi-Ethnic	8.8
Household Income	
\$0-\$30,000	20.6
\$30,001-\$60,000	25.7
\$60,001-\$100,000	33.8
Over \$100,000	19.9
Years of Education	15.99 ± 2.35 (9 – 19)
Obstetric Complications Sum Score	0.34 ± 0.56 (0 – 2)
Maternal Sensitivity Composite (6 months)	9.98 ± 1.06 (6.5 – 12.0)
Maternal Sensitivity Composite (12 months)	9.67 ± 1.67 (4.5 – 12.0)
Child Characteristics (N = 136)	
Sex (% Male)	58.1, n=79
Apgar Score (5 min)	9.00 ± 0.27 (8 – 10)
Gestational Age at Birth (GAB)	39.44 ± 1.26 (35.29 – 42.57)
Birth Weight Percentile**	52.77 ± 27.90 (1 – 99)
Mental Developmental Index (MDI)	97.71 ± 16.39 (54 – 140)
Negative Affectivity (ECBQ)	2.80 ± 0.58 (1.38 – 5.08)

* Values presented as (means ± SD) or (%) where applicable

** Birth weight percentile was calculated according to the infant's sex and gestational age at birth.

Table 2

Regression Model Examining the Association Between Prenatal Maternal Psychological Distress, Maternal Care and Child Cognitive Function

Variable	Model 1: Prenatal Distress		Model 2: Prenatal Distress and Sensitivity Composite			Model 3: Interaction			
	<i>B</i>	<i>SE B</i>	<i>B</i>	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	0.038	0.334	0.011	-0.062	0.333	-0.019	-0.042	0.327	-0.013
SES	2.159 [†]	1.139	0.225	1.771	1.138	0.185	1.839	1.120	0.192
Parity	-2.503	1.728	-0.125	-1.610	1.755	-0.080	-1.610	1.726	-0.080
Cohabitation with Child's Father	4.281	4.612	0.078	3.732	4.554	0.068	1.920	4.550	0.035
Index of Intelligence (WAIS)	0.166 [†]	0.097	0.174	0.144	0.096	0.151	0.130	0.094	0.137
Obstetric Risk	-1.925	2.831	-0.053	-1.795	2.792	-0.049	-1.921	2.746	-0.053
Non-Hispanic White	-1.764	3.191	-0.054	-1.951	3.148	-0.060	-2.611	3.109	-0.080
Latina	-6.728 [†]	3.747	-0.189	-6.048	3.709	-0.170	-6.544 [†]	3.654	-0.184
Gestational Age at Birth (GAB)	0.863	1.017	0.066	0.641	1.008	0.049	0.458	0.995	0.035
Child Sex (male)	-6.473 [*]	2.464	-0.195	-6.244 [*]	2.432	-0.188	-6.703 ^{**}	2.400	-0.202
Prenatal Distress	-1.376	1.616	-0.079	-1.301	1.594	-0.074	-1.425	1.568	-0.081
Sensitivity Composite				2.599 [*]	1.229	0.190	2.759 [*]	1.211	0.201
Prenatal Distress * Sensitivity Composite							2.813 [*]	1.245	0.161
<i>R</i> ²		0.379			0.401			0.426	
<i>F</i>		6.715 ^{***}			6.705 ^{***}			6.794 ^{***}	

B = unstandardized coefficient. β = standardized coefficient.

[†] $p < .10$

^{*} $p < .05$.

^{**} $p < .01$.

^{***} $p < .001$

Table 3

Regression Model Examining the Association Between Prenatal Maternal Psychological Distress, Maternal Care and Child Negative Emotionality

Variable	Model 1: Prenatal Distress			Model 2: Prenatal Distress and Sensitivity Composite			Model 3: Interaction		
	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β	<i>B</i>	<i>SE B</i>	β
Maternal Age	-0.012	0.011	-0.105	-0.012	0.011	-0.105	-0.014	0.011	-0.121
SES	-0.023	0.041	-0.065	-0.021	0.042	-0.061	-0.023	0.041	-0.065
Cohabitation with Child's Father	-0.172	0.193	-0.079	-0.173	0.194	-0.080	-0.122	0.193	-0.056
Index of Intelligence (WAIS)	-0.003	0.003	-0.100	-0.003	0.004	-0.097	-0.003	0.003	-0.087
Non-Hispanic White	-0.258*	0.127	-0.223	-0.258*	0.128	-0.224	-0.233 [†]	0.127	-0.201
Latina	0.152	0.152	0.117	0.148	0.154	0.114	0.182	0.153	0.141
Prenatal Distress	0.125*	0.060	0.201	0.124*	0.061	0.199	0.126*	0.060	0.204
Sensitivity Composite				-0.008	0.048	-0.016	-0.003	0.047	-0.006
Prenatal Distress * Sensitivity Composite							-0.105*	0.052	-0.164
<i>R</i> ²		0.280			0.280			0.306	
<i>F</i>		6.173***			5.358***			5.342***	

B = unstandardized coefficient. β = standardized coefficient.

[†]*p* < .10

**p* < .05.

***p* < .01.

****p* < .001