

UC Berkeley

UC Berkeley Previously Published Works

Title

Molecular Imaging of Labile Iron(II) Pools in Living Cells with a Turn-On Fluorescent Probe

Permalink

<https://escholarship.org/uc/item/6846t4bt>

Journal

Journal of the American Chemical Society, 135(40)

ISSN

0002-7863

Authors

Au-Yeung, Ho Yu
Chan, Jefferson
Chantarojsiri, Teera
[et al.](#)

Publication Date

2013-10-09

DOI

10.1021/ja4072964

Peer reviewed


Molecular Imaging of Labile Iron(II) Pools in Living Cells with a Turn-On Fluorescent Probe

Ho Yu Au-Yeung,[†] Jefferson Chan,[†] Teera Chantarojsiri,[†] and Christopher J. Chang^{*,†,‡,§}

[†]Departments of Chemistry and [‡]Molecular and Cell Biology and [§]Howard Hughes Medical Institute, University of California, Berkeley, California 94720, United States

S Supporting Information

ABSTRACT: Iron is an essential metal for living organisms, but misregulation of its homeostasis at the cellular level can trigger detrimental oxidative and/or nitrosative stress and damage events. Motivated to help study the physiological and pathological consequences of biological iron regulation, we now report a reaction-based strategy for monitoring labile Fe²⁺ pools in aqueous solution and living cells. Iron Probe 1 (IP1) exploits a bioinspired, iron-mediated oxidative C–O bond cleavage reaction to achieve a selective turn-on response to Fe²⁺ over a range of cellular metal ions in their bioavailable forms. We show that this first-generation chemical tool for fluorescence Fe²⁺ detection can visualize changes in exchangeable iron stores in living cells upon iron supplementation or depletion, including labile iron pools at endogenous, basal levels. Moreover, IP1 can be used to identify reversible expansion of labile iron pools by stimulation with vitamin C or the iron regulatory hormone hepcidin, providing a starting point for further investigations of iron signaling and stress events in living systems as well as future probe development.



INTRODUCTION

Iron is a required metal nutrient for life and the most abundant transition metal in the human body.^{1–5} In particular, potent redox capacity of iron, mainly in its ferrous (Fe²⁺) and ferric (Fe³⁺) forms, is utilized for a wide array of physiological functions ranging from O₂ binding and transport to electron transfer to catalysis. However, this same redox potency can also trigger damage to biological macromolecules and metabolites if cellular iron homeostasis is compromised.^{6–9}


The central importance of iron in biological systems provides a compelling need for new methods to track biological iron pools, particularly in exchangeable, labile forms, to help assess the contributions of this metal to healthy and diseased states. In this regard, molecular imaging with iron-responsive fluorescent probes can potentially provide a convenient and an effective means to study bioavailable iron in living specimens.^{10–15} However, the field of iron-responsive fluorescent indicators for living biological samples remains underdeveloped owing to significant challenges in overcoming issues of metal selectivity, specificity for different oxidation and spin states, and potent fluorescence quenching by iron in its open-shell forms, all while maintaining biocompatibility.^{10,11} Moreover, the reducing environment of the cell and greater aqueous solubility of Fe²⁺ versus Fe³⁺ presages that the ferrous state may be more relevant for assaying labile iron pools.^{16–18} As such, the vast majority of reported fluorogenic probes for iron imaging in living cells operate through a turn-off response,^{10,11,14,17,19–25} including the commercially available reagents Phen Green SK^{17,24} and calcein,¹⁹ which limits spatial resolution. There are few

examples of turn-on Fe³⁺ sensors that have been applied to cells,^{26–32} and Fe²⁺ fluorescent probes are even more rare. Indeed, only recently has one turn-on fluorescent probe for Fe²⁺ in cells based on an elegant N-oxide reduction been described,³³ and no endogenous biological stimulations were reported in this initial work.

In this report, we present the design, synthesis, characterization, and biological applications of Iron Probe 1 (IP1), a new type of reaction-based turn-on fluorescent probe for monitoring exchangeable Fe²⁺ pools in aqueous solution and living cells. Using a bioinspired Fe²⁺- and O₂-dependent C–O cleavage reaction to release a fluorescent dye product, IP1 features visible excitation and emission profiles and the dual requirement of metal coordination and bond-breaking reactivity promotes selectivity for Fe²⁺ over a range of cellularly relevant metal ions in their bioavailable forms. We demonstrate the utility of IP1 in biological settings by showing that this probe can detect changes in exchangeable iron stores in living cells upon iron supplementation or depletion. Moreover, this first-generation probe is capable of imaging endogenous, basal labile iron pools and their expansion by vitamin C or the iron regulatory hormone hepcidin, providing a unique tool that has the capacity to monitor changes in natural cellular iron stores.


Received: July 17, 2013

Published: September 24, 2013

Scheme 1. Design of Iron Probe 1 (IP1), a Reaction-Based Probe for Fe²⁺ Featuring a Biomimetic, Oxidative Bond-Cleavage Reaction^a

^aIP1 contains an expanded mimic of a mononuclear non-heme 2-His-1-carboxylate coordination motif with an open sixth apical site for O₂ activation.

Scheme 2. Synthesis of IP1



RESULTS AND DISCUSSION

Design and Synthesis of IP1. Our design strategy for turn-on detection of Fe²⁺ relies on a reaction-based approach in which a selective but transient Fe²⁺-triggered bond-cleavage reaction converts a weakly fluorescent caged probe into a highly fluorescent product.^{15,34–38} Inspired by the diverse range of oxidative chemical transformations catalyzed by iron centers in biology within heme P450 and mononuclear nonheme enzymes and models,^{39–46} we decided to exploit a biomimetic oxidation strategy for design of new fluorescent iron probes. Specifically, we sought to construct a fluorophore caged with a biomimetic ligand that can be cleaved by a Fe²⁺-mediated oxygenation reaction. Indeed, recent work from Taki⁴⁷ as well as from our

laboratory⁴⁸ shows that reaction-based probes exploiting such biomimetic oxidative cleavage can be used to selectively detect Cu⁺ and Co²⁺, respectively, in live-cell settings (Scheme 1), and work by Kodanko⁴⁹ and Nam^{50,51} has shown biomimetic oxidations for catalysis and sensing purposes.


Unlike traditional molecular recognition approaches where metal selectivity is solely based on preferential coordination of the target metal with a particular ligand, the reaction-based strategy adds an extra selectivity filter besides coordination to distinguish Fe²⁺ from other 3d metal ions, which are similar in size and charge, as both the geometry and the electronic structure of the metal complex have to be matched with the chemical reactivity in order to result in the bond breaking

reaction. Also, this reaction-based strategy separates the metal and the fluorophore after the bond cleavage reaction, circumventing the problem of fluorescence quenching by the paramagnetic metal ions, thus allowing turn-on detection that improves spatial resolution. Moreover, since the biomimetic ligand can be modularly coupled to different fluorophores or reporter groups, it allows versatile modification of the photophysical properties of the sensors to better suit the specific need of a particular imaging experiment.

IP1 was synthesized according to Scheme 2. To promote selective iron reactivity, we designed a pentadentate N_4O ligand to mimic the 2-His-1-carboxylate facial triad commonly found in the active sites of O_2 -activating mononuclear nonheme iron enzymes.^{44–46} The N_4O ligand **3** was coupled onto the reduced fluorescein alcohol derivative **4**, disrupting conjugation of the fused rings and thus giving the nonfluorescent IP1 (Scheme 2). Notably, precursor **3** allows for facile incorporation of the Fe^{2+} -responsive unit to other fluorescent scaffolds and imaging modalities via a simple nucleophilic substitution reaction.

To better understand the coordination of Fe^{2+} to the N_4O motif, a model ligand **6** and its Fe^{2+} complex were synthesized (Scheme 3). The neutral mononuclear Fe^{2+} complex **6-FeCl**

Scheme 3. Synthesis of **6-FeCl** (30% thermal ellipsoids)^a



^aSelected bond parameters (Angstroms and degrees): Fe1–N1 2.229(4), Fe1–N2 2.214(5), Fe1–N3 2.198(4), Fe1–N4 2.166(5), Fe1–O1 2.116(3), Fe1–Cl1 2.301(2), N1–Fe1–Cl1 175.07(11), N2–Fe1–N4 153.04(19), N3–Fe1–O1 152.94(16).

was prepared by reaction of **6** with $FeCl_2$ and K_2CO_3 in MeCN. A single crystal was obtained from slow evaporation of a MeCN solution of the complex. As anticipated, the Fe^{2+} center is coordinated by the pentadentate ligand in a distorted octahedral geometry, with a chloride ligand occupying the apical position (Scheme 3). The $Fe-N_{py}$ and $Fe-O_{RCOO}$ distances are comparable to related N_3O_2 pyridine–carboxylate Fe^{2+} complexes.⁵² Moreover, FT-IR analysis of the complex in the solid state showed absorptions at 1600 and 1360 cm^{-1} , which are assigned to the symmetric and asymmetric vibrations of the carboxylate ligand, while the 1H NMR spectrum of **6-FeCl** (CD_3CN , 298 K, 400 MHz) showed paramagnetically shifted resonances comparable to similar $S = 2$ Fe^{2+} complexes.⁵¹ The relatively simple spectrum suggests the mononuclear structure persists in solution, which is also supported by detection of the $[M + H]^+$ peak at m/z 425.0 by ESI-MS. The UV–vis spectrum of the Fe^{2+} complex showed two absorptions at $\lambda = 404$ ($\epsilon = 390$ $cm^{-1} M^{-1}$) and 496 nm ($\epsilon = 360$ $cm^{-1} M^{-1}$), similar to other mononuclear, high-spin Fe^{2+} complexes supported by related N_3O_2 pyridine–carboxylate ligands.⁵²

Fe^{2+} Reactivity and Metal-Ion Selectivity of IP1.

Spectroscopic evaluation of IP1 was performed in aqueous buffer (50 mM Tris, pH 7.6) in the presence of 5 mM glutathione to mimic the reducing intracellular environment. As

expected, a 1 μM solution of IP1 is weakly fluorescent, and addition of 20 equiv of Fe^{2+} leads to a 6-fold emission turn-on response within 1 h of reaction (Figure 1).⁵³ In addition to




Figure 1. Fluorescence response of 1 μM IP1 (50 mM Tris, pH 7.6, 5 mM GSH) with $\lambda_{ex} = 470$ nm. (A) Emission spectra collected after reaction with 20 μM Fe^{2+} at time = 0 (bottom black trace), 10, 20, 30, 40, 50, and 60 (top black trace) min. (B) Relative emission intensity at 508 nm after 1 h reaction with various s-block (1 mM) and d-block (20 μM) metals as well as vitamin B₁₂ (20 μM).

these spectral signatures, release of the fluorophore **4** upon reaction of IP1 with Fe^{2+} was directly identified by ESI-MS, showing the dye product at m/z 333.0, whereas the peak at m/z 449.0 suggests the ligand has been oxidized to form the iron complex $[Fe(L-COO)]^+$ (Figure S1, Supporting Information). Furthermore, no significant emission increase was observed when IP1 was treated with Fe^{2+} under anaerobic conditions nor when IP1 was treated with 100 equiv of various ROS, confirming the requirement of O_2 in the C–O bond cleavage reaction and that it is Fe^{2+} mediated (Figures S2 and S3, Supporting Information). A detailed mechanism of the bond cleavage reaction is yet to be determined, but it presumably involves high-valent iron species analogous to those found in the nonheme iron systems.^{44–46} We were unable to further characterize intermediates of this O_2 -dependent process, as no significant changes in spectral features above 400 nm were observed upon reaction of an acetonitrile solution of the model compound **6-FeCl** with O_2 at temperatures ranging from -40 to 25 $^{\circ}C$.

The fluorescence turn-on response of IP1 is selective for Fe^{2+} over abundant cellular alkali and alkaline earth metal ions, like Na^+ , K^+ , Mg^{2+} , and Ca^{2+} , as well as other biologically relevant 3d metal ions like Mn^{2+} , Ni^{2+} , Cu^+ , Cu^{2+} , and Zn^{2+} (Figure 1). In addition, IP1 shows redox selectivity for Fe^{2+} over Fe^{3+} . Of the 3d transition metal series, only Co^{2+} at high concentrations can also give a turn-on response; however, ionic Co^{2+} at these levels is not physiologically relevant within mammalian systems.⁵⁴ As an additional control experiment, we therefore tested the fluorescence response of IP1 toward cobalamin




Figure 2. Representative confocal microscopy images of HepG2/C3A cells stained with 20 μM IP1-AM for 2 h. (A) Control cells. (B) Cells enriched with 100 μM ferrous ammonium sulfate (FAS) for 20 h. (C) Iron-enriched cells treated with 50 μM DFO for 40 min. (D) Cells treated with 50 μM desferoxamine (DFO) for 40 min. (E–H) Overlays of bright field images and Hoechst stain in A–D. (I) Quantification of relative fluorescence intensity of confocal microscopy images of HepG2/C3A cells obtained under conditions A–D. Error bars are $\pm\text{SD}$ ($n = 3$). Statistical analysis was performed with a two-tailed Student's t test. Asterisk (*): $p < 0.001$. Fluorescent intensity in images A–D is represented in a pseudocolor scale for better visual contrast.

(vitamin B₁₂), which is the dominant form of cobalt in mammalian systems under physiological conditions.⁵⁵ As expected, because the cobalt ion in cobalamin is tightly bound, no change in emission of IP1 was observed after reaction with 20 μM cobalamin, a level which is 20-fold above the maximum limit found in the human body.⁵⁵ In addition to cobalamin, the fluorescence response of IP1 toward Fe²⁺ in the presence of various concentrations of Co²⁺ and Zn²⁺ was also evaluated under conditions that mimic cellular overload of these metal (Figures S4 and S5, Supporting Information). We discovered that in the presence of Co²⁺ up to 10% relative to Fe²⁺, a concentration far higher than the total available cobalt in a cell, did not result in a further enhancement of the fluorescence signal compared to just Fe²⁺ alone, showing that the presence of labile cobalt, if any, would not result in a false positive fluorescence enhancement when cells are imaged with IP1.⁵⁶ The presence of Zn²⁺ attenuates the fluorescence enhancement from IP1 by Fe²⁺ only at relatively high concentrations (a 30% decrease at 10 μM Zn²⁺). For most mammalian and down to microbial/yeast cell types, labile zinc concentrations are in the femtomolar to nanomolar range, which is many orders of magnitude less than the concentrations used in these in vitro experiments.

Cellular Evaluation of IP1 for Imaging Labile Iron Pools. We next established the ability of IP1 to detect changes in exchangeable, labile iron pools within living cells. We chose a liver cell line model, HepG2/C3A, as the liver is a major iron storage organ in the body.⁵⁷ To improve cellular loading, IP1 was masked as an acetoxymethyl ester.^{58–61} and IP1-AM was employed in all cellular imaging studies. First, we tested if IP1 can detect exogenously elevated levels of labile iron in live HepG2/C3A cells. To this end, HepG2/C3A cells were supplemented with 100 μM ferrous ammonium sulfate (FAS) for 20 h and then stained with 20 μM IP1-AM staining for an additional 2 h. As shown in Figure 2B, a significant increase in intracellular fluorescence in iron-supplemented cells compared to control cells was observed (Figure 2A). To verify that the fluorescence increase was due to iron-dependent events, iron-enriched cells were treated with 50 μM of the clinically used iron chelator desferoxamine (DFO) for 40 min to reduce intracellular levels of labile iron. As expected, DFO chelation

attenuated cellular fluorescence (Figure 2C), establishing that the observed fluorescence increase was indeed correlated to elevated levels of exchangeable Fe²⁺ in the iron-supplemented cells.

With these data in hand, we next tested if IP1 could also detect labile iron pools at basal, endogenous levels within cells. Accordingly, HepG2/C3A cells were treated with 50 μM DFO for 40 min to produce iron-depleted specimens. Subsequent IP1-AM staining showed less fluorescent signal than untreated control samples (Figure 2D), suggesting that IP1 is sensitive enough to image labile iron pools at endogenous, basal levels. Co-staining of IP1-AM with LysoTracker-Red showed good colocalization of fluorescence signals (Figure S6, Supporting Information), demonstrating the applicability of IP1-AM in the study of labile iron in the lysosome, where Fe(II) predominates due to the acidic and reducing environment and the organelle is involved in the degradation and recycling of the metal from iron-containing components such as ferritin, mitochondria, and endocytosed erythrocyte in macrophages.⁶² Finally, the cellular toxicity of IP1-AM and its products was assessed by nuclear staining with Hoechst 33342 and WST-1 assays. Both sets of control experiments confirmed that the cells were viable throughout the imaging studies (Figures 2E–H and S9–S11, Supporting Information) and thus establish the suitability of IP1 for cellular imaging studies in live specimens.

Imaging Labile Iron Pools in Hepcidin-Stimulated Cells. After establishing that IP1 can reliably detect changes in labile Fe²⁺ pools within living cells, including exchangeable iron at endogenous, basal levels, we turned our attention to monitoring changes in cellular labile Fe²⁺ stores triggered by physiological stimulation. In this context, we first chose to explore the effects of hepcidin, a key iron regulatory hormone produced by the liver. Hepcidin controls iron homeostasis by regulating iron absorption in the intestine and release of the metal from hepatic stores and macrophages that recycle erythrocytes.^{63–65} Previous work has shown that coinubation of iron with hepcidin triggers a greater increase in the total cellular iron pool compared to control cells supplemented with iron alone, as measured by ferritin content and ⁵⁹Fe labeling, most likely due to the shutdown of the iron export pathway by degrading ferroportin.⁶⁶ To test whether hepcidin stimulation




Figure 3. Representative confocal microscopy images of HepG2/C3A cells stained with 20 μM IP1-AM for 2 h. (A) Control cells. (B) Cells enriched with 100 μM ferrous ammonium sulfate (FAS) for 20 h. (C) Cells treated with 100 FAS for 1 h, followed by addition of 0.5 μM hepcidin and coincubated for a further 19 h. (D) Cells treated with FAS and hepcidin for 20 and 19 h, respectively, and followed by a 40 min treatment with 50 μM desferoxamine (DFO). (E–H) Overlays of bright field images and Hoechst stain in A–D. (I) Quantification of relative fluorescence intensities of confocal microscopy images of HepG2/C3A cells obtained under conditions A–D. Error bars are $\pm\text{SD}$ ($n = 3$). Statistical analysis was performed with a two-tailed Student's t test. Asterisk (*): $p < 0.001$. Fluorescent intensity in images A–D is represented in a pseudocolor scale for better visual contrast.




Figure 4. Representative confocal microscopy images of HepG2/C3A cells stained with 20 μM IP1-AM for 2 h. (A) Control cells. (B) Cells treated with 1 mM ascorbic acid (Asb) for 40 min. (C) Cells treated with 1 mM ascorbic acid for 40 min and then 50 μM desferoxamine (DFO) for a further 40 min. (D–F) Overlays of bright field images and Hoechst stain in A–C. (G) Quantification of relative fluorescence intensities of confocal microscopy images of HepG2/C3A cells obtained under conditions A–C. Error bars are $\pm\text{SD}$ ($n = 3$). Statistical analysis was performed with a two-tailed Student's t test. Asterisk (*): $p < 0.001$. Fluorescent intensity in images A–C is represented in a pseudocolor scale for better visual contrast.

can also expand the labile cellular iron pool in addition to the total cellular iron content, we used IP1-AM to image HepG2/C3A cells treated with hepcidin and/or iron. As shown in Figure 3, we observed elevations in intracellular fluorescence in hepcidin-treated, iron-supplemented cells compared to cells supplemented with iron alone, suggesting that the labile iron pool does expand under hepcidin stimulation. Moreover, the observed IP1 fluorescence increases were attenuated by DFO chelation, showing that the fluorescence increase was due to an increase in the level of labile iron. To eliminate the possibility of any cobalt-dependent fluorescence enhancements, cells were also stained with the cobalt-specific indicator CPI as an additional control.⁴⁸ As expected, no observable differences in fluorescence between control and hepcidin-treated cells were observed, further supporting that the fluorescence enhancement in hepcidin-stimulated cells was iron dependent and due

to an increase in labile Fe^{2+} pool (Figures S7 and S8, Supporting Information). Finally, ICP-OES experiments on cell lysates independently show the cells treated with both hepcidin and FAS have more total iron compared to cells supplemented with FAS alone (Figure S12, Supporting Information), in agreement with the anticipated increase in total iron content after hepcidin stimulation. Taken together, the IP1 imaging data show that the iron regulatory hormone hepcidin not only leads to an increase in total cellular iron pool via ferritin storage⁶⁶ but also triggers a rise in the labile cellular iron pool.

Imaging Labile Iron in Vitamin C-Stimulated Cells. As a second stimulation model in addition to hepcidin supplementation, we also examined the effects of ascorbic acid (vitamin C) on labile cellular iron pools. Previous work establishes that vitamin C can promote redox cycling and mobilization of iron, therefore enhancing cellular uptake of

nontransferrin-bound iron as well as the efficiency of iron chelation in treatment for iron-overload disorders.^{67–70} To probe this relationship by molecular imaging, HepG2/C3A cells were stained with IP1-AM and compared to cells treated with ascorbic acid or ascorbic acid/DFO. As shown in Figure 4, we observed a marked increase in IP1 fluorescence in vitamin C treated cells (Figure 4B) over untreated control samples (Figure 4A), suggesting an expansion of labile Fe²⁺ pools in the presence of ascorbic acid. Moreover, the observed fluorescence increases triggered by vitamin C treatment were attenuated by DFO chelation, confirming that the fluorescence enhancement was due to an elevated level of intracellular labile Fe²⁺ (Figure 4C). As an additional control experiment, no fluorescence increases in vitamin C treated cells were observed by staining with the cobalt-sensitive dye CPI (Figures S7 and S8, Supporting Information), providing further evidence for an iron-dependent effect. Interestingly, ICP-OES analyses of cell lysates show no significant difference in the levels of total iron between ascorbic acid treated cells and control cells (Figure S12, Supporting Information), suggesting that the increases in intracellular fluorescence imaging observed with IP1-AM are likely due to either a mobilization of iron stores into a more labile form and/or a shift in intracellular Fe²⁺/Fe³⁺ ratio within the labile pool, rather than a global change in the total iron pool.

CONCLUDING REMARKS

To summarize, we presented the design, synthesis, spectroscopy, and molecular imaging applications of IP1, a first-generation probe for turn-on detection of Fe²⁺ in aqueous solution and in living cells. IP1 utilizes a bioinspired Fe²⁺- and O₂-dependent oxidative C–O bond cleavage reaction to provide a selective and sensitive design strategy for fluorescence Fe²⁺ detection. Notably, the reaction-based approach described here circumvents fluorescence quenching events commonly associated with open-shell transition metal ions and allows for selective detection of the redox-active metal in a specific oxidation state (i.e., Fe²⁺ over Fe³⁺) as well as selectivity over other s- and d-block metal ions in their bioavailable forms. IP1 is capable of being loaded into live cells and can report both increases and decreases in exchangeable iron pools upon iron supplementation and/or depletion with visible excitation and emission profiles. Moreover, this probe is sensitive enough to detect endogenous, basal labile iron pools in living cells and can identify and visualize expansions in these labile iron pools upon stimulation with either the hormone hepcidin or vitamin C, establishing a starting point for further investigations of dynamic, bioavailable iron stores in cellular iron homeostasis and their physiological and pathological consequences. Efforts to modify both the sensor and the dye portions of this platform to achieve higher signal-to-noise responses, multicolor imaging, and extension to other imaging modalities, along with application of IP1 and related chemical tools for studying the cell and animal biology of iron, are currently underway.

EXPERIMENTAL SECTION

General Methods. All solvents were of reagent grade. Acetonitrile, THF, and dichloromethane (DCM) were dried by passing through activated alumina. *N,N*-Di-isopropylethylamine was dried by distilling over KOH. All commercially purchased chemicals were used as received. Di(2-pyridyl)methylamine **1** was synthesized according to literature procedures.⁷¹ ¹H and ¹³C NMR spectra were obtained from a Bruker AVB-400 NMR spectrometer at the College of Chemistry

NMR facility at UC Berkeley. Signals were internally referenced to solvent residues. Low-resolution mass spectral analyses were carried out using a LCMS (Agilent Technology 6130, Quadrupole LC/MS). High-resolution mass spectral analyses (ESI-MS) were carried out at the College of Chemistry Mass Spectrometry Facility at UC Berkeley. Single-crystal X-ray diffraction was conducted at the College of Chemistry X-ray crystallography facility, UC Berkeley. Crystals were mounted on nylon loops in Paratone-N hydrocarbon oil. Data integration was performed using SAINT. Preliminary data analysis and absorption correction were performed using XPREP and SADABS. Structure solution and refinement was performed using the SHELX software package. X-ray crystallographic data collection was carried out on a Bruker three-circle diffractometer mounted with a SMART 1000 detector using monochromated Mo K α radiation (0.71073 Å) outfitted with a low-temperature, N₂-stream aperture and an APEXII CCD detector and equipped with an Oxford Cryostream 700. The structure was solved using direct methods in conjunction with standard difference Fourier techniques and refined by full-matrix least-squares procedures. A semiempirical absorption correction (SADABS) was applied to the diffraction data. All non-hydrogen atoms were refined anisotropically, and hydrogen atoms were treated as idealized contributions and refined isotropically. All software used for diffraction data processing and crystal-structure solution and refinement are contained in the APEX2 program suite (Bruker AXS, Madison, WI).

Synthesis of 2. To a THF solution of di(2-pyridyl)methylamine (2.4 g, 13 mmol, 50 mL) cooled in an ice bath was added dropwise a solution of *tert*-butyl bromoacetate (2.5 g, 12.8 mmol) and DIPEA (2.3 mL, 13.2 mmol) in 50 mL of THF. The mixture was stirred in the melting ice bath overnight. THF was removed by a rotary evaporator; the residue was dissolved in DCM (50 mL) and washed with saturated NaHCO₃ (20 mL), water (20 mL), and brine (20 mL). The DCM extract was dried over Na₂SO₄ and evaporated to dryness. Residue was purified by basic alumina column (ethyl acetate/hexanes = 8:2). Yield = 2.5 g, 65%. ¹H NMR (400 MHz, CDCl₃, 298 K) δ (ppm): 8.57 (dd, *J* = 0.6 Hz, 4.8 Hz, 2 H), 7.62 (dt, *J* = 2.0 Hz, 9.6 Hz, 2H), 7.44 (d, *J* = 8.0 Hz, 2 H), 7.14 (ddd, *J* = 1.0 Hz, 4.8 Hz, 7.4 Hz, 2 H), 5.12 (s, 1 H), 3.34 (s, 2 H), 1.44 (s, 9 H). ¹³C{¹H} NMR (100.6 MHz, CDCl₃, 298 K) δ (ppm): 171.4, 161.1, 149.4, 136.8, 122.4, 81.3, 68.9, 49.9, 28.2. LRMS (ESI): calcd for C₁₇H₂₂N₃O₂ [M + H]⁺ *m/z* 300.2, found 300.2.

Synthesis of 3. A mixture of **2** (0.99 g, 3.3 mmol), 2,6-dibromomethylpyridine (1.1 g, 4.0 mmol), and Cs₂CO₃ (1.3 g, 4.0 mmol) in 50 mL of MeCN was heated at 40 °C overnight. Insoluble material was removed by filtration. Filtrate was concentrated and purified by basic alumina column (ethyl acetate/hexanes = 8:2 → 9:1). Yield = 0.5 g, 31%. ¹H NMR (400 MHz, CD₃OD, 298 K) δ (ppm): 8.44 (d, *J* = 0.8 Hz, 2 H), 7.84–7.75 (m, 5 H), 7.68 (d, *J* = 7.6 Hz, 1 H), 7.38 (d, *J* = 7.6 Hz, 1 H), 7.25 (ddd, *J* = 1.2 Hz, 4.8 Hz, 7.4 Hz, 2 H), 5.63 (s, 1 H), 4.53 (s, 2 H), 4.03 (s, 2 H), 3.34 (s, 2 H), 1.39 (s, 9 H). ¹³C{¹H} NMR (100.6 MHz, CD₃OD, 298 K) δ (ppm): 171.9, 161.1, 160.7, 157.6, 149.9, 139.3, 138.5, 125.6, 124.1, 123.7, 123.4, 82.2, 74.5, 58.3, 53.4, 33.9, 28.4. LRMS (ESI): calcd for C₂₄H₂₈N₄O₂⁷⁹Br [M + H]⁺ *m/z* 483.1, found 483.2; calcd for C₂₄H₂₈N₄O₂⁸¹Br [M + H]⁺ *m/z* 485.1, found 485.2.

Synthesis of 5. A mixture of **3** (116 mg, 0.24 mmol), **4** (88 mg, 0.26 mmol), and Cs₂CO₃ (78 mg, 0.24 mmol) in 40 mL of DMF was heated at 90 °C overnight. DMF was removed by a rotary evaporator, and residue was redissolved in 40 mL of DCM and filtered. The filtrate was concentrated and purified by a basic alumina column (ethyl acetate/hexanes = 1:1 → 100% ethyl acetate). Yield = 85 mg, 52%. ¹H NMR (400 MHz, CD₃OD, 298 K) δ (ppm): 8.41 (d, *J* = 5.2 Hz, 2 H), 7.81 (d, *J* = 7.2 Hz, 2 H), 7.75–7.71 (m, 3 H), 7.65 (d, *J* = 7.6 Hz, 1 H), 7.38–7.31 (m, 3 H), 7.22–7.16 (m, 3 H), 6.82–6.78 (m, 3 H), 6.73–6.66 (m, 3 H), 6.58 (dd, *J* = 2.4 Hz, 8.8 Hz, 1 H), 5.64 (s, 1 H), 5.22 (s, 2 H), 5.10 (s, 2 H), 4.05 (s, 2 H), 3.73 (s, 3 H), 3.34 (s, 2 H), 1.36 (s, 9 H). ¹³C{¹H} NMR (100.6 MHz, CD₃OD, 298 K) δ (ppm): 171.9, 161.9, 161.2, 160.5, 160.4, 157.4, 152.6, 149.9, 146.2, 140.1, 139.0, 138.5, 131.2, 131.0, 129.5, 129.3, 125.5, 124.7, 124.1, 123.5, 121.9, 121.4, 118.8, 118.1, 112.7, 112.1, 102.6, 101.3, 85.0, 82.1, 74.6,

72.9, 71.5, 58.4, 55.9, 53.4, 28.4. LRMS (ESI): calcd for $C_{45}H_{43}N_4O_6$ $[M + H]^+$ m/z 735.3, found 735.3.

Synthesis of Iron Probe 1 (IP1). Compound 5 (85 mg, 0.11 mmol) was dissolved in 4 mL of DCM under N_2 followed by addition of TFA (0.9 mL, 12 mmol). The mixture was stirred at room temperature under N_2 overnight. Solvent and excess acid were removed by a rotary evaporator; residue was redissolved in DCM and washed with diluted $NaHCO_3$ and water until the pH of the aqueous phase was between 7–8. The organic component was then washed with brine and dried over Na_2SO_4 , and DCM was removed to give the product. Yield = 74 mg, quant. 1H NMR (400 MHz, CD_3OD , 298 K) δ (ppm): 8.50 (d, $J = 4.8$ Hz, 2 H), 7.78–7.73 (m, 3 H), 7.55 (d, $J = 8.0$ Hz, 2 H), 7.46–7.37 (m, 4 H), 7.32–7.28 (m, 2 H), 7.23 (t, $J = 7.6$ Hz, 1 H), 6.86–6.81 (m, 3 H), 6.75–6.69 (m, 3 H), 6.62 (dd, $J = 2.4$ Hz, 8.8 Hz, 1 H), 5.49 (s, 1 H), 5.25 (s, 2 H), 5.21 (s, 2 H), 4.22 (s, 2 H), 3.78 (s, 3 H), 3.56 (s, 2 H). $^{13}C\{^1H\}$ NMR (100.6 MHz, $CDCl_3$, 298 K) δ (ppm): 174.0, 160.4, 159.0, 158.8, 158.3, 156.4, 151.6, 149.0, 144.9, 139.3, 138.0, 137.1, 130.1, 129.9, 128.5, 125.4, 124.5, 124.0, 123.0, 121.8, 120.8, 120.2, 117.8, 117.1, 111.7, 111.3, 101.8, 100.5, 83.6, 72.1, 71.1, 70.4, 59.3, 55.7, 55.6. HRMS (ESI): calcd for $C_{41}H_{35}N_4O_6$ $[M + H]^+$ m/z 679.2557, found 679.2548.

Synthesis of IP1-AM. IP1 (74 mg, 0.11 mmol) was dissolved in 10 mL of DCM, and bromomethyl acetate (15 μ L, 0.15 mmol) and Et_3N (20 μ L, 0.15 mmol) were added. The mixture was stirred at room temperature overnight. Volatiles were removed by a rotary evaporator, and residue was purified by a basic alumina column (ethyl acetate/hexanes = 8:2 \rightarrow 100% ethyl acetate \rightarrow ethyl acetate/MeOH = 8:2). Yield = 20 mg, 24%. 1H NMR (400 MHz, CD_3OD , 298 K) δ (ppm): 8.43 (d, $J = 4.4$ Hz, 2 H), 7.80–7.76 (m, 3 H), 7.46–7.37 (m, 4 H), 7.25–7.23 (m, 2 H), 6.85–6.81 (m, 4 H), 6.79–6.77 (m, 2 H), 6.74–6.70 (m, 2 H), 6.64–6.61 (m, 2 H), 5.64 (s, 1 H), 5.26 (s, 2 H), 5.25 (s, 2 H), 5.15 (s, 2 H), 4.05 (s, 2 H), 3.80 (s, 3 H), 3.78 (s, 3 H), 3.53 (s, 3 H). $^{13}C\{^1H\}$ NMR (100.6 MHz, CD_3OD , 298 K) δ (ppm): 183.1, 171.3, 162.0, 160.6, 152.8, 151.2, 150.0, 140.1, 139.4, 139.1, 138.6, 131.3, 131.1, 129.5, 129.4, 125.6, 124.7, 124.2, 122.0, 121.6, 118.0, 112.7, 112.1, 102.7, 101.3, 72.9, 67.8, 61.5, 56.0. HRMS (ESI): calcd for $C_{44}H_{39}N_4O_8$ $[M + H]^+$ m/z 751.2768, found 751.2761.

Fluorescence Spectroscopy. Fluorescence spectra were recorded on a Photon Technology International Quanta Master 4 L-format scanning spectrofluorometer (Lawrenceville, NJ) equipped with an LPS-220B 75-W xenon lamp and power supply, A-1010B lamp housing with an integrated igniter, switchable 814 photon-counting/analog photomultiplier detection unit, and MDS020 motor driver. Samples for emission measurements were contained in a quartz cuvette with a path length of 1 cm and 1.5 mL cell volume (Starna, Atascadero, CA).

Millipore water was used to prepare all aqueous solutions. Spectroscopic measurements were performed in 50 mM Tris buffer at pH 7.6. A typical sample for fluorescence measurement of IP1 in the presence of 20 equiv of Fe^{2+} was prepared as follows: 1 μ L of a stock solution of IP1 (1 mM in DMSO) was added to 50 mM Tris, followed by 1 μ L of 20 mM $Fe(NH_4)_2(SO_4)_2 \cdot 6H_2O$ and 500 μ L of 10 mM glutathione (GSH) to a final volume of 1000 μ L to give a solution of IP1, Fe^{2+} , and GSH at final concentrations of 1 μ M, 20 μ M, and 5 mM, respectively. The solution was air oxidized and mixed well with a plastic disposable pipet after addition of each reagent and at 5 min intervals until the emission spectrum was recorded. For metal selectivity study, aqueous metal solutions of NaCl, KCl, $MgCl_2 \cdot 6H_2O$, $CaCl_2 \cdot 2H_2O$, $MnCl_2 \cdot 4H_2O$, $FeCl_3 \cdot 6H_2O$, $CoCl_2$, $NiCl_2 \cdot 6H_2O$, $CuCl_2 \cdot 2H_2O$, and $ZnCl_2$ and MeCN solution of $Cu(MeCN)_4PF_6$ at 20 mM and cobalamin at 2 mM were used. For fluorescence measurement in the absence of O_2 , Tris buffer was degassed by bubbling N_2 for 30 min, and all solutions were prepared from the degassed buffer. Excitation was at 470 nm, and emission spectra were collected from 480–600 nm.

ESI-MS Product Analysis. Samples for ESI-MS analysis were prepared as follows: 20 μ L of a 1 mM IP1 solution in DMSO was added to 440 μ L of 50 mM Tris at pH 7.6, followed by addition of 40 μ L of a 10 mM FAS solution in the same buffer. The mixture was allowed to react for 1 h, which was then injected into a LC-MS

(Agilent Technology 6130, Quadrupole LC/MS) and analyzed by flow injection analysis.

Cell Culture and Confocal Microscopy. Cells were grown in the UC Berkeley Tissue Culturing Facility. HepG2/C3A cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Invitrogen) supplemented with 10% fetal bovine serum (FBS, Hyclone) and incubated at 37 $^\circ$ C in 5% CO_2 . One or two days before imaging, cells were passaged and plated in phenol red-free medium on 4-well chamber slides (Corning, Corning, NY) and allowed to grow to 50–70% confluence. Iron enrichment was performed in the same medium with supplementation with 100 μ M FAS with an equal concentration of sodium citrate for 20 h. For cells that required hepcidin treatment, the medium was first supplemented with 100 μ M FAS as described above and incubated for 1 h, a stock solution of hepcidin (AnaSpec) in Milli-Q water was added, and cells were coincubated with FAS and hepcidin for a further 19 h. DFO (Sigma Aldrich) and ascorbic acid (Sigma Aldrich) treatment were performed at the last 40 min of the cell culture period. Immediately before staining or DFO treatment the cells were washed twice with 200 μ M EDTA in PBS. A solution of IP1-AM in DMSO (20 mM) was diluted into DMEM at 20 μ M, added to the cells, and incubated for 2 h. Before imaging, dye-containing medium was removed and cells were washed twice with PBS. For nuclear staining, cells were incubated with 1 μ M Hoechst at 37 $^\circ$ C for 15 min prior to imaging.

Confocal fluorescence imaging studies were performed with a Zeiss laser scanning microscope 710 with a 40x water-immersion objective lens, with Zen 2009 software (Carl Zeiss). The dye was excited at 488 nm with an Ar laser, and emission was collected using a META detector between 500 and 625 nm. Hoechst 33342 was excited with a 405 nm diode laser, and emission was collected using a META detector between 410 and 490 nm. Image analysis was performed using ImageJ (National Institute of Health). For quantitative analysis of fluorescence of the confocal images, the threshold of the images was set to 10, the area of fluorescence was selected with the "create selection" function, and the fluorescent intensity of the whole image was measured.

WST-1 Assay. Cells were grown in the UC Berkeley Tissue Culturing Facility. HepG2/C3A cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Invitrogen) supplemented with 10% fetal bovine serum (FBS, Hyclone) and incubated at 37 $^\circ$ C in 5% CO_2 . Two days before the experiment, the cells were passaged and plated in phenol red-free medium on clear-bottom, black 96-well plates and allowed to grow to 100% confluence. Iron enrichment and IP1-AM treatment were performed as described for confocal microscopy experiment. At the end of the incubation period, medium was removed and cells were washed with 200 μ M EDTA and PBS. WST-1 reagent (Roche Applied Science), diluted by DMEM (1:10 v/v, 100 μ M), was then added to cells and incubated for 2 h. Absorbance at 450 nm was recorded by a plate reader.

ICP-OES. Cells were grown in the UC Berkeley Tissue Culturing Facility. HepG2/C3A cells were cultured in Dulbecco's Modified Eagle Medium (DMEM, Invitrogen) supplemented with 10% fetal bovine serum (FBS, Hyclone) and incubated at 37 $^\circ$ C in 5% CO_2 . Two days before the experiment, the cells were passaged and plated in 10 cm culture dish and allowed to grow to 100% confluence. Iron enrichment and hepcidin and ascorbate treatment were performed as described for the confocal microscopy experiment. At the end of the incubation period, medium was removed and cells were washed twice with PBS. Cells were then scraped in PBS, transferred to a 15 mL Falcon tube, and resuspended in 5 mL of PBS. Aliquots of 100 μ L cell suspension were transferred to a 1.5 mL microtube for BCA protein assay. The cell suspension was pelleted by centrifugation, supernatant was carefully removed, and the cell pellet was digested in 200 μ L of 70% HNO_3 at 90 $^\circ$ C for 4 h and diluted with Milli-Q water to a total volume of 5 mL. Samples were analyzed by an ICP Optima 7000 DV instrument in the Microanalytical Facility at the College of Chemistry, UC Berkeley. Iron signal was detected at 238.204 nm with an internal 10 ppm Ga standard at 417.206 nm.

■ ASSOCIATED CONTENT

■ Supporting Information

Experimental details, including synthesis of **4**, **6**, and **6-FeCl**, ESI-MS spectra, ICP-OES data, and fluorescence and cell data. This material is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

chrischang@berkeley.edu

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

This work was supported by the NIH (GM79465), Packard Foundation, and UC Berkeley. H.Y.A. thanks the Croucher Foundation for a postdoctoral fellowship. J.C. thanks the Human Frontiers Science Program for a postdoctoral fellowship, and T.C. was supported by a scholarship from the Ministry of Science, Thailand. C.J.C. is an Investigator of the Howard Hughes Medical Institute. We thank Ann Fischer and Xiaozhu Zhang (UC Berkeley Tissue Culture Facility) and Elena Kreimer (UC Berkeley College of Chemistry) for expert technical assistance.

■ REFERENCES

- (1) Lippard, S. J.; Berg, J. M. *Principles of Bioinorganic Chemistry*; University Science Book: Mill Valley, CA, 1994.
- (2) Aisen, P.; Enns, C.; Wessling-Resnick, M. *Int. J. Biochem. Cell Biol.* **2001**, *33*, 940.
- (3) Hentze, M. W.; Muckenthaler, M. U.; Andrews, N. C. *Cell* **2004**, *117*, 285.
- (4) Kaplan, C. D.; Kaplan, J. *Chem. Rev.* **2009**, *109*, 4536.
- (5) Theil, E. C.; Goss, D. J. *Chem. Rev.* **2009**, *109*, 4568.
- (6) Crichton, R. R.; Wilmet, S.; Leggsyter, R.; Ward, R. J. *J. Inorg. Biochem.* **2002**, *91*, 9.
- (7) Salahudeen, A. A.; Bruick, R. K. *Ann. N.Y. Acad. Sci.* **2009**, *1177*, 30.
- (8) Schultz, I. J.; Chen, C.; Paw, B. H.; Hamza, I. J. *Biol. Chem.* **2010**, *285*, 26753.
- (9) Philpott, C. C. *J. Biol. Chem.* **2012**, *287*, 13518.
- (10) Domaille, D. W.; Que, E. L.; Chang, C. J. *Nat. Chem. Bio.* **2008**, *4*, 168.
- (11) Que, E. L.; Domaille, D. W.; Chang, C. J. *Chem. Rev.* **2008**, *108*, 1517.
- (12) McRae, R.; Bagchi, P.; Sumalekshmy, S.; Fahrni, C. J. *Chem. Rev.* **2009**, *109*, 4780.
- (13) Haas, K. L.; Franz, K. J. *Chem. Rev.* **2009**, *109*, 4921.
- (14) Sahoo, S. K.; Sharma, D.; Bera, R. K.; Crisponi, G.; Callan, J. F. *Chem. Soc. Rev.* **2012**, *41*, 7195.
- (15) Chan, J.; Dodani, S. C.; Chang, C. J. *Nat. Chem.* **2012**, *4*, 973.
- (16) Breuer, W.; Shvartsman, M.; Cabantchik, Z. I. *Int. J. Biochem. Cell Biol.* **2008**, *40*, 350.
- (17) Petrat, F.; de Groot, H.; Rauen, U. *Biochem. J.* **2001**, *356*, 61.
- (18) Richardson, D. R.; Lane, D. J. R.; Becker, E. M.; Huang, M. L.-H.; Whitnall, M.; Rahmanto, Y. S.; Sheftel, A. D.; Ponka, P. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 10775.
- (19) Breuer, W.; Epsztejn, S.; Millgram, P.; Cabantchik, I. Z. *Am. J. Physiol. Cell Physiol.* **1995**, *268*, C1354.
- (20) Petrat, F.; Weisheit, D.; Lensen, M.; de Groot, H.; Sustmann, R.; Rauen, U. *Biochem. J.* **2002**, *362*, 137.
- (21) Qu, X.; Liu, Q.; Ji, X.; Chen, H.; Zhou, Z.; Shen, Z. *Chem. Commun.* **2012**, *48*, 4600.
- (22) Mei, Q.; Jiang, C.; Guan, G.; Zhang, K.; Liu, B.; Liu, R.; Zhang, Z. *Chem. Commun.* **2012**, *48*, 7468.
- (23) For a ratiometric, turn-off Fe(II) probe, see: Li, P.; Fang, L.; Zhou, H.; Zhang, W.; Wang, X.; Li, N.; Zhong, H.; Tang, B. *Chem.—Eur. J.* **2011**, *17*, 10520.
- (24) Petrat, F.; Rauen, U.; de Groot, H. *Hepatology* **1999**, *29*, 1171.
- (25) Rauen, U.; Springer, A.; Weisheit, D.; Petrat, F.; Korth, H. G.; de Groot, H.; Sustmann, R. *ChemBioChem* **2007**, *8*, 341.
- (26) Bricks, J. L.; Kovalchuk, A.; Trieflinger, C.; Nofz, M.; Büschel, M.; Tolmachev, A. I.; Daub, J.; Rurack, K. *J. Am. Chem. Soc.* **2005**, *127*, 13522.
- (27) Kennedy, D. P.; Kormos, C. M.; Burdette, S. C. *J. Am. Chem. Soc.* **2009**, *131*, 8578.
- (28) Wang, B.; Hai, J.; Liu, Z.; Wang, Q.; Yang, Z.; Sun, S. *Angew. Chem., Int. Ed.* **2010**, *49*, 4576.
- (29) Lee, M. H.; Giap, T. V.; Kim, S. H.; Lee, Y. H.; Kang, C.; Kim, J. S. *Chem. Commun.* **2010**, *46*, 1407.
- (30) Yang, Z.; She, M.; Yin, B.; Cui, J.; Zhang, Y.; Sun, W.; Li, J.; Shi, Z. *J. Org. Chem.* **2012**, *77*, 1143.
- (31) Wang, R.; Yu, F.; Liu, P.; Chen, L. *Chem. Commun.* **2012**, *48*, 5310.
- (32) Wei, Y.; Aydin, Z.; Zhang, Y.; Liu, Z.; Guo, M. *ChemBioChem* **2012**, *13*, 1569.
- (33) Hirayama, T.; Okuda, K.; Nagasawa, H. *Chem. Sci.* **2013**, *4*, 1250.
- (34) Czarnik, A. W. *Acc. Chem. Res.* **1994**, *27*, 302.
- (35) Kim, H. N.; Lee, M. H.; Kim, H. J.; Kim, J. S.; Yoon, J. *Chem. Soc. Rev.* **2008**, *37*, 1465.
- (36) Cho, D. G.; Sessler, J. L. *Chem. Soc. Rev.* **2009**, *38*, 1647.
- (37) Jun, M. E.; Roy, B.; Ahn, K. H. *Chem. Commun.* **2011**, *47*, 7583.
- (38) Du, J.; Hu, M.; Fan, J.; Peng, X. *Chem. Soc. Rev.* **2012**, *41*, 4511.
- (39) Ortiz de Montellano, P. R. *Cytochrome P450: Structure, Mechanism, and Biochemistry*, 3rd ed.; Kluwer Academic, Plenum Publishers: New York, 2005.
- (40) Groves, J. T. *J. Inorg. Biochem.* **2006**, *100*, 434.
- (41) Nam, W. *Acc. Chem. Res.* **2007**, *40*, 522.
- (42) Kovaleva, E. G.; Neiberger, M. B.; Chakrabarty, S.; Lipscomb, J. D. *Acc. Chem. Res.* **2007**, *40*, 475.
- (43) Sono, M.; Roach, M. P.; Coulter, E. D.; Dawson, J. H. *Chem. Rev.* **1996**, *96*, 2841.
- (44) Costas, M.; Mehn, M. P.; Jensen, M. P.; Que, L., Jr. *Chem. Rev.* **2004**, *104*, 939.
- (45) Tshuva, E. Y.; Lippard, S. J. *Chem. Rev.* **2004**, *104*, 987.
- (46) Krebs, C.; Fujimori, D. G.; Walsh, C. T.; Bollinger, J. M., Jr. *Acc. Chem. Res.* **2007**, *40*, 484.
- (47) Taki, M.; Iyoshi, S.; Ojida, A.; Hamachi, I.; Yamamoto, Y. *J. Am. Chem. Soc.* **2010**, *132*, 5938.
- (48) Au-Yeung, H. Y.; New, E. J.; Chang, C. J. *Chem. Commun.* **2012**, *48*, 5268.
- (49) Ekkati, A. R.; Kodanko, J. J. *J. Am. Chem. Soc.* **2007**, *129*, 12390.
- (50) Song, D.; Lim, J. M.; Cho, S.; Park, S.-J.; Cho, J.; Kang, D.; Rhee, S. G.; You, Y.; Nam, W. *Chem. Commun.* **2012**, *48*, 5449.
- (51) Han, Y.; You, Y.; Lee, Y. M.; Nam, W. *Adv. Mater.* **2012**, *24*, 2748.
- (52) McDonald, A. R.; Guo, Y.; Vu, V. V.; Bominaar, E. L.; Münck, E.; Que, L. *Chem. Sci.* **2012**, *3*, 1680.
- (53) A pseudo-first-order rate constant of the oxidative release of the fluorophore has been estimated to be $1 \times 10^{-5} \text{ s}^{-1}$.
- (54) To date, only eight noncorrinoid cobalt proteins have been identified and characterized, which are all from bacterial systems. The loosely bound form of cobalt is even less known and studied. See: Kobayashi, M.; Shimizu, S. *Eur. J. Biochem.* **1999**, *261*, 1. Okamoto, S.; Eltis, L. D. *Metallomics* **2011**, *3*, 963.
- (55) Ståhlberg, K. G.; Radner, S.; Nordén, Å. *Scand. J. Haematol.* **1967**, *4*, 312.
- (56) Fluorescence was indeed attenuated by around 50% in the presence of 0.1–10% of Co^{2+} but enhanced by around 30% in a 1:1 mixture of Fe^{2+} and Co^{2+} , indicating the possibility of a different bimetallic mechanism that is dependent on the relative concentrations of the metals.
- (57) Anderson, E. R.; Shah, Y. M. *Comp. Physiol.* **2013**, *3*, 315.

- (58) Tsien, R. Y. *Nature* **1981**, *290*, 527.
- (59) Izumi, S.; Urano, Y.; Hanaoka, K.; Terai, T.; Nagano, T. *J. Am. Chem. Soc.* **2009**, *131*, 10189.
- (60) McQuade, L. E.; Ma, J.; Lowe, G.; Ghatpande, A.; Gelperin, A.; Lippard, S. J. *Proc. Natl. Acad. Sci. U.S.A.* **2010**, *107*, 8525.
- (61) Dickinson, B. C.; Peltier, J.; Stone, D.; Schaffer, D. V.; Chang, C. *J. Nat. Chem. Biol.* **2011**, *7*, 106.
- (62) Terman, A.; Kurz, T. *Antioxid. Redox Signaling* **2013**, *18*, 888.
- (63) Park, C. H.; Valore, E. V.; Waring, A. J.; Ganz, T. *J. Biol. Chem.* **2001**, *276*, 7806.
- (64) Nicolas, G.; Viatte, L.; Bennoun, M.; Beaumont, C.; Kahn, A.; Vaulont, S. *Blood Cells Mol. Dis.* **2002**, *29*, 327.
- (65) Ganz, T. *Blood* **2003**, *102*, 783.
- (66) Nemeth, E.; Tuttle, M. S.; Powelson, J.; Vaughn, M. B.; Donovan, A.; Ward, D. M.; Ganz, T.; Kaplan, J. *Science* **2004**, *306*, 2090.
- (67) Bridges, K. R.; Hoffman, K. E. *J. Biol. Chem.* **1986**, *261*, 14273.
- (68) Toth, I.; Rogers, J. T.; McPhee, J. A.; Elliott, S. M.; Abramson, S. L.; Bridges, K. R. *J. Biol. Chem.* **1995**, *270*, 2846.
- (69) Lane, D. J. R.; Lawen, A. J. *Biol. Chem.* **2008**, *283*, 12701.
- (70) Brewer, C.; Otto-Duessel, M.; Lykkesfeldt, J.; Nick, H.; Wood, J. C. *Exp. Hematol.* **2012**, *40*, 820.
- (71) Renz, M.; Hemmert, C.; Meunier, B. *Eur. J. Org. Chem.* **1998**, 1271–1273.