

UC Berkeley

UC Berkeley Previously Published Works

Title

Leaf venation networks of Bornean trees: images and hand-traced segmentations

Permalink

<https://escholarship.org/uc/item/6b1423rg>

Journal

Ecology, 100(11)

ISSN

0012-9658

Authors

Blonder, Benjamin

Both, Sabine

Jodra, Miguel

et al.

Publication Date

2019-11-01

DOI

10.1002/ecy.2844

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Ecology, 100(11), 2019, e02844

© 2019 The Authors. *Ecology* published by Wiley Periodicals, Inc. on behalf of The Ecological Society of America.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Leaf venation networks of Bornean trees: images and hand-traced segmentations

BENJAMIN BLONDER,^{1,4,6} SABINE BOTH,^{2,5} MIGUEL JODRA,¹ NOREEN MAJALAP,³ DAVID BURSLEM,²
YIT ARN TEH,² AND YADVINDER MALHI¹

¹*Environmental Change Institute, School of Geography and the Environment, University of Oxford, South Parks Road, Oxford OX1 3QY, UK*

²*School of Biological Sciences, University of Aberdeen, Zoology Building, Tillydrone Avenue, Aberdeen AB24 2TZ, UK*
³*Forest Research Centre, Sabah Forestry Department, Peti Surat 1407, 90715 Sandakan, Sabah, Malaysia*

Citation: Blonder, B., S. Both, M. Jodra, N. Majalap, D. Burslem, Y. A. Teh, and Y. Malhi. 2019. Leaf venation networks of Bornean trees: images and hand-traced segmentations. *Ecology* 100(11):e02844. 10.1002/ecy.2844

Abstract. The data set contains images of leaf venation networks obtained from tree species in Malaysian Borneo. The data set contains 726 leaves from 295 species comprising 50 families, sampled from eight forest plots in Sabah. Image extents are approximately 1 × 1 cm, or 50 megapixels. All images contain a region of interest in which all veins have been hand traced. The complete data set includes over 30 billion pixels, of which more than 600 million have been validated by hand tracing. These images are suitable for morphological characterization of these species, as well as for training of machine-learning algorithms that segment biological networks from images. Data are made available under the Open Data Commons Attribution License. You are free to copy, distribute, and use the database; to produce works from the database; and to modify, transform, and build upon the database. You must attribute any public use of the database, or works produced from the database, in the manner specified in the license. For any use or redistribution of the database, or works produced from it, you must make clear to others the license of the database and keep intact any notices on the original database.

Key words: botany; cleared leaf; ecology; image analysis; image segmentation; leaf venation; machine learning; plant ecophysiology; tropical forest; tropical forest; vein network; venation network.

The complete data sets corresponding to abstracts published in the Data Papers section in the journal are published electronically as Supporting Information in the online version of this article at <http://onlinelibrary.wiley.com/doi/10.1002/ecy.2844/supinfo>.

DATA AVAILABILITY

Because of file size, data are permanently available at the Oxford Research Archive at <https://ora.ox.ac.uk/objects/uuid:de65fc07-4b8f-4277-a6c4-82836afbdeb3>.

Manuscript received 21 June 2019; accepted 8 July 2019. Corresponding Editor: William K. Michener.

⁴ Present address: School of Life Sciences, Arizona State University, Tempe, Arizona, USA.

⁵ Present address: School of Environmental and Rural Science, University of New England, Armidale, New South Wales, Australia.

⁶ E-mail: bblonder@gmail.com