

UC San Diego

UC San Diego Previously Published Works

Title

Variants Associated with the Ankle Brachial Index Differ by Hispanic/Latino Ethnic Group: a genome-wide association study in the Hispanic Community Health Study/Study of Latinos

Permalink

<https://escholarship.org/uc/item/6bp8f1js>

Journal

Scientific Reports, 9(1)

ISSN

2045-2322

Authors

Sofer, Tamar
Emery, Leslie
Jain, Deepti
[et al.](#)

Publication Date

2019

DOI

10.1038/s41598-019-47928-5

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

OPEN

Variants Associated with the Ankle Brachial Index Differ by Hispanic/Latino Ethnic Group: a genome-wide association study in the Hispanic Community Health Study/Study of Latinos

Tamar Sofer^{1,2}, Leslie Emery³, Deepti Jain³, Alicia M. Ellis⁴, Cathy C. Laurie³, Matthew A. Allison⁵, Jiwon Lee¹, Nuzulul Kurniansyah¹, Kathleen F. Kerr³, Hector M. González⁶, Wassim Tarraf⁷, Michael H. Criqui⁵, Leslie A. Lange⁸, Walter R. Palmas⁹, Nora Franceschini¹⁰ & Christina L. Wassel¹¹

Lower extremity peripheral artery disease (PAD) burden differs by race/ethnicity. Although familial aggregation and heritability studies suggest a genetic basis, little is known about the genetic susceptibility to PAD, especially in non-European descent populations. Genome-wide association studies (GWAS) of the ankle brachial index (ABI) and PAD (defined as an ABI < 0.90) have not been conducted in Hispanics/Latinos. We performed a GWAS of PAD and the ABI in 7,589 participants aged >45 years from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). We also performed GWAS for ABI stratified by Hispanic/Latino ethnic subgroups: Central American, Mexican, and South American (Mainland group), and Cuban, Dominican, and Puerto Rican (Caribbean group). We detected two genome-wide significant associations for the ABI in *COMMD10* in Puerto Ricans, and at *SYBU* in the Caribbean group. The lead SNP rs4466200 in the *COMMD10* gene had a replication $p = 0.02$ for the ABI in Multi-Ethnic Study of Atherosclerosis (MESA) African Americans, but it did not replicate in African Americans from the Cardiovascular Health Study (CHS). In a regional look-up, a nearby SNP rs12520838 had Bonferroni adjusted $p = 0.05$ (unadjusted $p = 7.5 \times 10^{-5}$) for PAD in MESA Hispanics. Among three suggestive associations ($p < 10^{-7}$) in subgroup-specific analyses, *DMD* on chromosome X, identified in Central Americans, replicated in MESA Hispanics ($p = 2.2 \times 10^{-4}$). None of the previously reported ABI and PAD associations in whites generalized to Hispanics/Latinos.

The ankle brachial index (ABI) is the ratio of systolic blood pressure in the ankle to the arm, and reflects the degree of atherosclerotic obstruction in the lower extremity arteries. The ABI has been the major clinical diagnostic criterion for peripheral artery disease (PAD) for more than 40 years¹. PAD, defined either by ABI ≤ 0.90

¹Division of Sleep and Circadian Disorders, Brigham and Women's Hospital, Boston, MA, USA. ²Division of Sleep Medicine, Harvard Medical School, Boston, MA, USA. ³Department of Biostatistics, University of Washington, Seattle, WA, USA. ⁴Duke Clinical Research Institute, Duke University, Durham, NC, USA. ⁵Department of Family Medicine and Public Health, University of California – San Diego, La Jolla, CA, USA. ⁶Department of Neurosciences, Shiley-Marcos Alzheimer's Disease Research Center, University of California San Diego, San Diego, CA, United States of America. ⁷Institute of Gerontology, Wayne State University, Detroit, MI, USA. ⁸Division of Biomedical Informatics and Personalized Medicine, University of Colorado-Denver, Denver, CO, USA. ⁹Department of Medicine, Columbia University, New York, NY, USA. ¹⁰Department of Epidemiology, University of North Carolina, Chapel Hill, NC, USA. ¹¹Applied Sciences, Premier, Inc, Charlotte, NC, USA. Nora Franceschini and Christina L. Wassel contributed equally. Correspondence and requests for materials should be addressed to T.S. (email: tsofer@bwh.harvard.edu)

or via other criteria, is associated with an increased risk for incident cardiovascular disease (CVD) events and mortality^{2–6}. Even borderline ABI values (0.90–1.00) are associated with increased risk of mortality⁷, as well as mobility impairment, including inability to walk 1/4 mile or climb one flight of stairs, as well as inability to complete a 6 minute walk⁷. The burden of PAD is greater in African-Americans and Cuban Americans compared to non-Hispanic whites and other Hispanic groups such as Mexican Americans^{8,9}. However, these race/ethnic differences have not been explained by traditional, genetic, or novel risk factors to date^{10–12}. Although familial aggregation and heritability estimates suggest a significant genetic contribution, little is known about the genetic susceptibility to PAD in non-European populations^{13–16}.

GWAS conducted in European ancestry populations have identified the 9p21 locus significantly associated with ABI¹⁷, although associations were attenuated and no longer genome-wide significant after accounting for coronary artery disease (CAD). Additionally, a variant in *TCF7L2* was significantly associated with ABI in a large-scale candidate gene (~50 K SNPs) analysis of European ancestry (n = 21,000), but this association failed to replicate in independent samples¹². No significant associations were observed in more than 7,000 African-Americans in a candidate gene array study¹². In general, previous studies have been limited by discovery of loci that have lacked specificity for PAD (e.g. 9p21 is also associated with CHD), a lack of racial/ethnic diversity, or a lack of robust independent replication of initial findings. In particular, no previous studies have examined the genetic variants underlying the ABI and PAD in Hispanic/Latino populations.

In this study, we sought to identify novel loci associations with ABI and PAD in a large cohort of US Hispanics/Latinos from the Hispanic Community Health Study/Study of Latinos (HCHS/SOL) using GWAS, replicate novel associations in independent samples, and to study whether previously identified genetic variants in European ancestry generalize to Hispanics/Latinos. We also performed GWAS for ABI in subgroups based on participant background, using genetically estimated ancestry and self-reported ethnicity, while taking into account their genetic and environmental heterogeneity.

Methods

HCHS/SOL population. The HCHS/SOL^{18,19} is a community-based cohort study of 16,415 self-identified Hispanic/Latino individuals aged 18–74 years and selected from households in predefined census-block groups across four US field centers (Chicago, IL; Miami, FL; Bronx, NY; and San Diego, CA). The HCHS/SOL baseline clinical examination occurred between 2008 and 2011 and included comprehensive biological, behavioral, and sociodemographic assessments. This study was approved by the institutional review boards (IRBs) at each field center, where all participants gave written informed consent, and by the Non-Biomedical IRB at the University of North Carolina at Chapel Hill, to the HCHS/SOL Data Coordinating Center. See Supplementary Material Section 3 for a complete list of IRBs that approved this study. This research was performed in accordance with relevant guidelines and regulations. The current study included participants who were aged >45 with measured ABI, and who had given DNA consent.

The cohort includes participants who self-identified as having a Hispanic/Latino background. The largest subgroups are Central American, Mexican, South American (comprising the Mainland group), Cuban, Dominican, and Puerto Rican (comprising the Caribbean group). Genetic analysis groups were based on these self-identified subgroups and on genetic similarity between individuals, as described in detail in Conomos, *et al.*²⁰. In brief, clusters of individuals in the genetic principal components space were constructed while accounting for self-identified group. Therefore, these subgroups mostly overlap with self-reported background, while recovering classification in some individuals who did not report ethnicity, and in a few instances, assigning individuals who self-reported to be from one subgroup, to a different one²⁰. Thus, we used these genetic analysis subgroups consequently, rather than self-reported background.

Assessment of ABI and outcomes definitions. For each of the left and right side, the ABI was calculated as the maximum systolic blood pressure in the posterior tibial artery or the dorsalis pedis artery in the same leg, divided by the maximum systolic blood pressures in the left and right brachial arteries. The overall composite ABI was then calculated for each participant as the minimum of the left and right side ABI. A participant was classified as having PAD if the overall ABI was ≤ 0.90 . We also defined a “borderline PAD” as having either ABI ≤ 1.00 . These classifications are visualized in Fig. S1 in the Supplementary Information.

Inclusion and exclusions criteria and study sample. The study sample included individuals aged 45–74 years (N = 7,662 genotyped individuals). Further exclusions were: missing ABI components of blood pressure data (N = 60), and primarily Asian genetic ancestry (n = 13, genetic outliers²⁰), for a total of 7,589 individuals included in the primary ABI analysis. In the stratified analysis by Hispanic/Latino background group, additional 20 participants were excluded due to missing group, so final sample size numbers were: Central American (n = 760), Cuban (n = 1,500), Dominican (n = 700), Mexican (n = 2,619), Puerto Rican (n = 1,437), and South American (n = 553). For PAD and borderline PAD analyses, we excluded 192 individuals with arterial stiffness (ABI > 1.4), for N = 7,397 individuals included.

Genotyping and imputation. HCHS/SOL individuals who consented for participation in genetic studies were genotyped using a HCHS/SOL custom 15041502 B3 array at Illumina. This array consisted of the Illumina Omni 2.5 M array (HumanOmni2.5-8 v.1-1), and additional ~150,000 SNPs selected based on multiple 1000 Genomes²¹ phase 1 data groups (CLM: Colombian in Medellin, Colombia); MXL: Mexican Ancestry in Los Angeles, California; and PUR: Puerto Rican in Puerto Rico), to increase the captured Amerindian genetic variation²². We applied standardized quality-assurance and quality-control (QA/QC) methods²³ to generate recommended SNP- and sample-level quality filters, as previously described in Conomos, *et al.*²⁰, who also provide comprehensive details of imputation. Following filtering for quality and informativeness (polymorphic

and unduplicated), 2,232,944 SNPs were carried forward for imputation and downstream association analyses. Imputation was performed using the complete 1000 Genome phase 1 reference panel ($n = 1,092$)²¹. Genotypes were first pre-phased with SHAPEIT2 (v.2.r644)²⁴ and then imputed with IMPUTE2 (v.2.3.0)²⁵. Only variants with at least two copies of the minor allele present in any of the four 1000 Genomes continental panels were imputed. We performed downstream association analyses on the results 27,887,661 variants, and considered only variants with imputation quality $oevar > 0.3$ and $MAF \geq 1\%$.

Association testing. Since ABI had a highly skewed distribution (Fig. S1), and to prevent spurious associations, we winsorized ABI values so that the highest 1% of the values of ABI were set to the value of the 99th percentile of the distribution, and the lowest 1% of the values were set to the value of the 1st percentile of the distribution. After winsorization, the distribution was approximately normal. As a sensitivity analysis, for the top results we also report results from analysis that removed individuals with $ABI > 1.4$.

To examine the association between genetic variants and the ABI, we fit a linear mixed model adjusted for sex, age, study center, sampling weights (to prevent potential selection bias due to the study sampling scheme), and the five first principal components as fixed-effects. We also had random-effects corresponding to genetic relatedness, household, and block unit sharing. For the binary traits of PAD and borderline PAD, we used the GMMAT algorithm²¹, which calculates the score test for each genetic variant based on a logistic mixed model. We used the same fixed- and random-effects as in the analysis of ABI.

For ABI, we investigated association results for variants with $MAF \geq 0.01$, and (if imputed) imputation quality $oevar \geq 0.3$. For the binary traits of PAD and borderline PAD, we further restricted the association results to variants with effective number of counts of the minor alleles, defined as $effN = N \times MAF \times (1 - MAF) \times oevar$ of at least 50, in both cases and controls. For the X-chromosome association testing, we set the count of alleles to be either 0 or 2 in males, and we calculated the effect allele frequency by separately calculating the minor allele frequency in males to get p_m (here count of alleles was either 0 or 1), separately in females to get p_f , and combining the two as $(N_m p_m + 2N_f p_f)/(N_m + 2N_f)$, where N_m and N_f are the number of males and female individuals with measured/imputed genotypes at the locus, respectively. Associations were genome-wide significant if their p -value $< 5 \times 10^{-8}$, and we also investigated suggestive associations, with p -value $< 10^{-7}$. In addition, for an X-chromosome association, we performed analysis stratified by sex, to study whether it is evidently driven by one of the sexes, or both.

Stratified analysis by hispanic/latino ethnic background. To study potential differences in genetic association patterns between and within subgroups, we performed a secondary analysis of ABI in which a GWAS was conducted separately in each of the genetic analysis subgroups. We then meta-analyzed the summary statistics from each subgroup into a Mainland group and a Caribbean group separately. We used a fixed-effects meta-analysis that accounts for the correlations between the groups, due to relatives and shared environment²⁶, and provide the p -value from the Cochran's Q test of heterogeneity of effect sizes across the genetic analysis groups. To limit the potential number of false positive associations with the increased of the multiple testing burden due to performing a large number of GWAS, in the stratified analyses we considered only results from common variants, defined as those with approximate effective count of the minor allele $effN$ (defined above) of at least 250, where this threshold is defined separately in each of the subgroups, yielding potentially different number of SNPs in each group. For reporting subgroup results, we used the same genome-wide significance and suggestive thresholds as in the main analysis.

Generalization and replication analysis of discovered associations. To investigate the replication and generalization of findings from the stratified analysis to other populations, we tested the association of the index SNPs (SNPs with lowest p -value in a detected association region) with the ABI in Multi-Ethnic Study of Atherosclerosis (MESA)²⁷ African Americans ($N = 1,613$), Hispanics/Latinos ($N = 1,447$), and European Americans ($N = 2,527$). In the Hispanic subgroup, 54% self-reported as Mexican descent, 12% as Dominican, 13% as Puerto Rican, 4% as Cuban, and 14% as Other Hispanic (primarily Central and South American groups), with 3% not reporting. Briefly, the MESA is a prospective population based study of European descent, African-American, Hispanic and Asian men and women aged 45–84 at the baseline examination in 2000–02 and designed to study subclinical cardiovascular disease and its progression, and risk factors that predict progression to clinically overt cardiovascular disease²⁷. MESA was approved by the IRBs at each field center, where all participants gave written informed consent, and by the Human Subjects Division at the University of Washington, Seattle, WA, to the MESA Data Coordinating Center. See Supplementary Material Section 3 for complete list of IRBs approving this study.

All analyses were adjusted for age, sex, and five principal components. Since we interrogated five variants in three ethnic subgroups (as one of the detected SNPs was not available in MESA), the p -value criterion for replication was $0.05/15 = 0.003$. Because it was previously shown to have better control of type 1 error and higher power, under the assumption that an association replicates only if the direction of estimated association matches between the discovery and replication study, we used one-sided p -values for replication testing²⁸, with the expected direction of association in MESA depending on the direction of association in the HCHS/SOL. Therefore, if an association had the same direction in the HCHS/SOL and the replication study, the one-sided p -value is related to the usual two-sided p -value by $p/2$. However, if the directions of estimated associations do not match, the one-sided p -value is $(1 - p/2)$, where p is the two-sided p -value.

Additional replication tests were performed in the Cardiovascular Health Study (CHS) and the Atherosclerosis Risk In Communities (ARIC), in almost 6,000 European Americans and 749 African Americans using genotypes and phenotypes downloaded from dbGaP. See Section 1.8 in the Supplementary Information for description.

As an exploratory analysis, and due to the high correlation between the traits and the observed heterogeneity in genetic association patterns, we defined regions of 10^5 bp upstream and downstream of our top detected loci,

	Overall n = 7589	Central American n = 760	South American n = 553	Mexican n = 2619	Puerto Rican n = 1437	Cuban n = 1500	Dominican n = 700
Age	55 (7)	55 (7)	55 (7)	55 (7)	56 (8)	56 (8)	55 (7)
Ankle Brachial Index	1.06 (0.13)	1.06 (0.13)	1.07 (0.12)	1.08 (0.12)	1.07 (0.15)	1.04 (0.12)	1.06 (0.13)
Male Sex	2966 (39%)	265 (35%)	207 (37%)	963 (37%)	575 (40%)	709 (47%)	242 (35%)
Systolic BP	132 (21)	132 (22)	128 (21)	127 (20)	135 (21)	135 (20)	136 (21)
Diastolic BP	78 (12)	78 (11)	75 (12)	75 (11)	79 (12)	80 (12)	81 (12)
Hypertension	3075 (41%)	295 (39%)	165 (30%)	866 (33%)	702 (49%)	715 (48%)	323 (46%)
Body Mass Index	30 (6)	31 (6)	29 (5)	30 (6)	31 (6)	29 (5)	30 (5)
Current smoker	1495 (20%)	89 (12%)	78 (14%)	381 (15%)	417 (29%)	467 (31%)	62 (9%)
Diabetes	2162 (28%)	214 (28%)	103 (19%)	784 (30%)	502 (35%)	361 (24%)	192 (27%)

Table 1. Participant Characteristics Overall and by Hispanic Ethnic Subgroup*. *Mean \pm SD or n (%).

and recorded the lead associated SNP for each of the ABI, PAD, and borderline PAD in each of the MESA replicating populations in these loci. The SNPs in these loci were restricted to those with MAF \geq 0.05 and an imputation quality score of at least 0.8.

Results

Study population characteristics overall and by self-reported Hispanic ethnic subgroup are provided in Table 1. There were 7,589 individuals in the primary ABI analysis. In the PAD analysis, there were 382 affected individuals with PAD and 7,015 unaffected (non-PAD without arterial stiffness). In the borderline PAD, there were 2,104 affected individuals and 5,293 unaffected (non-PAD/borderline PAD, and without arterial stiffness). ABI distribution varied in ethnic subgroups with the lowest mean among Cuban Americans, and highest mean among Mexican Americans.

Heritability estimation. We estimated the heritability of ABI using Haseman-Elston regression (²⁹; Supplemental Information Section 1.3). The estimated heritability of ABI was 10% (95% CI: 3% to 18%) when including related individuals. The estimated contribution of the environmental factors household and block unit to the variance of ABI was 1% (95% CI: 0% to 6%). In the analysis including only unrelated individuals (n = 6,856), the estimated heritability was lower, 7%, (95% CI: 0% to 16%). The environmental contribution to the ABI variance had similarly a low estimate, with a 95% CI including zero.

Overall pooled analysis of Hispanics/Latinos. GWAS inflation factors λ_{gc} ³⁰, were all between 0.99 and 1.005, indicating good control of population stratification. Manhattan plots and qq-plots for overall pooled analyses are provided in the Supplementary Figs S2 and S3. When examining the whole cohort, none of the variants were significantly associated with ABI, PAD, or borderline PAD at the genome-wide significance level.

Stratified ABI analyses by Hispanic/Latino background. Supplemental Fig. S4 displays the Manhattan plots the ABI stratified by Hispanic background subgroup. Table 2 provides the lead variants from six loci with $p < 10^{-7}$ in subgroup stratified analyses for the ABI. Figures S8–S13 in the Supplementary Information provide regional association plots for all background subgroups for the variants reported in Table 2. For all loci identified in the stratified analysis, the heterogeneity test p was $< 10^{-3}$. Further, the power to detect these association in the pooled analysis at the genome-wide significance level, assuming that the effect size in the group in which they were originally detected was the same across the population, were all > 0.99 . Table S1 in the Supplementary Information presents the PAD results for the top ABI SNPs in the subgroup-stratified analyses. Table S2 presents the pooled results for the ABI and PAD for the top ABI SNPs in the subgroup-stratified analysis. There were two genome-wide significant associations identified for ABI. The first locus with index SNP rs3133941 ($p = 1.8 \times 10^{-8}$) in the *SYBU* region on chromosome 8 (Fig. 1), was identified in the meta-analysis of the three Caribbean subgroups: Cuban, Dominican, and Puerto Rican (n = 3,637, MAF ranging from 14% to 26%). The second locus, *COMMD10* on chromosome 5 (Fig. 2), was identified in the analysis of Puerto-Ricans only (n = 1,437), with the index SNP rs4466200 ($p = 3.4 \times 10^{-9}$, MAF = 0.35).

Among suggestive associations with the ABI (Supplemental Figs S5–S7, Fig. 3), rs6631478 in the *DMD* gene on the X-chromosome was identified in analysis of the Central American subgroup (Fig. 3; n = 760, $p = 7.0 \times 10^{-8}$, MAF = 0.31). A sex-stratified analysis at this locus suggested that the association exists in both sexes (males $p = 4.2 \times 10^{-4}$, females $p = 1.2 \times 10^{-4}$). There were three additional loci with $p < 10^{-7}$, driven by associations with common variants, including an imputed indel, and two genotyped SNPs (Table 2).

To help contextualize these findings, Figs S18 and S19 in the Supplementary Information provide power analysis for the main (pooled) analysis, as well the group- and subgroup-specific GWAS, for a range of MAFs and effect sizes. One can see that the effect sizes required for the two genome-wide significant findings, given the analyses' sample sizes and SNPs' MAFs, are quite high: between 0.2–0.3 standard deviations (SDs) of ABI in the Caribbean group for the *SYBU* locus, and a little bit higher for the *COMMD10* locus detected in the Puerto-Ricans only. Indeed, the effect sizes reported for these loci in Table 2 are about 0.2 SDs of ABI (SDs of ABI in the HCHS/SOL range from 0.090 in Mexicans and South Americans to 0.105 in Puerto-Ricans). Sensitivity analysis removing a handful of individuals with ABI > 1.4 from the analysis reports similar results, as reported in Table S4 in the Supplementary Information.

Subgroup	Index SNP	Chr	Position	Allele A	Allele B	Type	EAF	Beta	SE	p-value	Het p-value	Gene
Dominican	rs6750426	2	15131582	G	A	G	0.75	0.031	0.006	7.39E-08	1.44E-06	
Puerto Rican	rs4466200	5	115596152	G	A	G	0.35	0.024	0.004	3.40E-09	<1E-06	COMMD10
Puerto Rican	rs7755533	6	157626334	G	A	G	0.59	0.022	0.004	9.28E-08	6.67E-06	TMEM242
Cuban	rs113916643	7	68415037	GA	G	I	0.73	-0.024	0.004	6.47E-08	4.03E-05	
Central American	rs6631478	X	32092380	T	C	G	0.69	-0.024	0.004	7.02E-08	2.41E-05	DMD
Caribbean	rs3133941	8	110715932	A	G	I	0.74-0.86	-0.018	0.003	1.78E-08	5.24E-01	SYBU

Table 2. Significant or Suggestive Loci for the Ankle Brachial Index in Stratified Subgroup Analyses*. *Significant or suggestive SNPs with $p < 10^{-7}$ in any of the subgroup-specific analyses, or in the meta-analysis of the Caribbean and the Mainland subgroups. Positions are provided in genome build 37. Allele A is the effect allele, EAF is the effect allele frequency, and Beta is the estimated effect of an increase in the allele count/dosage of allele A. Type is G when the variant is genotyped, and I when it is imputed. Het p-value is the p-values from the Cochran’s test of heterogeneity, adapted to the settings where there are correlated individuals between strata.




Figure 1. LocusZoom plot (top) and forest plot (bottom) of the SYBU locus, detected as associated with the ankle brachial index (ABI) in the Caribbean group. In the locusZoom plots, each point represents a variant, with location marked on the x-axis, and p-values marked as the location on the y-axis. The lead SNP is represented by the triangle, indicating that it is imputed. The colors of the variants correspond to the strength of their LD (r^2) with the lead SNP, with LD estimated using the combined population of the HCHS/SOL Caribbean group. Circles correspond to genotyped variants, x symbols to imputed variants. The p-value of heterogeneity (across all HCHS/SOL genetic analysis groups) was 9.3×10^{-4} . The bottom of the forest plot provides results from MESA replication groups.




Figure 2. LocusZoom plot (top) and forest plot (bottom) of the *COMMD10* locus, detected as associated with ankle brachial index (ABI) in the Puerto Rican subgroup. In the locusZoom plot, each point represents a variant, with location marked on the x-axis, and p-values marked on the y-axis. The lead SNP is represented by the diamond, indicating that it is genotyped. The colors of the variants correspond to the strength of their LD (r^2) with the lead SNP, with LD estimated using the Puerto Rican population of the HCHS/SOL. Circles correspond to genotyped variants, x symbols to imputed variants. The p-value of heterogeneity (across all HCHS/SOL genetic analysis groups) was 1×10^{-10} . The bottom of the forest plot provides results from MESA replication groups.

Replication and generalization results. Table 3 provides the replication results for both genome-wide significant and suggestive discovery results in the MESA cohort. Neither rs3133941 in *SYBU* nor rs4466200 in *COMMD10* replicated in MESA African American, European American or Hispanic groups, but rs4466200 had $p = 0.02$ in MESA African Americans (Table 3). However, this association did not replicate in African Americans from CHS (Table S5 in the Supplementary Information). Rs6631478 in the *DMD* gene replicated in MESA Hispanics, $p = 2.2 \times 10^{-4}$ (Table 3). Rs113916643 on chromosome 7 was not available in MESA. None of the associations replicated in European Americans (Table S6 in the Supplementary Information).

Supplementary Table S7 reports the results from the exploratory analysis that searched for the lead SNP in each of the MESA replicating populations and each of the traits. For the *DMD* locus identified for the ABI that replicated in MESA Hispanics, this analysis identified rs77460337, a SNP only 14 base-pairs away from the HCHS/SOL Central American lead SNP with a $p = 5.3 \times 10^{-7}$ in the PAD results of the MESA Hispanics. This SNP was not available in the HCHS/SOL data. At the *COMMD10* locus, rs12520838 was associated with PAD in MESA Hispanics ($p = 7.5 \times 10^{-5}$), a SNP that is 99,000 base-pairs away from the HCHS/SOL Puerto Rican lead variant (rs4466200); these variants are not in LD, as estimated in the HCHS/SOL.

None of the previously report associations with ABI or PAD in European descent, African-Americans or Japanese generalized to Hispanics/Latinos (all r -values = 1, see Supplementary Information Section 1.4 for methods and Supplementary Tables S8 and S9 for results). Additionally, Supplementary Figs S15 and S16 compare the




Figure 3. LocusZoom plot (top) and forest plot (bottom) of the *DMD* locus, detected as suggestively associated with ABI in the Central American genetic analysis group. In the locusZoom plots, each point represents a variant, with location marked on the x-axis, and p-values marked on the y-axis. The lead SNP is represented by the diamond, indicating that it is genotyped. The colors of the variants correspond to the strength of their LD (r^2) with the lead SNP, with LD estimated using the Central American population of the HCHS/SOL. Circles correspond to genotyped variants, x symbols to imputed variants. The p-value of heterogeneity (across all HCHS/SOL genetic analysis groups) was 2.41×10^{-5} . The bottom of the forest plot provides results from MESA replication groups.

effect sizes between the Hispanics/Latinos from our ABI analysis to the effect sizes in two previously published ABI analyses in individuals of European descent.

SYTL3 gene region. A previous candidate gene study in European ancestry participants¹⁶ reported a significant association of rs2171209 in the *SYTL3* gene region on chromosome 6 with the ABI. Although this variant was not associated with ABI, PAD or borderline PAD in our analyses, we detected a suggestive association of rs317789, about 700 bp from rs2171209 and uncorrelated with rs317789, with borderline PAD ($p = 7.56 \times 10^{-7}$) in the overall pooled Hispanic/Latino HCHS/SOL analysis. To study whether this association is driven by the Mainland group (Central American, Mexican, South American), which has higher proportion of European ancestry on average, we tested the association of rs317789 with borderline PAD in the Mainland and Caribbean groups separately. However, there was no difference between the groups (Mainland $p = 0.001$, Caribbean $p = 0.0002$). Further, the association with the European descent index SNP rs2171209 was not significant in either the Mainland or Caribbean groups ($p > 0.5$ in both groups).

Functional annotation of replicated association regions. At the *COMMD10* locus, the lead variant rs4466200 is in LD with 261 intronic variants ($r^2 \geq 0.8$) based in HCHS/SOL data, including three variants which have evidence for biological functionality. These include (1) rs12654321 ($r^2 = 0.96$) which lies in DNaseI

HCHS/SOL group	Index SNP	Chr	Allele A	Allele B	HCHS/SOL		MESA Hispanic		MESA AA		MESA EA	
					Beta	p-value	Beta	p-value	Beta	p-value	Beta	p-value
Dominican	rs6750426	2	G	A	0.031	7.39E-08	-0.006	5.58E-01	-0.019	6.79E-01	-0.023	7.93E-01
Puerto Rican	rs4466200	5	G	A	0.024	3.40E-09	0.016	3.34E-01	0.074	2.16E-02	-0.037	8.81E-01
Puerto Rican	rs7755533	6	G	A	0.022	9.28E-08	0.014	3.73E-01	0.024	2.82E-01	0.043	1.02E-01
Central American	rs6631478	X	T	C	-0.024	7.02E-08	-0.118	2.22E-04	-0.012	3.52E-01	0.013	6.43E-01
Caribbean	rs3133941	8	A	G	-0.018	1.78E-08	-0.032	2.50E-01	-0.04	1.27E-01	-0.001	4.94E-01

Table 3. Replication Results for Significant and Suggestive Loci for the ABI in Stratified Subgroup Analyses.

*Allele A is the effect allele. All variants were imputed, and all were common (MAF range 0.11–0.48 across all variants and all ethnic groups) in MESA. MESA replication sample sizes were $n = 1,447$ (Hispanic), $n = 1,613$ African American (AA), and $n = 2,527$ European American (EA). Rs113916643 was not available in MESA and is omitted from this table.

hypersensitive genomic region and is bound by transcription factor CTCF in aortic adventitial fibroblast cells, (2) rs1382342, which is located in a DNaseI hypersensitive site in monocytes and inflammatory macrophage cells, and (3) rs4921067, which overlaps enhancer-binding transcription factor CEBPB binding site in IMR90 (normal human lung cell line) cells. Functional annotation in the heart tissues did not identify any strong candidates for likely casual regulatory variants at this locus. Most of the LD proxies are reported as eQTL for the *COMMD10* gene. At the chromosome X locus that replicated, the lead variant rs6631478 is located in the intronic region of the *DMD* gene and does not have any LD proxies ($r^2 \geq 0.8$ in HCHS/SOL data). Other significant and suggestive loci had no clear functional roles.

Colocalization analysis for the region around rs4466200. Using GTEx summary statistics we identified three genes in three tissues that had statistically significant eQTLs at the *COMMD10* gene region, and their list of significantly associated SNPs overlapped with the SNPs with p -value < 0.001 in the Puerto Rican ABI analysis. Sections 1.6 and 2.7 in the Supplementary Information provide complete description of the analysis and results, and Supplementary Table S10 in the Supplementary Information provides the results. There were four analyses, and two SNPs were identified. Rs4466200, the lead ABI Puerto Rican SNP had the highest posterior probability in the co-localization analysis with atrial appendage heart tissue, with the *COMMD10* gene, and in the analysis with the left ventricle heart tissue with the *SEMA6A* gene. However, these posterior probabilities were low: 0.08–0.09. An additional SNP in the region, rs4466200, also had higher posterior probabilities, although below our threshold of significance (0.75) – first, in the aortic artery tissue, with gene *CTB-118N6.3* (posterior probability 0.31), and second, in the left ventricle heart tissue with gene *COMMD10* (posterior probability 0.40).

Pathway enrichment analysis. We used GOrilla³¹ to search for enriched pathways based on gene scores from the pooled ABI analysis. Detailed methods are provided in Supplementary Information Section 1.7. After False Discovery Rate control (FDR), there were two pathways, both of type “function” (as annotated in GOrilla) that were significantly enriched (FDR q -value < 0.05): nucleosomal DNA binding (q -value = 0.002), and chromatin DNA binding (q -value = 0.002) pathways. Figure S17 in the Supplementary Information displays these pathways in a larger context of molecular function.

Discussion

We examined the genetic architecture underlying ABI and PAD in the Hispanic/Latino population using a large cohort of ancestrally diverse Hispanics/Latinos living in the US. We report results from GWAS of ABI, PAD, and borderline PAD, both between and within ethnic subgroups based on genetic background. We identified two genome-wide significant loci (*SYBU*, *COMMD10*) and three suggestive associations for the ABI in analyses stratified by Hispanic/Latino ethnic background. However, we found no genome wide significant results in the overall pooled analyses of HCHS/SOL Hispanics/Latinos, perhaps due to significant genetic heterogeneity in risk for PAD among Hispanic/Latino subgroups⁸, as well as substantial differences in local and global genetic admixture among these groups, as suggested by differences in allele frequencies between groups in the detected association regions.

The *COMMD10* locus on chromosome 5 had p -value = 0.02 in MESA African-Americans for ABI, and regionally replicated for PAD in MESA Hispanics (FDR $p = 0.05$). This SNP had smaller estimated effect size in 749 African-Americans from CHS and replication p -value = 0.43. This could be because this is not a real signal, or due to low power, and difference in age distribution (CHS participants were 65 and older). *COMMD10* is a member of the COMM domain containing proteins, and encodes a protein that inhibits and mediates NF- κ B, a transcription factor involved in innate and adaptive immune responses, and possibly with inflammatory processes^{32,33}. Variants in *COMMD10* have been previously associated with multiple sclerosis³³, inflammatory factors (tumor necrosis factor-(TNF)- α and monocyte chemoattractant protein (MCP)-1)³⁴, as well as chronic obstructive pulmonary disease (COPD) and asthma³⁵. The functional annotation results showed *COMMD10* variants overlapping putative regulatory regions in adventitial fibroblast cells, inflammatory macrophages and monocytes, suggesting that the regulatory variants in *COMMD10* locus may be influencing the ABI or PAD via inflammatory and/or coagulation pathways, which is consistent with previously hypothesized roles for *COMMD10*. However, functional evaluation is needed to confirm the postulated underlying mechanisms. The co-localization analysis pointed to rs10062588 as a potential functional SNP in aortic artery and left ventricle heart tissues, however, the posterior probabilities were 0.31–0.40.

This analysis is limited by the different populations between GTEx (European Ancestry) and HCHS/SOL, despite minor allele frequencies being quite similar (Supplementary Fig. S16).

The *DMD* locus on chromosome X replicated in MESA Hispanics for the ABI. Deletions, duplications, rearrangements, and point mutations in *DMD*, the dystrophin gene, have been strongly linked to Duchenne muscular dystrophy (DMD), Becker muscular dystrophy (BMD), and cardiomyopathy^{36–39}. However, our functional annotation results did not identify any regulatory features associated with rs6631478.

None of the previously reported associations in populations of European, African, or East Asian (Japanese) ancestries generalized to Hispanics/Latinos, which could be due to the persisting low coverage of imputed genotypes using 1000 Genomes data, as well as the presence of genetic admixture. However, the stratified analysis revealed multiple associations with ABI, in specific Hispanic/Latino subgroups. These associations comprised largely common variants, and are therefore less likely to be statistical artifacts driven by outliers. In addition, these subgroup-specific associations were highly heterogeneous, as observed by comparing effect sizes and allele frequencies across Hispanic ethnic subgroups.

However, there is at least some evidence of common genetic basis at the regional level for ABI and PAD across race/ethnicities, with significant heterogeneity within each region. At the *SYTL3* locus, we identified a SNP associated with borderline PAD (rs317789, $p < 10^{-6}$) in a close proximity (700 bp) to a previously reported association with the ABI in Europeans (rs2171209), although rs317789 is not correlated with rs2171209 in European or Hispanic populations. Variants in *SYTL3* have been previously associated with circulating lipoprotein(a) levels (Lp(a))⁴⁰ which is a largely genetically regulated biomarker⁴¹. We have previously hypothesized that the association of *SYTL3* with the ABI and PAD may be mediated through Lp(a) levels¹²; however, circulating Lp(a) is unfortunately also not available in the HCHS/SOL to test this hypothesis. Our results for *SYTL3* also indicate that while the same loci may contribute to the underlying genetic architecture of PAD, heterogeneity within these loci and in ancestral background plays a substantial role.

This study has a few limitations. The ABI outcome is highly skewed, which may reduce power and increase type I error of genetic association studies⁴². Moreover, more than half of the study participants have ABI > 1. It is unclear which factors determine variation of ABI in the range of ABI > 1, and some of these factors are likely arterial elastic properties, height, and weight⁴³. However, in a sensitivity analysis reported in Supplementary Table S3, we tested all associations reported in Table 2, with added adjustment to height and weight, and the results were essentially the same. For PAD analysis, the low number of cases of this population-based sample, reduces power, compared to a potential case-control study that would specifically target PAD cases. This may reduce the power of ABI analysis as well. Another limitation is the small sample size, resulting in low statistical power. The largest, Mexican subgroup, had 2,619 individuals, and the pooled analysis had 7,589 individuals, which is considered low for a GWAS. Future analyses should leverage the admixture pattern for genetic discoveries via admixture mapping, which has reduced multiple testing burden and was useful in other genetic analyses in this cohort^{44–46}, and combine the HCHS/SOL GWAS with analyses in other cohorts in meta-analysis. Another limitation is the large number of tests, considering the different traits and subgroups: we performed ABI (pooled and by subgroup), PAD, and borderline PAD analyses. Still, the analyses were well controlled in terms of inflation, resulting in only a handful of findings among all analyses.

Additional evidence of regional heterogeneity in common genetic loci was observed in our MESA replication analysis. Specifically, an exploratory regional replication analysis of all common SNPs available in MESA around the HCHS/SOL lead SNPs from the stratified ABI analysis, detected two PAD SNPs different from the lead in HCHS/SOL: in the *DMD* locus, rs77460337 was 14 bp away from the HCHS/SOL Central American ABI lead SNP, and at the *COMMD10* locus, rs12520838 was 99,000 bp away from the HCHS/SOL Puerto Rican lead SNP. Overall, our generalization and replication results suggest there may be additional associations existing in both HCHS/SOL and MESA Hispanic populations, which may be detected once denser imputation panels are used, or whole genome sequencing data is available.

The current analysis detected multiple novel genetic variants associated with ABI in Hispanic/Latino ethnic subgroups from the HCHS/SOL. One of the lead SNPs in the *DMD* gene replicated in MESA Hispanics, despite the discovery being specific to Central Americans, and both the *DMD* and *COMMD10* loci regionally replicated in MESA Hispanics, albeit with different MESA-specific lead SNPs. Our results suggest regional commonality, but significant heterogeneity at the regional level for genetic architecture underlying the ABI and PAD. Care should be taken in interpreting these results due to the large multiple testing burden and low sample sizes within the Hispanic/Latino subgroups, and additional efforts for replication are needed, ideally, in diverse Hispanic/Latino populations. Another possible explanation for these results is that associated genotypes have different allele frequencies, and even different effect sizes, between ancestral populations (Europeans, Africans, Amerindians), and different admixture patterns in subpopulations of Hispanics/Latinos lead to different tag SNPs for the causal variants, and differences in power to discover these associations. Admixture mapping may help in revealing such insights. Future whole genome sequencing studies in diverse populations are important to facilitate better understanding and fine mapping of ABI/PAD loci, and may inform the mystery of lack of generalization of ABI/PAD loci across populations, including loci reported here, and in previous studies of European Ancestry and Japanese individuals.

Data Availability

Genotype data of the HCHS/SOL and summary statistics from all discovery GWAS for ABI, PAD, borderline PAD, and arterial stiffness can be requested via dbGaP study accession phs000880. Complete meta-data related to these analyses is recorded for reproducibility in the HCHS/SOL Genetic Analysis Center's Integrated Computing and Tracking system (unique analysis IDs are provided in Supplementary Table S11 in the Supplementary Information). Phenotype data can be requested via dbGaP study accession phs000810.

References

- Hiatt, W. R. *et al.* Atherosclerotic Peripheral Vascular Disease Symposium II: nomenclature for vascular diseases. *Circulation* **118**, 2826–2829, <https://doi.org/10.1161/CIRCULATIONAHA.108.191171> (2008).
- Ankle Brachial Index, C. *et al.* Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA* **300**, 197–208, <https://doi.org/10.1001/jama.300.2.197> (2008).
- Steg, P. G. *et al.* One-year cardiovascular event rates in outpatients with atherothrombosis. *JAMA* **297**, 1197–1206, <https://doi.org/10.1001/jama.297.11.1197> (2007).
- Criqui, M. H. *et al.* Mortality over a period of 10 years in patients with peripheral arterial disease. *N Engl J Med* **326**, 381–386, <https://doi.org/10.1056/NEJM199202063260605> (1992).
- Resnick, H. E. *et al.* Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality: the Strong Heart Study. *Circulation* **109**, 733–739, <https://doi.org/10.1161/01.CIR.0000112642.63927.54> (2004).
- Fowkes, F. G. *et al.* Development and validation of an ankle brachial index risk model for the prediction of cardiovascular events. *Eur J Prev Cardiol* **21**, 310–320, <https://doi.org/10.1177/2047487313516564> (2014).
- McDermott, M. M. *et al.* Associations of borderline and low normal ankle-brachial index values with functional decline at 5-year follow-up: the WALCS (Walking and Leg Circulation Study). *J Am Coll Cardiol* **53**, 1056–1062, <https://doi.org/10.1016/j.jacc.2008.09.063> (2009).
- Allison, M. A. *et al.* Cuban Americans have the highest rates of peripheral arterial disease in diverse Hispanic/Latino communities. *J Vasc Surg* **62**, 665–672, <https://doi.org/10.1016/j.jvs.2015.03.065> (2015).
- Allison, M. A. *et al.* Ethnic-specific prevalence of peripheral arterial disease in the United States. *Am J Prev Med* **32**, 328–333, <https://doi.org/10.1016/j.amepre.2006.12.010> (2007).
- Allison, M. A. *et al.* The effect of novel cardiovascular risk factors on the ethnic-specific odds for peripheral arterial disease in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Am Coll Cardiol* **48**, 1190–1197, <https://doi.org/10.1016/j.jacc.2006.05.049> (2006).
- Ix, J. H., Allison, M. A., Denenberg, J. O., Cushman, M. & Criqui, M. H. Novel cardiovascular risk factors do not completely explain the higher prevalence of peripheral arterial disease among African Americans. The San Diego Population Study. *J Am Coll Cardiol* **51**, 2347–2354, <https://doi.org/10.1016/j.jacc.2008.03.022> (2008).
- Wassel, C. L. *et al.* Genetic determinants of the ankle-brachial index: a meta-analysis of a cardiovascular candidate gene 50K SNP panel in the candidate gene association resource (CARE) consortium. *Atherosclerosis* **222**, 138–147, <https://doi.org/10.1016/j.atherosclerosis.2012.01.039> (2012).
- Murabito, J. M., Guo, C. Y., Fox, C. S. & D'Agostino, R. B. Heritability of the ankle-brachial index: the Framingham Offspring study. *Am J Epidemiol* **164**, 963–968, <https://doi.org/10.1093/aje/kwj295> (2006).
- Carmelli, D. *et al.* Contribution of genetic and environmental influences to ankle-brachial blood pressure index in the NHLBI Twin Study. National Heart, Lung, and Blood Institute. *Am J Epidemiol* **151**, 452–458 (2000).
- Wassel, C. L. *et al.* Family history of peripheral artery disease is associated with prevalence and severity of peripheral artery disease: the San Diego population study. *J Am Coll Cardiol* **58**, 1386–1392, <https://doi.org/10.1016/j.jacc.2011.06.023> (2011).
- Khaleghi, M., Isseh, I. N., Bailey, K. R. & Kullo, I. J. Family history as a risk factor for peripheral arterial disease. *Am J Cardiol* **114**, 928–932, <https://doi.org/10.1016/j.amjcard.2014.06.029> (2014).
- Murabito, J. M. *et al.* Association between chromosome 9p21 variants and the ankle-brachial index identified by a meta-analysis of 21 genome-wide association studies. *Circ Cardiovasc Genet* **5**, 100–112, <https://doi.org/10.1161/CIRCGENETICS.111.961292> (2012).
- Lavange, L. M. *et al.* Sample design and cohort selection in the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol* **20**, 642–649, <https://doi.org/10.1016/j.annepidem.2010.05.006> (2010).
- Sorlie, P. D. *et al.* Design and implementation of the Hispanic Community Health Study/Study of Latinos. *Ann Epidemiol* **20**, 629–641, <https://doi.org/10.1016/j.annepidem.2010.03.015> (2010).
- Conomos, M. P. *et al.* Genetic Diversity and Association Studies in US Hispanic/Latino Populations: Applications in the Hispanic Community Health Study/Study of Latinos. *Am J Hum Genet* **98**, 165–184, <https://doi.org/10.1016/j.ajhg.2015.12.001> (2016).
- Genomes Project, C. *et al.* An integrated map of genetic variation from 1,092 human genomes. *Nature* **491**, 56–65, <https://doi.org/10.1038/nature11632> (2012).
- Rosenberg, N. A., Li, L. M., Ward, R. & Pritchard, J. K. Informativeness of genetic markers for inference of ancestry. *Am J Hum Genet* **73**, 1402–1422, <https://doi.org/10.1086/380416> (2003).
- Laurie, C. C. *et al.* Quality control and quality assurance in genotypic data for genome-wide association studies. *Genet Epidemiol* **34**, 591–602, <https://doi.org/10.1002/gepi.20516> (2010).
- Delaneau, O., Zagury, J. F. & Marchini, J. Improved whole-chromosome phasing for disease and population genetic studies. *Nat Methods* **10**, 5–6, <https://doi.org/10.1038/nmeth.2307> (2013).
- Howie, B. N., Donnelly, P. & Marchini, J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* **5**, e1000529, <https://doi.org/10.1371/journal.pgen.1000529> (2009).
- Sofer, T. *et al.* Meta-Analysis of Genome-Wide Association Studies with Correlated Individuals: Application to the Hispanic Community Health Study/Study of Latinos (HCHS/SOL). *Genet Epidemiol* **40**, 492–501, <https://doi.org/10.1002/gepi.21981> (2016).
- Bild, D. E. *et al.* Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol* **156**, 871–881 (2002).
- Sofer, T. *et al.* A powerful statistical framework for generalization testing in GWAS, with application to the HCHS/SOL. *Genet Epidemiol* **41**, 251–258, <https://doi.org/10.1002/gepi.22029> (2017).
- Sofer, T. Confidence intervals for heritability via Haseman-Elston regression. *Stat Appl Genet Mol Biol* **16**, 259–273, <https://doi.org/10.1515/sagmb-2016-0076> (2017).
- Devlin, B. & Roeder, K. Genomic control for association studies. *Biometrics* **55**, 997–1004 (1999).
- Eden, E., Navon, R., Steinfeld, I., Lipson, D. & Yakhini, Z. GOrilla: a tool for discovery and visualization of enriched GO terms in ranked gene lists. *BMC Bioinformatics* **10**, 48, <https://doi.org/10.1186/1471-2105-10-48> (2009).
- Burstein, E. *et al.* COMMD Proteins, a Novel Family of Structural and Functional Homologs of MURR1. *Journal of Biological Chemistry* **280**, 22222–22232 (2005).
- Beck, J. *et al.* Serum DNA motifs predict disease and clinical status in multiple sclerosis. *The Journal of molecular diagnostics: JMD* **12**, 312–319 (2010).
- Aslibekyan, S. *et al.* A genome-wide association study of inflammatory biomarker changes in response to fenofibrate treatment in the Genetics of Lipid Lowering Drug and Diet Network. *Pharmacogenet Genomics* **22**, 191–197, <https://doi.org/10.1097/FPC.0b013e32834fdd41> (2012).
- Smolonska, J. *et al.* Common genes underlying asthma and COPD? Genome-wide analysis on the Dutch hypothesis. *Eur Respir J* **44**, 860–872, <https://doi.org/10.1183/09031936.00001914> (2014).
- Soltanzadeh, P. *et al.* Clinical and genetic characterization of manifesting carriers of DMD mutations. *Neuromuscular disorders: NMD* **20**, 499–504 (2010).
- Tsuda, T. *et al.* Early-progressive dilated cardiomyopathy in a family with Becker muscular dystrophy related to a novel frameshift mutation in the dystrophin gene exon 27. *Journal of human genetics* **60**, 151–155 (2015).
- Juan-Mateu, J. *et al.* DMD Mutations in 576 Dystrophinopathy Families: A Step Forward in Genotype-Phenotype Correlations. *PloS one* **10**, e0135189 (2015).
- Flanigan, K. M. *et al.* Nonsense mutation-associated Becker muscular dystrophy: interplay between exon definition and splicing regulatory elements within the DMD gene. *Human mutation* **32**, 299–308 (2011).

40. Ober, C. *et al.* Genome-wide association study of plasma lipoprotein(a) levels identifies multiple genes on chromosome 6q. *J Lipid Res* **50**, 798–806 (2009).
41. Utermann, G. Genetic architecture and evolution of the lipoprotein(a) trait. *Current opinion in lipidology* **10**, 133–141 (1999).
42. Sofer, T. *et al.* A Fully-Adjusted Two-Stage Procedure for Rank Normalization in Genetic Association Studies. *Genetic Epidemiology*, <https://doi.org/10.1002/gepi.22188> (2019).
43. Aboyans, V. *et al.* Intrinsic contribution of gender and ethnicity to normal ankle-brachial index values: the Multi-Ethnic Study of Atherosclerosis (MESA). *J Vasc Surg* **45**, 319–327, <https://doi.org/10.1016/j.jvs.2006.10.032> (2007).
44. Brown, L. A. *et al.* Admixture Mapping Identifies an Amerindian Ancestry Locus Associated with Albuminuria in Hispanics in the United States. *J Am Soc Nephrol* **28**, 2211–2220, <https://doi.org/10.1681/ASN.2016091010> (2017).
45. Burkart, K. M. *et al.* A Genome-Wide Association Study in Hispanics/Latinos Identifies Novel Signals for Lung Function. The Hispanic Community Health Study/Study of Latinos. *Am J Respir Crit Care Med* **198**, 208–219, <https://doi.org/10.1164/rccm.201707-1493OC> (2018).
46. Sofer, T. *et al.* Admixture mapping in the Hispanic Community Health Study/Study of Latinos reveals regions of genetic associations with blood pressure traits. *PLoS One* **12**, e0188400, <https://doi.org/10.1371/journal.pone.0188400> (2017).

Acknowledgements

The authors thank the staff and participants of HCHS/SOL for their important contributions. The Hispanic Community Health Study/Study of Latinos was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The following Institutes/Centers/Offices contribute to the HCHS/SOL through a transfer of funds to the NHLBI: National Institute on Minority Health and Health Disparities, National Institute on Deafness and Other Communication Disorders, National Institute of Dental and Craniofacial Research, National Institute of Diabetes and Digestive and Kidney Diseases, National Institute of Neurological Disorders and Stroke, NIH Institution-Office of Dietary Supplements. This study was also supported by the Intramural Program of the National Institute of Diabetes and Digestive and Kidney Diseases. The Genetic Analysis Center at the University of Washington was supported by NHLBI and NIDCR contracts (HHSN268201300005C AM03 and MOD03). This study was supported in part by NIMHD R01 MD012765, NIDDK R56 DK104806-01A1 and NHLBI R21HL123677-01 to Nora Franceschini. Christina Wassel was supported in part by R01HL110955. Tamar Sofer was supported by NHLBI R01 HL120393-03S1, 1R35HL135818, and NIA 7R01AG048642-04, 1RF1AG054548-01. Genotype and imputed data of the HCHS/SOL can be requested via dbGaP study accession phs000880. Phenotype data can be requested via dbGaP study accession phs000810. The authors thank the other investigators, the staff, and the participants of the MESA study for their valuable contributions. A full list of participating MESA investigators and institutions can be found at <http://www.mesa-nhlbi.org>. MESA and the MESA SHARe project are conducted and supported by the National Heart, Lung, and Blood Institute (NHLBI) in collaboration with MESA investigators. Support for MESA is provided by contracts HHSN268201500003I, N01-HC-95159, N01-HC-95160, N01-HC-95161, N01-HC-95162, N01-HC-95163, N01-HC-95164, N01-HC-95165, N01-HC-95166, N01-HC-95167, N01-HC-95168, N01-HC-95169, UL1-TR-000040, UL1-TR-001079, UL1-TR-001420, UL1-TR-001881, and DK063491. Funding for SHARe genotyping was provided by NHLBI Contract N02-HL-64278. Genotyping was performed at Affymetrix (Santa Clara, California, USA) and the Broad Institute of Harvard and MIT (Boston, Massachusetts, USA) using the Affymetrix Genome-Wide Human SNP Array 6.0.

Author Contributions

T.S. drafted the manuscript. T.S., L.E., A.E., C.W., J.L. and N.K. performed statistical data analysis. T.S. and L.E. prepared figures. D.J. performed functional annotation of findings. C.L., H.G., W.T., L.L. and C.W., contributed to either phenotype or genetic data collection and quality control. C.W., M.A., M.C., W.P. and N.F. conceptualize the manuscript. All authors critically reviewed and edited the manuscript.

Additional Information

Supplementary information accompanies this paper at <https://doi.org/10.1038/s41598-019-47928-5>.

Competing Interests: The authors declare no competing interests.

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2019