

# UCLA

## UCLA Previously Published Works

### Title

HENSELIAN VALUED FIELDS AND inp-MINIMALITY

### Permalink

<https://escholarship.org/uc/item/6d6066b5>

### Journal

Journal of Symbolic Logic, 84(4)

### ISSN

0022-4812

### Authors

CHERNIKOV, ARTEM  
SIMON, PIERRE

### Publication Date

2019-12-01

### DOI

10.1017/jsl.2019.56

Peer reviewed

# Henselian valued fields and inp-minimality

Artem Chernikov<sup>\*†‡</sup> and Pierre Simon<sup>\*†</sup>

January 28, 2016

## Abstract

We prove that every ultraproduct of  $p$ -adics is inp-minimal (i.e., of burden 1). More generally, we prove an Ax-Kochen type result on preservation of inp-minimality for Henselian valued fields of equicharacteristic 0 in the RV language.

## 1 Introduction

In his work on the classification of first-order theories [She90] Shelah has introduced a hierarchy of combinatorial properties of families of definable sets, so called dividing lines, which includes stable theories, simple theories, NIP, NSOP, etc. An important line of research in model theory is to characterize various algebraic structures depending on their place in this classification hierarchy (this knowledge can later be used to analyze various algebraic objects definable in such structures using methods of generalized stability theory). Here we will be concerned with valued fields and Ax-Kochen-type statements, i.e. statements of the form “a certain property of the valued field can be determined by looking just at the value group and the residue field”. For example, a classical theorem of Delon [Del78] shows that given a Henselian valued field of equicharacteristic 0, if the residue field is NIP, then the whole valued field is NIP. More recent result of similar type are [Bél99] demonstrating preservation of NIP for certain valued fields of positive characteristic, [She05] demonstrating that the field of  $p$ -adics is strongly dependent, and [DGL<sup>+</sup>11] demonstrating that it is in fact dp-minimal.

A motivating example for this article is to determine the model-theoretic complexity of the theory of an ultraproduct of the fields of  $p$ -adics  $\mathbb{Q}_p$  modulo a non-principal ultrafilter on the set of prime numbers. Namely, let  $K = \prod \mathbb{Q}_p / \mathcal{U}$ , where  $\mathcal{U}$  is a non-principal ultrafilter on the set of prime numbers. Note that the residue field  $k$  is a pseudo-finite field of characteristic 0 and that the value group  $\Gamma$  is a  $\mathbb{Z}$ -group. Besides, both  $k$  and  $\Gamma$  are interpretable in  $K$  in the pure ring language (e.g. by a result of Ax [Ax65]). This implies that the theory of  $K$  is neither NIP, nor simple — the two classes of structures extensively studied in model theory. However it turns out that any ultraproduct of  $p$ -adics is NTP<sub>2</sub> [Che14]. The class of NTP<sub>2</sub> theories was introduced by Shelah [She90, Chapter III] and generalizes both simple and NIP theories. We recall the definition.

**Definition 1.** Let  $T$  be a complete first-order theory in a language  $L$ , and let  $\mathbb{M} \models T$  be a monster model. Let  $\kappa$  be a cardinal (finite or infinite).

1. An inp-pattern of depth  $\kappa$  is given by  $(\phi_i(x, y_i), \bar{a}_i, k_i : i \in \kappa)$ , where  $\phi_i(x, y_i)$  are  $L$ -formulas with a fixed tuple of free variables  $x$  and a varying tuple of parameter variables  $y_i$ ,  $\bar{a}_i = (a_{i,j} : j \in \omega)$  are sequences of tuples of elements from  $\mathbb{M}$ , and  $k_i$  are natural numbers such that:

---

<sup>\*</sup>The research leading to this paper has been (partially) supported by the European Research Council under the European Union’s Seventh Framework Programme (FP7/2007-2013)/ERC Grant Agreement No. 291111.

<sup>†</sup>Partially supported by ValCoMo (ANR-13-BS01-0006).

<sup>‡</sup>This work is (partially) supported by the Fondation Sciences Mathématiques de Paris (FSMP) and by a public grant overseen by the French National Research Agency (ANR) as part of the Investissements d’Avenir program (reference: ANR-10-LABX-0098).

- (a) For every  $i \in \kappa$ , the set  $\{\phi_i(x, a_{i,j})\}_{j \in \omega}$  is  $k_i$ -inconsistent (i.e. no subset of size  $\geq k_i$  is consistent).
- (b) For every  $f : \kappa \rightarrow \omega$ , the set  $\{\phi_i(x, a_{i,f(i)})\}_{i \in \kappa}$  is consistent.

2.  $T$  is  $\text{NTP}_2$  if there is a (cardinal) bound on the depths of inp-patterns.

Other algebraic examples of  $\text{NTP}_2$  structures were identified recently, including bounded pseudo real closed and pseudo  $p$ -adically closed fields [Mon14], certain model complete multi-valued fields [Joh13] and certain valued difference fields, e.g. the theory  $\text{VFA}_0$  of a non-standard Frobenius on an algebraically closed valued field of characteristic zero [CH14]. See also [CKS15] and [HO15] for some general results about groups and fields definable in  $\text{NTP}_2$  structures.

The notion of *burden* was introduced by Adler [Adl07] based on Shelah's cardinal invariant  $\kappa_{\text{inp}}$  and provides a quantitative refinement of  $\text{NTP}_2$ . In the special case of simple theories burden corresponds to preweight, and in the case of NIP theories to dp-rank (e.g. see [Che14, Section 3] for the details and references).

**Definition 2.** 1.  $T$  is *strong* if there are no inp-patterns of infinite depth.

2.  $T$  is of *finite burden* if there are no inp-patterns of arbitrary large finite depth, with  $x$  a singleton.

3.  $T$  is *inp-minimal* if there is no inp-pattern of depth 2, with  $x$  a singleton.

Note that inp-minimality implies finite burden implies strong (the last implication uses submultiplicativity of burden from [Che14]). All the examples mentioned above have been demonstrated to be strong of finite burden, with the exception of  $\text{VFA}_0$ : it remains open if  $\text{VFA}_0$  is strong, see [CH14, Question 5.2]. Some results about strong groups and fields can be found in [CKS15, Section 4] and [DG15].

Returning to ultraproducts of  $p$ -adics, we have the following more general result.

**Fact 3.** [Che14] *Let  $\bar{K} = (K, k, \Gamma, \text{val}, \text{ac})$  be a Henselian valued field of equicharacteristic 0, considered as a three-sorted structure in the Denef-Pas language  $L_{\text{ac}}$  (i.e. there is a sort  $K$  for the field itself, as well as sorts  $k$  for the residue field and  $\Gamma$  for the value group, together with the maps  $v : K \rightarrow \Gamma$  for the valuation and  $\text{ac} : K \rightarrow k$  for an angular component).*

1. *If  $k$  is  $\text{NTP}_2$ , then  $\bar{K}$  is  $\text{NTP}_2$ .*

2. *If both  $k$  and  $\Gamma$  are strong (of finite burden) then  $\bar{K}$  is strong (respectively, of finite burden).*

Any pseudofinite field is supersimple of SU-rank 1, so in particular is inp-minimal. Any ordered  $\mathbb{Z}$ -group is dp-minimal, so in particular is inp-minimal. It follows that any ultraproduct of  $p$ -adics is strong, of finite burden. However, Fact 3(2) gives a finite bound on the burden of  $\bar{K}$  in terms of the burdens of  $k$  and  $\Gamma$  via a certain Ramsey number, and is far from optimal in general. It was conjectured in [Che14, Problem 7.13] that all ultraproducts of  $p$ -adics in the pure ring language are inp-minimal (note that in the Denef-Pas language, no valued field with an infinite residue field can be inp-minimal as  $\{\text{ac}(x) = a_i\}, \{\text{val}(x) = v_i\}$  with  $(a_i), (v_i)$  pairwise different give an inp-pattern of depth 2).

In this paper we establish an Ax-Kochen type result for inp-minimality in the RV language for valued fields, in particular confirming that conjecture.

**Theorem 4.** *Let  $\bar{K} = (K, \text{RV}, \text{rv})$  be a Henselian valued field of equicharacteristic 0, viewed as a structure in the RV-language (see Section 2). Assume that both the residue field  $k$  and the value group  $\Gamma$  are inp-minimal, and that moreover  $k^\times / (k^\times)^p$  is finite for all prime  $p$ . Then  $\bar{K}$  is inp-minimal.*

**Corollary 5.** *Any ultraproduct of  $p$ -adics is inp-minimal.*

Recall that an NIP theory is dp-minimal if and only if it is inp-minimal. Johnson [Joh15] shows that a dp-minimal not strongly minimal field admits a definable Henselian valuation. It follows that if  $K$  is dp-minimal, then  $K^\times/(K^\times)^p$  is finite for all prime  $p$  (a fact which Johnson states and uses). Combining this with Delon's result on preservation of NIP we have the following corollary (which also appears in Johnson [Joh15]).

**Corollary 6.** *Under the same assumptions on  $\bar{K}$ , if both  $k$  and  $\Gamma$  are dp-minimal, then  $\bar{K}$  is dp-minimal.*

There are three steps in the proof of the main theorem, corresponding to the sections of the paper. First, we recall some facts about the RV setting and show that the whole valued field is inp-minimal if and only if the RV sort is inp-minimal. Second, we show that the RV sort eliminates quantifiers down to the residue field  $k$  and the value group  $\Gamma$ . Using this quantifier elimination, in the last section we show that the RV sort is inp-minimal if and only if both  $k$  and  $\Gamma$  are inp-minimal. Finally, we discuss some problems and future research directions.

## 2 Reduction to RV

We recall some basic facts about the RV setting, we are going to use [Fle11] as a reference. Fix a valued field  $K$ , with value group  $\Gamma$  and residue field  $k$ . Let  $\text{RV}$  be the quotient group  $K^\times/(1 + \mathfrak{m})$  where  $\mathfrak{m} = \{x \in K : \text{val}(x) > 0\}$  is the maximal ideal of the valuation ring. We have a short exact sequence  $1 \rightarrow k^\times \rightarrow \text{RV} \xrightarrow{\text{val}_{\text{rv}}} \Gamma \rightarrow 0$ .

Consider now the two-sorted structure  $\bar{K} = (K, \text{RV}, \text{rv})$  in the language  $L_{\text{RV}+}$  consisting of:

- the quotient map  $\text{rv} : K \rightarrow \text{RV}$ ,
- on the sort  $K$ , the ring structure,
- on the sort  $\text{RV}$ , the structure  $\cdot, 1$  of a multiplicative group, a symbol  $0$ , a symbol  $\infty$  and a ternary relation  $\oplus$ .

The multiplicative group structure is interpreted as the group structure induced from  $K^\times$  and  $0 \cdot x = x \cdot 0 = 0$ ,  $\infty = \text{rv}(0)$ . The relation  $\oplus$  is interpreted as the partially defined addition inherited from  $K$ :  $\oplus(a, b, c) \iff \exists x, y, z \in K (a = \text{rv}(x) \wedge b = \text{rv}(y) \wedge c = \text{rv}(z) \wedge x + y = z)$ .

*Remark 7.* 1. One can define the set  $\text{WD}(x, y)$  of pairs of elements for which the sum is well-defined as  $\forall z, z' (\oplus(x, y, z) \wedge \oplus(x, y, z') \implies z = z')$ . Given a pair of elements  $x, y \in \text{RV}$  such that  $\text{WD}(x, y)$  holds, we write  $x + y$  to denote the unique element  $z \in \text{RV}$  satisfying  $\oplus(x, y, z)$ .

2. We have  $\text{WD}(\text{rv}(a), \text{rv}(b)) \iff \text{val}(a + b) = \min\{\text{val}(a), \text{val}(b)\}$ , in which case  $\text{rv}(a + b) = \text{rv}(a) + \text{rv}(b)$  (see [Fle11, Proposition 2.4]).
3. The relation  $\text{val}_{\text{rv}}(x) \leq \text{val}_{\text{rv}}(y)$  on  $\text{RV}$  is definable in this language [Fle11, Proposition 2.8(1)]. Namely, let  $d \in \text{RV}$  be arbitrary with  $\text{val}_{\text{rv}}(d) = 0$ . Then  $\text{val}_{\text{rv}}(x) > 0 \iff dx + 1 = 1$ , and  $\text{val}_{\text{rv}}(x) = 0 \iff \neg \text{val}_{\text{rv}}(x) > 0 \wedge \exists y (x \cdot y = 1 \wedge \neg \text{val}_{\text{rv}}(y) > 0)$ . Then  $\text{val}_{\text{rv}}(x) = \text{val}_{\text{rv}}(y) \iff \exists u (\text{val}_{\text{rv}}(u) = 0 \wedge x = u \cdot y)$  and  $\text{val}_{\text{rv}}(x) < \text{val}_{\text{rv}}(y) \iff x \neq \infty \wedge x + dy = x$ .

Let  $\bar{\mathbb{K}} \succ \bar{K}$  be a monster model. We may always assume that  $\bar{\mathbb{K}}$  admits a cross-section map  $\text{ac} : K \rightarrow k^\times$ , so we can view  $\bar{\mathbb{K}}$  also as a structure in the language  $L_{\text{ac}}$  with  $\text{ac}$  added to the language.

**Fact 8.** [Fle11, Proposition 5.1]

1. Let  $K$  be a Henselian valued field with  $\text{char}(k) = 0$ , and suppose that  $S \subseteq K$  is definable. Then there are  $\alpha_1, \dots, \alpha_k$  and a definable subset  $D \subseteq \text{RV}^k$  such that

$$S = \{x \in K : (\text{rv}(x - \alpha_1), \dots, \text{rv}(x - \alpha_k)) \in D\}.$$

2. The RV sort is fully stably embedded (i.e. the structure on RV induced from  $\bar{K}$ , with parameters, is precisely the one described above).

The following two lemmas are easy to verify (see [Che10] for the details).

**Lemma 9.** *Let  $(a_i)_{i \in I}$  be an  $L_{\text{ac}}$ -indiscernible sequence of singletons in  $\mathbb{K}$ , and consider the function  $(i, j) \mapsto \text{val}(a_j - a_i)$  for  $i < j \in I$ . Then one of the following cases occurs:*

1. It is strictly increasing depending only on  $i$  (so the sequence is pseudo-convergent).
2. It is strictly decreasing depending only on  $j$  (so the sequence taken in the reverse direction is pseudo-convergent).
3. It is constant (we'll refer to such a sequence as a "fan").

**Lemma 10.** *Let  $(a_i)_{i \in I}$  be an  $L_{\text{ac}}$ -indiscernible pseudo-convergent sequence from  $\mathbb{K}$ . Then for any  $d \in \mathbb{K}$  there is some  $i_* \in \bar{I} \cup \{+\infty, -\infty\}$  (where  $\bar{I}$  is the Dedekind closure of  $I$ ) such that (taking  $a_\infty$  from  $\mathbb{K}$  such that  $I \frown a_\infty$  is indiscernible):*

For  $i < i_*$ :  $\text{val}(a_\infty - a_i) < \text{val}(d - a_\infty)$ ,  $\text{val}(d - a_i) = \text{val}(a_\infty - a_i)$  and  $\text{ac}(d - a_i) = \text{ac}(a_\infty - a_i)$ .

For  $i > i_*$ :  $\text{val}(a_\infty - a_i) > \text{val}(d - a_\infty)$ ,  $\text{val}(d - a_i) = \text{val}(d - a_\infty)$  and  $\text{ac}(d - a_i) = \text{ac}(d - a_\infty)$ .

Note also that for any non-zero  $x, y \in K$ ,  $\text{rv}(x) = \text{rv}(y)$  if and only if  $\text{val}(x - y) > \text{val}(y)$  and for any  $z \in K$  and  $x, y \in K \setminus \{z\}$ ,  $\text{rv}(x - z) = \text{rv}(y - z)$  if and only if  $\text{val}(x - y) > \text{val}(y - z)$ .

In the remainder of this section we will reduce inp-minimality of  $\bar{K}$  to inp-minimality of the RV sort with the induced structure.

First we treat a key special case. Assume that there is an inp-pattern consisting of formulas  $\psi(x, yz) = \phi(\text{rv}(x - y), z)$  and  $\psi'(x, yz') = \phi'(\text{rv}(x - y), z')$  and mutually  $L_{\text{ac}}$ -indiscernible sequences  $(c_i)_{i \in \mathbb{Z}}, (c'_i)_{i \in \mathbb{Z}}$  with  $c_i = a_i \tilde{b}_i$  and  $c'_i = a'_i b'_i$  where  $\phi$  and  $\phi'$  are RV-formulas,  $b_i \in \text{RV}^{|z|}, b'_i \in \text{RV}^{|z'|}$  and  $a_i, a'_i \in K$ . Without loss of generality both  $\{\phi(\text{rv}(x - a_i), b_i)\}_{i \in \mathbb{Z}}$  and  $\{\phi'(\text{rv}(x - a'_i), b'_i)\}_{i \in \mathbb{Z}}$  are  $k$ -inconsistent, and let  $d \models \phi(\text{rv}(x - a_0), b_0) \wedge \phi'(\text{rv}(x - a'_0), b'_0)$ . We may also add to the base elements  $a_\infty, a_{-\infty}, a'_\infty, a'_{-\infty}$  continuing our sequences on the left and on the right.

*Claim 11.*  $\text{val}(d - a_i) \leq \text{val}(d - a'_0)$  and  $\text{val}(d - a'_j) \leq \text{val}(d - a_0)$  for all  $i$  and  $j$ . In particular,  $\text{val}(d - a_0) = \text{val}(d - a'_0) = \gamma$ .

*Proof.* Assume that  $\text{val}(d - a_i) > \text{val}(d - a'_0)$  for some  $i$ . Then  $\text{rv}(d - a'_0) = \text{rv}(a_i - a'_0)$ . So  $\models \phi'(\text{rv}(a_i - a'_0), b'_0)$ , and by mutual indiscernibility  $a_i \models \{\phi'(\text{rv}(x - a'_j), b'_j)\}_{j \in \omega}$  — a contradiction. The other part is by symmetry.  $\square$

*Claim 12.*  $\gamma \leq \text{val}(a_0 - a'_0)$ .

*Proof.* As otherwise  $\text{val}(d - a_0) = \text{val}(d - a'_0) = \gamma > \text{val}(a_0 - a'_0) = \text{val}(d - a_0)$  — a contradiction.  $\square$

We now consider several cases separately.

**Case A:**  $\text{val}(a_i - a'_j)$  is constant, equal to some  $\gamma' \in \Gamma$ .

As in this case the two sequences are mutually indiscernible over  $\gamma'$ , we may add it to the base. Note that  $\gamma \leq \gamma'$  by Claim 12. The following subcases cover all the possible situations, using mutual indiscernibility of the sequences over  $\gamma'$ .

**Subcase 1:**  $\gamma < \gamma'$ .

Then  $\text{rv}(d - a_i) = \text{rv}(d - a'_i) = \alpha$  for all  $i, j$ , for some some  $\alpha \in \text{RV}$  with  $\text{val}_{\text{rv}}(\alpha) = \gamma$ . Note furthermore that for any  $\alpha^* \in \text{RV}$  such that  $\text{val}_{\text{rv}}(\alpha^*) < \gamma'$  we can find some  $d^* \in K$  such that  $\text{rv}(d^* - a_i) = \text{rv}(d^* - a'_i) = \alpha^*$ .

But then consider the array  $\tilde{\phi}(\tilde{x}, b_i) = \phi(\tilde{x}, b_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) < \gamma'$ ,  $\tilde{\phi}'(\tilde{x}, b'_i) = \phi(\tilde{x}, b_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) < \gamma'$ , where  $\tilde{x}$  and  $b_i, b'_i$  are ranging over the RV sort and  $\phi, \phi'$  are RV-formulas (we are abusing the notation by writing  $\text{val}_{\text{rv}}(\tilde{x}) < \gamma'$  as a shortcut for  $\text{val}_{\text{rv}}(\tilde{x}) < \text{val}_{\text{rv}}(a_\infty - a'_\infty)$ ). We have  $\models \tilde{\phi}(\alpha, b_0) \wedge \tilde{\phi}'(\alpha, b'_0)$  and  $\left\{ \tilde{\phi}(\tilde{x}, b_i) \right\}_{i \in \mathbb{Z}}, \left\{ \tilde{\phi}'(\tilde{x}, b'_i) \right\}_{i \in \mathbb{Z}}$  are both inconsistent by the previous observation as the original array was inconsistent. This gives us an inp-pattern in the structure induced on the RV sort, and so implies that RV is not inp-minimal.

**Subcase 2:**  $\gamma = \gamma'$ ,  $\text{val}(a_i - a_j) > \gamma$  and  $\text{val}(a'_i - a'_j) > \gamma$  for all  $i < j$ .

It follows that there are  $\alpha, \alpha' \in \text{RV}$  with  $\text{val}_{\text{rv}}(\alpha) = \text{val}_{\text{rv}}(\alpha') = \gamma$  such that  $\text{rv}(d - a_i) = \alpha$  and  $\text{rv}(d - a'_i) = \alpha'$  for all  $i$ . Furthermore,  $\text{rv}(a_i - a'_j) = \alpha' - \alpha =: \beta$  for all  $i, j$ . It follows that our sequences are mutually indiscernible over  $\beta$  and we can add it to the base. Moreover, for any  $\alpha^* \in \text{RV}$  with  $\text{val}_{\text{rv}}(\alpha^*) = \gamma$  we can find some  $d^* \in K$  with  $\text{rv}(d^* - a_i) = \alpha^*$  and  $\text{rv}(d^* - a'_i) = \alpha^* + \beta$ .

Then let  $\tilde{\phi}(\tilde{x}, b_i) = \phi(\tilde{x}, b_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma$  and  $\tilde{\phi}'(\tilde{x}, b'_i) = \phi'(\tilde{x} - \beta, b'_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma$ . It follows that  $\alpha \models \tilde{\phi}(\tilde{x}, b_0) \wedge \tilde{\phi}'(\tilde{x}, b'_0)$  and that  $\left\{ \tilde{\phi}(\tilde{x}, b_i) \right\}_{i \in \mathbb{Z}}, \left\{ \tilde{\phi}'(\tilde{x}, b'_i) \right\}_{i \in \mathbb{Z}}$  are both inconsistent as the original array was inconsistent — contradicting inp-minimality of RV.

**Subcase 3:**  $\gamma = \gamma'$ ,  $\text{val}(a_i - a_j) > \gamma$  and  $\text{val}(a'_i - a'_j) = \gamma$  for all  $i < j$ .

In this case we still have some  $\alpha \in \text{RV}$  such that  $\text{rv}(d - a_i) = \alpha$  for all  $i$ . On the other hand, it follows that  $\text{rv}(d - a'_i) = \text{rv}(d - a_\infty) + \text{rv}(a_\infty - a'_i)$ . Moreover, for any  $\alpha^* \in \text{RV}$  with  $\text{val}_{\text{rv}}(\alpha^*) = \gamma$  and  $\text{WD}(\alpha^*, \text{rv}(a_\infty - a'_i))$  we can find some  $d^*$  such that  $\text{rv}(d^* - a_i) = \alpha^*$  and  $\text{rv}(d^* - a'_i) = \alpha^* + \text{rv}(a_\infty - a'_i)$ .

Then define  $\tilde{\phi}(\tilde{x}, b_i) = \phi(\tilde{x}, b_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma$  and  $\tilde{\phi}'(\tilde{x}, \tilde{b}'_i) = \phi'(\tilde{x} + \text{rv}(a_\infty - a'_i), b'_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma \wedge \text{WD}(\tilde{x}, \text{rv}(a_\infty - a'_i))$ , so  $\tilde{b}'_i = \text{rv}(a_\infty - a'_i) \tilde{b}'_i$ . Note that  $(b_i)_{i \in \mathbb{Z}}$  and  $(\tilde{b}'_i)_{i \in \mathbb{Z}}$  are mutually indiscernible sequences in RV. It follows that  $\alpha \models \tilde{\phi}(\tilde{x}, b_0) \wedge \tilde{\phi}'(\tilde{x}, \tilde{b}'_0)$  and that both  $\left\{ \tilde{\phi}(\tilde{x}, b_i) \right\}_{i \in \mathbb{Z}}, \left\{ \tilde{\phi}'(\tilde{x}, \tilde{b}'_i) \right\}_{i \in \mathbb{Z}}$  are inconsistent — contradicting inp-minimality of RV.

**Subcase 4:**  $\gamma = \gamma'$ ,  $\text{val}(a_i - a_j) = \text{val}(a'_i - a'_j) = \gamma$  for all  $i < j$ .

It then follows that  $\text{rv}(d - a_i) = \text{rv}(d - a_\infty) + \text{rv}(a_\infty - a_i)$  and  $\text{rv}(d - a'_i) = \text{rv}(d - a_\infty) + \text{rv}(a_\infty - a'_i)$  (as  $\text{val}(d - a'_i) = \text{val}(d - a_i) = \text{val}(d - a_\infty) = \text{val}(a_\infty - a'_i)$ ). Moreover, for any  $\alpha^* \in \text{RV}$  with  $\text{val}_{\text{rv}}(\alpha^*) = \gamma$  (and such that the corresponding sums are well-defined) there is  $d^*$  such that  $\text{rv}(d^* - a_i) = \alpha^* + \text{rv}(a_\infty - a_i)$ ,  $\text{rv}(d^* - a'_i) = \alpha^* + \text{rv}(a_\infty - a'_i)$ .

We define  $\tilde{\phi}(\tilde{x}, \tilde{b}_i) = \phi(\tilde{x} + \text{rv}(a_\infty - a_i), b_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma \wedge \text{WD}(\tilde{x}, \text{rv}(a_\infty - a_i))$  and  $\tilde{\phi}'(\tilde{x}, \tilde{b}'_i) = \phi'(\tilde{x} + \text{rv}(a_\infty - a'_i), b'_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) = \gamma \wedge \text{WD}(\tilde{x}, \text{rv}(a_\infty - a'_i))$ , so  $\tilde{b}_i = \text{rv}(a_\infty - a_i) \tilde{b}_i$  and  $\tilde{b}'_i = \text{rv}(a_\infty - a'_i) \tilde{b}'_i$ . Note that  $(\tilde{b}_i)_{i \in \mathbb{Z}}$  and  $(\tilde{b}'_i)_{i \in \mathbb{Z}}$  are mutually indiscernible sequences in RV. It follows that  $\alpha \models \tilde{\phi}(\tilde{x}, \tilde{b}_0) \wedge \tilde{\phi}'(\tilde{x}, \tilde{b}'_0)$  and that both  $\left\{ \tilde{\phi}(\tilde{x}, \tilde{b}_i) \right\}_{i \in \mathbb{Z}}, \left\{ \tilde{\phi}'(\tilde{x}, \tilde{b}'_i) \right\}_{i \in \mathbb{Z}}$  are inconsistent — contradicting inp-minimality of RV.

**Subcase 5:**  $\gamma = \gamma'$ ,  $\text{val}(a_i - a_i) = \gamma$  and  $\text{val}(a'_i - a'_j) > \gamma$  for all  $i < j$ .

Follows from Subcase 3 by symmetry.

**Case B:** Not Case A.

*Claim 13.* At least one of the sequences  $(a_i)_{i \in \mathbb{Z}}, (a'_i)_{i \in \mathbb{Z}}$  is not a fan.

*Proof.* Assume that both are, say  $\text{val}(a_i - a_j) = \alpha$  and  $\text{val}(a'_i - a'_j) = \alpha'$  for all  $i < j$ . It follows by mutual indiscernibility that  $\text{val}(a_i - a'_j) \leq \min\{\alpha, \alpha'\}$  for all  $i, j$ . But then  $\text{val}(a_i - a'_j) = \text{val}(a_0 - a'_0)$  for all  $i, j$ , thus putting us in Case A.  $\square$

So we may assume that  $(a_i)_{i \in \mathbb{Z}}$  is a pseudo-convergent sequence (by Lemma 9, possibly exchanging  $(a_i)$  with  $(a'_i)$  and reverting the ordering of the sequence).

**Subcase 1:** Some (equivalently, every)  $a'_i$  is a pseudo-limit of  $(a_i)_{i \in \mathbb{Z}}$ .

Then  $\text{rv}(d - a'_i) = \text{rv}(d - a_\infty)$  for all  $i$  (by Claim 12).

We define  $\tilde{\phi}'(\tilde{x}, b'_i) = \phi'(\tilde{x}, b'_i) \wedge \text{val}_{\text{rv}}(\tilde{x}) < \text{val}(a_\infty - a'_\infty)$ .

By Lemma 10 it follows that there is some  $i^* \in \{-\infty\} \cup \mathbb{Z} \cup \{\infty\}$  such that  $\text{rv}(d - a_i) = \text{rv}(d - a_\infty)$  for  $i > i^*$  and  $\text{rv}(d - a_i) = \text{rv}(a_\infty - a_i)$  for  $i < i^*$ . Again by Claim 12,  $i^* \leq 0$ . Let's restrict  $(a_i)_{i \in \mathbb{Z}}$  to  $(a_i)_{i \in \omega}$ .

If  $\text{val}(d - a_\infty) < \text{val}(a_\infty - a_0)$  then  $\text{rv}(d - a_i) = \text{rv}(d - a_\infty)$  for all  $i$ . If  $\text{val}(d - a_\infty) = \text{val}(a_\infty - a_0)$  then  $\text{rv}(d - a_i) = \text{rv}(d - a_\infty)$  for all  $i > 0$  and  $\text{rv}(d - a_0) = \text{rv}(d - a_\infty) + \text{rv}(a_\infty - a_0)$ . We thus define

$$\begin{aligned} \tilde{\phi}(\tilde{x}, \tilde{b}_i) &= (\text{val}(a_i - a_\infty) > \text{val}_{\text{rv}}(\tilde{x}) \wedge \phi(\tilde{x}, b_i)) \vee \\ &\vee (\text{val}(a_i - a_\infty) = \text{val}_{\text{rv}}(\tilde{x}) \wedge \text{WD}(\tilde{x}, \text{rv}(a_\infty - a_i)) \wedge \phi(\tilde{x} + \text{rv}(a_\infty - a_i), b_i)) \end{aligned}$$

with  $\tilde{b}_i = b_i \hat{\text{rv}}(a_i - a_\infty)$ . Then  $(\tilde{b}_i), (b'_i)$  are mutually indiscernible sequences in RV and  $\text{rv}(d - a_\infty) \models \tilde{\phi}(\tilde{x}, \tilde{b}_0) \wedge \tilde{\phi}'(\tilde{x}, b'_0)$ . By inp-minimality of RV we have that either there is some  $\alpha^* \models \left\{ \tilde{\phi}'(\tilde{x}, b'_i) \right\}_{i \in \omega}$ , in which case we can find  $d^*$  with  $\text{rv}(d^* - a_\infty) = \alpha^*$  and thus  $d^* \models \left\{ \phi'(\text{rv}(x - a'_i), b'_i) \right\}_{i \in \omega}$ , or that  $\alpha^* \models \left\{ \tilde{\phi}(\tilde{x}, \tilde{b}_i) \right\}_{i \in \omega}$ . Then it follows from the definition of  $\tilde{\phi}$  that there is  $d^*$  satisfying  $\text{rv}(d^* - a_\infty) = \alpha^*$  and such that that  $d^* \models \left\{ \phi(\text{rv}(x - a_i), b_i) \right\}_{i \in \omega}$  — a contradiction.

**Subcase 2:** Not Subcase 1.

By inspection (remembering that we are not in the Case A), the only possibility is that  $(a'_i)$  is pseudo-convergent and that any  $a_i$  is a pseudo-limit of it. But then reversing the roles of the two sequences we are back to Subcase 1.

Now we reduce the case of a general inp-pattern to the special case treated above. Assume that there is an inp-pattern of depth 2. By Ramsey and compactness we may assume that the rows are mutually indiscernible in the  $L_{\text{ac}}$ -language. Though in Fact 8 the formula defining  $D$  may depend on the formula defining  $S$ , by indiscernibility, Ramsey and compactness we may assume that the formulas in our inp-pattern are in fact of the form  $\phi(\text{rv}(x - y_1), \dots, \text{rv}(x - y_n), z_n)$  and  $\phi'(\text{rv}(x - y_1), \dots, \text{rv}(x - y_n), z'_n)$ , for some  $n \in \omega$ , where  $\phi$  and  $\phi'$  are RV-formulas. Let  $d$  realize the first column of the inp-pattern.

**Case 1:**  $\text{val}(d - a_{0,0}) < \text{val}(a_{0,n} - a_{0,0})$ . Then  $\text{rv}(d - a_{0,0}) = \text{rv}(d - a_{0,n})$  and we define  $\tilde{\phi}(x, a_i \tilde{b}_i) = \phi(\text{rv}(x - a_{i,0}), \dots, \text{rv}(x - a_{i,n-1}), \text{rv}(x - a_{i,0})) \wedge \text{val}(x - a_{i,0}) < \text{val}(a_{i,n} - a_{i,0})$  with  $\tilde{b}_i = b_i \hat{\text{rv}}(a_{i,n} - a_{i,0})$ .

**Case 2:**  $\text{val}(d - a_{0,0}) > \text{val}(a_{0,n} - a_{0,0})$ . Then  $\text{rv}(d - a_{0,n}) = \text{rv}(a_{0,n} - a_{0,0})$  and we define  $\tilde{\phi}(x, a_i \tilde{b}_i) = \phi(\text{rv}(x - a_{i,0}), \dots, \text{rv}(x - a_{i,n-1}), \text{rv}(a_{i,n} - a_{i,0})) \wedge \text{val}(x - a_{i,0}) > \text{val}(a_{i,n} - a_{i,0})$  with  $\tilde{b}_i = b_i \hat{\text{rv}}(a_{i,n} - a_{i,0})$ .

**Case 3:**  $v(d - a_{0,n}) < v(a_{0,n} - a_{0,0})$  and **Case 4:**  $v(d - a_{0,n}) > v(a_{0,n} - a_{0,0})$  are symmetric to **Case 1** and **Case 2** respectively.

**Case 5:**  $v(d - a_{0,0}) = v(d - a_{0,n}) = v(a_{0,n} - a_{0,0})$ . Then  $\text{rv}(d - a_{0,0}) = \text{rv}(d - a_{0,n}) + \text{rv}(a_{0,n} - a_{0,0})$ . We define

$$\begin{aligned} \tilde{\phi}(x, a_i \tilde{b}_i) &= \phi(\text{rv}(x - a_{i,n}) + \text{rv}(a_{i,n} - a_{i,0}), \dots, \text{rv}(x - a_{i,n-1}), \text{rv}(x - a_{i,n})) \\ &\wedge v(x - a_{i,n}) = v(a_{i,n} - a_{i,0}) \wedge \text{WD}(\text{rv}(x - a_{i,n}), \text{rv}(a_{i,n} - a_{i,0})) \end{aligned}$$

with  $\tilde{b}_i = b_i \hat{\text{rv}}(a_{i,n} - a_{i,0})$ .

In any of the cases, we still have that  $(\tilde{b}_i)_{i \in \mathbb{Z}}, (b'_i)_{i \in \mathbb{Z}}$  are mutually indiscernible, that  $d \models \tilde{\phi}(x, a_0 \tilde{b}_0) \wedge \phi'(x, a'_0 b'_0)$  and that  $\left\{ \tilde{\phi}(x, a_i \tilde{b}_i) \right\}_{i \in \mathbb{Z}}$  is inconsistent. Thus we get a new inp-pattern

replacing  $\{\phi(x, a_i b_i)\}$  by  $\{\tilde{\phi}(x, a_i \tilde{b}_i)\}$ , with  $\tilde{\phi}$  involving one less term of the form  $\text{rv}(x - y_i)$ . Repeating the same operation  $n$  times for  $\phi$ , and then for  $\phi'$ , we reduce the situation to the special case of formulas considered before.

### 3 Relative quantifier elimination for RV

Now it will be more convenient to consider a valued field  $K$  in a slightly weaker language  $L_{\text{RV}}$ . Namely, we associate with it a three-sorted structure  $\bar{K} = (K, \text{RV}, \Gamma, \text{val}_{\text{rv}})$  such that on  $\text{RV}$  we have the multiplicative group structure  $\cdot, 1$ , a constant  $0$ , a predicate for the residue field  $k \subseteq \text{RV}$  along with addition  $\tilde{+}$  on  $k$ , and a map  $\text{val}_{\text{rv}} : \text{RV} \rightarrow \Gamma$ .

The partial addition relation  $\oplus$  on  $\text{RV}$  is definable in  $L_{\text{RV}}$  (using [Fle11, Proposition 2.7]):

$$\begin{aligned} \oplus(x, y, z) \iff & (\text{val}_{\text{rv}}(x) < \text{val}_{\text{rv}}(y) \wedge z = x) \vee (\text{val}_{\text{rv}}(y) < \text{val}_{\text{rv}}(x) \wedge z = y) \vee \\ & \vee \left( \text{val}_{\text{rv}}(x) = \text{val}_{\text{rv}}(y) \wedge \left( \left( \frac{x}{y} \tilde{+} 1 = 0 \wedge \text{val}_{\text{rv}}(z) > \text{val}_{\text{rv}}(x) \right) \vee \left( \left( \frac{x}{y} \tilde{+} 1 \right) y = z \wedge z \neq 0 \right) \right) \right). \end{aligned}$$

The conclusion is that in particular if  $(\text{RV}, \Gamma, \text{val}_{\text{rv}})$  is inp-minimal as an  $L_{\text{RV}}$ -structure, then  $(\text{RV}, \cdot, \oplus)$  is inp-minimal as an  $L_{\text{RV}^+}$ -structure. In the next section we are going to demonstrate the former under the assumptions of the main theorem, but in order to do that we prove a relative quantifier elimination result for (a certain expansion of) the  $L_{\text{RV}}$  language.

#### Assumptions

- $G$  is an abelian group such that  $G/nG$  is finite for all  $n < \omega$ .
- $K \subseteq G$  is a subgroup, with quotient  $H = G/K$ . Let  $\pi : G \rightarrow H$  denote the projection map.
- $M$  is the two-sorted structure with sorts  $G$  and  $H$ , and the following language.
  - On  $G$ : we have the group structure  $+, -, 0$ , a predicate  $K(x)$  for the subgroup  $K$ , predicates  $(P_n(x) : n < \omega)$  interpreted as  $P_n(x) \leftrightarrow \exists y ny = x$ , and constants naming a countable subgroup  $G_0$  containing representatives of each class of  $G/nG$ , for each  $n < \omega$  (such that moreover all classes of elements from  $K$  are represented by elements from  $G_0 \cap K$ ).
  - On  $H$ : we have some language  $L_H$  (containing the induced group structure) and we assume that the structure  $(H, L_H)$  eliminates quantifiers.
  - On  $K$ : we have some language  $L_K$  such that  $(K, L_K)$  eliminates quantifiers and contains the language induced from  $G$  (via the group structure and predicates  $P_n$ ).
  - We have the projection group homomorphism  $\pi : G \rightarrow H$ .
- Moreover, we assume that the language contains no other function symbols apart from  $\pi$  and the group structures on  $G$  and  $H$ .
- Finally,  $H$  is torsion-free.

**Proposition 14.**  *$M$  has quantifier elimination.*

*Proof.* We prove it by back-and-forth. So assume that  $M$  is  $\omega$ -saturated and we have two substructures  $A$  and  $B$  from  $M$  and a partial isomorphism  $f : A \rightarrow B$ . So  $A, B \supseteq G_0$  contain elements from both  $G$  and  $H$ , both are closed under the group operations, inverse and  $\pi$ .

Let  $\alpha \in M$  be arbitrary, and we want to extend  $f$  to be defined on  $A_1 = A(\alpha)$ , the substructure generated by  $\alpha A$ . We assume that  $\alpha \notin A$ .

**Step 1:** If  $\alpha \in H$ , then we can extend  $f$ .



As  $f|_{A \cap H}$  is  $L_H$ -elementary by quantifier elimination in  $(H, L_H)$ , there is  $\beta \in H$  and a partial  $L_H$ -automorphism  $g$  extending  $f|_{A \cap H}$  and sending  $A(\alpha) \cap H$  to  $B(\beta) \cap H$ . Then we extend  $f$  to  $F$  defined on  $A(\alpha)$  by taking  $F = f \cup g$  (note that, as there are no functions from  $H$  to  $G$  in the language,  $A(\alpha) \cap G = A \cap G$ ).

So by iterating Step 1 we may assume that  $\alpha \in G$  and that  $\pi(a + n\alpha) \in A$  for all  $a \in A$  and  $n \in \mathbb{Z}$ .

**Step 2:** Assume that  $\alpha \in K$ . Then we can extend  $f$ .

As  $f|_{A \cap K}$  is  $L_K$ -elementarity by quantifier elimination, we can find  $\beta \in K$  and a partial  $L_K$ -automorphism  $g$  extending it and sending  $A(\alpha) \cap K$  to  $B(\beta) \cap K$ . Then we define  $F$  on  $A(\alpha)$  by setting  $F(a + n\alpha) = f(a) + g(n\alpha) = f(a) + ng(\alpha)$  for all  $a \in A$ ,  $n \in \mathbb{Z}$  (note that  $n\alpha \in A(\alpha) \cap K$  for all  $n \in \mathbb{Z}$  by the assumption) and  $F$  acts like  $f$  on  $A(\alpha) \cap H = A \cap H$ .

- $F$  is well-defined: Assume that  $a + n\alpha = a' + n'\alpha$ , so  $A \ni a - a' = (n' - n)\alpha$ , and thus  $f(a) - f(a') = f(a - a') = f((n' - n)\alpha) = \dots$  as  $(n' - n)\alpha \in K \cap A$  and  $g|_{A \cap K} = f|_{A \cap K} \dots = g((n' - n)\alpha) = ng(\alpha) - n'g(\alpha)$ . Then we have  $F(a + n\alpha) - F(a' + n'\alpha) = f(a) + g(n\alpha) - f(a') - g(n'\alpha) = 0$ .
- $F$  extends  $f$ : immediate from the definition.
- Note that  $F|_{A(\alpha) \cap K} = g$ , as given  $a + n\alpha \in A(\alpha) \cap K$  it follows that  $a \in A \cap K$ , and as  $f|_{A \cap K} = g|_{A \cap K}$  we have  $F(a + n\alpha) = f(a) + g(n\alpha) = g(a) + g(n\alpha) = g(a + n\alpha)$ .
- $F|_G$  is a group homomorphism: indeed,  $F(a + n\alpha + a' + n'\alpha) = F((a + a') + (n + n')\alpha) = f(a + a') + g((n + n')\alpha) = f(a) + f(a') + g(n\alpha) + g(n'\alpha) = F(a + n\alpha) + F(a' + n'\alpha)$ .
- $F$  is onto  $B(\beta)$ : every element of  $B(\beta)$  is of the form  $b + n\beta$ , so  $F(f^{-1}(b) + n\alpha) = b + n\beta$ .
- $F$  preserves  $\pi$ : On one hand  $\pi(F(a + n\alpha)) = \pi(f(a) + ng(\alpha)) = \pi(f(a)) + n\pi(g(\alpha)) = \dots$  as  $g(\alpha) \in K \dots = \pi(f(a)) + 0 = f(\pi(a)) = F(\pi(a))$  (recall that  $\pi(a) \in A$ ). On the other hand we have  $F(\pi(a + n\alpha)) = F(\pi(a) + n\pi(\alpha)) = F(\pi(a) + 0) = F(\pi(a))$ .
- In particular,  $F$  preserves  $K(x) = \{x \in G : \pi(x) = 0\}$ .
- $F$  preserves  $P_k$ :  $P_k(F(a + n\alpha)) \Leftrightarrow P_k(f(a) + ng(\alpha)) \Leftrightarrow P_k(a + ng(\alpha))$  (as  $f(a) = a \pmod{kG} \Leftrightarrow P_k(a + n\alpha)$  (as  $g(\alpha) = \alpha \pmod{kG}$  because all representatives of classes of  $\alpha \in K$  are in  $G_0 \cap K \subseteq A \cap K$ ,  $P_k \cap K$  is  $L_K$ -definable and  $g|_{A(\alpha) \cap K}$  is  $L_K$ -elementary).
- $F$  preserves every  $\phi(x_1, \dots, x_k) \in L_K$ : As  $F|_{A(\alpha) \cap K} = g$  and  $g$  is an  $L_K$ -elementary map.
- $F$  preserves every  $\psi \in L_H$ : As  $\pi(a + n\alpha) = \pi(a) + n\pi(\alpha) \in A \cap H$  (as  $\pi(\alpha) \in A$  by the assumption), and  $F|_{A \cap H} = f|_{A \cap H}$  is  $L_H$ -elementary.

So  $F$  is a partial isomorphism as wanted.

By iterating Step 2 we may assume that  $a + n\alpha \in K \Rightarrow a + n\alpha \in A$  for all  $a \in A$  and  $n \in \omega$ .

**Step 3:** Assume that  $m\alpha \in A$  for some  $m \geq 1$ . Then we can extend  $f$ .

Let  $m$  be minimal with this property.

*Claim 15.* There is  $\beta \in G$  satisfying  $m\beta = f(m\alpha)$  and  $\beta = \alpha \pmod{kG}$  for all  $k \in \omega$ .

*Proof.* By  $\omega$ -saturation it suffices to show this one  $k$  at a time. By assumption there is some  $g \in G_0$  such that  $P_k(\alpha - g)$ , then  $P_k(\alpha - g) \Rightarrow P_{mk}(m\alpha - mg) \Rightarrow P_{mk}(f(m\alpha) - mg)$  (as  $m\alpha, mg \in A$ ,  $f(mg) = mf(g) = g$  and  $f$  preserves  $P_l$  for all  $l < \omega$ )  $\Rightarrow \exists \gamma \in G$  such that  $mk\gamma = f(m\alpha) - mg$ . Let  $\beta = k\gamma + g$ . Then  $m\beta = f(m\alpha)$  and  $\beta = g = \alpha \pmod{kG}$ , and the claim is proved.  $\square$

We define  $F$  on  $A(\alpha) \cap G$  by setting  $F(a + n\alpha) = f(a) + n\beta$  and  $F|_{A(\alpha) \cap H} = f|_{A(\alpha) \cap H}$  as  $A(\alpha) \cap H = A \cap H$ .

- $F$  is well-defined: If  $a + n\alpha = a' + n'\alpha$  with  $a, a' \in A$ , then  $(n - n')\alpha = a' - a \in A$ . It follows that  $m$  divides  $(n - n')$  by minimality (assume that  $n - n' = km + m_1$ ,  $|m_1| < m$ , then  $m_1\alpha = a' - a - km\alpha \in A$ , contradiction), say  $(n - n') = km$ . Thus  $f(a') - f(a) = f(a' - a) = f((n - n')\alpha) = f(km\alpha) = kf(m\alpha) = km\beta = (n - n')\beta$ . But then  $F(a + n\alpha) - F(a' + n'\alpha) = f(a) + n\beta - f(a') - n'\beta = 0$ .
- $F$  extends  $f$  is obvious from the definition.
- $F$  is a group homomorphism from  $A(\alpha)$  to  $B(\beta)$ :  

$$F((a + n\alpha) + (a' + n'\alpha)) = F((a + a') + (n + n')\alpha) = f(a + a') + (n + n')\beta = (f(a) + n\beta) + (f(a') + n'\beta) = F(a + n\alpha) + F(a' + n'\alpha).$$
- $F$  preserves  $\pi$ : First observe that  $\pi(m\beta) = \pi(f(m\alpha))$ , so  $m\pi(\beta) = \pi(f(m\alpha)) \stackrel{\text{as } m\alpha \in A}{=} f(\pi(m\alpha)) = f(m\pi(\alpha)) = mf(\pi(\alpha))$ , and as  $H$  is torsion free this implies that  $\pi(\beta) = f(\pi(\alpha))$ . But then  $F(\pi(a + n\alpha)) = f(\pi(a + n\alpha)) = f(\pi(a) + n\pi(\alpha)) = f(\pi(a)) + nf(\pi(\alpha)) = \pi(f(a)) + n\pi(\beta) = \pi(f(a) + n\beta) = \pi(F(a + n\alpha))$ .
- In particular,  $F$  preserves  $K(x) = \{x \in G : \pi(x) = 0\}$ .
- $F$  preserves  $P_k(x)$ : By the choice of  $\beta$  we have  $\alpha = \beta \pmod{kG}$  for all  $k$ , and for any  $a \in A$  we have  $f(a) = a \pmod{kG}$  for all  $k$  (as  $G_0 \subseteq A$  and  $f$  preserves  $P_k$ ), hence  $P_k(F(a + n\alpha)) \Leftrightarrow P_k(f(a) + n\beta) \Leftrightarrow P_k(a + n\alpha)$ .
- $F$  preserves  $L_K$ -formulas: As  $a + n\alpha \in K \Rightarrow a + n\alpha \in A$  by the assumption and  $F|_{A \cap K} = f|_{A \cap K}$  is  $L_K$ -elementary by elimination of quantifiers in  $(K, L_K)$ .
- $F$  preserves  $L_H$ -formulas: As  $F|_{A(\alpha) \cap H} = f|_{A(\alpha) \cap H = A \cap H}$  by definition, and  $f$  is  $L_H$ -elementary.

So we may assume that:

1.  $A \cap H$  is a relatively divisible subgroup of  $H$  (iterating Step 1);
2.  $A \cap G$  is a relatively divisible subgroup of  $A(\alpha) \cap G$  (iterating Step 3);
3.  $\pi(a + n\alpha) \in A$  for all  $a \in A, n \in \mathbb{Z}$  (iterating Step 1);
4.  $a + n\alpha \notin K$  for all  $a \in A, n \in \mathbb{Z} \setminus \{0\}$  (as  $a + n\alpha \in K \Rightarrow a + n\alpha \in A$  by Step 2, so  $n\alpha \in A$ , so  $\alpha \in A$  by divisibility of  $A$  — contradicting the assumption).

**Step 4:** General case.

*Claim 16.* There is some  $\beta \in G$  such that  $\pi(\beta) = f(\pi(\alpha))$  and  $\alpha = \beta \pmod{kG}$  for all  $k \in \omega$ .

*Proof.* By  $\omega$ -saturation we only need to consider one value of  $k$  at a time. Let  $g \in G_0$  be such that  $P_k(g + \alpha)$  holds, then  $\pi(g + \alpha)$  is  $k$ -divisible as well. As  $g \in A \Rightarrow g + \alpha \in A(\alpha) \Rightarrow \pi(g + \alpha) \in A \cap H$  and  $f|_{A \cap H}$  is  $L_H$ -elementary, it follows that  $f(\pi(g + \alpha))$  is  $k$ -divisible as well. Take  $\beta$  to be  $k\beta' - g$  where  $\pi(\beta') = \frac{f(\pi(g + \alpha))}{k}$  (recall that  $H$  is torsion free). Now we have  $P_k(g + \beta)$  and  $\pi(\beta) = k\pi(\beta') - \pi(g) = f(\pi(g + \alpha)) - \pi(g) = f(\pi(g)) + f(\pi(\alpha)) - \pi(g) = f(\pi(\alpha))$  as  $f(\pi(g)) = \pi(f(g))$  and  $f(g) = g$ , so the claim is proved.  $\square$

We define  $F(a + n\alpha) = f(a) + n\beta$  and  $F|_{A(\alpha) \cap H = A \cap H} = f|_{A \cap H}$ .

- $F$  is well-defined: If  $a + n\alpha = a' + n'\alpha$ , then  $(a - a') + (n - n')\alpha = 0 \in A$ , which implies by the assumption that  $n = n'$  and  $a = a'$ .
- $F$  is a homomorphism: clear from definition and as  $f$  is a homomorphism on  $A$ .
- $F$  preserves  $\pi$  (so in particular  $K$ ):  $\pi(F(a + n\alpha)) = \pi(f(a) + n\beta) = \pi(f(a)) + n\pi(\beta) = f(\pi(a)) + nf(\pi(\alpha)) = f(\pi(a) + n\pi(\alpha)) = f(\pi(a + n\alpha)) = F(\pi(a + n\alpha))$ .

- $F$  preserves  $P_k$ :  $P_k(F(a + n\alpha)) \Leftrightarrow P_k(f(a) + n\beta) \Leftrightarrow P_k(a + n\beta)$  (as  $f(a) = a \pmod{kG}$  because we have all the representatives in  $G_0$ )  $\Leftrightarrow P_k(a + n\alpha)$  (as  $\alpha = \beta \pmod{kG}$  by the choice of  $\beta$ ).
- $F$  preserves  $L_K$ -formulas and  $L_H$ -formulas: as in Step 3. □

**Corollary 17.**  *$H$  and  $K$  are fully stably embedded, i.e. any subset of  $H$  (resp.  $K$ ) definable with external parameters is already definable with internal parameters in  $L_H$  (resp.,  $L_K$ ) — this follows directly from the elimination of quantifiers.*

## 4 Reduction from RV to $k$ and $\Gamma$

**Proposition 18.** *Let  $M = (G, K, H)$  be a structure satisfying the assumptions from the previous section. Assume moreover that:*

1.  $K$  (viewed as an  $L_K$  structure) and  $H$  (viewed as an  $L_H$  structure) are both inp-minimal;
2. for every  $n$ , there are only finitely many  $x \in G$  for which  $nx = 0$  (since  $H$  is torsion-free, such elements are in fact in  $K$ ).

Then  $M$  is inp-minimal.

*Proof.* First, we note that we may assume that  $K$  is infinite. To see this, assume that  $K$  is finite, and let  $b \in G$  be given along with two mutually indiscernible sequences  $(a_i : i < \omega)$  and  $(a'_i : i < \omega)$ . By dp-minimality of  $H$ , we may assume that  $(\pi(a_i) : i < \omega)$  is indiscernible over  $\pi(b)$ . But then as  $a_i \in \text{acl}(\pi(a_i))$ , we have that  $(a_i : i < \omega)$  is indiscernible over  $\text{acl}(\pi(b))$ , hence over  $b$ . This shows that  $G$  is dp-minimal. Hence from now on, we assume that  $K$  is infinite.

We are working in a saturated extension of  $M$ . Assume that the conclusion fails, then we have an inp-pattern  $\phi(x, y), \phi'(x, y'), \bar{a} = (a_i), \bar{a}' = (a'_i)$  witnessing this, with  $\bar{a}$  and  $\bar{a}'$  mutually indiscernible. In particular they are mutually indiscernible over  $G_0 \subseteq \text{acl}(\emptyset)$  which contains representatives of each class of  $G/nG$  and all torsion of  $G$ , and rows are  $k_*$ -inconsistent. It follows from quantifier elimination that  $\phi(x, a_i)$  is equivalent to a disjunction of conjuncts of the form  $\theta(t_{i,0}(x), \dots, t_{i,l-1}(x), \alpha_i) \wedge \psi(\pi(x), b_i) \wedge \chi(x, c_i) \wedge \rho(x, e_i)$  where:

- the  $t_{i,j}$  are terms with parameters in  $G$ ,  $\alpha_i \in K$  and  $\theta$  is an  $L_K$ -formula;
- $\psi$  is an  $L_H$ -formula and  $b_i \in H$ ;
- $\chi(x, c_i)$  is of the form  $\bigwedge_{j < k} n_j x + c_{i,j} = 0 \wedge \bigwedge_{j < k} m_j x + d_{i,j} \neq 0$  with  $c_i = (c_{i,j})_{j < k} \hat{\ } (d_{i,j})_{j < k}$  from  $G$ ;
- $\rho(x, e_i)$  is of the form

$$\bigwedge_{j < k} P_{m'_j}(n'_j x + e'_{i,j}) \wedge \bigwedge_{j < k} \neg P_{m''_j}(n''_j x + e''_{i,j})$$

with  $e_i = (e'_{i,j})_{j < k} \hat{\ } (e''_{i,j})_{j < k}$ .

Forgetting all but one disjunct satisfied by  $b$ , we may assume that  $\phi(x, a_i)$  is equal to such a conjunction.

Any term  $t_{i,j}$  is of the form  $n_{i,j}x - g_{i,j}$  and the formula makes sense only when  $n_{i,j}x - g_{i,j} \in K$ , that is when  $\pi(x) = \pi(g_{i,j})/n_{i,j}$ . Choose some  $h_i$  such that  $\pi(h_i) = \pi(g_{i,j})/n_{i,j}$  for some/all  $j$ . We can then replace  $n_{i,j}x - g_{i,j}$  with  $n(x - h_i) + h'_{i,j}$  with  $h'_{i,j} \in K$ . Adding  $h'_{i,j}$  to  $\alpha_i$  and changing the formula  $\theta$ , we replace  $\theta$  by a formula  $\theta'(x - h_i, \alpha'_i)$ ,  $\theta' \in L_K$ .

Recalling that  $G/nG$  is finite for every  $n < \omega$ ,  $\rho(x, e_i)$  is equivalent to some finite disjunction of the form  $\bigvee_{i < N} P_{k_i}(x - g_i)$  where  $g_i \in G_0$  (so for example to express  $\neg P_k(nx + e)$  we have to say that  $x$  belongs to one of the finitely many classes  $\pmod{kG}$  satisfying this, and to

express  $P_k(nx + e) \wedge P_l(n'x + e')$  we have to say that  $x$  belongs to a certain subset of the classes mod  $klG$ .

Note that  $\chi(x, c_0)$  is infinite as  $\chi(x, c_0) \wedge \phi'(x, a_i)$  is consistent for every  $i \in \omega$ , while  $\{\phi'(x, a_i)\}_{i \in \omega}$  is  $k_*$ -inconsistent. Thus  $\chi(x, c_i)$  can only be of the form  $\bigwedge_{j < k} n_j x + c_{i,j} \neq 0$  (as every equation of the form  $nx + c = 0$  has only finitely many solutions by assumption (2)).

Thus we may assume that  $\phi(x, a_i) = \theta(x - h_i, \alpha_i) \wedge \psi(\pi(x), b_i) \wedge \chi(x, c_i) \wedge P_l(x - g)$  where:

- $\alpha_i \in K$  and  $\theta$  is an  $L_K$ -formula,
- $\psi$  is an  $L_H$ -formula and  $b_i \in \Gamma$ ,
- $\chi(x, c_i) = \left( \bigwedge_{j < k} n_j x + c_{i,j} \neq 0 \right)$
- $l \in \omega, g \in G_0$ .

Similarly, we may assume that  $\phi'(x, a'_i) = \theta'(x - h'_i, \alpha'_i) \wedge \psi'(\pi(x), b'_i) \wedge \chi'(x, c'_i) \wedge P_{l'}(x - g')$  with the same properties.

Let  $b \models \phi(x, a_0) \wedge \phi'(x, a'_0)$ .

**Case 1:**  $b \in H$ . Then by full stable embeddedness of  $H$  we can replace our array by  $\tilde{\phi}(x, \tilde{a}_i)$  and  $\tilde{\phi}'(x, \tilde{a}'_i)$  where  $\tilde{\phi}, \tilde{\phi}' \in L_H$  and  $\tilde{a}_i, \tilde{a}'_i \in H$  are such that  $\tilde{\phi}(x, \tilde{a}_i) \cap H(x) = \phi(x, a_i) \cap H(x)$ , and similarly for  $\tilde{\phi}'$ . But this contradicts inp-minimality of  $(H, L_H)$ .

**Case 2:**  $b \in K$ . Similarly, by full stable embeddedness of  $K$  we can replace our array by  $\tilde{\phi}(x, \tilde{a}_i)$  and  $\tilde{\phi}'(x, \tilde{a}'_i)$  where  $\tilde{\phi}, \tilde{\phi}' \in L_K$  and  $\tilde{a}_i, \tilde{a}'_i \in K$  are such that  $\tilde{\phi}(x, \tilde{a}_i) \cap K(x) = \phi(x, a_i) \cap K(x)$ , and similarly for  $\tilde{\phi}'$ . But this contradicts inp-minimality of  $(K, L_K)$ .

**Case 3:**  $b \notin K \cup H$ .

**Subcase 3.1** Neither  $\theta$  occurs in  $\phi$  nor  $\theta'$  occurs in  $\phi'$ .

Then we have  $\phi(x, a_i) = \psi(\pi(x), b_i) \wedge \chi(x, c_i) \wedge P_l(x - g)$  and  $\phi'(x, a'_i) = \psi'(\pi(x), b'_i) \wedge \chi'(x, c'_i) \wedge P_{l'}(x - g')$ .

Consider  $\tilde{\psi}(x', b_i) := \psi(x', b_i) \wedge \text{“}x' - \pi(g) \text{ is } l\text{-divisible”}$  and  $\tilde{\psi}'(x', b'_i) := \psi'(x', b'_i) \wedge \text{“}x' - \pi(g') \text{ is } l'\text{-divisible”}$  — this is an array in the structure induced on  $H$ . Note that  $\pi(b) \models \tilde{\psi}(x', b_0) \wedge \tilde{\psi}'(x', b'_0)$ .

As  $H$  is inp-minimal, it follows without loss of generality that the set

$$\{\psi(x', b_i) \wedge \text{“}x' - \pi(g) \text{ is } l\text{-divisible”} : i < \omega\}$$

has a solution  $h$  in  $H$ .

Say  $h - \pi(g) = l\gamma$ . Take  $\beta \in G$  such that  $\pi(\beta) = \gamma$ . As  $K$  is infinite, there is an infinite sequence  $(\beta_i)_{i \in \omega}$  in  $K$  such that all the differences  $\beta_i - \beta_j$  are pairwise different. Let  $e'_i = \beta + \beta_i$ . Then we still have that  $e'_i - e'_j$  are all pairwise different, and that  $\pi(e'_i) = \pi(\beta) + \pi(\beta_i) = \gamma$ . Note that as by assumption there are only finitely many  $l$ -torsion elements in  $G$ , we may assume that  $e'_i - e'_j$  is not  $l$ -torsion, for any  $i \neq j$ .

Finally, define  $e_i = le'_i + g$ . We have:

- all  $e_i$ 's are pairwise different (as  $e_i = e_j \Rightarrow (e'_i - e'_j)$  is  $l$ -torsion, contradicting the choice of the elements  $b'_i$ ).
- $\pi(e_i) = l\pi(e'_i) + \pi(g) = l\gamma + \pi(g) = h$ .
- $P_l(e_i - g)$  holds as  $e_i - g = le'_i$ .

As the set  $\bigvee_{i < k_* + 1} \left( \bigvee_{j < k} n_j x + c_{i,j} = 0 \right)$  is finite, then one of the  $e_i$ 's realizes  $k_*$  elements of the first row — a contradiction.

So let now  $(e_i : i \in \omega)$  be an infinite list of pairwise different solutions of  $\{\tilde{\psi}(x', b_i) : i \in \omega\}$  in  $H$ . In particular  $e_i - \pi(g) = l\gamma_i$  for some  $\gamma_i \in H$  with  $(\gamma_i : i \in \omega)$  pairwise different. Let  $\beta_i \in G$  be arbitrary such that  $\pi(\beta_i) = \gamma_i$ . As all fibers of  $\pi$  are finite, we may assume that all of  $\beta_i$ 's are pairwise different as well. Finally, let  $f_i := l\beta_i + g$ . We have:

- $(f_i : i \in \omega)$  are pairwise different,
- $P_l(f_i - g)$  holds for all  $i \in \omega$ , as  $f_i - g = l\beta_i$ ,
- $\pi(f_i) = l\pi(\beta_i) + \pi(g) = l\gamma_i + \pi(g) = e_i$ .

As the set  $\bigvee_{i < k_* + 1} \left( \bigvee_{j < k} n_j x + c_{i,j} = 0 \right)$  is finite, then one of the  $f_i$ 's realizes at least  $k_*$  elements of the first row — a contradiction.

**Subcase 3.2**  $\theta$  occurs in  $\phi$  and  $\theta'$  occurs in  $\phi'$ .

Syntactically, this is only possible if  $b - g_0 \in K, b - g'_0 \in K$ , hence both  $\pi(b) \in \text{dcl}(g_0)$  and  $\pi(b) \in \text{dcl}(g'_0)$ . By mutual indiscernibility of the rows it follows that  $\bar{a}, \bar{a}'$  are mutually indiscernible over  $\pi(b)$  and we can add it to the base.

Then by mutual indiscernibility of  $\bar{a}, \bar{a}'$  over  $\pi(b)$ , Ramsey, compactness and automorphism we can find some  $f \in G$  such that  $\pi(f) = \pi(b)$  and  $\bar{a}, \bar{a}'$  are mutually indiscernible over  $f$ . So we can add  $f$  to the base as well.

Taking  $c := b - f$  we have  $c \in K$ . Translating by  $f$ , we can consider a new array  $\tilde{\phi}(x, \tilde{a}_i), \tilde{\phi}'(x, \tilde{a}'_i)$  where  $\tilde{\phi}(x, a_i) = \theta(x + f - h_i, \alpha_i) \wedge \psi(\pi(x + f), b_i) \wedge \chi(x + f, c_i) \wedge P_l(x + f - g)$ , and analogously for  $\tilde{\phi}'$ . Note that the first column is realized by  $c \in K$ . By Case 2, we can find some  $c'$  realizing, say, the first row of the new array. But then taking  $b' := c' + f$  clearly  $b'$  realizes the first row of the old array.

**Subcase 3.3**  $\theta$  occurs in  $\phi$ , but  $\theta'$  does not occur in  $\phi'$  (and the symmetric case by permuting the rows).

By assumption  $\phi'(x, a'_i) = \psi'(\pi(x), b'_i) \wedge \chi'(x, c'_i) \wedge P_{l'}(x - g')$ . As in Subcase 3.1, it follows that  $\pi(b) \in \text{dcl}(a_0)$ , say  $\pi(b) = f(a_0)$  for some  $\emptyset$ -definable function  $f$ . We have  $b \models \phi'(x, a'_0)$ . In particular,  $\models \psi'(f(a_0), b'_0) \wedge \text{“}f(a_0) - \pi(g') \text{ is } l'\text{-divisible”}$ . By mutual indiscernibility of  $\bar{a}, \bar{a}'$  it follows that  $\models \psi'(f(a_i), b'_j) \wedge \text{“}f(a_i) - \pi(g') \text{ is } l'\text{-divisible”}$  for all  $i, j \in \omega$ .

We may also assume that all of  $\{f(a_i) : i \in \omega\}$  are pairwise different. Otherwise, if  $f(a_i) = f(a_j)$  for some  $i < j$ , by indiscernibility  $\pi(b) = f(a_0) = f(a_\infty)$ , and so  $\bar{a}, \bar{a}'$  are mutually indiscernible over  $\pi(b)$  — and we can conclude as in Subcase 3.1. It follows that the partial type  $\{\psi'(x', b'_j) \wedge \text{“}x' - \pi(g') \text{ is } l'\text{-divisible”}\}$  has infinitely many solutions in  $H$ , witnessed by  $\{f(a_i) : i \in \omega\}$ . Now this implies that the second row of the original array  $\{\phi'(x, a'_i) : i \in \omega\}$  is consistent as in Case 3.1(a).  $\square$

**Proof of Theorem 4.** Given a valued field  $\bar{K}$  satisfying the assumption of Theorem 4, via the reductions in Sections 2 and 3 it is enough to demonstrate that  $(\text{RV}, k, \Gamma)$  is inp-minimal. For this it is enough to show that the assumptions of Proposition 18 are satisfied for  $G = \text{RV}$ ,  $K$  a Morleyzation of  $k$  and  $H$  a Morleyzation of  $\Gamma$ . Both  $K$  and  $H$  are inp-minimal as Morleyzation obviously preserves inp-minimality,  $H$  is torsion-free since  $\Gamma$  is an ordered abelian group.

As  $\Gamma$  is an inp-minimal ordered group, it follows from [Sim11, Lemma 3.2] that  $\Gamma/n\Gamma$  is finite for all  $n \in \omega$ . Besides, we have that  $k^\times/(k^\times)^p$  is finite for all prime  $p$  by assumption. Therefore also  $\text{RV}/n\text{RV}$  is finite for all  $n$ . Finally,  $k^\times$  has finite  $n$ -torsion for all  $n$ .

## Remarks and questions

We do not know if the assumption that  $k^\times/(k^\times)^p$  is finite for all  $p$  is in fact necessary. It follows from the proof of [CKS15, Corollary 4.6] that if  $k$  is an inp-minimal field, then there can be at most one prime  $p$  for which  $k^\times/(k^\times)^p$  is infinite.

**Problem 19.** *Let  $k$  be an inp-minimal field. Is it true that  $k^\times/(k^\times)^p$  is finite for all prime  $p$ ? Or at least, can we omit this extra assumption from Theorem 4?*

The answer is positive for a dp-minimal field by the results of Johnson [Joh15] (so under the assumptions of Theorem 4, we have that  $\bar{K}$  is dp-minimal if and only if both  $k$  and  $\Gamma$  are dp-minimal), but the proof relies on the construction of a valuation which doesn't seem to be available in the general inp-minimal case.

Another natural direction is to generalize Theorem 4 from the case of burden 1 to a general burden calculation.

**Problem 20.** Let  $\bar{K} = (K, \text{RV}, \text{rv})$  be a Henselian valued field of equicharacteristic 0, viewed as a structure in the RV-language. Is it true that  $\text{bdn}(\bar{K}) = \text{bdn}(k) \times \text{bdn}(\Gamma)$ ?

## References

- [Adl07] Hans Adler. Strong theories, burden, and weight. *Preprint*, 2007. <http://www.logic.univie.ac.at/~adler/docs/strong.pdf>.
- [Ax65] James Ax. On the undecidability of power series fields. *Proceedings of the American Mathematical Society*, 16(4):846, 1965.
- [Bél99] Luc Bélair. Types dans les corps valués munis d’applications coefficients. *Illinois Journal of Mathematics*, 43(2):410–425, 1999.
- [CH14] Artem Chernikov and Martin Hils. Valued difference fields and NTP2. *Israel Journal of Mathematics*, 204(1):299–327, 2014.
- [Che10] Artem Chernikov. Indiscernible sequences and arrays in valued fields. *RIMS Kokyuroku*, 1718:127–131, 2010. <http://www.kurims.kyoto-u.ac.jp/~kyodo/kokyuroku/contents/pdf/1718-14.pdf>.
- [Che14] Artem Chernikov. Theories without the tree property of the second kind. *Annals of Pure and Applied Logic*, 165(2):695–723, 2014.
- [CKS15] Artem Chernikov, Itay Kaplan, and Pierre Simon. Groups and fields with NTP<sub>2</sub>. *Proceedings of the American Mathematical Society*, 143(1):395–406, 2015.
- [Del78] Françoise Delon. Types sur  $C((x))$ . *Study Group on Stable Theories (Bruno Poizat), Second year: 1978/79 (French)*, 1978.
- [DG15] Alfred Dolich and John Goodrick. Strong theories of ordered abelian groups. *Preprint, arXiv:1511.08274*, 2015.
- [DGL<sup>+</sup>11] Alfred Dolich, John Goodrick, David Lippel, et al. Dp-minimality: basic facts and examples. *Notre Dame Journal of Formal Logic*, 52(3):267–288, 2011.
- [Fle11] Joseph Flenner. Relative decidability and definability in Henselian valued fields. *The Journal of Symbolic Logic*, 76(04):1240–1260, 2011.
- [HO15] Nadja Hempel and Alf Onshuus. Groups in NTP2. *Preprint, arXiv:1510.01049*, 2015.
- [Joh13] Will Johnson. Forking and dividing in fields with several orderings and valuations. 2013. Preprint, <https://math.berkeley.edu/~willij/drafts/combined.pdf>.
- [Joh15] Will Johnson. On dp-minimal fields. *Preprint, arXiv:1507.02745*, 2015.
- [Mon14] Samaria Montenegro. Pseudo real closed field, pseudo p-adic closed fields and NTP2. *Preprint, arXiv:1411.7654*, 2014.
- [She90] S. Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, second edition, 1990.
- [She05] Saharon Shelah. Strongly dependent theories. *Preprint, arXiv:math/0504197*, 2005.
- [Sim11] Pierre Simon. On dp-minimal ordered structures. *The Journal of Symbolic Logic*, 76(02):448–460, 2011.