

UC Irvine

UC Irvine Previously Published Works

Title

The Brain, Seizures and Epilepsy Throughout Life: Understanding a Moving Target

Permalink

<https://escholarship.org/uc/item/6j88x1fq>

Journal

Epilepsy Currents, 12(Suppl 3)

ISSN

1535-7511

Author

Baram, Tallie Z.

Publication Date

2012

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



The Brain, Seizures and Epilepsy Throughout Life: Understanding a Moving Target

Tallie Z. Baram, MD, PhD

Departments of Anatomy/Neurobiology, Pediatrics and Neurology, University of California–Irvine, Irvine, CA
Address correspondence to Tallie Z. Baram, MD, PhD, Departments of Anatomy/Neurobiology, Pediatrics and Neurology, University of California–Irvine, Med Sci I; ZOT 4475, Irvine, CA 92697-4475. E-mail: allie@uci.edu

The human brain is a tremendously complex and still enigmatic three-dimensional structure, composed of countless interconnected neurons and glia. The temporal evolution of the brain throughout life provides a fourth dimension, one that influences every element of the brain's function in health and disease. This temporal evolution contributes to the probability of seizure generation and to the type and the nature of these seizures. The age-specific properties of the brain also influence the consequences of seizures on neuronal structure and behavior. These, in turn, govern epileptic activity and cognitive and emotional functions, contributing to the diverse consequences of seizures and epilepsy throughout life.

The Scope of the Problem

The incidence of seizures and of epilepsy varies throughout life, peaking in neonates and children and increasing again after the age of 50 (1–4). In addition to the quantitative measure of the incidence of seizures, the type of seizures (and of epilepsy syndromes) varies with age: for example, febrile seizures take place in infants and children (1, 5–7), whereas the incidence of post-stroke epilepsy increases in adults compared with children (1, 3, 8). What is the basis of these age-dependent variations? What are the age-specific properties of the brain that contribute to the probability of seizure generation and to the nature of the resulting seizure?

A second, reciprocal aspect of the interaction of the age of the brain and seizures involves the effects of a seizure on the structure and function of neurons and neuronal networks. Here, again, clinical evidence suggests that the consequences of a seizure might vary with age. For example, the consequences of status epilepticus (SE) in young children seem to depend on the inciting etiology (9, 10), and, in general, mortality and cognitive outcomes are more favorable than in the adult and elderly individual (11–14). In contrast, status epilepticus in adult and aging individuals can have major detrimental effects on cognitive function and even survival (14–16). What is the basis of these age-dependent variations? What are the age-specific properties of the brain that modulate the effects of seizures on neuronal integrity and function?

These questions are of paramount clinical importance because they should provide clues for developing age-specific diagnostic tools, prognostic models, and intervention strate-

gies. Thus, while the answers to these questions are not fully elucidated, the questions provide a framework for discussing the salient elements of the evolving interaction between the brain and seizures throughout life.

The following paragraphs highlight a few of the many facts that influence age-specific seizures and the age-specific consequences of seizures. Because most of the direct information about seizures throughout the life-span often derives from animal models, these are discussed. The author regrets that this overview is not comprehensive and likely omits a number of important publications. Nonetheless, this review serves to illustrate crucial features of the topic.

Age-Specific Brain Properties Influence Seizures That Are Generated During Each Age

Whereas brain development from embryonic life to senescence is a continuum, here it is divided into four stages: the neonatal, childhood, adulthood, and aging epochs.

The Neonatal Brain and Age-Specific Neonatal Seizures

The neonatal brain is in a stage of rapid flux from both structural and functional perspectives. Neurons are still being born, circuits are being formed, and synapses are being established (17, 18). Synaptic currents are often slower (19), neurotransmitters play trophic roles (20), and circuits are not fully mature (21). These facts predict that seizures may propagate poorly and may thus remain subclinical, or manifest as fragmented motor activity (22, 23).

From a molecular standpoint, there are developmental changes in both excitatory and inhibitory neurotransmission (24–34) excitatory and inhibitory components of synaptic transmission do not develop concurrently (33, 35). GABAergic synapses seem to be the first to function (36). In addition, GABA receptor subunit expression is developmentally regulated and is relatively low during development (30), whereas



glutamate receptor expression is high, favoring robust neurotransmission in excitatory synapses.

Indeed, both presynaptic and postsynaptic elements of glutamatergic synapses undergo developmental changes that favor hyperexcitability in the neonatal period: glutamate transporter expression is low, increasing glutamate levels at synapses. In addition, the developmental characteristics of all three types of the major glutamate receptors, NMDA, AMPA, and metabotropic, favor augmented glutamatergic neurotransmission (31–33). Together, all of the facts described above contribute to the observed facility in which insults such as hypoglycemia or hypoxia generate seizures in the neonate, and to the fragmentary nature of many neonatal seizures.

The Child's Brain and Age-Specific Seizures

Animal models have been instrumental in elucidating the changes in excitability and seizure vulnerability during the developmental transition from the neonatal to the childhood periods. In this context, it is helpful to note the complexity of comparing developmental status of human and rodent brain. A useful approach is to focus on a single region or circuit and assess multiple developmental processes, such as neurogenesis, synaptogenesis, dendritic growth, and functional maturity across species (see table in Avishai-Eliner et al. [37] for such comparison of rat, monkey, and human hippocampus). This approach enables studying the role of specific stages of maturation of a given brain region on the susceptibility to seizures and the generation of specific seizure types (38–42). Insight from non-human work has suggested that the child's limbic circuit is characterized by an overshoot of axons in some hippocampal pathways (43), and an accompanying overshoot of excitatory synapses and glutamate receptors (31, 32). This apparent augmented excitability is combined with the susceptibility of young children to sustain fever and febrile seizures as well as trauma and infections. Hence, typical seizures in the child include febrile (5, 44, 45) and traumatic (46) seizures, as well as other reactive or provoked seizures. Increasing maturation of neuronal circuits supports rapid propagation and often severe seizures in this age group (41, 42).

The Mature Brain: Sex-Specific Properties and Seizures

The spurt in the release of sex hormones that accompanies puberty and adolescence contributes to structural and functional neuronal changes and further maturation of neuronal circuits. Sex-dependent dimorphism further arises in numerous brain regions and circuits (47–49). In addition to the arrival of sex hormones from the gonad and other peripheral sources, sex hormones are also synthesized within brain regions including the hippocampus (50). The effects of the female sex hormones on neuronal excitability are complex. Estrogen is generally believed to promote excitability (51) as well as augmented synaptic plasticity (e.g., [52]). In contrast, progesterone-derived neurosteroids augment GABA_A receptor function (48, 53, 54). The cyclical nature of hormonal release during the menstrual cycle in women promotes cyclical or catamenial seizures and epilepsy, often complicating the management of seizures in women (47, 51). In analogy to the female sex hormones, the effects of testosterone—directly and indirectly—on neuronal excitability are complex (55, 56).

How Aging Influences Seizures

The incidence of seizures increases in the middle-aged and the elderly (57, 58). This is a result, in part, of the increased incidence of tumors and cerebrovascular diseases including stroke (3). In addition, there are likely intrinsic processes within the aging brain that might promote susceptibility to seizures. For example, an increasingly recognized source of the increased incidence of seizures in the aging brain may derive from increased accumulation of the molecules Tau and amyloid beta. It has been shown that these compounds may interact to influence excitability at several levels (59, 60), thus influencing excitability in both normal and dementing aging individuals. Much more work remains to be done to explain and hopefully prevent the human observational studies on increased seizure susceptibility during aging (61). The nature of seizures in the elderly might differ from those observed earlier in the life cycle. Partial complex seizures may be confused with “senior” moments, and a number of additional factors contribute to the complexity of diagnosing seizures in the elderly (62).

Age-Specific Brain Properties Influence the Consequences of Seizures at Different Ages

The Neonatal Brain: Specific Vulnerabilities to the Effects of Seizures

The neonatal brain is in a stage of rapid flux from both a structural and a functional perspective. As mentioned above, neurons are still being born, circuits are being formed, and synapses continue to be established and mature. Many of these processes are activity dependent, in that they are influenced by synaptic communication. For example, activation of certain synapses is believed to strengthen them at the expense of nonactive synapses that are pruned (63, 64). Clearly, then, a burst of abnormal neuronal activity (a seizure) has the potential to disrupt this activity-dependent process, leading to aberrant hyperexcitable or hypo-excitable circuit. At the structural level, excitatory synapses reside on dendritic spines (65); spine size and integrity, as well as the integrity of the dendrites that carry the spines, are activity dependent (66–69). These observations suggest that seizures might influence the structure of spines and dendrites during development, resulting in stunted, dysfunctional neurons (70–73).

At the molecular level, seizures during the neonatal and infancy periods modify neurotransmission through several mechanisms. Neonatal seizures may directly influence the effects of activation of GABA-A receptors (74–77) and alter receptor expression at both excitatory and inhibitory synapses. For example, such seizures modify the expression of glutamate receptors of various types (78–81), GABA receptors (82), and a number of additional important ion channels (83–86). Notably, both the structural and molecular changes provoked by neonatal seizures might be persistent (87, 88).

Excitotoxicity Throughout the Life Cycle

In both human and animal models, prolonged seizures and status epilepticus appear to result in loss of specific neuronal populations in hippocampus and the hilus of the dentate gyrus (89–94), as well as other vulnerable brain regions (for examples, see Cavalheiro et al. [90], Motte et al. [93], Pitkänen



et al. [95], and Kubová et al. [96]). However, the majority of evidence suggests that even prolonged seizures and SE are less likely to provoke cell death during development (70, 71, 97–102). Reorganization of neuronal connectivity is another common outcome of seizures in the mature brain. Typical is axonal “sprouting” and the formation of new synapse (103–107). This process, which might be driven in part by the loss of neuronal targets of the affected axons, is also more attenuated during development (108, 109). Subtle changes in neuronal structure, including loss of dendritic spines and synapses (70, 110), with potential loss of hippocampal volume (111) are found. Thus, during the developmental stages of infancy and childhood, rodent brain seems more resilient to seizure-induced excitotoxicity. However, the age-specific mechanisms responsible for this relative resilience are not fully understood.

Understanding why neurons in hippocampus and other vulnerable regions do not die, when the inciting seizures are prolonged and severe is of paramount clinical importance. The underlying mechanisms might enable future protection of neurons in the adult brain from seizure-provoked death. One potential mechanism for this resilience of neurons in immature brain to long seizures and status epilepticus is the relatively mild inflammation in response to seizures during development (112, 113). This is in contrast to the adult, where cytokines and related mediators are both released from injured cells and contribute to neuronal death (113). A second mechanism might involve the resilience of mitochondria in the immature brain to accumulation of reactive oxygen species (ROS), which promotes mitochondrial injury and cell death (114, 115). During adult SE, metabolic demand in neurons results in the formation of ROS and overwhelming of mitochondrial function. In contrast, mitochondria in immature brain are partially uncoupled due to the high basal expression of the mitochondrial uncoupling protein 2 (UCP2). This protein reduces mitochondrial membrane gradient, prevents ROS accumulation, and protects from seizure-induced cell loss (116, 117). Other properties of the developing brain, potentially including higher levels of BDNF and other growth factors might protect neurons from excitotoxic injury.

Notably, the paucity of cell loss does not indicate that seizures during infancy or childhood do not have significant sequelae. Indeed, a large body of existing and emerging literature suggests that functional (45, 118) as well as structural changes in neurons may be induced by seizures throughout life, and may contribute to cognitive deficits and emotional dysfunction (119).

Cognitive Vulnerability During Aging

An important correlate (comorbidity) of seizures and epilepsy involves cognitive deficits (110, 120–122). In humans, it is more difficult to discern the relationship of the epilepsy and the cognitive defects (122). Temporal lobe or limbic seizures propagate through the same neuronal pathways that are engaged in learning and memory processes (111). Thus, seizures might directly cause such problems; alternatively, the abnormalities that result in the seizures might independently provoke cognitive dysfunction. Work in experimental animal models demonstrated that seizures, and even interictal activity, reduce cognitive performance (118). Indeed, in both humans and ani-

mals, rapid seizure-induced injury to dendritic spines and the synapses that they carry might underlie some of the cognitive effects of epileptic activity (123).

During aging, whereas there does not seem to be a major reduction in the numbers of hippocampal neurons, spine density is reduced in pyramidal cells. This finding, reported in aging monkeys, suggests that the number of synapses is also reduced (124). This loss of functional excitatory synapses might be a basis for reduced cognitive reserve in the aging brain. The aging brain may thus be more vulnerable to seizure-induced loss of additional synapses. Other factors that contribute to vulnerability to cognitive loss that might be provoked by seizures include an established, age-dependent accumulation of mitochondrial injury (125). In addition, there is an accumulation of beta amyloid and tau, as described above. These factors, and probably many additional ones that are as yet unknown, combine to exacerbate the potential consequences of seizures on the aging brain.

Summary

The brain evolves throughout life, and this evolution contributes to the probability of the occurrence of seizures, as well as to their type. In addition, the consequences of seizures are significantly influenced by age-specific properties of the brain. Understanding the unique features of the neonatal, child, adult, and aging brain is crucial for optimal diagnosis and treatment of both seizures and their consequences.

Acknowledgments

The author thanks Mrs. Barbara Cartwright for excellent editorial assistance.

The author's research has been supported by National Institutes of Health (NIH) Grants R37 NS35439; R21 HD57563.

References

1. Hauser WA. Epidemiology of epilepsy. *Adv Neurol* 1978;19:313–339.
2. Hesdorffer DC, Logroscino G, Cascino G, Annegers JF, Hauser WA. Incidence of status epilepticus in Rochester, Minnesota, 1965–1984. *Neurology* 1998;50:735–741.
3. Jallon P, Smadja D, Cabre P, Le Mab G, Bazin M. EPIMART: Prospective incidence study of epileptic seizures in newly referred patients in a French Caribbean island (Martinique). *Epilepsia* 1999;40:1103–1109.
4. Olafsson E, Ludvigsson P, Gudmundsson G, Hesdorffer D, Kjartansson O, Hauser WA. Incidence of unprovoked seizures and epilepsy in Iceland and assessment of the epilepsy syndrome classification: A prospective study. *Lancet Neurol* 2005;4:627–634.
5. Shinnar S, Glauser TA. Febrile seizures. *J Child Neurol* 2002;17(suppl):S44–S52.
6. Dube C, Brewster AL, Richichi C, Zha QQ, Baram TZ. Fever, febrile seizures and epilepsy. *Trends Neurosci* 2007;30:490–496.
7. Hesdorffer DC, Benn EK, Bagiella E, Nordli D, Pellock J, Hinton V, Shinnar S; for the FEBSTAT Study Team. Distribution of febrile seizure duration and associations with development. *Ann Neurol* 2011;XX:XX–XX.
8. Dhanuka AK, Misra UK, Kalita J. Seizures after stroke: A prospective clinical study. *Neurol India* 2001;49:33–36.
9. Maytal J, Shinnar S, Moshé SL, Alvarez LA. Low morbidity and mortality of status epilepticus in children. *Pediatrics* 1989;83:323–331.
10. Arzimanoglou A. Outcome of status epilepticus in children. *Epilepsia* 2007;48(suppl):91–93.



11. Lowenstein DH, Alldredge BK. Status epilepticus. *N Engl J Med* 1998;338:970–976.
12. Shinnar S, Pellock JM, Berg AT, O'Dell C, Driscoll SM, Maytal J, Moshe SL, DeLorenzo RJ. Short-term outcomes of children with febrile status epilepticus. *Epilepsia* 2001;42:47–53.
13. Raspall-Chaure M, Chin RF, Neville BG, Bedford H, Scott RC. The epidemiology of convulsive status epilepticus in children: A critical review. *Epilepsia* 2007;48:1652–1663.
14. Towne AR. Epidemiology and outcomes of status epilepticus in the elderly. *Int Rev Neurobiol* 2007;81:111–127.
15. Treiman DM, Walker MC. Treatment of seizure emergencies: Convulsive and non-convulsive status epilepticus. *Epilepsy Res* 2006;68(suppl):S77–S82.
16. Leppik IE, Kelly KM, deToledo-Morrell L, Patrylo PR, DeLorenzo RJ, Mathern GW, White HS. Basic research in epilepsy and aging. *Epilepsy Res* 2006;68(suppl):S21–S37.
17. Owens DF, Kriegstein AR. Is there more to GABA than synaptic inhibition? *Nat Rev Neurosci* 2002;3:715–727.
18. Ben-Ari Y, Spitzer NC. Nature and nurture in brain development. *Trends Neurosci* 2004;27:361.
19. Hollrigel GS, Soltesz I. Slow kinetics of miniature IPSCs during early postnatal development in granule cells of the dentate gyrus. *J Neurosci* 1997;17:5119–5128.
20. Represa A, Ben-Ari Y. Trophic actions of GABA on neuronal development. *Trends Neurosci* 2005;28:278–283.
21. Ribak CE, Navetta MS. An immature mossy fiber innervation of hilar neurons may explain their resistance to kainate-induced cell death in 15-day-old rats. *Brain Res Dev Brain Res* 1994;79:47–62.
22. Mizrahi EM. Neonatal seizures: Problems in diagnosis and classification. *Epilepsia* 1987;28(suppl):S46–S55.
23. Scher MS. Controversies regarding neonatal seizure recognition. *Epileptic Disord* 2002;4:139–158.
24. Morrisett RA, Mott DD, Lewis DV, Wilson WA, Swartzwelder HS. Reduced sensitivity of the N-methyl-D-aspartate component of synaptic transmission to magnesium in hippocampal slices from immature rats. *Brain Res Dev Brain Res* 1990;56:257–262.
25. Hestrin S. Activation and desensitization of glutamate-activated channels mediating fast excitatory synaptic currents in the visual cortex. *Neuron* 1992;9:991–999.
26. Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABAA receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 1992;12:4151–4172.
27. Wisden W, Seeburg PH. GABAA receptor channels: From subunits to functional entities. *Curr Opin Neurobiol* 1992;2:263–269.
28. Monyer H, Burnashev N, Laurie DJ, Sakmann B, Seeburg PH. Developmental and regional expression in the rat brain and functional properties of four NMDA receptors. *Neuron* 1994;12:529–540.
29. Rivera C, Voipio J, Payne JA, Ruusuvuori E, Lahtinen H, Lamsa K, Pirvola U, Saarna M, Kaila K. The K⁺/Cl⁻ co-transporter KCC2 renders GABA hyperpolarizing during neuronal maturation. *Nature* 1999;397:251–255.
30. Brooks-Kayal AR. Rearranging receptors. *Epilepsia* 2005;46(suppl):29–38.
31. Talos DM, Fishman RE, Park H, Folkerth RD, Follett PL, Volpe JJ, Jensen FE. Developmental regulation of alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptor subunit expression in forebrain and relationship to regional susceptibility to hypoxic/ischemic injury. I. Rodent cerebral white matter and cortex. *J Comp Neurol* 2006;497:42–60.
32. Talos DM, Follett PL, Folkerth RD, Fishman RE, Trachtenberg FL, Volpe JJ, Jensen FE. Developmental regulation of alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid receptor subunit expression in forebrain and relationship to regional susceptibility to hypoxic/ischemic injury. II. Human cerebral white matter and cortex. *J Comp Neurol* 2006;497:61–77.
33. Ben-Ari Y, Khazipov R, Leinekugel X, Caillard O, Gaiarsa JL. GABAA, NMDA and AMPA receptors: A developmentally regulated 'ménage à trois.' *Trends Neurosci* 1997;20:523–529.
34. Flint AC, Maisch US, Weishaupt JH, Kriegstein AR, Monyer H. NR2A subunit expression shortens NMDA receptor synaptic currents in developing neocortex. *J Neurosci* 1997;17:2469–2476.
35. Pickard L, Noël J, Henley JM, Collingridge GL, Molnar E. Developmental changes in synaptic AMPA and NMDA receptor distribution and AMPA receptor subunit composition in living hippocampal neurons. *J Neurosci* 2000;20:7922–7931.
36. Ben-Ari Y. Excitatory actions of gaba during development: The nature of the nurture. *Nat Rev Neurosci* 2002;3:728–739.
37. Avishai-Eliner S, Brunson KL, Sandman CA, Baram TZ. Stressed-out, or in (utero)? *Trends Neurosci* 2002;25:518–524.
38. Mares P, Velíšek L. N-methyl-D-aspartate (NMDA)-induced seizures in developing rats. *Brain Res Dev Brain Res* 1992;65:185–189.
39. Baram TZ, Hirsch E, Snead OC III, Schultz L. Corticotropin releasing hormone induced seizures in the infant rats in the amygdala. *Ann Neurol* 1992;31:488–494.
40. Baram TZ, Gerth A, Schultz L. Febrile seizures: An appropriate-aged model suitable for long-term studies. *Brain Res Dev Brain Res* 1997;98:265–270.
41. Holmes GL, Ben-Ari Y. The neurobiology and consequences of epilepsy in the developing brain. *Pediatr Res* 2001;49:320–325.
42. Coppola A, Moshé SL. Why is the developing brain more susceptible to status epilepticus? *Epilepsia* 2009;50(suppl):25–26.
43. Gomez-Di Cesare CM, Smith KL, Rice FL, Swann JW. Axonal remodeling during postnatal maturation of CA3 hippocampal pyramidal neurons. *J Comp Neurol* 1997;384:165–180.
44. Stafstrom CE. The incidence and prevalence of febrile seizures. In: *Febrile Seizures*. (Baram TZ, Shinnar S, eds.) San Diego: Academic Press, 2002:1–25.
45. Dubé CM, Brewster AL, Baram TZ. Febrile seizures: Mechanisms and relationship to epilepsy. *Brain Dev* 2009;31:366–371.
46. Giza CC, Mink RB, Madikians A. Pediatric traumatic brain injury: Not just little adults. *Curr Opin Crit Care* 2007;13:143–152.
47. Scharfman HE, MacLusky NJ. The influence of gonadal hormones on neuronal excitability, seizures, and epilepsy in the female. *Epilepsia* 2006;47:1423–1440.
48. Lawrence C, Martin BS, Sun C, Williamson J, Kapur J. Endogenous neurosteroid synthesis modulates seizure frequency. *Ann Neurol* 2010;67:689–693.
49. McCarthy MM, Arnold AP. Reframing sexual differentiation of the brain. *Nat Neurosci* 2011;14:677–683.
50. Zhou L, Fester L, von Blittersdorff B, Hassu B, Nogens H, Prange-Kiel J, Jarry H, Wegscheider K, Rune GM. Aromatase inhibitors induce spine synapse loss in the hippocampus of ovariectomized mice. *Endocrinology* 2010;151:1153–1160.
51. Herzog AG, Harden CL, Liporace J, Pennell P, Schomer DL, Sperling M, Fowler K, Nikolov B, Shuman S, Newman M. Frequency of catamenial seizure exacerbation in women with localization-related epilepsy. *Ann Neurol* 2004;56:431–434.
52. Kramár EA, Chen LY, Brandon NJ, Rex CS, Liu F, Gall CM, Lynch G. Cytoskeletal changes underlie estrogen's acute effects on synaptic transmission and plasticity. *J Neurosci* 2009;29:12982–12993.



53. Reddy DS, Rogawski MA. Enhanced anticonvulsant activity of neuroactive steroids in a rat model of catamenial epilepsy. *Epilepsia* 2001;42:337–344.
54. Maguire JL, Stell BM, Rafizadeh M, Mody I. Ovarian cycle-linked changes in GABA(A) receptors mediating tonic inhibition alter seizure susceptibility and anxiety. *Nat Neurosci* 2005;8:797–804.
55. Harden C, MacLusky NJ. Aromatase inhibitors as add-on treatment for men with epilepsy. *Expert Rev Neurother* 2005;5:123–127.
56. Reddy DS, Jian K. The testosterone-derived neurosteroid androstenediol is a positive allosteric modulator of GABAA receptors. *J Pharmacol Exp Ther* 2010;334:1031–1041.
57. Hauser WA. Seizure disorders: The changes with age. *Epilepsia* 1992;33(suppl):S6–S14.
58. Leppik IE, Birnbaum AK. Drug treatment in the elderly. In: *Epilepsy: A Comprehensive Textbook*. Vol 2. 2nd ed. (Engel J Jr, Pedley TA, eds.) Philadelphia: Lippincott Williams & Wilkins, 2008:1269–1278.
59. Palop JJ, Mucke L. Epilepsy and cognitive impairments in Alzheimer disease. *Arch Neurol* 2009;66:435–440.
60. Roberson ED, Halabisky B, Yoo JW, Yao J, Chin J, Yan F, Wu T, Hamto P, Devidze N, Yu GQ, Palop JJ, Noebels JL, Mucke L. Amyloid- β /Fyn-induced synaptic, network, and cognitive impairments depend on tau levels in multiple mouse models of Alzheimer's disease. *J Neurosci* 2011;31:700–711.
61. Kelly KM. Aging models of acute seizures and epilepsy. *Epilepsy Curr* 2010;10:15–20.
62. Jetter GM, Cavazos JE. Epilepsy in the elderly. *Semin Neurol* 2008;28:336–41.
63. Constantine-Paton M, Cline HT. LTP and activity-dependent synaptogenesis: The more alike they are, the more different they become. *Curr Opin Neurobiol* 1998;8:139–148.
64. Akerman CJ, Cline HT. Refining the roles of GABAergic signaling during neural circuit formation. *Trends Neurosci* 2007;30:382–389.
65. Segal M. Rapid plasticity of dendritic spine: Hints to possible functions? *Prog Neurobiol* 2001;63:61–70.
66. Nikonenko I, Jourdain P, Alberi S, Toni N, Muller D. Activity-induced changes of spine morphology. *Hippocampus* 2002;12:585–591.
67. Lin B, Kramár EA, Bi X, Brucher FA, Gall CM, Lynch G. Theta stimulation polymerizes actin in dendritic spines of hippocampus. *J Neurosci* 2005;25:2062–2069.
68. Penzes P, Cahill ME, Jones KA, Srivastava DP. Convergent CaMK and RacGEF signals control dendritic structure and function. *Trends Cell Biol* 2008;18:405–413.
69. Saneyoshi T, Fortin DA, Soderling TR. Regulation of spine and synapse formation by activity-dependent intracellular signaling pathways. *Curr Opin Neurobiol* 2010;20:108–115.
70. Swann JW, Al-Noori S, Jiang M, Lee CL. Spine loss and other dendritic abnormalities in epilepsy. *Hippocampus* 2000;10:617–625.
71. Holmes GL, Khazipov R, Ben-Ari Y. Seizure-induced damage in the developing human: Relevance of experimental models. *Prog Brain Res* 2002;135:321–334.
72. Baram TZ. Long-term neuroplasticity and functional consequences of single versus recurrent early-life seizures. *Ann Neurol* 2003;54:701–705.
73. McNamara JO, Huang YZ, Leonard AS. Molecular signaling mechanisms underlying epileptogenesis. *Sci STKE* 2006;2006:re12.
74. Baram TZ, Snead OC III. Bicuculline induced seizures in infant rats: Ontogeny of behavioral and electrocortical phenomena. *Brain Res Dev Brain Res* 1990;57:291–295.
75. Velísková J, Velíšek L, Mares P, Rokyta R, Micianiková D. Ethosuximide suppresses seizures and lethality induced by picrotoxin in developing rats. *Pharmacol Biochem Behav* 1993;44:975–979.
76. Dzhala VI, Talos DM, Sdrulla DA, Brumback AC, Mathews GC, Benke TA, Delpire E, Jensen FE, Staley KJ. NKCC1 transporter facilitates seizures in the developing brain. *Nat Med* 2005;11:1205–1213.
77. Dzhala VI, Kuchibhotla KV, Glykys JC, Kahle KT, Swiercz WB, Feng G, Kuner T, Augustine GJ, Bacskai BJ, Staley KJ. Progressive NKCC1-dependent neuronal chloride accumulation during neonatal seizures. *Neurosci* 2010;30:11745–11761.
78. Zhang G, Raol YS, Hsu FC, Brooks-Kayal AR. Long-term alterations in glutamate receptor and transporter expression following early-life seizures are associated with increased seizure susceptibility. *J Neurochem* 2004;88:91–101.
79. Sanchez RM, Dai W, Levada RE, Lippman JJ, Jensen FE. AMPA/kainate receptor-mediated downregulation of GABAergic synaptic transmission by calcineurin after seizures in the developing rat brain. *J Neurosci* 2005;25:3442–3451.
80. Swann JW, Le JT, Lee CL. Recurrent seizures and the molecular maturation of hippocampal and neocortical glutamatergic synapses. *Dev Neurosci* 2007;29:168–178.
81. Rakhade SN, Zhou C, Aujla PK, Fishman R, Sucher NJ, Jensen FE. Early alterations of AMPA receptors mediate synaptic potentiation induced by neonatal seizures. *J Neurosci* 2008;28:7979–7990.
82. Raol YH, Zhang G, Lund IV, Porter BE, Maronski MA, Brooks-Kayal AR. Increased GABA(A)-receptor alpha1-subunit expression in hippocampal dentate gyrus after early-life status epilepticus. *Epilepsia* 2006;47:1665–1673.
83. Anderson AE, Hrachovy RA, Antalffy BA, Armstrong DL, Swann JW. A chronic focal epilepsy with mossy fiber sprouting follows recurrent seizures induced by intrahippocampal tetanus toxin injection in infant rats. *Neuroscience* 1999;92:73–82.
84. Brewster A, Bender RA, Chen Y, Dube C, Eghbal-Ahmadi M, Baram TZ. Developmental febrile seizures modulate hippocampal gene expression of hyperpolarization-activated channels in an isoform- and cell-specific manner. *J Neurosci* 2002;22:4591–4599.
85. Chen K, Ratzliff A, Hilgenberg L, Gulyás A, Freund TF, Smith M, Dinh TP, Piomelli D, Mackie K, Soltesz I. Long-term plasticity of endocannabinoid signaling induced by developmental febrile seizures. *Neuron* 2003;39:599–611.
86. Brewster AL, Bernard JA, Gall CM, Baram TZ. Formation of heteromeric hyperpolarization-activated cyclic nucleotide-gated (HCN) channels in the hippocampus is regulated by developmental seizures. *Neurobiol Dis* 2005;19:200–207.
87. Cornejo BJ, Mesches MH, Coultrap S, Browning MD, Benke TA. A single episode of neonatal seizures permanently alters glutamatergic synapses. *Ann Neurol* 2007;61:411–426.
88. McClelland S, Dubé CM, Yang J, Baram TZ. Epileptogenesis after prolonged febrile seizures: Mechanisms, biomarkers and therapeutic opportunities. *Neurosci Lett* 2011;497:155–162.
89. Ben-Ari Y. Limbic seizures and brain damage produced by kainic acid: Mechanisms and relevance to human temporal lobe epilepsy. *Neuroscience* 1985;14:375–403.
90. Cavalheiro EA, Silva DF, Turski WA, Calderazzo-Filho LS, Bortolotto ZA, Turski L. The susceptibility of rats to pilocarpine-induced seizures is age-dependent. *Brain Res* 1987;465:43–58.
91. Sloviter RS. The functional organization of the hippocampal dentate gyrus and its relevance to the pathogenesis of temporal lobe epilepsy. *Ann Neurol* 1994;35:640–654.



92. Buckmaster PS, Dudek FE. 1997. Neuron loss, granule cell axon reorganization and functional changes in the dentate gyrus of epileptic kainate-treated rats. *J Comp Neurol* 1997;385:385–404.
93. Motte J, Fernandes MJ, Baram TZ, Nehlig A. Spatial and temporal evolution of neuronal activation, stress and injury in lithium-pilocarpine seizures in adult rats. *Brain Res* 1998;793:61–72.
94. de Lanerolle NC, Kim JH, Williamson A, Spencer SS, Zaveri HP, Eid T, Spencer DD. A retrospective analysis of hippocampal pathology in human temporal lobe epilepsy: Evidence for distinctive patient subcategories. *Epilepsia* 2003;44:677–687.
95. Pitkänen A, Immonen RJ, Gröhn OH, Kharatishvili I. From traumatic brain injury to posttraumatic epilepsy: What animal models tell us about the process and treatment options. *Epilepsia* 2009;50(suppl):21–29.
96. Kubová H, Druga R, Lukasiuk K, Suchomelová L, Haugvicová R, Jirmanová I, Pitkänen A. Status epilepticus causes necrotic damage in the mediodorsal nucleus of the thalamus in immature rats. *J Neurosci* 2001;21:3593–3599.
97. Toth Z, Yan XX, Heftoglu S, Ribak CE, Baram TZ. Seizure-induced neuronal injury: Vulnerability to febrile seizures in an immature rat model. *J Neurosci* 1998;18:4285–4294.
98. Sperber EF, Stanton PK, Haas K, Ackermann RF, Moshé SL. Developmental differences in the neurobiology of epileptic brain damage. *Epilepsy Res Suppl* 1992;9:67–80; discussion 80–81.
99. Stafstrom CE, Thompson JL, Holmes GL. Kainic acid seizures in the developing brain: Status epilepticus and spontaneous recurrent seizures. *Dev Brain Res* 1992;65:227–236.
100. Raol YS, Budreck EC, Brooks-Kayal AR. Epilepsy after early-life seizures can be independent of hippocampal injury. *Ann Neurol* 2003;53:503–511.
101. Porter BE, Maronski M, Brooks-Kayal AR. Fate of newborn dentate granule cells after early life status epilepticus. *Epilepsia* 2004;45:13–19.
102. Baram TZ, Jensen FE, Brooks-Kayal A. Does acquired epileptogenesis in the immature brain require neuronal death. *Epilepsy Curr* 2011;11:21–26.
103. Sutula T, He XX, Cavazos J, Scott G. Synaptic reorganization in the hippocampus induced by abnormal functional activity. *Science* 1988;239:1147–1150.
104. Scharfman HE, Sollas AL, Berger RE, Goodman JH. Electrophysiological evidence of monosynaptic excitatory transmission between granule cells after seizure-induced mossy fiber sprouting. *J Neurophysiol* 2003;90:2536–2547.
105. Buckmaster PS, Zhang GF, Yamawaki R. Axon sprouting in a model of temporal lobe epilepsy creates a predominantly excitatory feedback circuit. *J Neurosci* 2002;22:6650–6658.
106. Zappone CA, Sloviter RS. Translaminar disinhibition in the rat hippocampal dentate gyrus after seizure-induced degeneration of vulnerable hilar neurons. *J Neurosci* 2004;24:853–864.
107. Thind KK, Yamawaki R, Phanwar I, Zhang G, Wen X, Buckmaster PS. Initial loss but later excess of GABAergic synapses with dentate granule cells in a rat model of temporal lobe epilepsy. *J Comp Neurol* 2010;518:647–667.
108. Holmes GL, Sarkisian M, Ben-Ari Y, Chevassus-Au-Louis N. Mossy fiber sprouting after recurrent seizures during early development in rats. *J Comp Neurol* 1999;404:537–553.
109. Bender RA, Dubé C, Gonzalez-Vega R, Mina EW, Baram TZ. Mossy fiber plasticity and enhanced hippocampal excitability, without hippocampal cell loss or altered neurogenesis, in an animal model of prolonged febrile seizures. *Hippocampus* 2003;13:399–412.
110. Ben-Ari Y, Holmes GL. Effects of seizures on developmental processes in the immature brain. *Lancet Neurol* 2006;5:1055–1063.
111. Dube C, Zhou J, Hamamura M, Zhao Q, Ring A, Abrahams J, McIntyre K, Nalcioglu O, Shatskih T, Baram TZ. Cognitive dysfunction after experimental febrile seizures. *Exp Neurol* 2009;1:167–177.
112. Dubé CM, Ravizza T, Hamamura M, Zha Q, Keebaugh A, Fok K, Andres AL, Nalcioglu O, Obenaus A, Vezzani A, Baram TZ. Epileptogenesis provoked by prolonged experimental febrile seizures: Mechanisms and biomarkers. *J Neurosci* 2010;30:7484–7494.
113. Vezzani A, French J, Bartfai T, Baram TZ. The role of inflammation in epilepsy. *Nat Rev Neurol* 2011;7:31–40.
114. Yin HZ, Sensi SL, Ogoshi F, Weiss JH. Blockade of Ca²⁺-permeable AMPA/kainate channels decreases oxygen-glucose deprivation-induced Zn²⁺ accumulation and neuronal loss in hippocampal pyramidal neurons. *J Neurosci* 2002;22:1273–1279.
115. Waldbaum S, Patel M. Mitochondria, oxidative stress, and temporal lobe epilepsy. *Epilepsy Res* 2010;88:23–45.
116. Sullivan PG, Dubé C, Dorenbos K, Steward O, Baram TZ. Mitochondrial uncoupling protein-2 protects the immature brain from excitotoxic neuronal death. *Ann Neurol* 2003;53:711–717.
117. Sullivan PG, Rippon NA, Dorenbos K, Concepcion RC, Agarwal AK, Rho JM. The ketogenic diet increases mitochondrial uncoupling protein levels and activity. *Ann Neurol* 2004;55:576–580.
118. Khan OI, Zhao Q, Miller F, Holmes GL. Interictal spikes in developing rats cause long-standing cognitive deficits. *Neurobiol Dis* 2010;39:362–371.
119. Swann JW, Baram TZ, Jensen FE, Moshé SL. Seizure mechanisms and vulnerability in the developing brain. In: *Epilepsy: A Comprehensive Textbook*. Vol 1. 2nd ed. (Engel J Jr, Pedley TA, eds). Philadelphia: Lippincott Williams & Wilkins, 2008:469–480.
120. Dreifuss FE. Cognitive function-victim of disease or hostage to treatment? *Epilepsia* 1992;33(suppl):S7–S12.
121. Hermann B, Seidenberg M, Jones J. The neurobehavioural comorbidities of epilepsy: Can a natural history be developed? *Lancet Neurol* 2008;7:151–160.
122. Hermann B, Meador KJ, Gaillard WD, Cramer JA. Cognition across the lifespan: Antiepileptic drugs, epilepsy, or both? *Epilepsy Behav* 2010;17:1–5.
123. Zeng LH, Xu L, Rensing NR, Sinatra PM, Rothman SM, Wong M. Kainate seizures cause acute dendritic injury and actin depolymerization in vivo. *J Neurosci* 2007;27:11604–11613.
124. Duan H, Wearne SL, Rocher AB, Macedo A, Morrison JH, Hof PR. Age-related dendritic and spine changes in corticocortically projecting neurons in macaque monkeys. *Cereb Cortex* 2003;13:950–961.
125. Roubertoux PL, Sluyter F, Carlier M, Marcet B, Maarouf-Veray F, Chérif C, Marican C, Arrechi P, Godin F, Jamon M, Verrier B, Cohen-Salmon C. Mitochondrial DNA modifies cognition in interaction with the nuclear genome and age in mice. *Nat Genet* 2003;35:65–69.