

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

Low subjective social status is associated with daily selection of fewer healthy foods and more high-fat/high sugar foods

Permalink

<https://escholarship.org/uc/item/6n05k8fv>

Authors

Rahal, Danny

Chiang, Jessica J

Huynh, Virginia W

et al.

Publication Date

2023

DOI

10.1016/j.appet.2022.106338

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Published in final edited form as:

Appetite. 2023 January 01; 180: 106338. doi:10.1016/j.appet.2022.106338.

Low subjective social status is associated with daily selection of fewer healthy foods and more high-fat/high sugar foods

Danny Rahal¹, Jessica Chiang², Virginia W. Huynh³, Julienne E. Bower^{4,5,6}, Heather McCreath⁷, Andrew Fuligni^{4,5,6}

¹Department of Biobehavioral Health, Pennsylvania State University, State College, PA

²Department of Psychology, Georgetown University, Washington, DC

³Department of Child and Adolescent Development, California State University, Northridge, Northridge, CA

⁴Department of Psychiatry and Biobehavioral Sciences, University of California, Los Angeles, Los Angeles, CA

⁵Cousins Center for Psychoneuroimmunology, Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA

⁶Department of Psychology, University of California Los Angeles, Los Angeles, CA

⁷Division of Geriatrics, David Geffen School of Medicine, UCLA, Los Angeles, CA

1. Introduction

People from lower socioeconomic backgrounds show greater risk for obesity and poorer metabolic health (Cohen et al., 2013; Levine, 2011; Svastisalee et al., 2012). These patterns may be partially due to differences in food selection, as lower socioeconomic status has been consistently related to less healthy eating behavior (e.g., Kirkpatrick et al., 2012; Wilson et al., 2004). However, in addition to objective income, relative differences in income can also impact obesity. Indeed, countries with higher income inequality tend to have greater prevalence of obesity, and obesity is more related to country's degree of inequality than its absolute economic level (Due et al., 2009). It is possible that feelings of low status relative to local peers may similarly relate to obesity risk in daily life, although this has not been previously tested.

Perceptions of having lower status relative to others, also known as having lower subjective social status (SSS), may be another important yet understudied factor that may relate to metabolic health and food selection. Subjective evaluations of social status can account for aspects of social position that objective socioeconomic status cannot (e.g., relative income). Studies consistently suggest that SSS is only moderately related to indicators of objective socioeconomic status such as income and education, in part because people can account for other daily experiences such as stressors when evaluating their SSS (Adler et al., 2000;

*Corresponding author: Danny Rahal, Department of Psychology, University of California, Los Angeles, 2311 Franz Hall, Los Angeles, California, 90095; Tel: (908) 249-3976; danrahal@ucla.edu.

Goodman et al., 2001). Lower SSS is often uniquely predictive of poorer metabolic health, including higher obesity risk, body mass index (BMI), adiposity, and metabolic syndrome, over and above income and education in adolescents and adults (Kaiser et al., 2012; Quon & McGrath, 2014; Tang et al., 2016). Despite these associations, it remains unclear how lower SSS may influence food selection on a daily basis, and whether this association may be driven by differences in stress. Therefore, the present study examined how feeling of low status relates to stress and daily food selection during young adulthood, when individuals have newfound autonomy in their food choices and are particularly sensitive to concerns regarding social status (Bassett et al., 2008; Forbes & Dahl, 2010).

1.1 Subjective Social Status and Eating Behavior

Theoretical research suggests that the subjective experience of having low status relative to other people, or having lower SSS, may influence eating behavior. Specifically, the insurance hypothesis posits that humans increase energy intake over energy expenditure when they are uncertain about having an adequate food supply, resulting in fat storage (Nettle et al., 2017). This response may be adaptive as people can ensure they have physiological resources available in times of need. Likewise, the resource scarcity hypothesis posits that people who perceive low status or low resource access would be in a state of chronic positive energy balance (i.e., energy intake exceeds energy expenditure) and prioritize calorie intake when resources are available (Dhurandhar, 2016). People consequently capitalize on high-calorie foods as they are available, especially when status is unstable and when future access to resources is uncertain (Kaiser et al., 2012). For instance, a previous study suggested that providing food access to low-income adults resulted in increased prevalence of obesity shortly thereafter (Fernald et al., 2008). People who have low SSS may also feel uncertain about their circumstances and prioritize calorie intake.

Empirical research similarly suggests that low SSS may influence eating behaviors. In animal models, both unstable and low hierarchical rank can elicit changes in eating behaviors. Status loss and social subordination result in preference for calorie-rich diets and increased adiposity in animals, potentially so that these animals can capitalize on available resources (e.g., Gosler, 1996; Roman et al., 2019). Subordinate animals consume more energy-dense foods and more food overall than dominant animals (e.g., Tamashiro et al., 2007; Wilson et al., 2008). Animals of lower hierarchical position may naturally tend to capitalize on high-energy foods in case they require energetic mobilization for processes to promote survival (e.g., fighting or fleeing). Indeed, species of birds of more subordinate status have evolved to have greater fat reserves than dominant status birds (Ekman & Lilliendahl, 1993; Pravosudov & Lucas, 2001). It is possible that lower subjective status (i.e., perceived rank status relative to other people) may better account for this form of social position than objective socioeconomic status in humans.

In humans, experimentally inducing people to feel of lower status or having lower resource access has been found to increase people's preferences for high-energy foods compared to fruits or vegetables (Cardel et al., 2016) and consumption of more calories and larger portions in subsequent snacks and meals (Cheon & Hong, 2017). Acute manipulations that induce lower perceived status can elicit physiological changes (e.g., increases in active

ghrelin) that stimulate appetite (Sim et al., 2018) and promote sensitivity to the caloric load of beverages, such that people can identify and prioritize high-calorie foods (Lim et al., 2020). However, despite results from these experimental studies, it remains unclear how chronic feeling of lower status in society or in one's local community can also influence food selection in everyday life.

Eating behavior in daily life may differ from the laboratory context because participants in experimental studies may not have access to preferred foods, may be more conscious of their eating in the laboratory setting, and may be influenced by other factors in their daily life (e.g., daily stress, food cost; e.g., Reichenberger et al., 2018; Robinson et al., 2015). To our knowledge, only one study to date has examined associations between low SSS and *daily* food preferences. In this study, lower SSS in society was associated with consumption of more calories per day among 17 college students across 14 days (Wijayatunga et al., 2019). However, additional investigation is needed to determine whether similar associations between lower SSS in society and food selection are found in a larger, more socioeconomically and ethnically diverse sample. Further, it remains unclear whether SSS relates to types of foods selected (e.g., fruits, fast foods), and whether SSS in both local and distal settings (e.g., college versus society) relates to daily food selection.

Food choices may be particularly influenced by SSS for young adults. Developmentally, youth are sensitive to social status concerns during adolescence and the transition to adulthood (Forbes & Dahl, 2010). Social comparison is prominent during the college transition, and lower SSS is related to poorer mental health at the start of this transition (Rahal et al., 2020; Yang et al., 2018). Also, as they age, youth transition from relying on their parents for meals to having greater autonomy over their own diets (Bassett et al., 2008). This adjustment can be challenging for youth, as college students often struggle to manage their weight (e.g., Nelson et al., 2008).

1.2 Associations between Subjective Social Status, Stress, and Eating Behavior

It is possible that feeling of low status may be stressful and thereby promote altered daily eating behavior (Bratanova et al., 2016). People of lower status generally live in circumstances which promote exposure to chronic and daily stressors, such as demands from other people, and higher perceived stress (Cundiff et al., 2020). Further, experimental and correlational studies suggest that lower SSS is related to greater psychological and physiological stress (e.g., Habersaat et al., 2018; Pieritz et al., 2016; Steen et al., 2020). Heightened stress among people with lower SSS may contribute to poorer daily eating. People often eat more when they feel stressed, potentially as a means of emotion regulation (e.g., Araiza & Lobel, 2018). Likewise, people who experience more daily hassles engage in more snacking, greater consumption of high-fat and high-sugar foods, and lower consumption of fruits and vegetables (e.g., Reichenberger et al., 2018; Zenk et al., 2014). Consequently, differences in stress may explain associations between SSS and eating behaviors. For instance, people with lower SSS were more likely to report eating as a means of managing emotions (Kauffman et al., 2020). Yet, no study has examined whether stress explains associations between SSS and food selection.

1.3 Present Study

The present study examined associations between low SSS in society and in college and daily food selection in young adults, as well as whether these associations were explained by higher levels of stress. Young adults reported their SSS in college and in American society as well as their daily servings of fruits, vegetables, fried foods, fast foods, desserts, and sweet drinks across 15 days. They also reported perceived stress and frequency of daily stressors, including conflicts and demands. In line with previous studies (e.g., Wijayatunga et al., 2019), we predicted that young adults with low SSS in society and low SSS in college would report fewer daily servings of healthy foods (i.e., fruits, vegetables) and more daily servings of high-fat/high-sugar foods (i.e., fried foods, fast foods, desserts, sweet drinks). Given evidence relating both lower SSS to higher stress and higher stress to daily eating behavior (Cundiff et al., 2020; Reichenberger et al., 2018; Zenk et al., 2014), we tested higher stress as one pathway relating low SSS to daily food selection. Models were repeated controlling for perceived stress over the past month and daily stressful events to determine whether associations between low SSS and fewer daily servings of healthy foods and more daily servings of high-fat/high-sugar foods were explained by stress.

2. Method

2.1 Participants

Data from a community sample of 131 young adults ($M_{age} = 20.3$, $SD = 0.8$; 60% female) who were part of a larger three-wave longitudinal examination of the transition from adolescence to adulthood were analyzed in this study. Participants were initially recruited from the 10th and 11th grades from four high schools in the greater Los Angeles area and had the option to participate in subsequent waves of data collection. Participants from the larger study were included in the analytic sample if they completed reports of daily food selection and reported society SSS ($N = 129$) or were enrolled at college and reported SSS at college ($N = 106$; $N = 104$ reported both). Participants were ethnically diverse (46% Latinos; 34% European American; 15% Asian American; 5% of other ethnic backgrounds including Middle Eastern, African American, and biracial). A primary caregiver reported each parent's level of education, and education was averaged across both parents when available (30% did not pursue education beyond high school, 45% completed vocational school or some college, 34% completed a college degree or higher). Caregivers also reported annual income ($M = \$81,745$, $SD = \$62,638$, range \$4,750-\$410,000). If caregivers did not report income, reports were used from data collection either two ($N = 15$) or four years prior ($N = 3$). Participants reported their personal annual income from a job, although many participants were students and without current employment ($M = \$1,559$, $SD = \$8,149$, range \$0-\$36,000).

2.2 Procedures

Participants learned about the study through flyers and in-class presentations. They had the option to continue data collection two and four years later. Analyses were limited to the third and final wave of data collection because participants reported daily food servings only at this wave. Young adults completed a psychosocial survey, in which they reported their SSS in society, SSS at their college, and perceived stress.

Young adults completed up to 15 physical, paper daily checklists ($M = 14.3$ days completed per participant, 95.3% possible days completed). At the end of each day before bed, participants reported their daily food servings and various daily events, including whether they had a meal with a family member and whether they experienced any conflicts or demands each day (yes/no). Participants received \$120 as compensation for completing the survey and two movie theater passes for on-time completion of the daily checklists in this wave. Procedures were approved by the University of California, Los Angeles Institutional Review Board and were in accordance with the Declaration of Helsinki, and all participants gave informed consent.

2.3 Measures

2.3.1 Society Subjective Social Status.—Young adults reported their SSS in society using the MacArthur Scale of Subjective Social Status—Youth Version (Adler et al., 2000; Goodman et al., 2001). Participants viewed a 10-rung ladder with the following prompt:

“Imagine that this ladder pictures how American society is set up. At the top of the ladder are the people who are the best off—they have the most money, the highest amount of schooling, and the jobs that bring the most respect. At the bottom are people who are the worst off—they have the least money, little or no education, no job or jobs that no one wants or respects. Now think about your family. Please tell us where you think your family would be on this ladder.”

SSS is a well-established indicator of status that is consistently related to objective indicators of status, relates to perceptions of status from mixed-methods research, shows test-retest reliability, and robustly shows unique associations with health (Goodman et al., 2001; Mistry et al., 2015; Operario et al., 2004; Quon & McGrath, 2014). This validated measure asks about the family’s socioeconomic status because individuals have not necessarily had enough time to develop their own socioeconomic status. Higher scores suggested higher society SSS.

2.3.2 College Subjective Social Status.—Participants viewed a second ladder with this prompt:

“Now assume that the ladder is a way of picturing your school. At the top of the ladder are the people in your school with the most respect, the highest grades, and the highest standing. At the bottom are the people who no one respects, no one wants to hang around with, and have the worst grades. Where would you place yourself on this ladder?”

Again, higher scores represented higher SSS. This scale has been well-validated, and high scores are consistently associated with better health (Goodman et al., 2001; Quon & McGrath, 2014).

2.3.3 Daily Food Selection.—Each day, participants reported how many servings they consumed of each of six types of food: fruits, vegetables, desserts, sweet drinks, fast foods, and fried foods. Items included examples of each type of food (e.g., for fast foods, “e.g., one burger, hot dog, burrito, slice of pizza, etc.”). Participants were asked to specify the number

of servings of each food type they had each day. Previous large-scale and daily studies have used similar items regarding daily servings of fruits and vegetables, and have found that greater selection of fruits and vegetables relates to greater positive affect and well-being (Conner et al., 2015; Russell et al., 1999; White et al., 2013). Desserts, sweet drinks, fast foods, and fried foods have also been examined in prior studies (e.g., Chan et al., 2015). Therefore, for this study, we created items for each food category to mirror the framing of the items regarding fruits and vegetables.

We used two exploratory factor analyses (EFA) of the daily food items, first at the daily level, and second at the person level after calculating an average for each person across all 15 days. Both EFAs suggested two factors: vegetable and fruit daily servings loaded onto one factor (healthy foods) and desserts, sweet drinks, fast foods, and fried foods loaded onto a second factor (high-sugar/high-fat foods). Items showed sufficient loading onto their respective factors (Table 1). Similar categorizations have been used in prior studies (e.g., Liao et al., 2018; O'Connor et al., 2008; White et al., 2013).

2.3.4 Daily Stressors.—Each day, young adults reported whether they experienced conflicts using five items and daily demands using six items. Separate items assessed whether participants argued with their mother or father, argued with another family member, argued with a close friend or partner, argued with or were punished by an adult at school, and were punished or disciplined by parents. Similar items regarding arguments have been used to index emotional reactivity to daily stress in married couples (Almeida et al., 2002). Participants also completed items regarding daily demands. Separate items assessed whether they had a lot of work at school, had a lot of work at home, had a lot of demands made by teachers, had a lot of demands made by friends, had a lot of demands made by family, and had a lot of demands made by a work supervisor. Prior research has used these items as indices of daily stressors and found that young adults experience shorter sleep duration and poorer mood on days when they experience more demands, and that young adults who experience more demands tend to have poorer academic performance and greater low-grade inflammation (Flook & Fuligni, 2008; Fuligni & Hardway, 2006; Levine et al., 2017). The sum number of conflicts and demands were calculated per day, and at least one conflict or demand occurred on 30.2% of days. The same pattern of results emerged when assessing daily conflicts and demands separately, as well as when dichotomizing days with respect to whether any conflict or demand was experienced that day (0 = none, 1 = any conflict or demand experienced that day).

2.3.5 Perceived Stress Scale.—As part of the psychosocial survey, young adults rated their subjective feelings of stress over the past month using the 10-item Perceived Stress Scale (Cohen et al., 1983). They rated how often they felt stressed (e.g., “How often have you found that you could not cope with all the things that you had to do”) using a five-point scale from 0 (*Never*) to 4 (*Very Often*). There were four reverse-coded items, and an average was computed such that higher scores represented more perceived stress. Items showed good reliability ($\alpha = .88$).

3. Analytic plan

Multilevel models with days (Level 1) nested within young adults (Level 2) were used to test whether SSS was related to daily food selection in Stata 16 software. Analyses were limited to the third and final wave of data collection because daily food servings were measured only at this wave. First, models tested the main effects of society SSS and college SSS on daily food selection, with each form of SSS tested in separate models. All models controlled for age (grand-mean centered), ethnicity (dummy-coded for Latino, Asian American, and other ethnic groups with European American as the reference group), and gender (effect-coded, -1 = male, 1 = female). Models were repeated additionally adjusting for parents' education, family income, and personal income to determine whether there was a unique effect of feeling of low status after controlling for objective socioeconomic status (e.g., Hoebel & Lampert, 2020). When associations emerged between SSS and healthy and high-fat/high-sugar foods, we tested whether SSS was especially related to a specific type of food. Society SSS, college SSS, parents' education, family income, and personal income were grand-mean centered.

Next, models examined how stress related to SSS and daily food servings. Society SSS and college SSS were tested as predictors of perceived stress in linear regressions and as predictors of daily stressors in multilevel models, controlling for demographic covariates. These models were also repeated controlling for parents' education, family income, and personal income. Perceived stress and daily stressors were then tested as predictors of daily food servings when controlling for demographic covariates. In these models, perceived stress was grand-mean centered, and the number of daily stressors was centered at the adolescent-mean. Finally, in order to determine whether associations between SSS and daily food selection may be explained by differences in stress, multilevel models tested whether SSS was related to young adults' daily food selection over and above demographic factors, indicators of objective socioeconomic status, and stress.

Because we were interested in young adults' own daily food choices, we examined only days on which participants did not eat with a family member in order to rule out the possibility that other people were selecting young adults' meals. Therefore, participants were included in analyses if they had reported daily servings on at least one day when they did not eat with a family member, and had reported either SSS in society ($N=129$) or reported SSS at their college ($N=106$). There were 1,035 daily observations of food choices for society SSS and 882 observations for college SSS.

4. Results

As shown in Table 2, young adults reported being above the mid-point for both society SSS and college SSS, in line with prior studies (e.g., Goodman et al., 2001). Young adults reported having about one serving each of fruits, vegetables, and sweet drinks daily and one serving each of desserts, fried foods, and fast foods every other day. As expected, young adults who had more servings of fast foods also tended to have more servings of sweet drinks, fried foods, and fast foods, and fewer vegetables. Young adults who reported having more fruits also had more vegetables and, interestingly, desserts. There was a marginally

significantly association between greater daily reports of sweet drinks, fast foods, and fried foods and larger waist-to-hip ratios (Table 3).

The distributions of college SSS and society SSS were normally distributed (skewness of -0.45 and -0.05 , respectively), although the distribution of caregiver-reported annual family income was positively skewed (skewness of 3.73). There were two outliers for college SSS (both 3.1 standard deviations below the mean) and one outlier for caregiver-reported family income (7.6 standard deviations above the mean). We repeated all analyses winsorizing these values to three standard deviations and observed no change in the reported pattern of results. Therefore, all models are presented using the unadjusted values.

First, multilevel models tested whether society SSS and college SSS related to daily food selection. Lower college SSS was associated with fewer daily servings of healthy foods ($B = 0.30$, $SE = 0.11$, $p = .006$) and more daily servings of high-fat/high-sugar foods ($B = -0.24$, $SE = 0.09$, $p = .007$). These associations remained significant over and above parents' education, family income, and personal income (Table 4, Figure 1). When disaggregating across food groups, associations were found between lower college SSS and fewer servings of both fruits ($B = 0.13$, $SE = 0.05$, $p = .017$) and vegetables ($B = 0.17$, $SE = 0.06$, $p = .006$) in fully adjusted models. The association between lower college SSS and more servings of high-fat/high-sugar foods was primarily driven by greater selection of sweet drinks ($B = -0.10$, $SE = 0.04$, $p = .028$), and marginally significantly greater selection of fast food ($B = -0.08$, $SE = 0.04$, $p = .065$). In contrast, society SSS was consistently not associated with daily food servings when adjusting for demographic factors and when additionally adjusting for parents' education, family income, and personal income, all $ps > .38$ (Table S1).

Interestingly, when examining model covariates, we observed that male participants had about 0.35 more servings of high-fat/high-sugar foods than female participants across models. Therefore, we also tested whether the degree to which associations between society SSS and college SSS differed by gender by testing interactions between SSS and gender. There was no evidence that associations between either society SSS or college SSS and daily servings were moderated by gender, all $ps > .07$.

Next, models examined whether SSS was related to perceived stress and daily stressors. Associations with stress were tested in hierarchical regressions, adjusting first for demographics and then adjusting for family income, personal income, and parents' education. Although society SSS was not related to perceived stress ($B = -0.02$, $SE = 0.03$, $p = .49$), lower college SSS was related to higher perceived stress as hypothesized ($B = -0.11$, $SE = 0.04$, $p = .003$) and this association remained significant after adjusting for objective socioeconomic status (Table S2). Associations between SSS and experiencing daily demands or conflicts were tested with multilevel models. Neither society nor college SSS were related to daily demands or conflicts, $ps > .50$ (Table S3).

Finally, models examined whether associations between SSS and daily food serving were explained by differences in stress. Models tested whether perceived stress and daily stressors were related to daily food selection, and suggested that neither perceived stress nor daily stressors were related to either healthy or high-fat/high-sugar foods when controlling for

demographic factors and when additionally controlling for socioeconomic status, $ps > .06$ (Table S4). Importantly, when perceived stress and daily stressors were included in the model, college SSS remained a significant predictor of daily selection of healthy foods ($B = 0.28$, $SE = 0.11$, $p = .015$) and high-fat/high-sugar foods ($B = -0.19$, $SE = 0.09$, $p = .043$). Overall, there was no evidence that perceived stress or daily stressors explained associations between college SSS and daily food selection.

5. Discussion

Although there are socioeconomic status-based disparities in obesity and metabolic health (e.g., Cohen et al., 2013; Levine, 2011), it remains unclear whether feeling of low status may contribute to poorer daily eating behavior. The present study examined how feeling of low status relative to others in society and college relates to daily food selection. Results suggested that young adults with lower college SSS had more daily servings of high-fat/high-sugar foods and fewer servings of healthy foods, even after controlling for objective socioeconomic status and stress, suggesting that feeling of low status in college may uniquely relate to poorer food selection.

As hypothesized, lower college SSS was related to daily selection of fewer healthy foods and more high-fat/high-sugar foods, particularly sweet drinks. In line with the resource scarcity hypothesis, young adults who have low SSS may capitalize on eating high-calorie foods including high-fat/high-sugar foods rather than healthy foods including fruits and vegetables (Dhurandhar, 2016). Research suggests that individuals tend to consume high-energy foods to promote physiological mobilization when perceiving uncertainty in food or other resources (Caldwell & Sayer, 2019). Academic performance and peer belonging are of high priority for college students (Tinto, 1975), and individuals who report low college SSS may feel insecure or uncertain about their social or academic rank relative to peers specifically. Uncertainty associated with psychologically feeling of low status relative to peers may promote poor eating behaviors, even without food insecurity or low objective socioeconomic status.

Our findings align with prior experimental studies that have found that temporarily feeling of low status is related to preference for higher-calorie meals (Cardel et al., 2016; Cheon & Hong, 2017) and heightened sensitivity to calorie-richness (Lim et al., 2020). Still, other experimental studies have found that individuals who are assigned to temporarily have lower social status (i.e., as a follower versus a leader; more challenging versus easier rules for earning money in Monopoly) do not show differences in lunchtime daily energy needs and energy intake (Cardel et al., 2020b; Pavela et al., 2017). Findings from the present study suggest that manipulating aspects of an individual's status relative to peers (e.g., academic rank, perceived reputation) may influence eating behavior.

Furthermore, our findings extend prior experimental research by examining how persistent low SSS in society and college relates to daily food selection. One prior study found that college students with lower SSS in society consume more calorie-rich foods (Wijayatunga et al., 2019). However, we observed that low SSS in college, as opposed to in society, was related to the types of foods that young adults select, which is important in light

of the protective health benefits of consuming fruits and vegetables (e.g., Van Duyn & Pivonka, 2000). It is possible that associations for society SSS would have emerged if measures assessed selection of more calorie-rich foods (e.g., grains, meat). These findings also suggest that low SSS relative to peers may position young adults for poorer health and may contribute to socioeconomic disparities in obesity and eating fruits and vegetables (e.g., Fisman et al., 2016; Sweeting et al., 1994).

The association between low college SSS and high-fat/high-sugar foods was driven by greater servings of sweet drinks. Low college SSS may position young adults for poorer metabolic health, as having sweet drinks increases risks for obesity (Luger et al., 2017; Te Morenga et al., 2013), and greater selection of sweet drinks by people of low socioeconomic status has been posited to contribute to socioeconomic status-based disparities in obesity (Bolt-Evensen et al., 2018; Hu, 2013). Young adults with lower college SSS may have more high-sugar beverages because they are more sensitive to the energy density of drinks. Prior research has demonstrated that people induced to feel of low status show greater preference for high-calorie foods and heightened ability to differentiate high-from low-calorie beverages, potentially through attentiveness to energy cues (e.g., sweetness; Cheon & Hong, 2017; Lim et al., 2020). As a result, people with chronically lower SSS in daily life may have more sweet drinks, as opposed to sugar-free substitutes or healthier alternatives.

Interestingly, lower college SSS, but not society SSS, was related to selection of fewer daily servings of healthy foods and more daily servings of high-sugar/high-fat foods. It is possible that individuals develop a unique sense of status across social contexts, as individuals tend to show only a moderate association between their standing in society and their standing in local contexts (Goodman et al., 2001; Rahal et al., 2020). Local SSS is often more strongly related to health outcomes than society SSS, potentially because local standing is more salient than societal standing (e.g., Habersaat et al., 2018; Rahal et al., 2020; Zell et al., 2018). For instance, female adolescents with lower school SSS have been previously found to show greater increases in BMI the following year (Goodman et al., 2003). Young adults may be more affected by their local status because they have more perceived control over their college status than their status in society. Perceptions of family's standing may become less tied to food selection as youth transition to adulthood and develop their own sense of status (e.g., Goodman et al., 2001). Also, social relationships and status are particularly salient during adolescence and young adulthood (e.g., Forbes & Dahl, 2010). As a result, young adults may be more invested in their status relative to peers than relative to society, and consequently more affected on a daily basis by their college SSS than their society SSS.

We also found that male participants reported daily selection of more high-fat/high-sugar foods, but not healthy foods, than female participants. This difference aligns with prior evidence that male young adults are more likely to have fast food and soft drinks compared to female young adults (Lee & Allen, 2021; Park et al., 2014). Yet, in contrast to a prior experimental study (Cardel et al., 2020b), we did not observe differences in associations between SSS and daily food selection between male and female participants. It is possible that gender differences may emerge in acute, but not chronic, feelings of low status may relate to diet-related outcomes. Another non-experimental study found that higher SSS was

related to lower severity of metabolic syndrome among women but not men in a sample of adults (Cardel et al., 2020a), suggesting that gender differences in associations may emerge later in adulthood or with respect to diet-related outcomes beyond food selection.

Associations between low college SSS and food selection appeared to be independent of stress. Lower college SSS was related to higher perceived stress, but not daily stress, and society SSS was not related to perceived or daily stress. SSS is inconsistently related to perceived stress (e.g., Steen et al., 2020; Ursache et al., 2015), and associations may be weaker for young adults, who experience stressors in varied domains including jobs and academics, than for older adults (Eccles et al., 2003). Also, neither perceived stress over the past month nor daily stressors related to food selection across the full sample in this study. This may be because the effect of stress on food selection has been found most robustly in the context of experimental rather than naturalistic stressors (e.g., Oliver et al., 2000), and the effect of stress on daily food selection often varies with dispositional factors, such as emotion regulation or stress management (e.g., Errisuriz et al., 2016).

Given that stress did not explain associations between low college SSS and poorer food selection, it is possible that low SSS may promote a sense of relative deprivation—or feeling lesser and worse off relative to other people—and this feeling may contribute to poorer food selection. For instance, low status has been associated with lower sense of control, such that people may capitalize on resources and high-calorie foods when available (Kraus et al., 2009). Young adults may also compensate for their low status by selecting high-calorie foods that are associated with higher status (Briers & Laporte, 2013). Certain foods such as meat are viewed as symbols of high status (Chan & Zlatevska, 2019), whereas plant-based diets can be more stigmatized or viewed more negatively (Markowski & Roxburgh, 2019).

This study has strengths including the socioeconomic and ethnic diversity of the sample, the rigorous sampling of food selection across 15 days, and the high levels of data completion across days. However, results must be interpreted in the context of limitations. First, participants were at different colleges. Both access to healthy foods and grocery stores and the effects of low college SSS may vary across college campuses. Future studies could be improved by measuring participants' perceptions of availability and barriers to access of different types of food. Second, generalizability of the results is limited by aspects of the sample. College SSS was assessed only among participants who were enrolled in college, and future research can assess whether college SSS with respect to the local community or workplace may also relate to daily food selection. Participants also reported generally high levels of college SSS on average, and no participants endorsed the scale minimum. Although the mean level of college SSS is comparable to values found in other studies of college students (e.g., Rahal et al., 2020), future studies should replicate these associations with greater representation of students with low college SSS. Further, although the present sample is ethnically diverse and has an ethnic breakdown comparable to the larger county, results should be replicated with nationally representative samples that include larger numbers of other ethnic groups (e.g., African Americans). Third, young adults reported foods via self-report. To reduce the burden of rigorously reporting food selection daily over two weeks, participants were asked about servings of only certain food groups. Although this scale was developed to mirror existing scales, the present study did not use a validated

measure of daily food servings, and the administered scale omitted daily servings of meat, grains, and snacks. Therefore, we were unable to determine whether society SSS and college SSS are related to daily selection of these foods.

Fourth, this study was embedded within a larger longitudinal study, such that attrition from earlier waves may have influenced participant characteristics (i.e., participants with lower society SSS may be less likely to continue to complete subsequent waves of the study). This also resulted in a relatively low sample size which may limit statistical power, and future studies may be better positioned to identify associations between low college SSS and perceived stress by including larger sample sizes. Still, the present study included up to 15 daily checklists per participant, which enabled assessment of associations at the daily level (i.e., associations between daily food selection and daily stressors), and included a larger and more diverse sample compared to previous studies (Wijayatunga et al., 2019). Fifth, although family and personal income were covaried in analyses, there was no measure of participants' subjective financial concerns or economic hardship. Finally, this study was correlational. Although low SSS may causally influence daily food choices in line with prior experiments, it is also possible that aspects of the home environment or personality factors (e.g., neuroticism) influence both food selection and college SSS. There is also the possibility for the reverse causal pathways, as individuals who have poorer eating habits may be mistreated by peers based on their appearance and therefore report lower college SSS. Further research is needed to identify the specific psychophysiological mechanisms that may explain associations between low college SSS and poorer food selection.

6. Conclusions

Taken together, results suggested that young adults with lower college SSS tend to have fewer daily servings of fruits and vegetables and more servings of high-sugar/high-fat foods, particularly sweet drinks. Feeling of low status or relative inequality may permeate one's daily life and contribute to dietary choices. Colleges can consider means of mitigating status-based differences in obesity by addressing factors that may influence students' perceived status, such as by promoting social belonging and reducing mistreatment for individuals from marginalized backgrounds. To reduce existing disparities in obesity, interventions may need to address both low objective socioeconomic status as well as feeling of relatively lower status.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Funding:

This research was supported by NIH National Center for Advancing Translational Science (NCATS) UCLA CTSA (UL1TR001881) and funding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (R01-HD062547), the UCLA California Center for Population Research (P2C-HD041022), the UCLA Older Americans Independence Center (P30-AG028748), and the USC/UCLA Center for Biodemography and Population Health (P30-AG017265). Danny Rahal was supported by a National Institutes of Health grant 1 F31 DA051181-01A1 and by funding from the Prevention and Methodology Training Program (T32 DA017629).

Data Availability Statement:

Data were collected as part of a longitudinal study. Data are available upon request.

References

- Adler NE, Epel ES, Castellazzo G, & Ickovics JR (2000). Relationship of subjective and objective social status with psychological and physiological functioning: Preliminary data in healthy, White women. *Health Psychology, 19*(6), 586. 10.1037/0278-6133.19.6.586 [PubMed: 11129362]
- Almeida DM, McGonagle KA, Cate RC, Kessler RC, & Wethington E (2002). Psychosocial moderators of emotional reactivity to marital arguments: Results from a daily diary study. *Marriage & Family Review, 34*(1–2), 89–113. 10.1300/J002v34n01_05
- Araiza AM, & Lobel M (2018). Stress and eating: Definitions, findings, explanations, and implications. *Social & Personality Psychology Compass, 12*(4), e12378. 10.1111/spc3.12378
- Bassett R, Chapman GE, & Beagan BL (2008). Autonomy and control: The co-construction of adolescent food choice. *Appetite, 50*(2–3), 325–332. 10.1016/j.appet.2007.08.009 [PubMed: 17936413]
- Bolt-Evensen K, Vik FN, Stea TH, Klepp KI, & Bere E (2018). Consumption of sugar-sweetened beverages and artificially sweetened beverages from childhood to adulthood in relation to socioeconomic status—15 years follow-up in Norway. *International Journal of Behavioral Nutrition & Physical Activity, 15*(1), 8. 10.1186/s12966-018-0646-8 [PubMed: 29343247]
- Bratanova B, Loughnan S, Klein O, Claassen A, & Wood R (2016). Poverty, inequality, and increased consumption of high calorie food: Experimental evidence for a causal link. *Appetite, 100*, 162–171. 10.1016/j.appet.2016.01.028 [PubMed: 26809142]
- Briers B, & Laporte S (2013). A wallet full of calories: The effect of financial dissatisfaction on the desire for food energy. *Journal of Marketing Research, 50*(6), 767–781. 10.1509/jmr.10.0513
- Caldwell AE, & Sayer RD (2019). Evolutionary considerations on social status, eating behavior, and obesity. *Appetite, 132*, 238–248. 10.1016/j.appet.2018.07.028 [PubMed: 30078673]
- Cardel MI, Guo Y, Sims M, Dulin A, Miller D, Chi X, ... & Gurka MJ (2020a). Objective and subjective socioeconomic status associated with metabolic syndrome severity among African American adults in Jackson Heart Study. *Psychoneuroendocrinology, 117*. 10.1016/j.psyneuen.2020.104686
- Cardel MI, Johnson SL, Beck J, Dhurandhar E, Keita AD, Tomczik AC, ... & Piff PK (2016). The effects of experimentally manipulated social status on acute eating behavior: A randomized, crossover pilot study. *Physiology & Behavior, 162*, 93–101. 10.1016/j.physbeh.2016.04.024 [PubMed: 27094920]
- Cardel MI, Pavela G, Janicke D, Huo T, Miller D, Lee AM, ... & Allison DB (2020b). Experimentally manipulated low social status and food insecurity alter eating behavior among adolescents: a randomized controlled trial. *Obesity, 28*(11), 2010–2019. 10.1002/oby.23002 [PubMed: 33150744]
- Chan EY, & Zlatevska N (2019). Jerkies, tacos, and burgers: Subjective socioeconomic status and meat preference. *Appetite, 132*, 257–266. doi.org/10.1016/j.appet.2018.08.027 [PubMed: 30172366]
- Chan TC, Yen TJ, Fu YC, & Hwang JS (2015). ClickDiary: Online tracking of health behaviors and mood. *Journal of Medical Internet Research, 17*(6), e147. 10.2196/jmir.4315 [PubMed: 26076583]
- Cheon BK, & Hong YY (2017). Mere experience of low subjective socioeconomic status stimulates appetite and food intake. *Proceedings of the National Academy of Sciences, 114*(1), 72–77. 10.1073/pnas.1607330114
- Cohen AK, Rai M, Rehkopf DH, & Abrams B (2013). Educational attainment and obesity: A systematic review. *Obesity Reviews, 14*(12), 989–1005. 10.1111/obr.12062 [PubMed: 23889851]
- Cohen S, Kamarck T, & Mermelstein R (1983). A global measure of perceived stress. *Journal of Health and Social Behavior, 24*, 385–396. 10.2307/2136404 [PubMed: 6668417]
- Conner TS, Brookie KL, Richardson AC, & Polak MA (2015). On carrots and curiosity: Eating fruit and vegetables is associated with greater flourishing in daily life. *British Journal of Health Psychology, 20*(2), 413–427. 10.1111/bjhp.12113 [PubMed: 25080035]

- Cundiff JM, Boylan JM, & Muscatell KA (2020). The pathway from social status to physical health: Taking a closer look at stress as a mediator. *Current Directions in Psychological Science*, 29(2), 147–153. 10.1177/0963721420901596
- Dhurandhar EJ (2016). The food-insecurity obesity paradox: A resource scarcity hypothesis. *Physiology & Behavior*, 162, 88–92. 10.1016/j.physbeh.2016.04.025 [PubMed: 27126969]
- Due P, Damsgaard MT, Rasmussen M, Holstein BE, Wardle J, Merlo J, ... & Lynch, J. (2009). Socioeconomic position, macroeconomic environment and overweight among adolescents in 35 countries. *International Journal of Obesity*, 33(10), 1084–1093. 10.1038/ijo.2009.128 [PubMed: 19621018]
- Eccles J, Templeton J, Barber B, & Stone M (2003). Adolescence and emerging adulthood: The critical passage ways to adulthood. In Bornstein MH, Davidson L, Keyes CLM, & Moore KA (Eds.), *Crosscurrents in contemporary psychology. Well-being: Positive development across the life course* (p. 383–406). Lawrence Erlbaum Associates Publishers.
- Ekman JB, & Lilliendahl K (1993). Using priority to food access: Fattening strategies in dominance-structured willow tit (*Parus montanus*) flocks. *Behavioral Ecology*, 4(3), 232–238. 10.1093/beheco/4.3.232
- Errisuriz VL, Pasch KE, & Perry CL (2016). Perceived stress and dietary choices: The moderating role of stress management. *Eating Behaviors*, 22, 211–216. [PubMed: 27310611]
- Fernald LC, Gertler PJ, & Hou X (2008). Cash component of conditional cash transfer program is associated with higher body mass index and blood pressure in adults. *The Journal of Nutrition*, 138(11), 2250–2257. 10.3945/jn.108.090506 [PubMed: 18936227]
- Fismen AS, Smith ORF, Torsheim T, Rasmussen M, Pedersen Pagh T, Augustine L, ... & Samdal O (2016). Trends in food habits and their relation to socioeconomic status among Nordic adolescents 2001/2002–2009/2010. *PLoS One*, 11(2), e0148541. 10.1371/journal.pone.0148541 [PubMed: 26859568]
- Flook L, & Fuligni AJ (2008). Family and school spillover in adolescents' daily lives. *Child Development*, 79(3), 776–787. 10.1111/j.1467-8624.2008.01157.x [PubMed: 18489427]
- Forbes EE, & Dahl RE (2010). Pubertal development and behavior: Hormonal activation of social and motivational tendencies. *Brain & Cognition*, 72(1), 66–72. 10.1016/j.bandc.2009.10.007 [PubMed: 19942334]
- Fuligni AJ, & Hardway C (2006). Daily variation in adolescents' sleep, activities, and psychological well-being. *Journal of Research on Adolescence*, 16(3), 353–378. 10.1111/j.1532-7795.2006.00498.x
- Goodman E, Adler NE, Daniels SR, Morrison JA, Slap GB, & Dolan LM (2003). Impact of objective and subjective social status on obesity in a biracial cohort of adolescents. *Obesity Research*, 11(8), 1018–1026. 10.1038/oby.2003.140 [PubMed: 12917508]
- Goodman E, Adler NE, Kawachi I, Frazier AL, Huang B, & Colditz GA (2001). Adolescents' perceptions of social status: Development and evaluation of a new indicator. *Pediatrics*, 108(2), e31–e31. 10.1542/peds.108.2.e31 [PubMed: 11483841]
- Gosler AG (1996). Environmental and social determinants of winter fat storage in the great tit *Parus major*. *Journal of Animal Ecology*, 1–17. 10.2307/5695
- Habersaat S, Abdellaoui S, Geiger AM, Urben S, & Wolf JM (2018). Low subjective social status in the police is linked to health-relevant changes in diurnal salivary alpha-amylase activity in Swiss police officers. *Stress*, 21(1), 11–18. 10.1080/10253890.2017.1389882 [PubMed: 29037115]
- Hoebel J, & Lampert T (2020). Subjective social status and health: Multidisciplinary explanations and methodological challenges. *Journal of Health Psychology*, 25(2), 173–185. 10.1177/1359105318800804 [PubMed: 30230391]
- Hu FB (2013). Resolved: There is sufficient scientific evidence that decreasing sugar-sweetened beverage consumption will reduce the prevalence of obesity and obesity-related diseases. *Obesity Reviews*, 14(8), 606–619. 10.1111/obr.12040 [PubMed: 23763695]
- Kaiser KA, Smith DL Jr., & Allison D (2012). Conjectures on some curious connections among social status, calorie restriction, hunger, fatness, and longevity. *Annals of the New York Academy of Sciences*, 1264(1), 1. 10.1111/j.1749-6632.2012.06672.x [PubMed: 22834696]

- Kauffman BY, Bakhshaie J, Manning K, Rogers AH, Shepherd JM, & Zvolensky MJ (2020). The role of emotion dysregulation in the association between subjective social status and eating expectancies among college students. *Journal of American College Health*, 68(1), 97–103. 10.1080/07448481.2018.1515761 [PubMed: 30570445]
- Kirkpatrick SI, Dodd KW, Reedy J, & Krebs-Smith SM (2012). Income and race/ethnicity are associated with adherence to food-based dietary guidance among US adults and children. *Journal of the Academy of Nutrition and Dietetics*, 112(5), 624–635. 10.1016/j.jand.2011.11.012 [PubMed: 22709767]
- Kraus MW, Piff PK, & Keltner D (2009). Social class, sense of control, and social explanation. *Journal of Personality & Social Psychology*, 97(6), 992. 10.1037/a0016357 [PubMed: 19968415]
- Lee J, & Allen J (2021). Gender differences in healthy and unhealthy food consumption and its relationship with depression in young adulthood. *Community Mental Health Journal*, 57(5), 898–909. 10.1007/s10597-020-00672-x [PubMed: 32602082]
- Levine CS, Hoffer LC, & Chen E (2017). Moderators of the relationship between frequent family demands and inflammation among adolescents. *Health Psychology*, 36(5), 493. 10.1037/hea0000469 [PubMed: 28192001]
- Levine JA (2011). Poverty and obesity in the U.S. *Diabetes* 60(11), 2667–2668. 10.2337/db11-1118 [PubMed: 22025771]
- Liao Y, Schembre SM, O'Connor SG, Belcher BR, Maher JP, Dzibur E, & Dunton GF (2018). An electronic ecological momentary assessment study to examine the consumption of high-fat/high-sugar foods, fruits/vegetables, and affective states among women. *Journal of Nutrition Education & Behavior*, 50(6), 626–631. 10.1016/j.jneb.2018.02.003 [PubMed: 29573964]
- Lim EX, Forde CG, & Cheon BK (2020). Low subjective socioeconomic status alters taste-based perceptual sensitivity to the energy density of beverages. *Physiology & Behavior*, 223, 112989. 10.1016/j.physbeh.2020.112989 [PubMed: 32502527]
- Luger M, Lafontan M, Bes-Rastrollo M, Winzer E, Yumuk V, & Farpour-Lambert N (2017). Sugar-sweetened beverages and weight gain in children and adults: A systematic review from 2013 to 2015 and a comparison with previous studies. *Obesity Facts*, 10(6), 674–693. 10.1159/000484566 [PubMed: 29237159]
- Markowski KL, & Roxburgh S (2019). “If I became a vegan, my family and friends would hate me:” Anticipating vegan stigma as a barrier to plant-based diets. *Appetite*, 135, 1–9. 10.1016/j.appet.2018.12.040 [PubMed: 30605705]
- Mistry RS, Brown CS, White ES, Chow KA, & Gillen-O'Neel C (2015). Elementary school children's reasoning about social class: A mixed-methods study. *Child Development*, 86(5), 1653–1671. 10.1111/cdev.12407 [PubMed: 26300338]
- Nelson MC, Story M, Larson NI, Neumark-Sztainer D, & Lytle LA (2008). Emerging adulthood and college-aged youth: An overlooked age for weight-related behavior change. *Obesity*, 16(10), 2205. 10.1038/oby.2008.365 [PubMed: 18719665]
- Nettle D, Andrews C, & Bateson M (2017). Food insecurity as a driver of obesity in humans: The insurance hypothesis. *Behavioral & Brain Sciences*, 40. 10.1017/S0140525X16000947
- O'Connor DB, Jones F, Conner M, McMillan B, & Ferguson E (2008). Effects of daily hassles and eating style on eating behavior. *Health Psychology*, 27(1S), S20. 10.1037/0278-6133.27.1.S20 [PubMed: 18248102]
- Oliver G, Wardle J, & Gibson EL (2000). Stress and food choice: A laboratory study. *Psychosomatic Medicine*, 62(6), 853–865. 10.1097/00006842-200011000-00016 [PubMed: 11139006]
- Operario D, Adler NE, & Williams DR (2004). Subjective social status: Reliability and predictive utility for global health. *Psychology & Health*, 19(2), 237–246. 10.1080/08870440310001638098
- Park S, Pan L, Sherry B, & Blanck HM (2014). Consumption of sugar-sweetened beverages among US adults in 6 states: Behavioral Risk Factor Surveillance System, 2011. *Preventing Chronic Disease*, 11, E65. 10.5888/pcd11.130304 [PubMed: 24762529]
- Pavela G, Lewis DW, Dawson JA, Cardel M, & Allison DB (2017). Social status and energy intake: A randomized controlled experiment. *Clinical Obesity*, 7(5), 316–322. 10.1111/cob.12198 [PubMed: 28877558]

- Pieritz K, Süssenbach P, Rief W, & Euteneuer F (2016). Subjective social status and cardiovascular reactivity: An experimental examination. *Frontiers in Psychology*, 7, 1091. 10.3389/fpsyg.2016.01091 [PubMed: 27486426]
- Pravosudov VV, & Lucas JR (2001). Daily patterns of energy storage in food-caching birds under variable daily predation risk: A dynamic state variable model. *Behavioral Ecology & Sociobiology*, 50(3), 239–250. 10.1007/s002650100361
- Quon EC, & McGrath JJ (2014). Subjective socioeconomic status and adolescent health: A meta-analysis. *Health Psychology*, 33(5), 433. 10.1037/a0033716 [PubMed: 24245837]
- Rahal D, Chiang JJ, Fales M, Fuligni AJ, Haselton MG, Slavich GM, & Robles TF (2020). Early life stress, subjective social status, and health during late adolescence. *Psychology & Health*, 1–19. 10.1080/08870446.2020.1761977
- Reichenberger J, Kuppens P, Liedlgruber M, Wilhelm FH, Tiefengrabner M, Ginzinger S, & Bleichert J (2018). No haste, more taste: An EMA study of the effects of stress, negative and positive emotions on eating behavior. *Biological Psychology*, 131, 54–62. 10.1016/j.biopsycho.2016.09.002 [PubMed: 27654506]
- Robinson E, Hardman CA, Halford JC, & Jones A (2015). Eating under observation: A systematic review and meta-analysis of the effect that heightened awareness of observation has on laboratory measured energy intake. *American Journal of Clinical Nutrition*, 102(2), 324–337. 10.3945/ajcn.115.111195 [PubMed: 26178730]
- Roman KM, Wilson ME, & Michopoulos V (2019). Social status predicts response to dietary cycling in female rhesus monkeys. *Appetite*, 132, 230–237. 10.1016/j.appet.2018.07.019 [PubMed: 30032952]
- Russell D, Parnell W, Wilson N, Faed J, Ferguson E, Herbison P, ... Tukuitonga C (1999). NZ food: NZ people. Key results of the 1997 National Nutrition Survey Wellington, New Zealand: Ministry of Health.
- Sim AY, Lim EX, Leow MK, & Cheon BK (2018). Low subjective socioeconomic status stimulates orexigenic hormone ghrelin—A randomised trial. *Psychoneuroendocrinology*, 89, 103–112. 10.1016/j.psyneuen.2018.01.006 [PubMed: 29358119]
- Steen PB, Poulsen PH, Andersen JH, & Biering K (2020). Subjective social status is an important determinant of perceived stress among adolescents: A cross-sectional study. *BMC Public Health*, 20, 1–9. 10.1186/s12889-020-08509-8 [PubMed: 31898494]
- Svastisalee CM, Holstein BE, & Due P (2012). Fruit and vegetable intake in adolescents: Association with socioeconomic status and exposure to supermarkets and fast food outlets. *Journal of Nutrition and Metabolism*, 2012. 10.1155/2012/185484
- Sweeting H, Anderson A, & West P (1994). Socio-demographic correlates of dietary habits in mid to late adolescence. *European Journal of Clinical Nutrition*, 48(10), 736–748. [PubMed: 7835328]
- Tamashiro KL, Nguyen MM, Ostrander MM, Gardner SR, Ma LY, Woods SC, & Sakai RR (2007). Social stress and recovery: Implications for body weight and body composition. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 293(5), R1864–R1874. 10.1152/ajpregu.00371.2007 [PubMed: 17855491]
- Tang KL, Rashid R, Godley J, & Ghali WA (2016). Association between subjective social status and cardiovascular disease and cardiovascular risk factors: A systematic review and meta-analysis. *BMJ*, 6(3). 10.1136/bmjopen-2015-010137
- Te Morenga L, Mallard S, & Mann J (2013). Dietary sugars and body weight: Systematic review and meta-analyses of randomised controlled trials and cohort studies. *BMJ*, 346. 10.1136/bmj.e7492
- Tinto V (1975). Dropout from higher education: A theoretical synthesis of recent research. *Review of Educational Research*, 45(1), 89–125. 10.3102/00346543045001089
- Ursache A, Noble KG, & Blair C (2015). Socioeconomic status, subjective social status, and perceived stress: Associations with stress physiology and executive functioning. *Behavioral Medicine*, 41(3), 145–154. 10.1080/08964289.2015.1024604 [PubMed: 26332932]
- Van Duyn MAS, & Pivonka E (2000). Overview of the health benefits of fruit and vegetable consumption for the dietetics professional: selected literature. *Journal of the American Dietetic Association*, 100(12), 1511–1521. 10.1016/S0002-8223(00)00420-X [PubMed: 11138444]

- White BA, Horwath CC, & Conner TS (2013). Many apples a day keep the blues away—Daily experiences of negative and positive affect and food consumption in young adults. *British Journal of Health Psychology*, 18(4), 782–798. 10.1111/bjhp.12021 [PubMed: 23347122]
- Wijayatunga NN, Ironuma B, Dawson JA, Rusinovich B, Myers CA, Cardel M, ... & Dhurandhar EJ (2019). Subjective social status is associated with compensation for large meals—A prospective pilot study. *Appetite*, 132, 249–256. 10.1016/j.appet.2018.07.031 [PubMed: 30059770]
- Wilson DK, Kirtland KA, Ainsworth BE, & Addy CL (2004). Socioeconomic status and perceptions of access and safety for physical activity. *Annals of Behavioral Medicine*, 28(1), 20–28. 10.1207/s15324796abm2801_4 [PubMed: 15249256]
- Wilson M, Fisher J, Fischer A, Lee V, Harris RB, & Bartness TJ (2008). Quantifying food intake in socially housed monkeys: Social status effects on caloric consumption. *Physiology & Behavior*, 94(4), 586–594. 10.1016/j.physbeh.2008.03.019 [PubMed: 18486158]
- Yang CC, Holden SM, Carter MD, & Webb JJ (2018). Social media social comparison and identity distress at the college transition: A dual-path model. *Journal of Adolescence*, 69, 92–102. 10.1016/j.adolescence.2018.09.007 [PubMed: 30278321]
- Zell E, Strickhouser JE, & Krizan Z (2018). Subjective social status and health: A meta-analysis of community and society ladders. *Health Psychology*, 37(10), 979–987. 10.1037/hea0000667 [PubMed: 30234357]
- Zenk SN, Horoi I, McDonald A, Corte C, Riley B, & Odoms-Young AM (2014). Ecological momentary assessment of environmental and personal factors and snack food intake in African American women. *Appetite*, 83, 333–341. 10.1016/j.appet.2014.09.008 [PubMed: 25239402]




Figure 1. Modeled Values of Daily Servings of Healthy Foods (a) and High-Fat/High-Sugar Foods (b) as a Function of College Subjective Social Status, Adjusting for Demographic Factors and Objective Socioeconomic Status (i.e., Parents' Education, Family Income, and Personal Income). Note: Shaded region indicates 95% confidence interval.

Table 1.

Exploratory Factor Analyses for Daily Serving Variables.

	Average Across All Days			Daily		
	High-Sugar/High-Fat Foods	Healthy Foods	High-Sugar/High-Fat Foods	High-Sugar/High-Fat Foods	Healthy Foods	Healthy Foods
Fruit	.01	.78	-.03			.42
Vegetable	-.08	.77	-.02			.40
Sweet Drinks	.49	.01	.28		.15	.05
Desserts	.41	.26	.25			
Fried Foods	.57	-.05	.42			-.05
Fast Foods	.54	-.14	.31			-.09
Variance	1.03	1.29	0.41			0.37
Proportion	0.59	0.73	0.60			0.53

Note: Values with a factor loading over .25 are bolded. Oblique factor analyses were run across participants' mean daily servings (Average Across All Days) and across all observations (Daily).

Table 2.

Descriptive Statistics of Study Variables.

Variable	<i>N</i>	<i>M</i>	<i>SD</i>	<i>Min</i>	<i>Max</i>
Age	131	20.3	0.8	16.5	22.1
Parents' Education	131	7.4	1.8	1.5	11.0
Family Annual Income	131	\$81774.9	\$62637.5	\$4750.0	\$410000.0
Personal Annual Earnings	131	\$1559.1	\$8149.4	\$0.0	\$36000.0
Society Subjective Social Status	129	5.9	1.6	1.0	10.0
College Subjective Social Status	106	7.0	1.5	2.0	10.0
Perceived Stress	131	1.8	0.5	0.0	3.4
Daily Fruit Servings	131	0.9	0.8	0.0	3.4
Daily Vegetable Servings	131	0.8	1.0	0.0	4.9
Daily Sweet Drink Servings	131	1.0	0.8	0.0	3.5
Daily Dessert Servings	131	0.7	0.6	0.0	3.3
Daily Fried Foods Servings	131	0.6	0.5	0.0	2.3
Daily Fast Foods Servings	131	0.7	0.8	0.0	6.0

Note: Averages for each participant are calculated across all days for daily servings.

Table 3.Correlations between Mean Daily Servings ($N=131$).

	1.	2.	3.	4.	5.	6.
1. Fruit Mean	—					
2. Vegetable Mean	.68***	—				
3. Sweet Drinks Mean	-.02	-.06	—			
4. Desserts Mean	.19*	.08	.19*	—		
5. Fried Foods Mean	-.01	-.12	.31***	.15	—	
6. Fast Foods Mean	-.12	-.20*	.24**	.25**	.39***	—
7. Waist-Hip Ratio	-.12	-.06	.18*	-.06	.16 [†]	.16 [†]

Note:[†] = $p < .1$ * = $p < .05$ ** = $p < .01$ *** = $p < .001$.

Table 4. Selection of Healthy Foods (Left) and High-Fat/High Sugar Foods (Right) as a Function of College Subjective Social Status and Stress.

	Healthy Foods						High-Fat/High-Sugar Foods					
	Unadjusted Model		Adjusted for SES		Unadjusted Model		Adjusted for SES		Unadjusted Model		Adjusted for SES	
	B	SE	B	SE	B	SE	B	SE	B	SE	B	SE
Constant	1.98***	0.30	2.03***	0.32	3.46***	0.25	3.41***	0.27	0.25	0.09	-0.24**	0.09
College SSS	0.30**	0.11	0.30**	0.11	-0.24**	0.09	-0.24**	0.09	0.02	0.38	0.14	0.40
Asian American	-0.68	0.46	-0.81	0.48	0.02	0.38	0.14	0.40	0.31	0.31	-0.36	0.34
Latino	-0.34	0.37	-0.43	0.40	-0.45	0.31	-0.36	0.34	0.71	0.71	-0.71	0.71
Other Ethnicity	-0.81	0.81	-0.89	0.81	-0.75	0.71	-0.71	0.71	-0.73***	0.14	-0.73***	0.15
Gender	-0.12	0.17	-0.10	0.17	-0.73***	0.14	-0.73***	0.15	0.10	0.18	0.10	0.19
Age	0.17	0.22	0.17	0.22	0.10	0.18	0.10	0.19	—	—	-0.01	0.09
Parents' Education	—	—	0.04	0.10	—	—	—	0.02	—	—	0.02	0.02
Family Income	—	—	-0.02	0.02	—	—	—	0.02	—	—	0.07	0.22
Personal Earnings	—	—	-0.20	0.27	—	—	—	0.07	—	—	—	—

Note.

* $p < .05$

** $p < .01$

*** $p < .001$

SSS= subjective social status; SES = socioeconomic status. College SSS, Age, Parents' Education, Family Income, and Personal Earnings were grand mean-centered. Family Income and Personal Income were divided by \$10,000. Ethnicity was dummy-coded with European American as the reference group. Gender was effect-coded (-1 = male, 1 = female).