

# Lawrence Berkeley National Laboratory

## Recent Work

**Title**

ELECTRONIC STRUCTURE AND CHEMICAL DYNAMICS

**Permalink**

<https://escholarship.org/uc/item/6p2002rt>

**Author**

Mahan, Bruce H.

**Publication Date**

1974-11-01

0 0 0 0 4 2 0 0 9 4 5

Submitted to Accounts of Chemical  
Research

LBL-3525  
Preprint c. |

ELECTRONIC STRUCTURE AND CHEMICAL DYNAMICS


Bruce H. Mahan

November, 1974

Prepared for the U. S. Atomic Energy Commission  
under Contract W-7405-ENG-48

**For Reference**

Not to be taken from this room



LBL-3525  
c. |

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

## ELECTRONIC STRUCTURE AND CHEMICAL DYNAMICS

Bruce H. Mahan

Department of Chemistry and Inorganic Materials  
Research Division of the Lawrence Berkeley Laboratory,  
University of California, Berkeley, California 94720




The connection between the measurable properties of isolated molecules and the course, mechanisms, and rates of reactions is a central problem in chemistry. While an enormous number and variety of empirical, semiempirical, and fundamental approaches to this problem have and are being explored, the successes of orbital and state symmetry and correlation methods have been particularly outstanding and widespread. In addition, the importance of intersections and avoided intersections of potential energy surfaces in a variety of thermal as well as photochemical reactions has become increasingly evident.<sup>1</sup> To provide a firm basis for the understanding of the behavior of complex chemical systems, it is valuable to determine the degree to which one can rationalize or predict detailed dynamical behavior in systems comprised of a small number of atoms, but which have several low-lying, strongly interacting potential energy surfaces. The area of gaseous ion-molecule reactions provides a particularly effective vehicle for this type of study, since even in systems with very few electrons, the surfaces corresponding to the different charge exchange states of reactants and products generally lie close to each other, and often interact strongly. Moreover, detailed information on the reaction dynamics of many of these systems

is available from ion beam scattering experiments.

In this Account, I shall describe how, by using the known electronic properties of reactants, products, and intermediates, much of the dynamical behavior of these simple systems can be understood. In particular, electronic state correlation diagrams will be used to deduce the qualitative nature of the potential surfaces which control the reaction dynamics. Molecular orbital correlation diagrams have, of course, been applied extensively and successfully<sup>2</sup> to problems of this type, but much less use has been made of electronic state correlations.<sup>3,4</sup> One of the major purposes of this paper is to demonstrate the value of making electronic state correlations and illuminating them by orbital correlations.


#### Noble Gas-Hydrogen Systems

The reactions of the noble gas ions and atoms with respectively the hydrogen molecule and molecule ion together constitute an intriguing example of the relation between reactivity and electronic configuration. Despite the simplicity of these systems, there are several possible product channels. Letting A stand for any noble gas atom, we have




These processes, in addition to simple collisional dissociation without charge transfer, constitute the possible results of low energy collisions. The relative importance of the different channels varies rather spectacularly in the noble gas family.

The reaction



is endoergic, but proceeds readily if the necessary energy is supplied either as relative translational energy of the reactants, or as vibrational energy of  $\text{H}_2^+$ . Indeed, the photoionization experiments of Chupka<sup>5,6</sup> have shown that vibrational energy is particularly effective in promoting reaction. In contrast, the reaction



has never been observed, and has a rate constant that is four orders of magnitude<sup>7</sup> smaller than that expected for an exoergic ion-molecule reaction.

The explanation of these results, and a rationalization of what does happen when  $\text{He}^+$  collides with  $\text{H}_2$  can be gleaned from Figure 1. There are plotted the potential energy curves that correspond to cuts through the potential energy surfaces in the asymptotic reactant and product regions. The separated atoms with lowest total energy are  $\text{He} + \text{H}^+ + \text{H}$ . Bringing together  $\text{H}^+$  and  $\text{H}$  on the lowest surface merely generates the  ${}^2\Sigma_g^+$  potential energy curve of  $\text{H}_2^+$ , if  $\text{He}$  is far away. Bringing together  $\text{He}$  and  $\text{H}^+$  generates the lowest  ${}^1\Sigma^+$  potential energy curve of  $\text{HeH}^+$ , and this constitutes a

cut through the lowest surface for the system in the product region. Because the reactants  $\text{H}_2^+ + \text{He}$  and the products  $\text{HeH}^+ + \text{H}$  both correlate adiabatically to  $\text{He} + \text{H}^+ + \text{H}$ , we can expect that the system can pass adiabatically (i.e., on one Born-Oppenheimer surface) from reactants to products. The same conclusion can be reached for the other of the first four noble gas-hydrogen systems, as was pointed out by Kuntz and Roach.<sup>8</sup> Indeed, reaction with  $\text{H}_2^+$  to form the protonated noble gas molecular ion is observed experimentally in all these cases. The lowest potential energy surface for the  $\text{H}_2^+ - \text{He}$  system has been calculated by Brown and Hayes<sup>9</sup> for collinear conformations, and appears to have the qualitative features expected to be consistent with the observed<sup>5,6</sup> effectiveness of vibrational energy in increasing the reaction cross section.

Consider now the higher energy atomic state  $\text{He}^+ + \text{H} + \text{H}$ . Bringing together the two hydrogen atoms with paired spins generates the  $^1\Sigma_g^+$  ground state potential energy curve, which is accordingly a cut in the reactant region through the surface on which  $\text{He}^+ + \text{H}_2$  collisions occur. The apparent peculiarity of having the ground state of  $\text{H}_2$  lie above the states of  $\text{H}_2^+$  is of course a result of our (proper) inclusion of the energies of  $\text{He}$  and  $\text{He}^+$  in the diagram. To generate a cut through the product region of this upper surface, we bring together  $\text{He}^+$  and  $\text{H}$ . There are calculations<sup>10</sup> that demonstrate that the resulting excited  $^1\Sigma$  state of  $\text{HeH}^+$  is totally repulsive, and the corresponding  $^3\Sigma$  state is at best very weakly bound. Consequently, we have the first indication of the difficulty of forming a stable  $\text{HeH}^+$  molecule from  $\text{He}^+$  and  $\text{H}_2$ .

Figure 1 shows that in the asymptotic reactant region, the potential energy curve for  $\text{H}_2$  crosses the repulsive  ${}^2\Sigma_u^+$  state of  $\text{H}_2^+$  at approximately  $1\text{\AA}$ , and may intersect the repulsive wall of the  ${}^2\Sigma_g^+$  ground state of  $\text{H}_2^+$  at a smaller distance. These potential surface crossings do occur when helium is far away, and not interacting with the hydrogen moiety. In effect, the crossings occur because there is no electron readily available to convert  $\text{H}_2^+$  to  $\text{H}_2$ . However, since all the surfaces involved are of  ${}^2\Sigma - {}^2A'$  symmetry, these intersections become avoided when the helium-hydrogen separation decreases. As a result, the surface which has asymptotically the character of  $\text{He}^+ + \text{H}_2$  may upon contraction of the H-H distance adiabatically acquire the characteristic of  $\text{He} + \text{H}_2^+$  ( ${}^2\Sigma_g^+$ ), and upon expansion of the H-H distance, assume the characteristics of  $\text{He} + \text{H}_2^+$  ( ${}^2\Sigma_u^+$ ). In the latter instance, the surface leads directly to dissociation to  $\text{H}^+ + \text{H} + \text{He}$ . In the former case, the shape of the potential surface is affected, but the system remains bound with respect to  $\text{H}^+ + \text{H}$ .

There are two important questions to be considered. First, is the interaction between the primitive crossing states in fact strong enough to produce substantial mixing and avoided crossings, and thereby form an adiabatic path from  $\text{He}^+ + \text{H}_2$  to  $\text{He} + \text{H}^+ + \text{H}$ ? Second, are the avoided intersections reached in the course of a significant number of collision trajectories? For the  $\text{He}^+ - \text{H}_2$  system, these questions can only be answered by detailed calculation of the potential energy surfaces and exact trajectories on them. However, we can say that if these



avoided intersections are accessible, the result of  $\text{He}^+$  colliding with  $\text{H}_2$  should be to produce dissociative charge transfer to  $\text{H}^+ + \text{H} + \text{He}$ . It is unlikely that significant amounts of either  $\text{H}_2^+$  or  $\text{HeH}^+$  can be formed. This argument is consistent with recent experimental findings<sup>7</sup> which show that the cross section for disappearance of  $\text{He}^+$  in  $\text{H}_2$  is very small and that the product is predominantly and perhaps exclusively  $\text{He} + \text{H} + \text{H}^+$ .

A diagram which is very similar to Figure 1 can be drawn for the  $(\text{Ne} - \text{H}_2)^+$  system, with the major difference being that the reactant states which dissociate to  $\text{Ne}^+ + \text{H} + \text{H}$  are 3 eV lower than the corresponding curves in Figure 1. The conclusions to be reached are the same:  $\text{Ne} + \text{H}_2^+$  will react to produce  $\text{NeH}^+$ ,  $\text{Ne}^+ + \text{H}_2$  should be largely unreactive and produce no  $\text{NeH}^+$  or  $\text{H}_2^+$ . If anything, dissociative charge transfer to  $\text{H}^+ + \text{H}$  should occur, but with small cross section. These conclusions are consistent with the experimental facts that are known.<sup>11,12</sup>

Qualitatively different chemical behavior is observed in the  $(\text{Ar} - \text{H}_2)^+$  system, and can also be rationalized by a similar analysis of the asymptotic potential energy surfaces displayed in Figure 2. Again, since the reactants  $\text{H}_2^+ + \text{Ar}$  and the products  $\text{ArH}^+ + \text{H}$  dissociate to  $\text{H}^+ + \text{H} + \text{Ar}$ , it is not surprising that there is an adiabatic surface which leads to reaction. In the lowest approximation, one would not expect that there is a surface which permits reaction of  $\text{Ar}^+$  with  $\text{H}_2$  to produce  $\text{ArH}^+$  in its ground state. However, because of the

near coincidence of the ionization energies of Ar and  $H_2$  (15.76 and 15.43 eV, respectively), the potential curve for  $H_2^+$  (and Ar) intersects that for  $H_2$  (and  $Ar^+$ ) very near the equilibrium internuclear distance of  $H_2$ . As the argon-hydrogen distance decreases in a collision, this intersection becomes avoided, and lower and upper adiabatic noncrossing surfaces are generated.<sup>8,13</sup> The lower adiabatic surface has the character of  $Ar^+ + H_2$  at small H-H separation, and of  $Ar + H_2^+$  at larger H-H separations. Thus, during a collision of  $Ar^+$  and  $H_2$ , the system can pass adiabatically to regions closely related to  $Ar + H_2^+$  by expansion of the H-H distance. If the collision is of a grazing nature, the argon and hydrogen may separate in this electronic condition, and thereby produce the simple charge exchange products  $Ar + H_2^+$ . It is also easy to imagine that if the collision is of a more intimate nature, production of  $ArH^+$  and H can occur. Both these processes have been observed experimentally. Moreover, an exact classical trajectory analysis of  $Ar^+ - H_2$  collisions by Chapman and Preston<sup>13</sup> shows very clearly that grazing collisions lead principally to electron transfer, while more intimate collisions lead to atom transfer.

The foregoing analyses show how knowledge of the asymptotic properties of reactants and products can be used to rationalize or predict the chemical behavior of systems that can lead to more than one set of reaction products. It is seen that the central points of the analysis are delineation of the adiabatic correlations between reactants and products, and recognition

of surface intersections and of avoided surface crossings. An important lesson to be learned is that these avoided surface intersections can lead to products whose electronic orbital configurations do not correlate in lowest order with those of the reactants. This type of analysis can also be applied to rationalize the qualitative nature of the detailed reaction dynamics, as the following examples show.

#### The $C^+ - H_2$ and $N^+ - H_2$ Systems

The reaction of  $C^+$  with  $H_2$  provides a particularly instructive example of the correlation of electronic structure and reaction dynamics. Ion beam scattering experiments<sup>14</sup> show that the velocity vector distribution of the  $CH^+$  product has considerable symmetry about the  $\pm 90^\circ$  axis in the center-of-mass coordinate system. As has been extensively discussed,<sup>14,15</sup> this is consistent with but does not necessarily indicate, the occurrence of a  $CH_2^+$  collision intermediate whose lifetime is greater than a molecular rotational period. Superficially, such a "long-lived" collision complex might be expected from the fact, known from appearance potential measurements, that the  $^2A_1$  ground state of  $CH_2^+$  lies approximately 4.3 eV below the minimum energy of the reactants  $C^+ + H_2$ , and 4.7 eV below the products  $CH^+(^1\Sigma)$  and H. Such a potential energy well can produce the long-lived collision complex that is suggested by the experimental data. However, the question which should be answered is whether or not the potential energy well is accessible to reactants by a low energy adiabatic path. In other words, does  $C^+$  readily insert into  $H_2$  to form the symmetric  $CH_2^+$ ?

A first attempt to answer this question can be made by using a molecular orbital correlation diagram to determine how the various possible electron configurations of the reactants evolve to those of the intermediate  $\text{CH}_2^+$ . An orbital correlation diagram suitable for this analysis is shown in Fig. 3. We consider the part of the diagram in which  $\text{C}^+$  approaches  $\text{H}_2$  along the perpendicular bisector of the bond, so that the system remains in the  $\text{C}_{2v}$  point group. From Fig. 3, which shows only the valence shell orbitals, it is clear that in the  ${}^2\text{A}_1$  ground state of  $\text{CH}_2^+$ , the C-H bonding  $2a_1$  and  $1b_2$  orbitals are doubly occupied, and the largely non-bonding  $3a_1$  orbital is singly occupied. For  $\text{C}^+({}^2\text{P})$  approaching  $\text{H}_2$  in the  $\text{C}_{2v}$  (isocetes triangle) conformation, we have two electrons in the carbon  $2s(a_1)$  orbital, two more in the hydrogen  $\sigma_g(a_1)$  orbital, and one electron in the carbon  $2p$  orbital, which may be of  $b_1$ ,  $b_2$ , or  $a_1$  symmetry, depending on whether the symmetry axis of the orbital is perpendicular to the nuclear plane, or in-plane parallel or perpendicular to the internuclear axis of  $\text{H}_2$ . The latter choice generates a  ${}^2\text{A}_1$  potential surface, since all electrons are in orbitals of  $a_1$  symmetry. Thus, this  ${}^2\text{A}_1$  surface which has lowest energy in the reactant region does not have the same orbital occupancy as does the  ${}^2\text{A}_1$  electronic ground state of  $\text{CH}_2^+$  near its most stable equilibrium geometry. In fact, as Figure 3 shows, the orbital occupancy of the reactants corresponds to the configuration  $(2a_1)^2 (3a_1)^2 (4a_1^*)^1$ , in which the bonding  $1b_2$  orbital is unoccupied, and the antibonding  $4a_1^*$  orbital possesses one electron.

Consideration of the electron configuration leads to the conclusion that as  $C^+$  approaches  $H_2$  on the  ${}^2A_1$  surface, the potential energy must rise. Eventually this "reactant" surface must attempt to cross the  ${}^2A_1$  surface which has the ground state electronic configuration, and which dissociates in the lowest or diabatic approximation to  $C({}^1D)$  and  $H_2^+({}^2\Sigma_g^+)$ . The result is the behavior indicated on the electronic state correlation diagram, Figure 4. While a correlation diagram can only represent the true potential energy surface in a very limited and almost entirely qualitative manner, certain features can be displayed with considerable assurance. In Figure 4, we see that the  ${}^2A_1$  ground state of  $CH_2^+$  is accessible to the reactants  $C^+({}^2P)$  and  $H_2$ , but only if the system passes over what must be a considerable potential energy barrier. It is important to note how both state and orbital correlations contribute to this conclusion. Since states of the same symmetry do not cross, the lowest  ${}^2A_1$  state of reactants must correlate adiabatically to the lowest  ${}^2A_1$  state of the intermediate. However, this forces the electron orbital occupancy to change as the system evolves from reactants to the intermediate. Consideration of the orbitals occupied in the reactants leads to the conclusion that they are separated from the intermediate by a potential energy barrier.

Much the same argument can be constructed for the  ${}^2B_1$  surface of Fig. 4. On this surface, reactants have an electronic configuration which according to Figure 3 correlates to the highly excited  $(2a_1)^2 (3a_1)^2 (1b_1)^1$  configuration of

$\text{CH}_2^+$ . The configuration of the lowest  ${}^2\text{B}_1$  state of  $\text{CH}_2^+$  near its equilibrium conformation is  $(2a_1)^2 (1b_2)^2 (1b_1)^1$ , which corresponds to the highly excited reactants  $\text{C}({}^3\text{P})$  and  $\text{H}_2^+({}^2\Sigma_u^+)$ . Thus the  ${}^2\text{B}_1$  surface must have a potential energy barrier between the reactants and the potential energy well, as is indicated in Figure 4. Again, access of reactants with low kinetic energy to the strongly bound regions of  $\text{CH}_2^+$  is blocked.

The behavior of the  ${}^2\text{B}_2$  surface is qualitatively different from that of the two previous situations. The electronic configuration of the reactants on the  ${}^2\text{B}_2$  surface corresponds to  $(2a_1)^2 (1b_2)^1 (3a_1)^2$ , the expected configuration of the lowest  ${}^2\text{B}_2$  state of  $\text{CH}_2^+$ . Because there are three bonding electrons and no antibonding electrons present, the surface might be expected to be flat or slightly attractive at large  $\text{C}^+-\text{H}_2$  distances. In fact, ab initio calculations<sup>16</sup> show that the surface is rather flat initially, but rises at small  $\text{C}^+-\text{H}_2$  separations. The  ${}^2\text{B}_2$  surface therefore crosses the lowest  ${}^2\text{A}_1$  and  ${}^2\text{B}_1$  surfaces in the region where they are rapidly dropping in energy. These crossings are allowed in strict  $\text{C}_{2v}$  nuclear conformations because of the different symmetries of the states. It therefore appears that in  $\text{C}_{2v}$  conformations, reactants with low kinetic energy do not have access to the deep potential energy wells associated with the  ${}^2\text{B}_1$  and  ${}^2\text{A}_1$  states of  $\text{CH}_2^+$ . In the light of these considerations alone, it would be hard to understand the occurrence of a long-lived collision complex in the  $\text{C}^+(\text{H}_2, \text{H})\text{CH}^+$  reaction at low kinetic energies.

A significant new feature appears when scalene triangular geometries for  $\text{CH}_2^+$  are considered. In the  $C_s$  point group, the  ${}^2B_1$  state of  $\text{CH}_2^+$  becomes  ${}^2A''$ , but both the  ${}^2A_1$  and  ${}^2B_2$  states become  ${}^2A'$ . Thus the crossing of the  ${}^2B_2$  and  ${}^2A_1$  surfaces, allowed in  $C_{2v}$  symmetry, becomes avoided in  $C_s$  symmetry. Accordingly, reactants on the  ${}^2B_2 - {}^2A'$  surface can pass adiabatically by a low energy path to the  ${}^2A' - {}^2A_1$  ground state of  $\text{CH}_2^+$ . The deep potential energy well of  $\text{CH}_2^+$  is therefore accessible to the reactants even when they have low kinetic energy, and the occurrence of a long-lived collision complex in the  $\text{C}^+(\text{H}_2, \text{H})\text{CH}^+$  reaction is quite reasonable.

There is a rather pictorial description of the mixing of the  ${}^2A_1$  and  ${}^2B_2$  states that occurs when the  $\text{C}^+-\text{H}_2$  collision complex is distorted from  $C_{2v}$  to  $C_s$  geometry. The carbon 2p orbitals which lie in the (y-z) plane defined by the nuclei together produce in the isolated atomic ion an electron distribution which is cylindrically symmetric, and appears to be a doughnut in the y-z plane. As  $\text{C}^+$  approaches  $\text{H}_2$ , the doughnut-like electron distribution is distorted. If the nuclear conformation has  $C_{2v}$  symmetry, these in-plane orbitals can be resolved and expressed in terms of either the conventional  $2p_z(a_1)$  or  $2p_y(b_2)$  orbitals of carbon, which point respectively perpendicular and parallel to the axis of  $\text{H}_2$ . Occupation of one of these orbitals produces either the  ${}^2A_1$  or  ${}^2B_2$  surface. When the nuclear conformation departs from  $C_{2v}$  symmetry, both these surfaces contain contributions from what have been designated as the  $a_1$  and  $b_2$  2p orbitals,

both mixed with the  $\sigma$ -a' orbital of  $H_2$ . The lower of the two  ${}^2A'$  surfaces generated in this manner has the lowest lying a' orbitals occupied to the maximum extent possible. Its close relation to the  ${}^2A_1$  ground state of  $CH_2^+$  is displayed by writing the electron configuration as  $(2a_1-2a')^2 (1b_2-3a')^2 (3a_1-4a')^1$ , which shows the association of the orbitals in  $C_{2v}$  and  $C_s$  symmetry explicitly. In brief: on the lowest energy surface available to  $C^+ + H_2$ , all electrons have a' symmetry for  $C_s$  nuclear conformations. This surface evolves adiabatically to the ground state of  $CH_2^+$ , which has its minimum energy in the symmetric  $C_{2v}$  conformation, and is then designated  ${}^2A_1$ .

The foregoing arguments were constructed in anticipation and as rationalizations of the experimental results<sup>14</sup> on the reactive scattering of  $C^+$  by  $H_2$ . Subsequent to the completion of that work and construction of the arguments based on the correlation diagram, Liskow, Bender, and Schaefer<sup>16</sup> performed extensive SCF-CI calculations to determine certain profiles of the  $C^+ - H_2$  potential energy surfaces. Some of the results of these calculations are shown in Figure 5. The barrier on the  ${}^2A_1$  surface between reactants and the complex anticipated from the correlation diagram is nicely confirmed by the calculations. The barrier on the  ${}^2B_1$  surface, which was not anticipated in the original work, is also clearly demonstrated. As required by experiment, the calculations show that for  $C_s$  geometries, the  ${}^2B_2$  and  ${}^2A_1$  states interact to form a  ${}^2A'$  surface which provides a low energy path from reactants to the



$\text{CH}_2^+ (^2A_1)$  potential energy well. Thus the major qualitative features of the potential surface deduced from experiments, correlation diagrams, and the asymptotic properties of reactant, product, and intermediate states are substantially confirmed by the ab initio calculation.

Can the lessons learned from this analysis be profitably applied elsewhere? The reaction of  $\text{N}^+ (^3P)$  with  $\text{H}_2$  to give  $\text{NH}_2^+$  provides a test. This reaction is thermoneutral or slightly exoergic, and it is known that the potential energy of symmetric  $\text{NH}_2^+$  in its  $^3B_1$  ground state lies approximately 6 eV below that of reactants and products. If this state is accessible to low energy reactants, a long-lived collision complex should be observed.

Construction of the correlation diagram for the  $\text{N}^+ - \text{H}_2$  reaction is greatly aided by our knowledge of the  $\text{C}^+ - \text{H}_2$  system. As  $\text{N}^+ (^3P)$  approaches  $\text{H}_2$  in the  $\text{C}_{2v}$  conformation,  $^3B_1$ ,  $^3B_2$ , and  $^3A_2$  states are generated. The  $^3B_1$  state is very closely related to the  $^2A_1$  state of the  $\text{C}^+ - \text{H}_2$  system, since it can be generated by adding a non-bonding  $1b_1$  electron to the  $\text{C}^+ - \text{H}_2$  system and appropriately increasing the nuclear charge. The  $^3B_1$  surface should therefore be repulsive at large  $\text{N}^+ - \text{H}_2$  distances, and then drop abruptly to the deep ground state potential energy well of symmetrical  $\text{NH}_2^+$ . This behavior is indicated in Figure 6.

Similarly, the  $^3B_2$  state of  $\text{N}^+ - \text{H}_2$  is generated from the repulsive  $^2A_1$  state of  $\text{C}^+ - \text{H}_2$  by addition of a bonding  $1b_2$  electron. This surface should also be repulsive, but not as

much as the  ${}^3B_1$  state. As Figure 6 shows, the  ${}^3B_2$  surface does not provide access to the deep potential wells of  $NH_2^+$ . The  ${}^3A_2$  state of  $N^+-H_2$  can be obtained from the  ${}^2B_2$  state of  $C^+-H_2$ , again by addition of a non-bonding  $1b_1$  electron. Accordingly, the  ${}^3A_2$  surface of the  $N^+-H_2$  system is indicated to be rather flat in Figure 6.

The surfaces appropriate for strict  $C_{2v}$  symmetry offer no access to the deep potential energy well of  $NH_2^+$  by a low energy path. However, in  $C_s$  symmetry, the  ${}^3A_2$  and  ${}^3B_1$  surfaces both assume  ${}^3A''$  character. Thus the crossing of the  ${}^3A_2$  and  ${}^3B_1$  surfaces indicated in Figure 6 is actually avoided when the system is distorted from  $C_{2v}$  symmetry. The reactants do in fact have a low energy adiabatic path to the potential energy well, and from there to the products  $NH^+ + H$ . Do the experimental measurements of the product distribution indicate the occurrence of a long-lived collision complex, as might be expected from this analysis? The velocity vector distributions of  $NH^+$  measured at reactant energies of 2 eV and above<sup>17</sup> indicate that the reaction proceeds by a direct, short-lived interaction, not a long-lived complex. Only when the initial relative energy is reduced to the vicinity of 1 eV and below<sup>18</sup> does any indication of a long-lived complex appear.

This result is superficially surprising, since the  $NH_2^+$  well is very deep, and might be expected to produce a long-lived collision complex at energies as high as 3 eV. However, an explanation is available. In order to reach the  ${}^3B_1 - {}^3A''$  well of  $NH_2^+$ , the reactants on the lower  ${}^3A''$  surface must travel adiabatically past the region where the  ${}^3B_1$  and  ${}^3A_2$  surfaces

intersect. In the limit of vanishing relative velocity, systems on the lower  ${}^3A''$  surface will remain on this surface as it evolves to the  ${}^3B_1 - {}^3A''$  ground state well of  $NH_2^+$ . However, when the relative velocity is finite, the system may exhibit adiabatic behavior, and make a transition to the upper  ${}^3A''$  surface. At small  $N^+ - H_2$  separations this surface is related to the excited  ${}^3A_2$  state of  $NH_2^+$ , which has no deep potential energy well. As Figure 6 shows, the system can then evolve to the products  $NH^+ + H$ .

Motion on the diabatic  ${}^3A'' - {}^3A_2$  surface should lead to product formation by a direct, short-lived interaction. The velocity vector distribution of  $NH^+$  should therefore be asymmetric at the higher relative energies as is observed experimentally. At low relative energies, the system should behave adiabatically, and remain on the lower  ${}^3A'' - {}^3B_1$  surface, visit the deep potential energy well, and form product by a long-lived interaction. This adiabatic behavior will be more pronounced, the greater is the splitting between the upper and lower  ${}^3A''$  surfaces, and the smaller is the initial relative translational energy. Thus it is the separation between the two  ${}^3A''$  surfaces, rather than the depth of the potential energy well which controls formation of a long-lived complex in this system.

The foregoing argument provides an adequate explanation of the behavior of the  $N^+ - H_2$  system. However, its validity probably can only be established by an elaborate calculation of the exact potential energy surfaces in the region of the

${}^3A_2 - {}^3B_1$  intersection. While this example illustrates the limitations of the predictions that can be made from asymptotic state properties and correlation diagrams, it also shows how these techniques can lead to recognition of at least the possibility of certain types of otherwise unexpected behavior.

In his paper which first outlined the electronic state correlation technique, Schuler<sup>3</sup> emphasized the importance of correlating reactants and products through the least symmetric intermediate complex. The formation of long-lived intermediates in the  $C^+-H_2$  and  $N^+-H_2$  systems illustrates this point, since it is only the scalene triangular conformations that provide low energy adiabatic paths from reactants to the deep potential well of the intermediate. On the other hand, it is important to realize that if only  $C_s$  conformations had been considered, an important aspect of the  $N^+-H_2$  problem would have been overlooked. In highly symmetric nuclear conformations, the electronic states each belong to one of a variety of symmetry species. Some of these states of different symmetry may cross each other. In less symmetric conformations, fewer symmetry species are available, and some of the states which crossed in symmetric conformations may belong to the same symmetry species of the less symmetric conformation. Thus their crossings become avoided as the intermediate is distorted. However, because these states cross in symmetric conformations, they will approach each other closely even in less symmetric conformations, thereby facilitating diabatic behavior, or electronic transitions in the course of reaction. In short,

making correlation diagrams for symmetric conformations is a good way to discover the possibility of diabatic behavior.

A brief comment on the dynamics of the  $O^+(H_2,II)OH^+$  reaction<sup>19,20</sup> is in order because of its obviously close relation to the  $C^+-H_2$  and  $N^+-H_2$  systems. The  ${}^2B_1$  ground state of  $H_2O^+$  lies approximately 5 eV below the ground states of the reactants and products, and again one can entertain the possibility that a long-lived complex reaction mechanism will be followed. However, the correlation diagram<sup>17</sup> for this system shows that the reactants  $O^+({}^4S) + H_2({}^1\Sigma_g^+)$  move initially on a  ${}^4A_2 - {}^4A'' - {}^4\Sigma$  surface which leads adiabatically to products, and has no deep wells that can produce a long-lived collision complex. This  ${}^4A_2$  surface does cross the  ${}^2B_1$  surface that leads to the ground state of  $H_2O^+$ . Because electron spin is for the most part conserved in systems of low atomic number, systems on the  ${}^4A_2 - {}^4A'' - {}^4\Sigma$  surface can not reach the deep  ${}^2B_1$  well with any appreciable probability. Consequently, it is not surprising that this reaction proceeds by a direct, short-lived interaction mechanism even at the lowest energies.<sup>21</sup>

### Summary

Analysis of the behavior of a number of triatomic ion-molecule systems shows that certain properties of the reactants, products and intermediates can be used to deduce the feasibility, course, and detailed dynamic mechanism of a reaction. Correlation of the electronic states of reactants and products is important, for in the course of a reaction, the principal

contributing electronic orbital configuration may change substantially. In constructing these correlation diagrams, information from one system may be used to ascertain the qualitative nature of some of the potential energy surfaces in other systems. Surface crossings and avoided crossings are important. To ascertain the existence of these features, it is necessary not only to correlate states corresponding to highly symmetric nuclear conformations, but to consider the effects of distortions to lower symmetry. In addition, the effect of variation of coordinates other than what appears to be the reaction coordinate should be assessed. Both adiabatic and diabatic behavior near avoided crossings is possible and has been observed to affect the reaction dynamics profoundly. It is to be hoped that there will be continued development of methods which permit facile quantitative assessment of the nature of avoided crossings of potential energy surfaces without extensive computational effort.

Acknowledgement: This work was carried out during my tenure as a Professor in the Miller Institute for Basic Research at the University of California, Berkeley. The experimental work on ion-molecule reactions in my laboratory is supported by the U. S. Atomic Energy Commission.

## REFERENCES

- (1) T. Carrington, Acc. Chem. Res., 7, 20 (1974).
- (2) R. B. Woodward and R. Hoffmann, "The Conservation of Orbital Symmetry", Academic Press, New York, 1970.
- (3) K. E. Shuler, J. Chem. Phys., 21, 624 (1953).
- (4) R. J. Donovan and D. Husain, Chem. Rev., 70, 489 (1970).
- (5) W. A. Chupka and M. E. Russell, J. Chem. Phys., 49, 5426 (1968).
- (6) W. A. Chupka, J. Berkowitz, and M. E. Russell, Proc. of the 6<sup>th</sup> Int. Conf. on the Physics of Electronic and Atomic Collisions, Cambridge, Mass., 1969, p 71. The MIT Press, Cambridge, Massachusetts.
- (7) R. Johnsen and M. Biondi, J. Chem. Phys., 61, 2112 (1974).
- (8) P. J. Kuntz and A. C. Roach, Trans. Faraday Soc. II, 68, 259 (1972).
- (9) P. J. Brown and E. F. Hayes, J. Chem. Phys., 53, 922 (1971).
- (10) H. H. Michels, J. Chem. Phys., 44, 3834 (1966).
- (11) L. Friedman and T. F. Moran, J. Chem. Phys., 42, 2624 (1965).
- (12) B. H. Mahan and J. S. Winn, J. Chem. Phys., 57, 4321 (1972).
- (13) S. Chapman and R. Preston, J. Chem. Phys., 60, 650 (1974).
- (14) B. H. Mahan and T. M. Sloane, J. Chem. Phys., 59, 5661 (1973).

- (15) B. H. Mahan, Accounts Chem. Res., 3, 393 (1970).
- (16) D. H. Liskow, C. F. Bender, and H. F. Schaefer III, J. Chem. Phys., 61, 2507 (1974).
- (17) E. A. Gislason, B. H. Mahan, C. W. Tsao, and A. S. Werner, J. Chem. Phys., 54, 3897 (1971).
- (18) J. A. Fair and B. H. Mahan, J. Chem. Phys., in press.
- (19) K. T. Gillen, B. H. Mahan, and J. S. Winn, J. Chem. Phys., 58, 5373 (1973).
- (20) K. T. Gillen, B. H. Mahan, and J. S. Winn, J. Chem. Phys., 59, 6380 (1973).
- (21) H. H. Harris and J. J. Leventhal, J. Chem. Phys., 58, 2333 (1973).



Figure 1. Potential energy curves for the diatoms in the asymptotic reactant and product regions of the  $(\text{He-H}_2)^+$  system. Note the crossing of the curves for  $\text{H}_2(1\Sigma_g^+)$  and  $\text{H}_2^+(2\Sigma_u^+)$  which occurs in the reactant region of  $\text{He}^+ + \text{H}_2$ , but which becomes avoided when all three atoms are close to each other.


Figure 2. Potential energy curves for the diatomic reactants and products of the  $(\text{Ar} - \text{H}_2)^+$  system. Note that the crossing of the curves for  $\text{H}_2(1\Sigma_g^+)$  and  $\text{H}_2^+(2\Sigma_g^+)$  occurs very near the minimum energy of the former state. This crossing becomes avoided when all three atoms are close to each other, and permits formation of  $\text{ArH}^+$  from  $\text{H}_2 + \text{Ar}^+$ , as well as from  $\text{H}_2^+ + \text{Ar}$ .

Figure 3. A molecular orbital correlation diagram for the approach of a second row atom or ion to hydrogen. At the left, the correlations are made assuming  $C_{2v}$  conformations, whereas at the far right, a collinear approach is assumed. The mixing of  $a_1$  and  $b_2$  orbitals which occurs when the system is distorted to  $C_s$  symmetry is indicated by dashed lines.

Figure 4. A correlation diagram linking the electronic states of reactants, intermediates, and products of the  $(\text{C-H}_2)^+$  system. On the left of the diagram, states which arise for isosceles triangular conformations ( $C_{2v}$ ) are indicated, which on the right, correlations for collinear unsymmetrical conformations are shown. The crossing of the  $2B_2$  and  $2A_1$  states which occurs for strict  $C_{2v}$  symmetry is avoided in  $C_s$  symmetry, as is indicated by the dashed curves.


Figure 5. Calculations by Liskow, Bender, and Schaefer of the minimum energies on the lowest  ${}^2A_1$ ,  ${}^2B_1$ , and  ${}^2B_2$  states of  $CH_2^+$ . The coordinate  $R$  is the distance between the center-of-mass of  $H_2$  and the carbon nucleus. Note the barriers between reactants and the  $CH_2^+$  complex on the  ${}^2A_1$  and  ${}^2B_2$  surfaces. The crossing of the  ${}^2A_1$  and  ${}^2B_2$  surfaces shown here becomes avoided when the system is distorted to  $C_s$  symmetry.

Figure 6. A correlation diagram linking the relevant electronic states of reactants, products, and intermediates of the  $(N-H_2)^+$  system. The arrangement of the diagram is as in Fig. 4, with  $C_{2v}$  correlations on the left, and  $C_{\infty v}$  correlations on the right. Note the close resemblance to the correlation diagram for the  $(C-H_2)^+$  system, particularly the crossing of the  ${}^3A_2$  and  ${}^3B_1$  states, which is avoided in  $C_s$  symmetry.



XBL 748-6982

Fig. 1



XBL 748-6983

Fig. 2




Fig. 3

XBL 7411-7615





Fig. 4

XBL 7411-7616

00004206959



XBL 7310-4163

Figure 5





Fig. 6

XBL 7410-7487

00004206960





LEGAL NOTICE

*This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.*

TECHNICAL INFORMATION DIVISION  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720