

UCLA

UCLA Previously Published Works

Title

Subjective Response to Alcohol as a Research Domain Criterion.

Permalink

<https://escholarship.org/uc/item/6r2158df>

Journal

Alcoholism, clinical and experimental research, 40(1)

ISSN

1530-0277

Authors

Ray, Lara A
Bujarski, Spencer
Roche, Daniel J O

Publication Date

2016

Peer reviewed

Critical Review

Subjective Response to Alcohol as a Research Domain Criterion

Lara A. Ray, Spencer Bujarski, and Daniel J. O. Roche

Background: Individual differences in the subjective experience of the pharmacological effects of alcohol have long been implicated in the likelihood that one will drink heavily and develop alcoholism. The theme of this conceptual review and perspective article is to synthesize the literature on subjective responses to alcohol and to set an agenda for the next generation of research in the area. Specifically, we contend that in order for subjective response to alcohol to play a prominent role in alcoholism research, it is critical that it be studied as a multimodal phenotype.

Methods: First, we review the human research on subjective response to alcohol measured under controlled laboratory conditions and draw recommendations for the application of these findings to understanding alcoholism neurobiology in humans. Second, we highlight multimodal approaches, including studies of the genetic and neural substrates of individual differences in subjective response to alcohol. Third, we review treatment implications with a focus on subjective response to alcohol as an intervention target. Upon review of the research on subjective response to alcohol across levels of analyses, we provide recommendations for leveraging these phenotypes in a systematic and methodologically rigorous fashion that can address central questions about alcoholism etiology, disease progression, and personalized treatment.

Discussion: The approach recommended herein is largely consistent with the Research Domain Criteria (RDoC) initiative across the National Institute of Mental Health. The defining feature of such domains is that they inform behavior yet be amenable to examination through multiple units of analysis, such as molecular, genetic, circuit-level, and behavioral measurements. To that end, we contend that subjective response to alcohol represents a behaviorally and biologically plausible phenotype upon which to build using the RDoC framework for understanding alcohol use disorder.

Key Words: Subjective Response, Alcohol, Translational Phenotype, Research Domain Criteria.

THROUGH ITS PHARMACODYNAMIC interaction with a multitude of neurotransmitter systems, consumption of alcohol alters the subjective state of the consumer, termed, subjective response. In particular, alcohol is known to have a variety of effects on subjective states including mood, alertness, and mental clarity. Individual differences in subjective responses to alcohol have long been implicated in the likelihood that one will drink heavily and experience problems related to alcohol, including alcoholism. While the notion that individuals who experience more positive, or reinforcing, subjective effects from drinking alcohol are more likely to drink and those who

experience more negative effects are less likely to drink has a great deal of intuitive appeal, operationalizing and systematically testing these effects remains challenging. Multiple conceptualizations of subjective response have been proposed in the literature (reviewed in more detail below and in Morean and Corbin 2010; Quinn and Fromme, 2011). For this review, we are guided by factor analytic work by our research group (Ray et al., 2009) which, in an attempt to promote consilience between different alcohol challenge “camps,” identified 3 factors of subjective response as measured by validated and reliable self-report assessments: (i) stimulation and other pleasant effects; (ii) sedation and unpleasant effects; and (iii) alleviation of tension and negative mood. This factor structure was recently replicated in a larger independent sample (Bujarski et al., 2015b), though with the addition of a fourth “craving” dimension. Although alcohol craving is clearly important for alcoholism risk and treatment development, a detailed review of the vast craving literature is beyond the scope of this present review. Furthermore, while alcohol administration is also known to affect more objective indices (e.g., static ataxia, acetaldehyde levels, fine motor task performance), this review focuses on the effects of alcohol that

From the Department of Psychology (LAR, SB, DJOR), University of California, Los Angeles, Los Angeles, California.

Received for publication December 23, 2014; accepted October 12, 2015.

Reprint requests: Lara A. Ray, PhD, Psychology Department, University of California, Los Angeles, 1285 Franz Hall, Box 951563, Los Angeles, CA 90095-1563; Tel.: 310-794-5383; Fax: 310-206-5895; E-mail: lararay@psych.ucla.edu

Copyright © 2016 by the Research Society on Alcoholism.

DOI: 10.1111/acer.12927

are experienced at the level of individual subjectivity, and that are assessed through validated self-report scales.

The approach to studying subjective response to alcohol discussed herein is highly consistent with the Research Domain Criteria (RDoC) initiative across the National Institute of Mental Health (Cuthbert and Insel, 2013; Insel, 2014; Insel et al., 2010). According to the RDoC proposal, scientists are asked to identify key domains of function that can effectively describe the phenomenology of a disorder and offer generative insights into treatment. The defining feature of such domains is that they inform behavior yet are amenable to examination through multiple units of analysis, such as molecular, genetic, circuit-level, and behavioral measurements. Recently, compelling arguments for an Alcohol Addiction RDoC have been raised (Litten et al., 2015). In this paper, we review multimodal research on subjective response to alcohol, thereby strengthening the argument for subjective response as a domain criterion in the RDoC approach to alcohol use disorder.

Although many outcome variables can be measured in these alcohol challenge studies with varying levels of objectivity/subjectivity, subjective responses to alcohol consumption, as measured through validated self-report scales, have emerged as primary predictive factors in multiple prominent etiological models of alcoholism risk (King et al., 2014; Schuckit, 1984; Schuckit and Smith, 1996; Schuckit et al., 1988). Similarly, while alcohol craving represents an important etiological construct that is often measured in alcohol challenge studies (e.g., Bujarski and Ray, 2014; King et al., 2011, 2014; Ray et al., 2013), a detailed review of alcohol craving as an RDoC is deserving of its own critical review. As such, the theme of this review and perspective article is to describe what we have learned thus far about subjective responses to alcohol and to set an agenda for the next generation of research in the area. Specifically, we contend that in order for subjective response to alcohol to play a prominent role in alcoholism research, it is critical that it be studied as a multimodal phenotype.

Multimodal phenotypes, by definition, are those that can be examined across methods and levels of analyses, thereby informing etiological and treatment models of a disorder. To that end, applying findings to subclinical as well as clinical populations requires careful consideration of the samples under study and where they may fit within the progression of the disorder, from initial drinking, to heavy drinking, to early and later stages of alcohol dependence. This review intends to consolidate major findings in this line of research as well as to identify opportunities to enhance the value of subjective response phenotypes that in turn can be carried forward into the era of transdisciplinary addiction science. We begin by reviewing the human research on subjective response to alcohol measured under controlled laboratory conditions and draw recommendations for future studies in the field. Second, we highlight multimodal approaches, including studies of the genetic and neural substrates of individual differences in subjective response to alcohol. Third, we review treatment

implications with a focus on subjective response to alcohol as an intervention target. Finally, we provide a set of conclusions and recommendations to extend upon the rich history of research on subjective response to alcohol and move this line of inquiry into the RDoC era of psychiatric research.

HUMAN LABORATORY MODELS OF SUBJECTIVE RESPONSE TO ALCOHOL

Alcohol administration studies have documented the substantial variability in individuals' subjective responses to alcohol and have shown that such differences impact the predisposition to alcohol use and misuse (King et al., 2014; Schuckit, 1984; Schuckit and Smith, 1996). Schuckit and colleagues pioneered this line of research by assessing subjective response to alcohol during controlled oral alcohol administration in the laboratory (i.e., oral alcohol challenge) (Schuckit, 1984). Although Schuckit's body of research examined multiple indices of alcohol response (e.g., static ataxia and electroencephalography), an early multivariate analysis identified subjective responses to alcohol as measured by the Subjective High Assessment Scale (SHAS; Judd et al., 1977; Schuckit, 1984) as the single strongest predictor of alcoholism risk (Schuckit and Gold, 1988). In terms of alcoholism risk, Schuckit and colleagues demonstrated that a blunted response to the subjective effects of alcohol, measured by the SHAS, represents a risk factor for future alcohol misuse and dependence (Schuckit, 1984, 1994; Schuckit and Smith, 1996). Termed the low level of response model, this theory examined response to alcohol as a unidimensional construct with a particular focus on the more sedative or other unpleasant components of subjective response as evidenced by factor analytic work by Schuckit and Gold (1988). Our group has corroborated that the SHAS is most sensitive to the sedative/unpleasant effects of alcohol (Bujarski et al., 2015b; Ray et al., 2009), although it should be noted that the factor analytic work by our group has utilized intravenous, as opposed to oral, alcohol administration.

Newlin and Thomson (1990) provided a careful review of the alcohol challenge literature available at the time focusing on studies of family history of alcoholism as a risk factor for the disorder. They proposed the influential differentiator model for understanding psychobiological responses to alcohol as a function of family history. The differentiator model proposed that among individuals with a family history of alcoholism, responses to alcohol may be accentuated during the rising blood alcohol curve (BAC) (i.e., acute sensitization) and attenuated during the falling BAC (i.e., acute tolerance). To that end, sons of alcohol-dependent individuals may be both more sensitive to the rewarding and psychomotor stimulating (e.g., increased heart rate) effects of alcohol during the rising limb of the oral alcohol BAC and less sensitive to the unpleasant effects of alcohol when BAC is declining (Newlin and Thomson, 1990). While in broad agreement with the low level of response model in terms of sedative effects, the differentiator model argues for limb of BAC to be

considered along with various dimensions of subjective response.

Following upon the differentiator model, recent studies have challenged the notion of subjective responses to alcohol as a unitary construct. Instead, the pharmacological and behavioral effects of alcohol may be best described using a biphasic model. Specifically, when blood alcohol levels are rising (i.e., the ascending limb of intoxication), alcohol produces robust stimulatory and other pleasurable subjective effects, whereas when blood alcohol levels are declining (i.e., the descending limb of intoxication), alcohol's effects are largely sedative and unpleasant (Earleywine, 1994; Erblich et al., 2003; Martin et al., 1993). This conceptualization of the effects of alcohol argues for the construct of subjective response to be parsed into stimulant and sedative dimensions leading to the development of the Biphasic Alcohol Effects Scale (BAES) (Martin et al., 1993).

Alcohol administration studies have since routinely employed the BAES, including a briefer version which has been recently developed and validated (Rueger and King, 2012; Rueger et al., 2009). However, it was not until recently that a large longitudinal study established the predictive utility of alcohol-induced stimulation and sedation as measured with the BAES during a controlled and blinded oral alcohol challenge with regard to the propensity for heavy drinking and alcohol problems (King et al., 2011, 2014). Specifically, this landmark study demonstrated that individuals who engaged in regular, binge-drinking behavior experienced greater stimulant/pleasant effects during the rising BAC limb than light drinkers, and this heightened alcohol stimulation and reward sensitivity robustly predicted more alcohol use disorder symptoms over time and was associated with greater binge-drinking frequency at 2- (King et al., 2011) and 6-year follow-ups (King et al., 2014). Additionally, heavy drinkers reported lower sensitivity to the sedative effects of alcohol across the duration of BAC, which in turn was also associated with a higher number of alcohol use disorder symptoms at 2- and 6-year follow-ups (King et al., 2014). In sum, this study demonstrated that drinking history is highly related to acute subjective response to alcohol and, beyond the risk associated with a low response to the sedative effects of alcohol as described in the low-response model, enhanced sensitivity to the stimulant subjective effects of alcohol predicts heavier alcohol use and problems at follow-up. This recognition is consistent with a meta-analysis compiling decades of alcohol administration studies and reporting considerable support for the low level of response to alcohol in the context of familial risk as well as support for the differentiator model with regard to its association with drinking patterns (Quinn and Fromme, 2011).

Another important model of subjective response and alcoholism risk is the tension-reduction model which is based on the notion that individuals drink alcohol because of its ability to reduce tension. This theory was initially influenced by the drive reduction theory of the 1940s (Hull, 1943), which emphasized motivational aspects underlying drinking.

Although there is significant intuitive appeal to the notion that individuals drink to reduce tension, an early review by Cappell and Herman (1972) suggested that empirical support for the theory was limited. A related theory was developed in the 1980s, which focused on the stress-response dampening (SRD) effects of alcohol (Levenson et al., 1980; Sher & Levenson, 1982). The SRD model focused on refining the operational definition and laboratory manipulation of stressors (e.g., electric shock, public speaking task) and examining individual differences in the SRD effects of alcohol, making it a more focused and testable theory (Greeley & Oei, 1999). As stated by Greeley and Oei (1999), alcohol, at certain dosages, is capable of reducing some signs of tension in some humans, under certain contextual conditions.

More recent work on the tension-reduction model and related theories emphasize the pharmacological and neurobiological mechanisms by which alcohol may dampen a stress-response as well as the individual and contextual differences that may moderate those effects. Examples of such moderators include hostility (Zeichner et al., 1995), anxiety sensitivity (Stewart et al., 1997), gender (Sinha et al., 1998), type of social situation (Armeli et al., 2003), or life stressor (Hart and Faza, 2004). More recently, anxiety and negative urgency were found to operate through coping motives leading to increased risk of alcohol use disorders (Menary et al., 2015). Further, genetic predispositions to stress reactivity, via variations in hypothalamic-pituitary-adrenal axis genes, have been tested as possible mediators of the effects of stress on alcohol consumption (Clarke et al., 2007). Thus, alcohol-induced stress reduction may be especially salient for those individuals with genetically determined heightened stress reactivity. In short, the tension-reduction theory and its most widely researched offshoot, the SRD model, identify alcohol's ability to reduce tension and stress reactivity as central to the motivation to drink and the development of alcohol-related problems.

In summary, subjective response to alcohol represents a multifaceted (Bujarski et al., 2015b; Ray et al., 2009) and replicable construct (King et al., 2015; Roche et al., 2014). Regarding the ability of subjective response to alcohol, measured under controlled laboratory conditions, to predict one's risk to develop alcohol-related problems elegant longitudinal studies have suggested 2 distinct and relatively independent, pathways of risk. The first pathway suggests lower levels of response to the sedative and unpleasant effects of alcohol, and the second suggests higher sensitivity to the stimulant and pleasant effects of alcohol. However, subjective response to alcohol is highly variable between individuals and is influenced by numerous factors, including participant characteristics (e.g., family history of alcoholism, drinking history, recent alcohol exposure) and methodology (dose, measures, timing of measures, route of administration, etc.). As noted by King and colleagues (2011), careful consideration of factors such as the specific response being measured, the dose and rate of alcohol administered, BAC level and the limb of BAC, and the risk factors under examination

should be carefully described so that greater consilience can be reached in the literature.

We would add to this set of recommendations that sample characteristics and additional potential methodological factors, such as route and total duration of alcohol administration be carefully considered as well. For example, when predicting future risk for alcoholism or describing subjective response as a risk factor and treatment target, it is critical to consider that social drinkers that are past the peak age of risk for alcoholism (Grant et al., 2012) are unlikely to adequately inform the literature, ultimately aimed at high risk and/or affected individuals. Additionally, it is currently unclear whether subjective response reliably differs between oral and intravenous alcohol administration, as only a single study has systematically examined differences in alcohol responses between these 2 methods (Ray et al., 2007) and reported greater levels of intoxication, but lower alcohol craving, from intravenous alcohol (possibly due to the absence of alcohol cues). However, recent developments in computerized intravenous alcohol administration paradigms allow for the mimicking of oral BACs (Ramchandani et al., 2009), thus permitting rigorous comparison of subjective response variables across these routes of administration while controlling for BAC-related confounds. It is also worth noting that comparison of drinking groups with different alcohol use histories requires careful consideration of tolerance effects. Considering the limitations noted above, pharmacokinetic tolerance can be controlled through utilization of experimental paradigms that carefully control blood alcohol dose, such as intravenous alcohol administration (Bujarski and Ray, 2014; Ray and Hutchison, 2004; Ray et al., 2013; Zimmermann et al., 2008, 2009, 2013). Pharmacodynamic tolerance, however, is more difficult to account for. Examination of multiple different alcohol responses consistent with the multidimensionality of subjective response (Bujarski et al., 2015b; Martin et al., 1993; Ray et al., 2009) allows researchers to test whether responses to alcohol are blunted across dimensions of alcohol response, consistent with a generalized tolerance syndrome, or whether only specific dimensions of alcohol's effects are altered by drinking status. Recent work by our group has found replicable differences between alcohol dependents and nondependent heavy drinkers in the relationship between stimulatory responses to alcohol and alcohol craving (Bujarski and Ray, 2014; Bujarski et al., 2015a). Although this line of inquiry is early in development, these differences in the coupling of subjective response and craving are potentially less sensitive to tolerance effects than the raw magnitude of subjective responses themselves.

Taken together, decades of human laboratory studies of individual differences in the subjective responses to alcohol have advanced 2 translational phenotypes, namely subjective response to the (i) stimulant/pleasant and (ii) sedative/unpleasant effects of alcohol. A tension-reduction factor has emerged in factor analytic work (Bujarski et al., 2015b; Ray et al., 2009) and has been recently examined by our group (Bujarski and Ray, 2014) in relation to the proposed transi-

tion from positive to negative reinforcement that is thought to occur during the progression of alcoholism (Koob and Le Moal, 2008b). However, the proper evaluation of this potentially useful phenotype may require testing alcohol self-administration in samples with higher levels of alcoholism severity than those typically enrolled in alcohol administration studies, if in fact this construct can capture negative reinforcement as a maintenance, as opposed to a predisposing, factor for alcoholism. Further, longitudinal studies that can repeatedly measure subjective responses to alcohol, document patterns of change over time, and associate such changes with dynamic changes in alcohol use will ultimately advance these multidimensional phenotypes that can identify risk and also track the progression of alcoholism from the initiation of heavy drinking to its more severe chronic and relapsing forms.

MULTIMODAL APPROACHES TO STUDYING SUBJECTIVE RESPONSE TO ALCOHOL

Subjective response to alcohol has been studied using a host of experimental methods from diverse fields, including behavioral genetics and neuroimaging. The goal of this section is to highlight genetic and neuroimaging studies of subjective response to alcohol, while extensive reviews of these areas can be found elsewhere (Ray et al., 2010d; Roche and Ray, 2015). Using heritability as a starting point to genetic studies, subjective responses to alcohol, measured in an oral alcohol administration twin study, had a heritability estimate of 60% (Viken et al., 2003). This study used a 22-item measure called the Sensation Scale, which included items such as drowsy, lightheaded, and dizzy. Similar estimates, ranging between 40 and 60%, were obtained in an Australian oral alcohol challenge twin study in which subjective response to alcohol was measured across levels of BAC by a single item, namely "how drunk do you feel now" (Heath and Martin, 1991). A more recent laboratory study of the offspring of fathers who completed an alcohol challenge 20 years earlier revealed a significant positive parent-offspring association for subjective feelings of intoxication and body sway after consuming a standardized oral dose of alcohol among family history positive individuals (Schuckit et al., 2005). Although not providing direct evidence of heritability, this study is consistent with prior reports of the genetic influences on these phenotypes and provides support for its reliability. Notably, these studies examine subjective response from the perspective of the low-response model and are less informative about the genetic bases of the stimulant/rewarding phenotype.

A number of behavioral genetic studies have also been conducted using subjective response to alcohol as the outcome variable. Given the experimental nature of the phenotype itself, such studies typically rely on the candidate gene approach, thus borrowing the strengths and weaknesses of this method. An important line of research has dealt with the genes subserving the metabolism of alcohol, which in turn

impacts subjective response. When alcohol is consumed, its metabolic breakdown is a 3-step hepatic process in which the alcohol is first oxidized into acetaldehyde by the enzyme alcohol dehydrogenase (ADH) and is then further metabolized into acetate, and other byproducts, by the enzyme aldehyde dehydrogenase (ALDH). As both acetaldehyde and acetate appear to have pharmacological properties and behavioral effects (Correa et al., 2003, 2012), the genes responsible for the ADH and ALDH enzymes exert important influences on the subjective effects of alcohol because they determine the relative levels of these metabolites over the course of alcohol metabolism. Indeed, genes underlying the pharmacokinetics of alcohol are among the best characterized in terms of their influence on subjective responses to alcohol and alcoholism risk. The behavioral and subjective consequences of possession of genetic variants that affect ADH enzymatic activity to increase the presence of acetaldehyde are acutely aversive in nature, including flushing, headache, tachycardia, and nausea. Genes involved in this pathway include the ALDH gene (*ALDH2*), which has been widely linked to both alcohol use (Sun et al., 2002) and the development of alcohol use disorders (Luczak et al., 2004), as well as *ADH1B*, which has been linked to reduced pleasant subjective effects in Asian Americans (Cook et al., 2005) and greater unpleasant alcohol effects in Caucasians (Wall et al., 2005). A recent study by Peng and colleagues (2014) found that in a sample of Han Chinese men who ingested a small dose of alcohol (0.3 g/kg of alcohol), *ALDH2*2*, rather than *ADH1B*2*, was a causal variant allele for the accumulation of blood acetaldehyde and the associated facial flushing response. These studies exemplify the application of subjective response models to understanding genetic variance in the pharmacokinetics of alcohol.

Given the large number of neurotransmitter systems potentially involved in alcohol's subjective effects (Spanagel, 2009), numerous studies have also probed genetic variation affecting subjective responses to alcohol via various pharmacodynamics pathways (for recent reviews, see Matsushita and Higuchi, 2014; Roche and Ray, 2015). Several studies using oral alcohol challenge and retrospective assessments have implicated genetic polymorphisms in the gamma-aminobutyric acid (GABA) system, particularly variation in the GABA_A receptor (Pierucci-Lagha et al., 2005; Ray and Hutchison, 2009), as being associated with subjective response to alcohol. Several *GABRA2* variants, which code $\alpha 2$ subunit of the GABA_A receptor, were related to attenuated aversive subjective effects, particularly during the declining BAC limb (Uhart et al., 2013), reduced pleasant subjective effects during the ascending BAC limb (Pierucci-Lagha et al., 2005), and blunted levels of self-reported high and intoxication during intravenous administration (Kareken et al., 2010; Roh et al., 2011). Genetic variation in nicotinic acetylcholine receptors, particularly variants within the *CHRNA5-CHRNA3-CHRN4* gene cluster on chromosome 15 which encode the $\alpha 5$, $\alpha 3$, and $\beta 4$ subunits, have also been associated with an attenuated sedative/unpleasant subjective

response as measured by the SHAS during an oral alcohol administration (Joslyn et al., 2008). Finally, a polymorphism in the serotonin transporter gene (*SLC6A4*) resulting in increased transporter activity is associated with reduced SHAS sedation after acute oral alcohol administration (Hu et al., 2005; Schuckit et al., 1999) and lower retrospective self-reports of intoxication levels during drinking episodes (Hinckers et al., 2006).

While the aforementioned studies have focused primarily on the low level of response phenotype, genetic studies of the stimulant/pleasant effects of alcohol have met with some convergence when testing a single nucleotide polymorphism (SNP) in the mu opioid receptor gene (*OPRM1*), the Asn40Asp SNP (rs17799971) (for a review, see Ray et al., 2012). In particular, a series of studies focusing on the stimulant and pleasant subjective effects of alcohol have shown that, compared to Asn40 homozygotes, Asp40 carriers report greater positive subjective effects from an alcohol infusion in the laboratory (Ray and Hutchison, 2004) and in the natural environment, measured using Ecological Momentary Assessment (EMA) methods (Ray et al., 2010e). EMA data revealed that Asp40 carriers consumed more alcohol per drinking episode and that although craving was positively associated with alcohol use in general among Asp40 carriers, craving was less strongly related to alcohol use as compared to Asn40 homozygotes (Ray et al., 2010e). Furthermore, a variable number of tandem repeats in the dopamine transporter 1 gene (*SLC6A3*) was found to moderate the effects of the Asn40Asp SNP on subjective response to alcohol, with Asp40 carriers who were also homozygous for the 10-repeat allele of *SLC6A3* reporting heightened stimulation, vigor, and positive mood after an alcohol infusion (Ray et al., 2014). An interactive effect between *SLC6A3* and *OPRM1* genotypes has also been reported in behavioral (Anton et al., 2012) and neuroimaging (Schacht et al., 2013) studies of naltrexone response. However, a subset of laboratory studies have not supported a relationship between the Asp40 allele and subjective response. Recent studies have found that non-treatment-seeking alcohol-dependent individuals who were homozygous for Asn40 self-reported more alcohol-induced stimulation than Asp40 carriers in a bar-laboratory setting (Anton et al., 2012b) and *OPRM1* genotype was not related to subjective response to intravenous administration in young, heavy drinkers (Hendershot et al., 2014). Yet, the latter study did report that Asp40 carriers self-administer substantially more alcohol and reach a higher BAC than Asn40 homozygotes. This line of inquiry also has important pharmacogenetic implications regarding the prediction of behavioral and clinical response to naltrexone for alcoholism, as discussed in detail elsewhere (Kranzler and Edenberg, 2010; Ray et al., 2012; Roche and Ray, 2015).

A recent review by Jones and colleagues (2015) provides a comprehensive summary of pharmacogenetics of alcohol's subjective effects, including alcohol metabolizing genes, opioidergic genes, dopaminergic genes, and genetic variation in GABAergic, serotonergic, and neurosteroidergic genes,

which is highly consistent with the genetic studies reviewed above. Importantly, the aforementioned review focuses on pharmacogenetics of treatment of alcoholism and calls for increased methodological rigor (e.g., correction for multiple comparison, control for sample heterogeneity) to promote greater consistency across studies. Increased standardization of methods may be critical for genetic studies of subjective response to alcohol, as variation in sample characteristics, alcohol dose, route of administration, and assessment may contribute to the inconsistent findings in the field. Further, the requirement for larger sample sizes in genetic studies along with the realities of time-intensive alcohol administration procedures calls for multisite projects that can effectively ascertain a large, homogenous, and methodologically matched large sample from which genetic determinants of alcohol subjective response could be more effectively ascertained.

The aforementioned genetic studies are mostly consistent with the hypothesized role of endogenous opioids as mediators of the reinforcing effects of alcohol and suggest that favorable phenotypes to probe for the effect of the Asn40Asp SNP may involve assays of the rewarding subjective effects of alcohol in humans. To that end, a study combining intravenous alcohol administration with positron emission tomography (PET) measuring displacement of a radiolabeled mu opioid receptor agonist, [^{11}C]carfentanil, found that alcohol administration induces endogenous opioid release in the orbitofrontal cortex and nucleus accumbens (NAc) (Mitchell et al., 2012). This study provides a critical demonstration of the involvement of endogenous opioids in the effects of alcohol in humans. Further, this study reported a positive correlation between change in opioid binding in the left NAc and subjective ratings of “best ever” feeling during the alcohol infusion, suggesting that as endogenous ligand release increased in the NAc so did subjective reports of “feeling good” in response to alcohol (Mitchell et al., 2012). A study of intravenous alcohol and PET imaging focused on [^{11}C]raclopride displacement implicated the Asp40 allele of the *OPRM1* gene in greater striatal dopamine response to alcohol, although associations between striatal dopamine levels and subjective response to alcohol were not reported (Ramchandani et al., 2011). Regarding the relationship between neuroimaging and behavioral measures of response to alcohol, our group has recently reported that subjective responses to alcohol in the laboratory, namely craving, high, and the reinforcing properties of alcohol, predict neural response patterns during alcohol cue presentation in the scanner (Courtney and Ray, 2014).

Last, behavioral studies of the subjective effects of alcohol can help to extend hypotheses from preclinical models of alcoholism to human clinical samples. To that end, our group has recently conducted studies exploring the transition from positive to negative reinforcement, posited by the allostatic model of addiction (Koob and Le Moal, 2005, 2008a), using subjective responses to alcohol as markers of the posi-

tive (i.e., stimulant/pleasant effects) and negative (i.e., tension relief) rewarding effects of alcohol in humans. In a study comparing heavy drinkers to alcohol-dependent individuals who received an intravenous alcohol administration, alcohol-induced stimulation was associated with alcohol craving to a significantly greater degree in heavy drinkers, as compared to alcohol-dependent individuals (Bujarski and Ray, 2014). This study provided initial support to hypotheses derived from the allostatic model as individuals with alcohol dependence showed a weaker association between the stimulant effects of alcohol and craving than did heavy drinkers, as would be expected if in fact the stimulant and pleasant effects of alcohol become a weaker determinant of drinking as individuals transition from heavy drinking to alcohol dependence. The diminished salience of the positive/hedonic effects of alcohol has since been replicated and extended in a larger sample that includes social drinkers (Bujarski et al., 2015a).

In sum, a host of studies combining oral or intravenous alcohol administration with genetic, imaging, and pharmacology methods have provided unique insights into the biological bases of subjective responses to alcohol and demonstrate a promising level of consistency between findings in humans and predictions from preclinical studies of the neural and behavioral substrates of alcohol's reinforcing effects. And while the findings themselves are encouraging, these studies make valuable contributions by informing the ways in which methodological tools may be employed/combined to answer transformative questions about the nature of addiction in humans.

TREATMENT APPROACHES TARGETING SUBJECTIVE RESPONSE TO ALCOHOL

Because of its predictive relationship with alcohol consumption and alcoholism etiology, subjective response to alcohol has been identified as a potential therapeutic target in medication development for alcoholism (Ray et al., 2010b). As described earlier, individuals with heightened sensitivity to alcohol's rewarding and stimulatory effects or with decreased sensitivity to its sedative and unpleasant effects are at greater risk for the development of alcoholism (King et al., 2011, 2014b; Schuckit and Smith, 1996). Furthermore, in laboratory and naturalistic studies, alcohol-induced stimulation is positively associated with alcohol preference and consumption, whereas the sedative and unpleasant effects of alcohol are negatively associated with these outcomes (Chutuape and de Wit, 1994; Corbin et al., 2008; DeWit et al., 1989; King et al., 2011; de Wit and Doty, 1994). Accordingly, a medication's ability to acutely reduce the stimulant/pleasant or potentiate the sedative/unpleasant effects of alcohol has become a standard biobehavioral marker of efficacy in early phase human laboratory studies of drug development for alcoholism (Ray et al., 2010b).

Several approved or promising pharmacotherapies for alcoholism may reduce the motivation to drink by “blocking

the buzz” (Heilig et al., 2010), or in other words, attenuating the positively rewarding effects of alcohol. For example, naltrexone (Drobes et al., 2004; King et al., 1997; Ray et al., 2008; Swift et al., 1994), nalmefene (Drobes et al., 2004), and topiramate (Miranda et al., 2008) have all been reported to reduce alcohol-induced stimulation in human laboratory studies, while varenicline was shown to reduce the pleasant effects of alcohol (e.g., liking, wanting more on a visual analogue scale; McKee et al., 2009). Naltrexone appears to be particularly effective in blunting alcohol’s other pleasant effects; in addition to its mitigating effects on stimulation, it also attenuates reports of liking and wanting more as measured on the Drug Effects Questionnaire (Morean et al., 2013), and vigor, and positive mood as measured by the Profile of Mood States (McNair et al., 1971) after alcohol administration either orally or intravenously (King et al., 1997; McCaul et al., 2000; Ray et al., 2008). A medication that blunts the stimulatory/pleasant effects of alcohol, such as naltrexone, has been theorized to produce a reduction in alcohol consumption through several potential biological and behavioral pathways (Heilig et al., 2010; Ray et al., 2010a). Primarily, it is thought that, rather than promoting long-term abstinence, the blunting of stimulation and other pleasurable effects reduces the likelihood that a single drinking episode during abstinence (i.e., a “slip”) would escalate to a heavy drinking event (e.g., if an individual likes a drink less than they anticipated, they may be less likely to continue drinking within that episode). A consistent reduction in heavy drinking episodes would thereby reduce harmful alcohol consumption and potentially prevent a full relapse. In support of this notion, the results of meta-analyses and systematic reviews suggest that naltrexone’s predominant therapeutic mechanism of treating alcoholism is gained through a reduction in heavy drinking as opposed to promoting abstinence (Bouza et al., 2004; Pettinati et al., 2006; Rosner et al., 2008). Interestingly, re-analysis of the COMBINE Study focused on individuals who drank during the trial found that a subgroup of nonabstainers, composed primarily of very regular drinkers, benefited from naltrexone in reducing heavy drinking days (Ray et al., 2010c). In sum, several pharmacotherapies, both U.S. Food and Drug Administration approved and currently under development, blunt the rewarding and stimulatory subjective effects of alcohol and, through this behavioral mechanism, may decrease alcohol consumption by decreasing the frequency of heavy and harmful drinking episodes.

Conversely, other medications may contribute to reducing alcohol consumption by potentiating the sedative and unpleasant subjective effects of alcohol. Varenicline has been shown to enhance dysphoria during an oral alcohol challenge (Childs et al., 2012) and weekly retrospective assessments of sedation (Fucito et al., 2011), whereas naltrexone increased sedation, unpleasant feelings, nausea, fatigue, tension, and confusion after both oral and intravenous alcohol administration (King et al., 1997; McCaul et al., 2000; Ray et al., 2008; Swift et al., 1994). While the

mechanistic role that augmented sedative/unpleasant alcohol responses may play in reducing alcohol consumption is not well understood, it is plausible a medication potentiating such subjective effects could simultaneously reduce positive reinforcement (e.g., produce sedative or unpleasant effects that overpower any stimulant/pleasant effects) and block negative reinforcement (e.g., increase tension). This would, in turn, block the tension reduction that is often sought by drinkers during late stages of alcoholism (Heilig et al., 2010; Koob and Kreek, 2007), while also introducing a punishment aspect (e.g., feel tired and unpleasant after drinking), each of which could theoretically contribute to a decrease in drinking behavior, particularly in heavy drinking episodes. Yet, disulfiram provides a cautionary example of using a medication to treat alcoholism via aversive subjective, as well as physiological, responses to alcohol. By disrupting alcohol metabolism and causing an accumulation of acetaldehyde, disulfiram produces headaches, flushing, nausea, and vomiting after alcohol consumption. These effects were intended to lead patients to associate aversive symptoms with alcohol ingestion and avoid future consumption, but the severity of these adverse reactions, coupled with disulfiram’s inability to reduce alcohol craving or protracted withdrawal symptoms, instead lead to poor compliance rates and a limited use in clinical practice (Johnson, 2008; Peterson, 2007). Thus, while a medication’s efficacy in reducing drinking may occur in part by potentiating alcohol-induced sedation or other unpleasant subjective effects, one must provide a cautious and balanced consideration of how the severity of such subjective responses may affect both medication compliance and drinking outcomes.

Although reducing the pleasurable or increasing the sedative/unpleasant effects of alcohol is commonly used markers of efficacy in the development of alcoholism pharmacotherapies, further refinement in several related areas is still needed to confidently determine whether subjective response is truly a viable target for alcoholism pharmacotherapy. First, future studies must identify whether subjective responses to alcohol in the laboratory are clinically meaningful markers of medication efficacy. To our knowledge, no study has explicitly examined whether a medication’s ability to alter subjective response in the laboratory is directly predictive of pharmacotherapy treatment outcomes in a clinical trial setting. It is imperative that future alcoholism medication development studies employ combined laboratory-longitudinal research approaches, particularly in the context of a clinical trial framework, in order to establish the predictive utility of pharmacological manipulation of laboratory-based subjective response to alcohol as a robust and reliable indicator of medication efficacy in clinical settings.

Second, subjective response to alcohol is a multidimensional construct and a moving target that is observably distinct across the progressive stages of alcoholism (e.g., early vs. late stage addiction; Heilig et al., 2010) and between different populations of drinkers (e.g., light drinkers vs. social drinkers vs. heavy drinkers vs. individuals

with alcoholism; Drobles et al., 2004; King et al., 2011; Ray et al., 2013). Unfortunately, it is currently unclear how this complexity and variability may affect medication efficacy. For example, as noted above, the impact of tolerance is an important consideration that requires careful attention in medication development laboratory studies. Relatedly, several prominent theories of addiction theorize that alcohol's positive reinforcing effects convey the greatest influence on motivation to consume alcohol during the transition from heavy drinking to initial development of alcoholism (e.g., "reward drinking"), whereas late stage alcoholism is characterized primarily by negative reinforcement processes (e.g., "relief drinking"; Heilig et al., 2010; Koob and Kreek, 2007). Therefore, a medication that targets a single dimension of subjective response to alcohol may only be clinically effective during a particular stage of alcoholism or in a certain population of problem drinkers. For example, a medication that primarily targets a reduction in the stimulatory or pleasant effects of alcohol could potentially be most effective in heavy episodic drinkers who are either at risk for or still in the early stages of alcoholism, but less so for individuals in late stage alcoholism. Furthermore, novel medications have generally been screened in nontreatment-seeking and nonalcoholic populations due to concerns of ethics in administering alcohol to individuals with alcoholism, which may limit the generalizability and impact of many laboratory studies that reported pharmacotherapy effects on subjective response to alcohol. While recent human laboratory studies have lent initial credence to the transition of subjective responses across the progressive stages of alcoholism (Bujarski and Ray, 2014; Bujarski et al., 2015a), more research focused on both at-risk and affected populations of drinkers is needed to better characterize how subjective response may vary as a function of drinking behavior and disorder severity, as well as how this progression may relate to medication efficacy.

Finally, while subjective response to alcohol may ultimately be found to be a reliable and clinically meaningful treatment target for some alcoholism pharmacotherapies, it must be noted that a medication can be effective in treating alcoholism without affecting acute subjective response to alcohol. Several pharmacotherapies have demonstrated meaningful reductions in alcohol consumption in clinical trials, yet either showed no effect on subjective response to alcohol in the laboratory (e.g., acamprosate [Brasser et al., 2004] and gabapentin [Bisaga and Evans, 2006]) or provided results that were directionally opposite to those hypothesized (e.g., baclofen, increased stimulation; Leggio et al., 2013). When using human laboratory paradigms to screen novel medications for alcoholism, there are many potential biobehavioral markers of drug efficacy in addition to subjective response to alcohol, including, but not limited to, alcohol cue- or stress-induced craving, protracted or acute alcohol withdrawal, and alcohol self-administration. Therefore, despite subjective response to alcohol representing a promising

target for alcoholism treatment, it is recommended that a comprehensive approach utilizing multiple biobehavioral markers of medication efficacy be employed when screening pharmacotherapies for alcoholism to capture the medication-specific effects that may best inform future clinical trials.

CONCLUSIONS AND FUTURE DIRECTIONS

The goal of this review and perspective article was to synthesize a large body of research on subjective response to alcohol and to advance recommendations for a research agenda in this area. We began by advancing the argument that in order for subjective response to alcohol to play a prominent role in alcoholism research it is critical that it be studied as a multimodal phenotype. This was followed by a review of key findings using subjective response to alcohol in the human laboratory, in studies combining imaging and genetic methods, and ending with a review of treatment studies. During each section, we provided recommendations for future research which we summarize below. Figure 1 is provided for a summary and visual depiction of the utility of subjective response in alcoholism etiology and treatment development.

The human laboratory represents the starting point as subjective response to alcohol is best captured under controlled experimental conditions. An important and relatively recent shift has occurred in how subjective response to alcohol is conceptualized. Whereas in the past, subjective response was measured as a unitary construct, there is now increasing recognition that this is a multidimensional construction with distinct etiological and treatment implications. Three phenotypes are advanced for future research including the stimulant and other pleasant effects of alcohol (capturing positive reward), the sedative and unpleasant effects of alcohol, and the relatively less well studied but potentially important, the tension/dysphoria relief effects of alcohol (capturing negative reward). To advance research in this area, it is recommended that careful attention be paid to how these phenotypes are operationalized, including limb of BAC, measurements, alcohol dose, participant characteristics, and risk factors. Ultimately, large-scale longitudinal studies may be most informative in determining correlates of these phenotypes as well as their clinical significance across stages of alcohol use and alcoholism.

The treatment section reviewed ways in which subjective response to alcohol already informs treatment development for alcoholism, with a strong recommendation that studies linking responses to alcohol in the laboratory to clinical response in trials be conducted in order to fully ascertain which, if any, dimensions of subjective response serve as useful and predictive treatment targets.

In conclusion, this paper calls for research leveraging subjective response to alcohol in a systematic and methodologically rigorous fashion that can address central questions about alcoholism etiology, progression, and




Fig. 1. Subjective responses to alcohol in alcoholism etiology and treatment development research. Subjective response represents a multidimensional construct with the domains of: (i) stimulation, (ii) sedation, and (iii) tension. Etiological studies of subjective response have examined multiple indicators of alcoholism risk including: family history of alcoholism, genetic risk loci, and participants' level of drinking. Treatment studies utilizing subjective response have focused on medication development with a number of approved and promising medications having significant effects on subjective response outcomes. FH, family history; ADH, alcohol dehydrogenase; ALDH, aldehyde dehydrogenase; OPRM1, opioid receptor mu 1; GABA_AR, gamma-aminobutyric acid type A receptor; 5HTTLPR, serotonin-transporter-linked polymorphic region; nAChR, nicotinic acetylcholine receptor.

treatment development. The approach recommended herein is highly consistent with the RDoC initiative (Cuthbert and Insel, 2013; Insel, 2014; Insel et al., 2010). According to the RDoC approach, scientists are asked to identify key domains of function that can effectively describe the phenomenology of a disorder or set of disorders. RDoC domains are expected to inform behavior yet be amenable to examination through multiple units of analysis, such as molecular, genetic, circuit-level, and behavioral measurements. As reviewed above, subjective response to alcohol represents a behaviorally and biologically plausible phenotype upon which to build using the RDoC framework for understanding alcohol use disorder. And while this approach is not without its limitations and criticisms (Peterson, 2015), there is considerable enthusiasm for the development of an Alcohol Addiction RDoC that can ultimately advance personalized treatment for this disorder (Litten et al., 2015).

CONFLICT OF INTEREST

LAR is a paid consultant for GSK, has received study medications from Pfizer and Medicinova, and has received research support from Pfizer. None of the other authors have any conflicts of interest to disclose. SB was supported as a pre-doctoral trainee from the National Institute on Alcohol Abuse and Alcoholism (F31 AA022569). DJOR was supported as a post-doctoral fellow from the Tobacco-Related Disease Research Program (TRDRP 23FT-0102).

REFERENCES

- Anton RF, Voronin KK, Randall PK, Myrick H, Tiffany A (2012) Naltrexone modification of drinking effects in a subacute treatment and bar-lab paradigm: influence of OPRM1 and dopamine transporter (SLC6A3) genes. *Alcohol Clin Exp Res* 36:2000–2007.
- Armeli S, Tennen H, Todd M, Carney MA, Mohr C, Affleck G, Hromi A (2003) A daily process examination of the stress-response dampening effects of alcohol consumption. *Psychol Addict Behav* 17:266–276.

- Bisaga A, Evans SM (2006) The acute effects of gabapentin in combination with alcohol in heavy drinkers. *Drug Alcohol Depend* 83:25–32.
- Bouza C, Angeles M, Munoz A, Amate JM (2004) Efficacy and safety of naltrexone and acamprosate in the treatment of alcohol dependence: a systematic review. *Addiction* 99:811–828.
- Brasser SM, McCaul ME, Houtsmuller EJ (2004) Alcohol effects during acamprosate treatment: a dose-response study in humans. *Alcohol Clin Exp Res* 28:1074–1083.
- Bujarski S, Hutchison KE, Prause N, Ray LA (2015a) Functional significance of subjective response to alcohol across levels of alcohol exposure. *Addict Biol* doi: 10.1111/adb.12293 [Epub ahead of print].
- Bujarski S, Hutchison KE, Roche DJ, Ray LA (2015b) Factor structure of subjective responses to alcohol in light and heavy drinkers. *Alcohol Clin Exp Res* 39:1193–1202.
- Bujarski S, Ray LA (2014) Subjective response to alcohol and associated craving in heavy drinkers vs. alcohol dependents: an examination of Koob's allostatic model in humans. *Drug Alcohol Depend* 140:161–167.
- Cappell H, Heman CP (1972) Alcohol and tension reduction: a review. *Q J Stud Alcohol* 33:33–64.
- Childs E, Roche DJ, King AC, de Wit H (2012) Varenicline potentiates alcohol-induced negative subjective responses and offsets impaired eye movements. *Alcohol Clin Exp Res* 36:906–914.
- Chutuape MA, de Wit H (1994) Relationship between subjective effects and drug preferences: ethanol and diazepam. *Drug Alcohol Depend* 34:243–251.
- Clarke TK, Treutlein J, Zimmermann US, Kiefer F, Skowronek MH, Rietschel M, Mann K, Schumann G (2007) Review: HPA-axis activity in alcoholism: examples for a gene-environment interaction. *Addict Biol* 13:1–14.
- Cook TA, Luczak SE, Shea SH, Ehlers CL, Carr LG, Wall TL (2005) Associations of ALDH2 and ADH1B genotypes with response to alcohol in Asian Americans. *J Stud Alcohol* 66:196–204.
- Corbin WR, Gearhardt A, Fromme K (2008) Stimulant alcohol effects prime within session drinking behavior. *Psychopharmacology* 197:327–337.
- Correa M, Arizzi M, Betz A, Mingote S, Salamone J (2003) Open field locomotor effects in rats after intraventricular injections of ethanol and the ethanol metabolites acetaldehyde and acetate. *Brain Res Bull* 62:197–202.
- Correa M, Salamone JD, Segovia KN, Pardo M, Longoni R, Spina L, Peana AT, Vinci S, Acquas E (2012) Piecing together the puzzle of acetaldehyde as a neuroactive agent. *Neurosci Biobehav Rev* 36:404–430.
- Courtney KE, Ray LA (2014) Subjective responses to alcohol in the lab predict neural responses to alcohol cues. *J Stud Alcohol Drugs* 75:124–135.
- Cuthbert BN, Insel TR (2013) Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC Med* 11:126.
- DeWit H, Pierri J, Johanson CE (1989) Assessing individual differences in ethanol preference using a cumulative dosing procedure. *Psychopharmacology* 98:113–119.
- Drobes DJ, Anton RF, Thomas SE, Voronin K (2004) Effects of naltrexone and nalmefene on subjective response to alcohol among non-treatment-seeking alcoholics and social drinkers. *Alcohol Clin Exp Res* 28:1362–1370.
- Earleywine M (1994) Confirming the factor structure of the anticipated biphasic alcohol effects scale. *Alcohol Clin Exp Res* 18:861–866.
- Erblich J, Earleywine M, Erblich B, Bovbjerg DH (2003) Biphasic stimulant and sedative effects of ethanol: are children of alcoholics really different? *Addict Behav* 28:1129–1139.
- Fucito LM, Toll BA, Wu R, Romano DM, Tek E, O'Malley SS (2011) A preliminary investigation of varenicline for heavy drinking smokers. *Psychopharmacology* 215:655–663.
- Grant JD, Verges A, Jackson KM, Trull TJ, Sher KJ, Bucholz KK (2012) Age and ethnic differences in the onset, persistence and recurrence of alcohol use disorder. *Addiction* 107:756–765.
- Greeley J, Oei T (1999) Alcohol and tension reduction, in *Psychological Theories of Drinking and Alcoholism*, 2nd edn (Leonard KE, Blane HT eds), pp 14–53. The Guilford Press, New York, NY.
- Hart KE, Fazaan N (2004) Life stress events and alcohol misuse: distinguishing contributing stress events from consequential stress events. *Subst Use Misuse* 39:1319–1339.
- Heath AC, Martin NG (1991) Intoxication after an acute dose of alcohol: an assessment of its association with alcohol consumption patterns by using twin data. *Alcohol Clin Exp Res* 15:122–128.
- Heilig M, Thorsell A, Sommer WH, Hansson AC, Ramchandani VA, George DT, Hommer D, Barr CS (2010) Translating the neuroscience of alcoholism into clinical treatments: from blocking the buzz to curing the blues. *Neurosci Biobehav Rev* 35:334–344.
- Hendershot CS, Claus ED, Ramchandani VA (2014) Associations of OPRM1 A118G and alcohol sensitivity with intravenous alcohol self-administration in young adults. *Addict Biol* doi: 10.1111/adb.12165 [Epub ahead of print].
- Hinckes AS, Laucht M, Schmidt MH, Mann KF, Schumann G, Schuckit MA, Heinz A (2006) Low level of response to alcohol as associated with serotonin transporter genotype and high alcohol intake in adolescents. *Biol Psychiatry* 60:282–287.
- Hu X, Oroszi G, Chun J, Smith TL, Goldman D, Schuckit MA (2005) An expanded evaluation of the relationship of four alleles to the level of response to alcohol and the alcoholism risk. *Alcohol Clin Exp Res* 29:8–16.
- Hull CL (1943) Principles of behavior: an introduction to behavior theory. Available at: <http://doi.apa.org/psycinfo/1944-00022-000>. Accessed November 5, 2015.
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, Sanislow C, Wang P (2010) Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *Am J Psychiatry* 167:748–751.
- Insel TR (2014) The NIMH Research Domain Criteria (RDoC) Project: precision medicine for psychiatry. *Am J Psychiatry* 171:395–397.
- Johnson BA (2008) Update on neuropharmacological treatments for alcoholism: scientific basis and clinical findings. *Biochem Pharmacol* 75:34–56.
- Jones JD, Comer SD, Kranzler HR (2015) The pharmacogenetics of alcohol use disorder. *Alcohol Clin Exp Res* 39:391–402.
- Joslyn G, Brush G, Robertson M, Smith TL, Kalmijn J, Schuckit M, White RL (2008) Chromosome 15q25.1 genetic markers associated with level of response to alcohol in humans. *Proc Natl Acad Sci USA* 105:20368–20373.
- Judd LL, Hubbard RB, Huey LY, Attewell PA, Janowsky DS, Takahashi KI (1977) Lithium carbonate and ethanol induced highs in normal subjects. *Arch Gen Psychiatry* 34:463–467.
- Kareken DA, Liang T, Wetherill L, Dziedzic M, Bragulat V, Cox C, Talavage T, O'Connor SJ, Foroud T (2010) A polymorphism in GABRA2 is associated with the medial frontal response to alcohol cues in an fMRI study. *Alcohol Clin Exp Res* 34:2169–2178.
- King AC, Hasin D, O'Connor SJ, McNamara PJ, Cao D (2015) A prospective 5-year re-examination of alcohol response in heavy drinkers progressing in alcohol use disorder (AUD). *Biol Psychiatry* doi: 10.1016/j.biopsych.2015.05.007 [Epub ahead of print].
- King AC, McNamara PJ, Hasin DS, Cao D (2014) Alcohol challenge responses predict future alcohol use disorder symptoms: a 6-year prospective study. *Biol Psychiatry* 75:798–806.
- King AC, Volpicelli JR, Frazer A, O'Brien CP (1997) Effect of naltrexone on subjective alcohol response in subjects at high and low risk for future alcohol dependence. *Psychopharmacology* 129:15–22.
- King AC, de Wit H, McNamara PJ, Cao D (2011) Rewarding, stimulant, and sedative alcohol responses and relationship to future binge drinking. *Arch Gen Psychiatry* 68:389–399.
- Koob GF, Kreek MJ (2007) Stress, dysregulation of drug reward pathways, and the transition to drug dependence. *Am J Psychiatry* 164:1149–1159.
- Koob GF, Le Moal M (2005) Plasticity of reward neurocircuitry and the 'dark side' of drug addiction. *Nat Neurosci* 8:1442–1444.
- Koob GF, Le Moal M (2008a) Addiction and the brain anti-reward system. *Annu Rev Psychol* 59:29–53.

- Koob GF, Le Moal M (2008b) Review. Neurobiological mechanisms for opponent motivational processes in addiction. *Philos Trans R Soc Lond B Biol Sci* 363:3113–3123.
- Kranzler HR, Edenberg HJ (2010) Pharmacogenetics of alcohol and alcohol dependence treatment. *Curr Pharm Des* 16:2141–2148.
- Leggio L, Zywiak WH, McGeary JE, Edwards S, Fricchione SR, Shoaff JR, Addolorato G, Swift RM, Kenna GA (2013) A human laboratory pilot study with baclofen in alcoholic individuals. *Pharmacol Biochem Behav* 103:784–791.
- Levenson RW, Sher KJ, Grossman LM, Newman J, Newlin DB (1980) Alcohol and stress response dampening: pharmacological effects, expectancy, and tension reduction. *J Abnorm Psychol* 89:528–538.
- Litten RZ, Ryan ML, Falk DE, Reilly M, Fertig JB, Koob GF (2015) Heterogeneity of alcohol use disorder: understanding mechanisms to advance personalized treatment. *Alcohol Clin Exp Res* 39:579–584.
- Luczak SE, Wall TL, Cook TA, Shea SH, Carr LG (2004) ALDH2 status and conduct disorder mediate the relationship between ethnicity and alcohol dependence in Chinese, Korean, and White American college students. *J Abnorm Psychol* 113:271–278.
- Martin CS, Earleywine M, Musty RE, Perrine MW, Swift RM (1993) Development and validation of the Biphasic Alcohol Effects Scale. *Alcohol Clin Exp Res* 17:140–146.
- Matsushita S, Higuchi S (2014) Genetic differences in response to alcohol. *Handb Clin Neurol* 125:617–627.
- McCaul ME, Wand GS, Eissenberg T, Rohde CA, Cheskin LJ (2000) Naltrexone alters subjective and psychomotor responses to alcohol in heavy drinking subjects. *Neuropsychopharmacology* 22:480–492.
- McKee SA, Harrison EL, O'Malley SS, Krishnan-Sarin S, Shi J, Tetrault JM, Picciotto MR, Petrakis IL, Estevez N, Balchunas E (2009) Varenicline reduces alcohol self-administration in heavy-drinking smokers. *Biol Psychiatry* 66:185–190.
- McNair DM, Lorr M, Droppleman LF (1971) *Manual for the Profile of Mood States*. Educational & Industrial Testing Service, San Diego, CA.
- Menary KR, Corbin WR, Leeman RF, Fucito LM, Toll BA, DeMartini K, O'Malley SS (2015) Interactive and indirect effects of anxiety and negative urgency on alcohol-related problems. *Alcohol Clin Exp Res* 39:1267–1274.
- Miranda R Jr, MacKillop J, Monti PM, Rohsenow DJ, Tidey J, Gwaltney C, Swift R, Ray L, McGeary J (2008) Effects of topiramate on urge to drink and the subjective effects of alcohol: a preliminary laboratory study. *Alcohol Clin Exp Res* 32:489–497.
- Mitchell JM, O'Neil JP, Janabi M, Marks SM, Jagust WJ, Fields HL (2012) Alcohol consumption induces endogenous opioid release in the human orbitofrontal cortex and nucleus accumbens. *Sci Transl Med* 4:116ra6.
- Morean ME, de Wit H, King AC, Sofuoglu M, Rueger SY, O'Malley SS (2013) The drug effects questionnaire: psychometric support across three drug types. *Psychopharmacology* 227:177–192.
- Morean ME, Corbin WR (2010) Subjective response to alcohol: a critical review of the literature. *Alcohol Clin Exp Res* 34:385–395.
- Newlin DB, Thomson JB (1990) Alcohol challenge with sons of alcoholics: a critical review and analysis. *Psychol Bull* 108:383–402.
- Peng GS, Chen YC, Wang MF, Lai CL, Yin SJ (2014) ALDH2*2 but not ADH1B*2 is a causative variant gene allele for Asian alcohol flushing after a low-dose challenge: correlation of the pharmacokinetic and pharmacodynamic findings. *Pharmacogenet Genomics* 24:607–617.
- Peterson AM (2007) Improving adherence in patients with alcohol dependence: a new role for pharmacists. *Am J Health Syst Pharm* 64:S23–S29.
- Peterson BS (2015) Editorial: Research Domain Criteria (RDoC): a new psychiatric nosology whose time has not yet come. *J Child Psychol Psychiatry* 56:719–722.
- Pettinati HM, O'Brien CP, Rabinowitz AR, Wortman SP, Oslin DW, Kampman KM, Dackis CA (2006) The status of naltrexone in the treatment of alcohol dependence: specific effects on heavy drinking. *J Clin Psychopharmacol* 26:610–625.
- Pierucci-Lagha A, Covault J, Feinn R, Nellissery M, Hernandez-Avila C, Oncken C, Morrow AL, Kranzler HR (2005) GABRA2 alleles moderate the subjective effects of alcohol, which are attenuated by finasteride. *Neuropsychopharmacology* 30:1193–1203.
- Quinn PD, Fromme K (2011) Subjective response to alcohol challenge: a quantitative review. *Alcohol Clin Exp Res* 35:1759–1770.
- Ramchandani VA, Plawecki M, Li TK, O'Connor S (2009) Intravenous ethanol infusions can mimic the time course of breath alcohol concentrations following oral alcohol administration in healthy volunteers. *Alcohol Clin Exp Res* 33:938–944.
- Ramchandani VA, Umhau J, Pavon FJ, Ruiz-Velasco V, Margas W, Sun H, Damadzic R, Eskay R, Schoor M, Thorsell A, Schwandt ML, Sommer WH, George DT, Parsons LH, Herscovitch P, Hommer D, Heilig M (2011) A genetic determinant of the striatal dopamine response to alcohol in men. *Mol Psychiatry* 16:809–817.
- Ray LA, Barr CS, Blendy JA, Oslin D, Goldman D, Anton RF (2012) The role of the Asn40Asp polymorphism of the mu opioid receptor gene (OPRM1) on alcoholism etiology and treatment: a critical review. *Alcohol Clin Exp Res* 36:385–394.
- Ray LA, Bujarski S, MacKillop J, Courtney KE, Monti PM, Miotto K (2013) Subjective response to alcohol among alcohol-dependent individuals: effects of the mu-opioid receptor (OPRM1) gene and alcoholism severity. *Alcohol Clin Exp Res* 37(Suppl 1):E116–E124.
- Ray LA, Bujarski S, Squeglia LM, Ashenhurst JR, Anton RF (2014) Interactive effects of OPRM1 and DAT1 genetic variation on subjective responses to alcohol. *Alcohol Alcohol* 49:261–270.
- Ray LA, Chin PF, Miotto K (2010a) Naltrexone for the treatment of alcoholism: clinical findings, mechanisms of action, and pharmacogenetics. *CNS Neurol Disord Drug Targets* 9:13–22.
- Ray LA, Hutchison KE (2004) A polymorphism of the mu-opioid receptor gene (OPRM1) and sensitivity to the effects of alcohol in humans. *Alcohol Clin Exp Res* 28:1789–1795.
- Ray LA, Hutchison KE (2009) Associations among GABRG1, level of response to alcohol, and drinking behaviors. *Alcohol Clin Exp Res* 33:1382–1390.
- Ray LA, Hutchison KE, MacKillop J, Miranda R Jr, Audette A, Swift R, Monti PM (2008) Effects of naltrexone during the descending limb of the blood alcohol curve. *Am J Addict* 17:257–264.
- Ray LA, Hutchison KE, Tartter M (2010b) Application of human laboratory models to pharmacotherapy development for alcohol dependence. *Curr Pharm Des* 16:2149–2158.
- Ray LA, Krull JL, Leggio L (2010c) The effects of naltrexone among alcohol non-abstainers: results from the COMBINE Study. *Front Psychiatry* 1:26.
- Ray LA, MacKillop J, Leventhal A, Hutchison KE (2009) Catching the alcohol buzz: an examination of the latent factor structure of subjective intoxication. *Alcohol Clin Exp Res* 33:2154–2161.
- Ray LA, MacKillop J, Monti PM (2010d) Subjective responses to alcohol consumption as endophenotypes: advancing behavioral genetics in etiological and treatment models of alcoholism. *Subst Use Misuse* 45:1742–1765.
- Ray LA, Meskew-Staer S, Hutchison KE (2007) The relationship between prospective self-rating of alcohol sensitivity and craving and experimental results from two alcohol challenge studies. *J Stud Alcohol Drugs* 68:379–384.
- Ray LA, Miranda R Jr, Tidey JW, McGeary JE, MacKillop J, Gwaltney CJ, Rohsenow DJ, Swift RM, Monti PM (2010e) Polymorphisms of the mu-opioid receptor and dopamine D4 receptor genes and subjective responses to alcohol in the natural environment. *J Abnorm Psychol* 119:115–125.
- Roche DJ, Palmeri MD, King AC (2014) Acute alcohol response phenotype in heavy social drinkers is robust and reproducible. *Alcohol Clin Exp Res* 38:844–852.
- Roche DJ, Ray LA (2015) Subjective response as a consideration in the pharmacogenetics of alcoholism treatment. *Pharmacogenomics* 16:721–736.
- Roh S, Matsushita S, Hara S, Maesato H, Matsui T, Suzuki G, Miyakawa T, Ramchandani VA, Li TK, Higuchi S (2011) Role of GABRA2 in moderating subjective responses to alcohol. *Alcohol Clin Exp Res* 35:400–407.
- Rosner S, Leucht S, Leher P, Soyka M (2008) Acamprosate supports abstinence, naltrexone prevents excessive drinking: evidence from a meta-analysis with unreported outcomes. *J Psychopharmacol* 22:11–23.
- Rueger SY, King AC (2012) Validation of the brief Biphasic Alcohol Effects Scale (B-BAES). *Alcohol Clin Exp Res* 37:470–476.

- Rueger SY, McNamara PJ, King AC (2009) Expanding the utility of the Biphasic Alcohol Effects Scale (BAES) and initial psychometric support for the Brief-BAES (B-BAES). *Alcohol Clin Exp Res* 33:916–924.
- Schacht JP, Anton RF, Voronin KE, Randall PK, Li X, Henderson S, Myrick H (2013) Interacting effects of naltrexone and OPRM1 and DAT1 variation on the neural response to alcohol cues. *Neuropsychopharmacology* 38:414–422.
- Schuckit MA (1984) Subjective responses to alcohol in sons of alcoholics and control subjects. *Arch Gen Psychiatry* 41:879–884.
- Schuckit MA (1994) Low level of response to alcohol as a predictor of future alcoholism. *Am J Psychiatry* 151:184–189.
- Schuckit MA, Gold EO (1988) A simultaneous evaluation of multiple markers of ethanol/placebo challenges in sons of alcoholics and controls. *Arch Gen Psychiatry* 45:211–216.
- Schuckit MA, Mazzanti C, Smith TL, Ahmed U, Radel M, Iwata N, Goldman D (1999) Selective genotyping for the role of 5-HT 2A, 5-HT 2C, and GABA α 6 receptors and the serotonin transporter in the level of response to alcohol: a pilot study. *Biol Psychiatry* 45:647–651.
- Schuckit MA, Risch SC, Gold EO (1988) Alcohol consumption, ACTH level, and family history of alcoholism. *Am J Psychiatry* 145:1391–1395.
- Schuckit MA, Smith TL (1996) An 8-year follow-up of 450 sons of alcoholic and control subjects. *Arch Gen Psychiatry* 53:202–210.
- Schuckit MA, Smith TL, Kalmijn J, Danko GP (2005) A cross-generational comparison of alcohol challenges at about age 20 in 40 father-offspring pairs. *Alcohol Clin Exp Res* 29:1921–1927.
- Sher KJ, Levenson RW (1982) Risk for alcoholism and individual differences in the stress-response-dampening effect of alcohol. *J Abnorm Psychol* 91:350–367.
- Sinha R, Robinson J, O'Malley S (1998) Stress response dampening: effects of gender and family history of alcoholism and anxiety disorders. *Psychopharmacology* 137:311–320.
- Spanagel R (2009) Alcoholism: a systems approach from molecular physiology to addictive behavior. *Physiol Rev* 89:649–705.
- Stewart SH, Karp J, Pihl RO, Peterson RA (1997) Anxiety sensitivity and self-reported reasons for drug use. *J Subst Abuse* 9:223–240.
- Sun F, Tsuritani I, Yamada Y (2002) Contribution of genetic polymorphisms in ethanol-metabolizing enzymes to problem drinking behavior in middle-aged Japanese men. *Behav Genet* 32:229–236.
- Swift RM, Whelihan W, Kuznetsov O, Buongiorno G, Hsuing H (1994) Naltrexone-induced alterations in human ethanol intoxication. *Am J Psychiatry* 151:1463–1467.
- Uhart M, Weerts EM, McCaul ME, Guo X, Yan X, Kranzler HR, Li N, Wand GS (2013) GABRA2 markers moderate the subjective effects of alcohol. *Addict Biol* 18:357–369.
- Viken RJ, Rose RJ, Morzorati SL, Christian JC, Li TK (2003) Subjective intoxication in response to alcohol challenge: heritability and covariation with personality, breath alcohol level, and drinking history. *Alcohol Clin Exp Res* 27:795–803.
- Wall TL, Shea SH, Luczak SE, Cook TA, Carr LG (2005) Genetic associations of alcohol dehydrogenase with alcohol use disorders and endophenotypes in white college students. *J Abnorm Psychol* 114:456–465.
- de Wit H, Doty P (1994) Preference for ethanol and diazepam in light and moderate social drinkers: a within-subjects study. *Psychopharmacology* 115:529–538.
- Zeichner A, Giancola PR, Allen JD (1995) Effects of hostility on alcohol stress-response-dampening. *Alcohol Clin Exp Res* 19:977–983.
- Zimmermann US, Mick I, Laucht M, Vitvitskiy V, Plawecki MH, Mann KF, O'Connor S (2009) Offspring of parents with an alcohol use disorder prefer higher levels of brain alcohol exposure in experiments involving computer-assisted self-infusion of ethanol (CASE). *Psychopharmacology* 202:689–697.
- Zimmermann US, Mick I, Vitvitskiy V, Plawecki MH, Mann KF, O'Connor S (2008) Development and pilot validation of computer-assisted self-infusion of ethanol (CASE): a new method to study alcohol self-administration in humans. *Alcohol Clin Exp Res* 32:1321–1328.
- Zimmermann US, O'Connor S, Ramchandani VA (2013) Modeling alcohol self-administration in the human laboratory, in *Behavioral Neurobiology of Alcohol Addiction, Behavioral Neurobiology of Alcohol Addiction* (Sommer WH, Spanagel R eds), pp 315–353, Springer, Heidelberg, Germany.